

Utilization, Equity and Determinants of Full Antenatal Care in India: Analysis from the National Family Health Survey 4

Gunjan Kumar

Knowledge Intergration and Transaltional Platform at the Centre for Health Research and Development, Society for Applied Studies

Tarun Shankar Choudhary

Knowledge Integration and Translational Platform at the Centre for Health Research and Development, Society for Applied Studies

Akanksha Srivastava

Knowldege Integration and Translational Platform at the Centre for Health Research and Development, Society for Applied Studies

Ravi Praksah Updhyay

Knowledge Integration and Translational Platform at the Centre for Health Research and Development, Society for Applied Studies

Sunita Taneja

Knowledge Integration and Translational Platform at the Centre for Health Research and Development, Society for Applied Studies

Rajiv Bahl

Department of Maternal, Newborn, Child and Adolescent Health, World Health Organisation

Jose Martines

Centre for Intervention Sciences in Maternal and Child Health, University of Bergen

Maharaj Kishan Bhan

Indian Institute of Technology Delhi; Knowledge Integration and Translational Platform, Biotechnology Industry Research Assistance Council

Nita Bhandari

Knowledge Integration and Translational Platform at the Centre for Health Research and Development, Society for Applied Studies

Sarmila Mazumder (✉ sarmila.mazumder@sas.org.in)

Centre for Health Research and Development, Society for Applied Studies <https://orcid.org/0000-0003-4200-6112>

Keywords: Antenatal care, India, NFHS-4

Posted Date: September 17th, 2019

DOI: <https://doi.org/10.21203/rs.2.11599/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on September 5th, 2019. See the published version at <https://doi.org/10.1186/s12884-019-2473-6>.

Abstract

Objectives: We examined the utilisation, equity and determinants of full antenatal care (ANC), defined as 4 or more antenatal visits, at least one tetanus toxoid (TT) injection and consumption of iron folic acid (IFA) for a minimum of 100 days, in India. **Methods:** We analysed a sample of 190,898 women from India's National Family Health Survey 4. Concentration curves and concentration index were used to assess equity in full ANC utilisation. Multivariable logistic regression model was used to examine the factors associated with full ANC utilization. **Results:** In India, 21% of pregnant women utilised full ANC, ranging from 2.3% - 65.9% across states. Overall, 51.6% had 4 or more ANC visits, 30.8% consumed IFA for atleast 100 days, and 91.1% had one or more doses of tetanus toxoid. Registration of pregnancy, utilisation of government's Integrated Child Development Services (ICDS) and health insurance coverage were associated with higher odds of full ANC utilisation. Lower maternal education, lower wealth quintile(s), lack of father's participation during antenatal visits, higher birth order, teenage and unintended pregnancy were associated with lower odds of full ANC utilisation. Full ANC utilisation was inequitable across area of residence, caste and maternal education. **Conclusions:** Full ANC utilization in India was inadequate and inequitable. Although half of the women did not receive the minimum recommended ANC visits, the utilisation of TT immunisation was almost universal. The positive association of full ANC with ICDS utilisation and child's father involvement may be leveraged for increasing the uptake of full ANC. Strategies to address the socio-demographic factors associated with low and inequitable utilisation of full ANC are imperative for strengthening India's maternal health program. **Key words:** Antenatal care, India, NFHS-4

Background

Antenatal care (ANC) is an opportunity to promote a positive pregnancy experience and improved maternal and child survival. Care in the antenatal period is also important for supporting the long-term growth and development of the child as its part of the critical "1000 days" window [1,2,3]. It serves as a facilitating platform linking the woman and her family to the healthcare system and possibly promotes higher utilisation of essential services like breastfeeding and nutritional counselling, post-partum family planning and childhood vaccination [4,5].

Globally, 62% of pregnant women received the WHO recommended minimum 4 antenatal visits during 2010-2016 [6]. Latest research has shown a lower still birth rate, among women with a minimum of 8 antenatal visits, based on which the minimum recommended number of antenatal contacts has now been increased from 4 to 8 [7].

In India, the proportion of pregnant women receiving the minimum 4 antenatal visits has increased from 37.0% to 51.2% during 2006-2016 [8]. This is relatively modest when compared to increase in the rate of institutional delivery which has doubled from 38.7% to 79% during the same time period, largely driven by the conditional cash transfer schemes of the government [8,9,10]. This differential coverage reflects a missed opportunity, as about one fourth of the maternal deaths are attributable to pre-eclampsia,

eclampsia and antepartum haemorrhage, which could be identified and managed during the antenatal contacts [11].

Antenatal contacts also provide a window of opportunity to detect and possibly prevent adverse birth events. Evidence suggests that there is significant inequality in the access to essential health services, including services for pregnancy and childbirth. Disparities in utilisation of maternal care services have been reported in low and middle-income countries (LMIC) [12,13]. A comprehensive understanding of determinants of antenatal care utilisation in India is lacking. In context of the recent focus of the Indian government on antenatal services, through Pradhan Mantri Matru Vandana Yojana (PMMVY), a conditional cash transfer scheme, understanding the factors driving the utilisation is crucial to aid the development of an informed policy. In this scheme the pregnant woman is eligible if she registers her pregnancy at the Anganwadi centre (AWC) within four months of conception, attends at least one prenatal care session and is taking Iron-folic acid tablets and TT (tetanus toxoid) injection [14].

We conducted a secondary data analysis using the recent National Family Health Survey (NFHS-4) data, with the objective of studying the coverage and equity in antenatal care utilisation at national and state level in India. We also examined the predictors of full antenatal care utilization. The policy implications of these findings taking the new and pre-existing guidelines and national health programmes into consideration have also been discussed.

Methods

Data Source

This analysis utilizes individual level data from the fourth round of India's National Family Health Survey (NFHS 4), conducted during 2015-16. NFHS are a series of nationally representative, cross-sectional surveys that provides data on a range of demographic, socioeconomic, maternal and child health outcomes, reproductive health and family planning. In the fourth round, around 700,000 women, aged 15-49 years from 601,509 households were interviewed, using a two-stage stratified sampling design, with a response rate of 97%. Details of the sampling design and instruments used are available elsewhere [8].

Children recode file was used for this analysis (IAKR73FL). The recode file contains data in a standard, easy to analyse format with common variable names and coding categories for easy comparability between countries and includes summary variables and indices which are calculated post data collection [15]. Information on the utilisation of antenatal care was available for the most recent pregnancy, resulting in a live birth during the five years preceding the survey.

Definitions and analysis

The outcome variable was "*Full antenatal care*" defined as four or more antenatal visits, at least one tetanus toxoid (TT) injection and reported consumption of iron folic acid (IFA) tablets or syrup for a minimum of 100 days [8]. Place of residence, caste, wealth quintile, access to health insurance, age at

conception for current pregnancy, education, registration of pregnancy, timing of first ANC visit, utilisation of government's Integrated child development services (ICDS), pregnancy intendedness, participation of child's father in ANC visit and birth order of the child were used as explanatory variables. The details and sub-categories of these variables are available in Additional file 1.

The analysis was restricted to the antenatal care utilised during the last pregnancy that resulted in a live birth, in the five years preceding the survey. From the total sample of 259,627, information pertaining to last live birth was available for 190,898 births. The proportion of pregnant women who utilised full ANC was estimated at national and sub-national level (states and union territories). There are 36 states/union territories in India. We used binary logistic regression to examine the association between the explanatory variables and full ANC care.

The variables with a p-value of < 0.20 in univariable analysis and variables of known clinical or contextual importance were included in the multivariable model [16]. A p-value of < 0.05 was considered as statistically significant for all analyses. We used concentration index to examine the extent of wealth-based inequality in utilisation of full ANC in India and states/union territories. Concentration curve provide statistical inference for the comparison of inequality between groups. They capture the extent to which health differs across individuals ranked by some indicator of socioeconomic status [17].

Analysis was done using STATA[®] 15.1 (StataCorp LLC, College Station, TX, USA). Stata's survey command (*svyset*) was used to adjust for sampling weight, clustering and stratification in the sampling design. User written Stata package "*conindex*" was used for calculating the concentration index. The guidelines for data use as required by the DHS program were strictly followed [18].

Results

We report the findings for 190,898 women, who had at least one live birth in the five years preceding the survey. Background characteristics of these women are shown in Table 1

[Table 1 about here]

Full ANC utilisation

Twenty One percent of pregnant women utilised full ANC during their last pregnancy ranging from 2.3% - 65.9% within states (median 30.7, IQR 14.8 - 39.0). The proportion of women who had a minimum of 4 ANC visits was 51.6%, ranging from 14.4% - 96.7% within states, IFA was consumed for a minimum of 100 days by 30.8% of women, ranging from 4.5% - 85.5% within states. At least one dose of tetanus toxoid was received by 91.1% of women ranging from 70% - 98.6% within states (Figure 1 & Figure 2). The median number of antenatal visits in the country was 4 (IQR 2-7).

[Figure 1 & Figure 2 about here]

Determinants of Full ANC

Results from the multivariable logistic regression model are presented in Table 2. The model was also adjusted for respondent's state of residence. We observed a gradient in full ANC utilisation across the wealth quintiles, the highest utilisation was among the richest and vice-versa. Women residing in rural areas had lower odds (adjusted odds ratio, AOR 0.90; 95% CI 0.84-0.96) of full ANC utilisation. Odds of full ANC utilisation was higher among women covered by health insurance (AOR 1.18; 95% CI 1.11-1.26).

Maternal age below 19 years at conception, had lower odds (AOR 0.86; 95% CI 0.79-0.93) of full ANC utilisation. A gradient in full ANC utilisation was also observed across the levels of education, with lowest utilisation among those with no formal education. The odds of full ANC decreased with increasing birth order. Unintended pregnancy was associated with a lower odds of full ANC utilisation (AOR 0.83; 95% CI 0.76-0.92).

Unregistered pregnancy was associated with lower odds of full ANC utilisation (AOR 0.78; 95% CI 0.71-0.85). Women who had their first ANC visit in the second or third trimester had lower odds of full ANC utilisation. Lack of child's father participation during the ANC visits was associated with lower odds of full ANC utilisation, (AOR 0.72; 95% CI 0.68-0.77). Women who had not received ICDS benefits had lower odds of full ANC utilisation (AOR 0.93; 95% CI 0.88-0.98). Caste was not associated with full ANC utilisation.

[Table 2 about here]

Inequity in Full ANC utilisation

The inequity in full ANC utilisation was higher in rural areas, among women with no education and those belong to socially disadvantaged sections (SC/ST/OBC) (Figure 3a-c). The utilisation of full ANC, was highly inequitable with a concentration index of 0.31, ranging from -0.11 to 0.67 within states. In general, the states with lower utilisation of full ANC had higher level of inequity in utilisation (Figure 4).

[Figure 3a-c & Figure 4 about here]

Discussion

The findings of this analysis are of concern as despite efforts of the government of India during the last two decades, only one fifth of pregnant women utilised ANC to a full extent. Half of the women did not receive these minimum recommended 4 ANC visits, which are a conservative expectation when compared to the recent recommendations of the World Health Organisation (WHO), a minimum of 8 visits. In 17 out of 36 states/UTs, less than 30% of the pregnant women received full ANC. Inequity in full ANC utilisation was higher in states with low rates of full ANC coverage.

The proportion of women with 4 or more ANC visits is considerably lower than the global average of 61.8% and implementing the recent WHO recommendation of a minimum of 8 ANC visits will be a major challenge for the national programme in India [6]. The number of ANC visits may also be critical to the delivery of other components of ANC and to provide adequate follow-up of pregnant women closer to

delivery. The high proportion of mothers with at least one tetanus toxoid immunisation can be achieved even in a single visit during any trimester. In contrast, 100 days of IFA consumption is possible only if multiple visits are made as currently the supplies are given for 1 month at each visit and lower number of visits may be a reason for low utilisation for 100 days of IFA.

Registration of pregnancy, utilisation of benefits from the government's ICDS program and having health insurance were associated with higher odds of full ANC utilisation. Lower maternal education, lower wealth quintile(s), higher birth order, father not accompanying for the ANC visit, teenage pregnancy and unintended pregnancy were associated with lower odds of full ANC utilisation.

Economic status and maternal education were highly associated with utilisation of ANC, which corroborates with previous literature [19,20,21]. Younger maternal age at conception is also common among these women. In LMICs including South East Asia, women from the highest wealth quintile have higher financial and social access to health care services in general, which may lead to higher utilisation of full ANC as seen in the current analysis [6].

Unintended pregnancy, which may be due to lack of awareness or, a reflection of inadequate or inaccessible family planning services has been associated with lower ANC utilisation previously and our findings also suggest the same [22,23].

Lower ANC utilisation among women of higher parity can be due to increased confidence from previous pregnancy and childbirth experience, constraints of time and resources, poor prior experience with the health system and financial barriers to ANC utilisation [19].

Equity in states with low ANC utilisation is expected to be lower as those who are better-off are more likely to utilise the services. This finding is critical to account for while redesigning of strategies for increasing ANC utilisation at national and subnational level. As India prepares for achieving universal health coverage, equity in utilisation of essential maternal healthcare services like ANC should be ensured.

Child's father presence during any ANC visit was associated with higher utilisation of full ANC, this is of special interest in the context of a patriarchal society like India. Child's father presence during ANC visit, may reflect greater spousal care and support, joint decision making and a more caring environment at home [24,25]. Positive association between health insurance coverage and full ANC utilisation in the adjusted analysis is relevant as, despite provision of free ANC services by the government families incur out-of-pocket expenditure for availing these. This finding is relevant to the current health policy as majority of the private insurance players and even the recently launched National Health Protection Scheme ("Ayushman Bharat") do not cover ANC services [26]. Higher utilisation of government's ICDS services, implemented through Anganwadis which are the focal point of community outreach sessions for antenatal care, health education and nutrition support during pregnancy and lactation was associated with higher odds of full ANC as it acts as a bridging platform for different government schemes like Janani Suraksha Yojna (JSY) and Pradhan Mantri Matru Vandana Yojna (PMMVY) [14,27].

Strengths and limitations

The findings from this analysis should be interpreted considering the following strengths. Our analysis uses data from the most recent survey with a large, nationally representative sample, and is based on a more comprehensive indicator i.e. *full ANC* (minimum of 4 ANC visits, consumption of 100 or more days of IFA and at least 1 dose of tetanus toxoid), as against the total number of ANC visits used in previous studies. Some limitations are as follows. Lack of information on pregnancies which did not result in a live birth i.e. abortion, miscarriage or still birth restricts us from commenting on the utilisation of ANC in this subset with adverse outcomes. The responses pertaining to the individual components of full ANC, were self-reported and therefore prone to recall bias, although the subgroup analysis restricted to births in the one year preceding the survey yielded similar results. We are unable to comment on some factors like distance from the health care facility, provider discrimination, health literacy, care seeking behaviour and other system side factors for which information was not collected in NFHS-4.

Conclusions

Full ANC utilization in India was inadequate and inequitable. Although half of the women did not receive the minimum number of recommended ANC visits, the utilisation of TT immunisation was almost universal. In a country like India, where still birth, prematurity, Intrauterine growth restriction (IUGR) and maternal mortality are still high, the recent WHO recommendations of 8 ANC visits, emphasises the need to provide more number of ANC visits and (should force us to recognise the need) intensive care to pregnant women [28,29,30]. Taking into consideration the factors associated with low and inequitable utilisation, it is imperative to design strategies, in order to achieve the WHO recommendations at the earliest.

Counselling of young married couples may create a channel for greater information flow, for planned parenthood. Broadcasting important messages related to antenatal care at relevant health and educational institutes and through mass media, may help in increasing awareness at the population level. Future surveys should capture details of antenatal care with higher granularity to understand the quality of care received at each visit. The states with low full ANC utilization should give greater priority to improving ANC services and using the opportunities opening in the country [14]. Coordinated effort is needed from the different sections of the government involved in delivering maternal care during pregnancy is needed. We suggest, that the newly launched Ayushman Bharat scheme, should also prioritize covering outpatient-based antenatal care.

List Of Abbreviations

95%CI: 95% Confidence Interval

AOR: Adjusted Odds Ratio

ANC: Antenatal Care

AWC: Anganwadi Centre

ICDS: Integrated Child Development Scheme

IFA: Iron Folic Acid

IQR: Inter Quartile Range

IUGR: Intrauterine Growth Retardation

JSY: Janani Suraksha Yojana

LMIC: Low- and Middle-Income Countries

NFHS: National Family Health Survey

PMMVY: Pradhan Mantri Matru Vandana Yojana

TT: Tetanus Toxoid

WHO: World Health Organisation

Declarations

Ethics statement

Ethical clearance was not needed for this analysis as it was based on secondary data available freely, upon request in the public domain.

Consent for publication

Not Applicable

Availability of data and materials

Data was available through MEASURE DHS program (www.dhsprogram.com).

Competing interests

The authors declare that they have no competing interests

Funding

None

Author's contributions

GK and TSC conceptualized the study, developed the analytical strategy, performed the statistical analysis, interpreted the results, and wrote the first draft and revised the manuscript. AS, RPU contributed to obtaining data, statistical analysis, interpretation of the results, and revision of the manuscript. NB, JM, RB, SM, ST, MKB contributed to the conceptualization of the study, analytical strategy, and interpretation of results, and gave critical feedback throughout the preparation of manuscript.

Acknowledgements

The Society for Applied Studies acknowledges the core support provided by the Department of Maternal, Newborn, Child and Adolescent Health, World Health Organization, Geneva (WHO Collaborating Centre IND-158) and the Centre for Intervention Science in Maternal and Child Health (RCN Project No. 223269), Centre for International Health, University of Bergen (Norway). Knowledge Integration and Technology Platform (KnIT), a Grand Challenges Initiative of the Department of Biotechnology and Biotechnology Industry Research Assistance Council (BIRAC) of Government of India and Bill & Melinda Gates Foundation (USA). The authors would also like to acknowledge the MEASURE DHS program for providing the India DHS datasets.

References

1. Noonan K, Corman H, Schwartz-Soicher O, Reichman NE. Effects of prenatal care on child health at age 5. *Matern Child Health J.* 2013 Feb;17(2):189–99.
2. Kuhnt J, Vollmer S. Antenatal care services and its implications for vital and health outcomes of children: evidence from 193 surveys in 69 low-income and middle-income countries. *BMJ Open.* 2017 Nov 15;7(11).
[<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5695442/>] Accessed on 16 Jan 2019
3. Liu X, Behrman JR, Stein AD, Adair LS, Bhargava SK, Borja JB, et al. Prenatal care and child growth and schooling in four low- and medium-income countries. *PloS One.* 2017;12(2): e0171299.
4. Dixit P, Dwivedi LK, Ram F. Strategies to Improve Child Immunization via Antenatal Care Visits in India: A Propensity Score Matching Analysis. *PloS One.* 2013;8(6): e66175.
5. World Health Organization (WHO). 2012. From Evidence to Policy: Expanding Access to Family Planning. Strengthening Health System Response: No Opportunities Missed. Geneva: WHO Department of Reproductive Health and Research. Accessed November 15, 2017, at: [http://apps.who.int/iris/bitstream/10665/75162/1/WHO_RHR_HRP_12.22_eng.pdf]
6. Antenatal care. UNICEF DATA. [<https://data.unicef.org/topic/maternal-health/antenatal-care/>] Accessed on 16 Jan 2019
7. WHO | WHO recommendations on antenatal care for a positive pregnancy experience. WHO. [http://www.who.int/reproductivehealth/publications/maternal_perinatal_health/anc-positive-pregnancy-experience/en/]. Accessed on 21 Jan 2019
8. International Institute for Population Sciences (IIPS) and ICF. National Family Health Survey (NFHS-4), 2015-16. Mumbai: IIPS; 2017.
[<http://rchiips.org/nfhs/pdf/NFHS4/India.pdf>] Accessed on 21 Jan 2019
9. Gupta SK, Pal DK, Tiwari R, Garg R, Shrivastava AK, Sarawagi R, et al. Impact of Janani Suraksha Yojana on Institutional Delivery Rate and Maternal Morbidity and Mortality: An Observational Study in India. *J Health Popul Nutr.* 2012 Dec;30(4):464–71.
10. Kumar V, Misra SK, Kaushal SK, Gupta SC, Khan AM. A study on the effect of janani suraksha yojana on antenatal registration and institutional deliveries in the Agra district of Uttar

pradesh. *Indian J Public Health* 2015; 59:54-7. 11. Say L, Chou D, Gemmill A, Tunçalp Ö, Moller A-B, Daniels J, et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Glob Health*. 2014 Jun;2(6): e323-333. 12. Arsenault C, Jordan K, Lee D, Dinsa G, Manzi F, Marchant T, et al. Equity in antenatal care quality: an analysis of 91 national household surveys. *Lancet Glob Health*. 2018 Nov;6(11):e1186–95. 13. Yaya S, Uthman OA, Amouzou A, Ekholuenetale M, Bishwajit G. Inequalities in maternal health care utilization in Benin: a population based cross-sectional study. *BMC Pregnancy Childbirth*. 2018 May 31;18(1):194. 14. PMMVY | Ministry of Women & Child Development | Gol. [<http://www.wcd.nic.in/schemes/pradhan-mantri-matru-vandana-yojana>] Accessed on 21 Jan 2019 15. Program., T. D. Demographic and Health Surveys- Dataset-Types. [<https://dhsprogram.com/data/Dataset-Types.cfm>] Accessed on 25 Mar 2019 16. David W. Hosmer, Jr., Stanley Lemeshow, Rodney X. Sturdivant. *Applied Logistic Regression*. 3rd ed. Hoboken, New Jersey: John Wiley & Sons, Inc; 2013. 17. O'Donnell O, O'Neill S, Van Ourti T, Walsh B. conindex: Estimation of concentration indices. *Stata J*. 2016;16(1):112–38. 18. Program., T. D. Demographic and Health Surveys-Terms of use. [<https://dhsprogram.com/data/terms-of-use.cfm>] Accessed on 25 Mar 2019 19. Miteniece E, Pavlova M, Shengelia L, Rechel B, Groot W. Barriers to accessing adequate maternal care in Georgia: a qualitative study. *BMC Health Serv Res*. 2018 Aug 13;18. [<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6090778/>] Accessed on 21 Jan 2019 20. Hijazi HH, Alyahya MS, Sindiani AM, Saqan RS, Okour AM. Determinants of antenatal care attendance among women residing in highly disadvantaged communities in northern Jordan: a cross-sectional study. *Reprod Health*. 2018 Jun 7;15. [<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5992715/>] Accessed on 21 Jan 2019 21. Sakeah E, Okawa S, Rexford Oduro A, Shibanuma A, Ansah E, Kikuchi K, et al. Determinants of attending antenatal care at least four times in rural Ghana: analysis of a cross-sectional survey. *Glob Health Action*. 2017 Jan 27;10(1). [<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5496066/>] Accessed on 25 Mar 2019 22. Ameyaw EK. Prevalence and correlates of unintended pregnancy in Ghana: Analysis of 2014 Ghana Demographic and Health Survey. *Matern Health Neonatol Perinatol*. 2018 Sep 5;4. [<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6123900/>] Accessed on 21 Jan 2019 23. Birmeta K, Dibaba Y, Woldeyohannes D. Determinants of maternal health care utilization in Holeta town, central Ethiopia. *BMC Health Serv Res*. 2013 Jul 3; 13:256. 24. Lewis S, Lee A, Simkhada P. The role of husbands in maternal health and safe childbirth in rural Nepal: a qualitative study. *BMC Pregnancy Childbirth*. 2015 Aug 4 [<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4523911/>] Accessed on 21 Jan 2019 25. Bhatta DN. Involvement of males in antenatal care, birth preparedness, exclusive breast feeding and immunizations for children in Kathmandu, Nepal. *BMC Pregnancy Childbirth*. 2013 Jan 16; 13:14. 26. Ayushman Bharat - National Health Protection Mission | National Portal of India [<https://www.india.gov.in/spotlight/ayushman-bharat-national-health-protection-mission>] Accessed on 29 May 2019 27. Janani Suraksha Yojana (JSY) | National Health Portal Of India [https://www.nhp.gov.in/janani-suraksha-yojana-jsy-_pg] Accessed on 29 May 2019 28. Office of the Registrar General and Census Commissioner of India (ORGI) (2018) Sample Registration System Statistical Report 2014-16. New Delhi: Office of the Registrar General, India. 29. Lee AC, Kozuki N, Cousens S, Stevens GA, Blencowe H, Silveira MF, et al. Estimates of burden and consequences of infants born small for gestational age in low- and middle-income countries with INTERGROWTH-21st standard:

analysis of CHERG datasets. The BMJ. 2017 Aug 17;358.

[<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5558898/>] Accessed on 28 Mar 2019 30. Blencowe H, Cousens S, Jassir FB, Say L, Chou D, Mathers C, et al. National, regional, and worldwide estimates of stillbirth rates in 2015, with trends from 2000: a systematic analysis. Lancet Glob Health. 2016 Feb;4(2): e98–108.

Tables

Due to technical limitations, Tables 1 & 2 are only available as a download in the supplemental file section.

Figures

Figure 1: Utilisation (%) of full antenatal care and different components in India, NFHS-4 (2015-16)




Figure 1

Utilisation (%) of full antenatal care and different components in India, NFHS-4 (2015-16)

Figure 2: State-wise utilisation (%) of full antenatal care and its components in India, NFHS-4 (2015-16)




Figure 2

State-wise utilisation (%) of full antenatal care and its components in India, NFHS-4 (2015-16)

Figure 3: Concentration curves for full antenatal care utilisation across area of residence (a), maternal education (b), social caste (c)




Figure 3

Concentration curves for full antenatal care utilisation across area of residence (a), maternal education (b), social caste (c)

Figure 4: State-wise inequity in full antenatal care utilization in India, NFHS-4 (2015-16)




Figure 4

State-wise inequity in full antenatal care utilization in India, NFHS-4 (2015-16)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Tablesandfigures11062019.pdf](#)
- [Additionalfile111062019.pdf](#)