

# Radioisotopic Assay of L-carnitine (CN)

Wei Zou (✉ [wzou@dtsc.ca.gov](mailto:wzou@dtsc.ca.gov))

Zou's lab, DTSC, California State Government

---

## Method Article

**Keywords:** L-carnitine

**Posted Date:** July 25th, 2011

**DOI:** <https://doi.org/10.1038/protex.2011.240>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

L-Carnitine +  $[1-^{14}\text{C}]$ acetyl-CoA  $\longrightarrow$   $[^{14}\text{C}]$ acetyl-Carnitine + CoASH Sodium tetrathionate is used to remove CoASH by oxidation so that the reaction is toward the right direction. Then,  $[^{14}\text{C}]$ acetyl-carnitine is separated from  $[1-^{14}\text{C}]$ acetyl-CoA by passing the mixture through an anion-exchange resin AG2-X8 (5g resin/100 mL sample). The isotope content in the elute ( $[^{14}\text{C}]$ acetyl-carnitine) is determined by a scintillation counter.

# Introduction

L-Carnitine (CN) transfers long-chain fatty acids into the mitochondrial matrix for fatty acid  $\beta$ -oxidation. In addition, CN transfers the peroxisomal  $\beta$ -oxidation product, acetyl-CoA, to the mitochondria, maintaining a favorable acetyl-CoA:CoA ratio and stores energy in the form of acetylcarnitine.

# Reagents

Tip: D.I. water: fresh de-ionized water Put everything into the ice when making standard solutions and acetyl-CoA mixture.

- L-Carnitine standard stock solution (1.00 mM): add 19.77 mg L-carnitine-HCl into a 100 mL volumetric flask. Top to the mark with D.I. water.
- 0.02 mM standard solution (1:50): use a 500  $\mu\text{L}$  syringe to add 500  $\mu\text{L}$  1.00 mM solution into a 25 mL volumetric flask. Top to the mark with D.I. water.
- 0.04 mM standard solution (1:25): use a 500  $\mu\text{L}$  syringe to add 1000  $\mu\text{L}$  1.00 mM solution into a 25 mL volumetric flask. Top to the mark with D.I. water.
- 0.10 mM standard solution (1:10): use a 500  $\mu\text{L}$  syringe to add 1000  $\mu\text{L}$  1.00 mM solution into a 10 mL volumetric flask. Top to the mark with D.I. water.
- 0.30 mM standard solution (3:10): use a 500  $\mu\text{L}$  syringe to add 3000  $\mu\text{L}$  1.00 mM solution into a 10 mL volumetric flask. Top to the mark with D.I. water. Transfer the standard solutions into test tubes with Teflon cap. Wrap with Teflon tape. Storage: Put the solutions in the refrigerator (4  $^{\circ}\text{C}$ ). It is stable for several months. Calibration: for each experiment, standardize the stock solution spectrophotometrically at 349 nm by monitoring the disappearance of NADH in the enzyme system.
- Tris buffer (1L of 0.24 M, pH 7.3): dissolve 29.06 g Trizma base (Sigma T6066) in 750 mL of fresh deionized water (D.I. water) and adjust with 1N HCl (about 207 mL) to pH 7.3; pool to the 1 L volumetric flask, top to the mark. Calculation: Problem: to make 1 L of 0.24 M, pH 7.3 Tris buffer ( $\text{pK}_a=8.1$ ) Solution:  $7.3 = 8.1 + \log \left( \frac{[\text{RNH}_2]}{[\text{RNH}_3^+]} \right)$  (1)  $[\text{RNH}_2] + [\text{RNH}_3^+] = 0.24 \text{ M} \times 1 \text{ L}$  (2) then: (1) + (2):  $7.31 \times \text{RNH}_2 = 0.24 \text{ mol}$  (3) then:  $\text{RNH}_3^+ = \text{HCl amount}$ , and,  $\text{RNH}_3^+ + \text{RNH}_2 = \text{Tris base amount}$  Storage: Put the solution in the refrigerator (4  $^{\circ}\text{C}$ ).
- Sodium tetrathionate (7.06 mM): dissolve 1.08 g sodium tetrathionate powder (Sigma S5758) in 500 mL of Tris buffer. Calculation:  $7.06 \text{ mM} \times 500 \text{ mL} = 7.06 \text{ mM} \times 0.5 \text{ L} = 3.53 \text{ mmol} = 3.53 \text{ mmol} \times 306.2 \text{ mg/mmol} = 1080.89 \text{ mg} = 1.08 \text{ g}$  Storage: Put the solution in the refrigerator (4  $^{\circ}\text{C}$ ).
- Acetyl-CoA mixture (0.4 mCi/L + 0.4 mM, or 0.4  $\mu\text{Ci}$   $[1-^{14}\text{C}]$ acetyl-CoA + 0.35 mg acetylCoA/mL deionized water): Thaw  $^{14}\text{C}$ -acetyl-CoA (NEC-313L or ARC 276A) first. Add Tris buffer carefully into the acetyl-CoA bottle (25 mg, Sigma A2181). Pool the solution into a 100 mL of graduate cylinder. Wash the acetyl-CoA bottle several time, collect the washing in the 100 mL graduate cylinder. Add Tris buffer to 75

mL, transfer the solution to a bottle with a glass cap. Pipette 2000  $\mu\text{L}$  of NEN  $[1-^{14}\text{C}]$  acetyl-CoA (1  $\mu\text{Ci}/100 \mu\text{L}$ ) or 200  $\mu\text{L}$  of ARC  $[1-^{14}\text{C}]$  acetyl-CoA (10  $\mu\text{Ci}/100 \mu\text{L}$ ) into the glass bottle. Mix the solution gently in the bottle. Label the bottle as radioactive material. Storage: Put the solution in the refrigerator (4  $^{\circ}\text{C}$ ). It is stable for several days. Calibration:

- Ammonium sulfate (2.9 M): Add 9.58 g ammonium sulfate in 25 mL D.I. water. Storage: Put the solution in the refrigerator (4  $^{\circ}\text{C}$ ).
- Carnitine acetyltransferase stock solution (550 kU/L): the commercial product of carnitine acetyltransferase suspension (Sigma C8757) has 10.08 mg protein in 1.8 mL of 2.9 M ammonium sulfate (5.6 mg protein/mL solution) with the activity of 98 U/mg protein (988 U/10.08g protein/1.8 mL = 550 U/mL = 550 kU/L). Storage: Put the solution in the refrigerator (4  $^{\circ}\text{C}$ ). Usage: For each working day, diluted the stock solution: 1 volume of the stock solution + 10 volume of 2.9 M  $(\text{NH}_4)_2\text{SO}_4$ . Put the diluted enzyme solution in ice. 20  $\mu\text{L}$  diluted enzyme solution (50 U/mL) have an activity of 1.00 U.
- 1 mL reaction mixture contains: 0.5 mL 0.24 M Tris buffer (120  $\mu\text{mol}$ ), 0.25 mL 7.06 mM sodium tetrathionate (1.76  $\mu\text{mol}$ ), and 0.25 mL 0.4 mM acetyl-CoA mixture (100 nmol + 0.1  $\mu\text{Ci}$ )
- Perchloric acid ( $\text{HClO}_4$ ) (1 M or 10 %): Measure 25 mL 70%  $\text{HClO}_4$  solution with a 50 mL volumetric cylinder and pour it into a glass bottle. Use 100 mL volumetric cylinder to measure 300 mL D.I. water and add it into the bottle (dilution factor 1:7). Seal the bottle with a glass stopper and mix the solution well by shaking.
- Hydrochloric acid ( $\text{HCl}$ ) (1 N): Use 50 mL graduate cylinder to add 41.50 mL of 38%  $\text{HCl}$  into a 500 mL volumetric flask. Top to the mark with D.I. water. Mix the solution in the flask. Transfer the solution to a glass bottle with a glass stopper. Titrate the solution with 1N  $\text{KOH}$  if accurate concentration is needed.
- Potassium hydroxide ( $\text{KOH}$ ) (3.5 M): Weigh 39.2 g crystal and dissolve into 20 mL D.I. water in a beaker. Transfer into a 200 mL volumetric flask. Top to the mark with D.I. water.
- Potassium hydroxide ( $\text{KOH}$ ) (1 M): Weigh 11.22 g crystal and dissolve into 20 mL D.I. water in a beaker. Transfer into a 200 mL volumetric flask. Top to the mark with D.I. water.

## Equipment

Preparation of anion exchange columns

1. Calculate and weigh proper amount of AG2-X8 anion exchange resin (0.75 g/sample, i.e., 60 g in 400 mL of water/80 samples/day)
2. Wash AG2-X8 anion exchange resin material with D.I. water. Swirl resin in water. Allow the solution to settle several minutes. Pour off fine articles that do not settle. Repeat.
3. Add 1 N  $\text{HCl}$  and let sit in cold room overnight to charge the resin. Pour off  $\text{HCl}$  and rinse twice with D.I. water.
4. Add water to the resin to make slurry. Deliver equal amount of resin into Poly-Prep columns.
5. Flush columns first with 1.0 mL of water using Pasteur pipette. Don't dry the resin in the column. Then add the tip to the column to stop the flow. Add 1 mL of D.I. water. Cap the column.

## Procedure

Free CN assay procedure: Tip: Most steps should be conducted in ice.

1. Check the accuracy of 100  $\mu\text{L}$  automatic pipette. Run blanks to check the whole procedure. Normally blank is about 100 DPM. If the value is much higher,  $[1-^{14}\text{C}]$ acetyl-CoA may be degraded.
2. Prepare blanks, standards and samples in

duplicates. 3. Add 0.5 mL tris buffer to each sample. 4. Finely tune each sample to pH 7.3 with 1 M HCl or 1 M KOH. Critical step! 5. 0.25 mL sodium tetrathionate solution, and 0.25 mL acetyl-CoA reaction mixture (should be FRESH). Vortex. 6. Make enzyme solution (1:10). Example: Take the stock enzyme solution from the refrigerator and put into the ice. Gently shake it 3 times inversely. Use 20  $\mu$ L automatic pipette to transfer 20  $\mu$ L stock solution into a micro centrifuge tube in the ice. Use 200  $\mu$ L automatic pipette to add 200  $\mu$ L 2.9 M  $(\text{NH}_4)_2\text{SO}_4$  solution into the micro centrifuge tube. Gently mix inversely 3 times. 7. Add 20  $\mu$ L of enzyme into each tube EXCEPT FOR BLANKS. mix inversely for 3 times. 8. Incubate for 30 minutes at 37  $^{\circ}\text{C}$ . 9. Vortex the tube drastically and place in ice for 5 min. 10. Prepare a set of 13x100 mm disposable tubes to contain eluant. 11. Change tubes! Transfer 1.0 mL of the solution in the assay tube into the anion exchange columns. Mark the latest loaded column. 12. Flush columns with 0.5 mL D.I. water. Mix well the eluant. 13. Pipette 0.5 mL elutes into the vials with 10 mL scintillation cocktail and count 14C with Beckman LS 6500.

**I. Deproteinization of serum/bile/lymph**  
 1. Defreeze the tissue overnight. 2. In the morning of the working day, turn on the water bath and centrifuge. 3. Prepare columns, tubes, racks, and scintillation vials. Weigh microcentrifuge tubes. Prepare 6 sets of 2 mL of centrifuge tubes. Weigh two of the 6 sets. Write down the values. 4. Use 100  $\mu$ L automatic pipette to add 50  $\mu$ L of the serum/bile/lymph into a 2 mL microcentrifuge tube. Wash with 50  $\mu$ L of D.I. water. Duplicate. 5. Add 200  $\mu$ L ice-cold 1M  $\text{HClO}_4$ . Vortex drastically. Stand for 10 min. 6. Centrifuge for 10 minutes at  $1.2 \times 10^4$  g at 5  $^{\circ}\text{C}$ . To separate long chain acyl-CNs (acid-insoluble fraction) from short chain acyl-CNs and free CN (acid-soluble fraction) by deproteinization. Medium chain acyl-CNs are partitioned into the two fractions. 7. Use 1000  $\mu$ L automatic pipette to transfer the supernatant to another 2 mL microcentrifuge tube. 8. Add 100  $\mu$ L ice-cold 1M  $\text{HClO}_4$ . Vortex drastically. Stand for 10 min. 9. Centrifuge for 10 minutes at  $1.2 \times 10^4$  g at 5  $^{\circ}\text{C}$ . To repeat separation. 10. Use 1000  $\mu$ L automatic pipette to collect the supernatant to the other 2 mL microcentrifuge tube. 11. Resuspend the precipitate in 100  $\mu$ L 3.5 M KOH at pH 13. 12. Incubate the suspension at 60  $^{\circ}\text{C}$  for 30 min. 13. Neutralize the supernatant with 100  $\mu$ L 3.5 M KOH. 14. Centrifuge the supernatant for 10 min at  $1.2 \times 10^4$  g. To precipitate  $\text{KClO}_4$  15. During this time, prepare 4 sets of tubes for Free CN and acid-soluble CN duplicates. 16. Weigh another set of microcentrifuge tubes. Transfer the supernatant to this set of tubes. Measure the final volume of the supernatant (About 500  $\mu$ L) by weighing the tubes again.

**II. Preparation of free CN**  
 1. Pipette 150  $\mu$ L supernatant into a 12x75 mm tube. 2. Ready for CN assay.

**III. Preparation of acid-soluble CN (free + short chain acyl-CN)**  
 1. Pipette 150  $\mu$ L supernatant into a 12x75 mm tube. 2. Add 100  $\mu$ L 3.5 M KOH to pH 13. Mix well. 3. Incubate the solution at 60  $^{\circ}\text{C}$  for 1 h. To alkalinize the short chain carnitine to free carnitine 4. Cool down the solution and neutralize with 200  $\mu$ L 1 M HCl. 5. Ready for CN assay.

**IV. Preparation of acid-insoluble carnitine (long chain acyl-CN)**  
 1. (Already done in steps 11-12) a. Resuspend the precipitate in 200  $\mu$ L 3.5 M KOH at pH 13. b. Incubate the suspension at 60  $^{\circ}\text{C}$  for 30 min. 2. Vortex the tube drastically. 3. Incubate the suspension at 60  $^{\circ}\text{C}$  for another 30 min. 4. Cool down the suspension and add 400  $\mu$ L 1 M  $\text{HClO}_4$ . Stir drastically. Wait for 10 min. To precipitate the protein in the sample 5. Centrifuge the suspension for 10 min at  $1.2 \times 10^4$  g. 6. During this time, prepare 6 sets of 12x75 mm tubes for acid-insoluble CN duplicates. 7. Transfer the supernatant to another microcentrifuge tube and neutralize with 50  $\mu$ L 1 M KOH. 8. Centrifuge the suspension for 10 min at  $1.2 \times 10^4$  g. To precipitate  $\text{KClO}_4$  9. Weigh another set of microcentrifuge tubes. Transfer the supernatant to this set of tubes.

Measure the final volume of the supernatant (About 1000  $\mu\text{L}$ ) by weighing the tubes again. 10. Take 500  $\mu\text{L}$  supernatant for CN assay. V. Preparation of blanks and standards Blanks: add 200  $\mu\text{L}$  D.I. water and run the same as samples but DO NOT ADD ENZYME. Sample range: By literature, serum total CN is 20-50  $\mu\text{M}$ , i.e., 1-2.5 nmol/50  $\mu\text{L}$  serum. Free CN:acyl CN ratio is about 2:1. Standard curve range: 2 – 5 nmol. Standards for free CN fraction: add 50  $\mu\text{L}$  standard solutions contain 40, and 100  $\mu\text{M}$  L-CN (2, 5 nmol) + 150  $\mu\text{L}$  D.I. water. Standards for acid soluble and acid insoluble CN fractions: add 50  $\mu\text{L}$  standard solutions contain 40, and 100  $\mu\text{M}$  L-CN (2, 5 nmol) + 350  $\mu\text{L}$  D.I. water. VI. Deproteinization of liver 1. Defreeze the tissue overnight. 2. In the morning of the working day, turn on the water bath and centrifuge. 3. Prepare columns, tubes, racks, and scintillation vials. 4. Weigh the homogenizer tube (about 32 g). Use spatula to add 200-300 mg minced tissue into the tube. Weigh again. Write down the exact numbers. 5. Homogenize the tissue in 750  $\mu\text{L}$  D.I. water. Wash the pestle and the tube with 750  $\mu\text{L}$  D.I. water. Collect the washings. Vortex to mix well. To disrupt the cell membrane 6. Measure the volume of the homogenized solution (About 1500  $\mu\text{L}$ ). 7. Use 1000  $\mu\text{L}$  automatic pipette to add 750  $\mu\text{L}$  of the homogenizing solution into a 12x75 mm disposable tube. Duplicate. 8. Add 100  $\mu\text{L}$  ice-cold 1M  $\text{HClO}_4$  (200  $\mu\text{L}$  if not duplicated). Vortex drastically. Stand for 10 min. 9. Centrifuge for 10 minutes at  $1.2 \times 10^4$  g at 5  $^\circ\text{C}$ . To separate long chain acyl-CNs (acid-insoluble fraction) from short chain acyl-CNs and free CN (acid-soluble fraction) by deproteinization. Medium chain acyl-CNs are partitioned into the two fractions. 10. Use Pasteur pipette to transfer the supernatant to another 12x75 mm tube. Combine the supernatant from the duplicates. 11. Resuspend the precipitate in 200  $\mu\text{L}$  3.5 M KOH at pH 13. 12. Incubate the suspension at 60  $^\circ\text{C}$  for 30 min. 13. Neutralize the supernatant with 215  $\mu\text{L}$  1 M KOH. 14. Centrifuge the supernatant for 10 min at  $1.2 \times 10^4$  g. To precipitate  $\text{KClO}_4$  15. During this time, prepare 4 sets of tubes for Free CN and acid-soluble CN duplicates. 16. Weigh another set of tubes. Transfer the supernatant to this set of tubes. Measure the final volume of the supernatant (About 1250  $\mu\text{L}$ ) by weighing the tubes again. VII. Preparation of free CN 1. Pipette 300  $\mu\text{L}$  supernatant into a 12x75 mm tube. 2. Ready for CN assay. VIII. Preparation of acid-soluble CN (free + short chain acyl-CN) 1. Pipette 300  $\mu\text{L}$  supernatant into a 12x75 mm tube. Duplicate. 2. Add 50  $\mu\text{L}$  3.5 M KOH to pH 13. Mix well. 3. Incubate the solution at 40  $^\circ\text{C}$  for 1 h. To alkalinize the short chain carnitine to free carnitine 4. Cool down the solution and neutralize with 160  $\mu\text{L}$  1 M HCl. 5. Ready for CN assay. IX. Preparation of acid-insoluble carnitine (long chain acyl-CN) 1. (Already done in steps 11, 12, and 13) a. Wash the precipitate with deionized water and discard the washings. Duplicate. b. Resuspend the precipitate in 200  $\mu\text{L}$  3.5 M KOH at pH 13. c. Incubate the suspension at 60  $^\circ\text{C}$  for 30 min. 2. Vortex the tube drastically. 3. Incubate the suspension at 60  $^\circ\text{C}$  for another 30 min. 4. Cool down the suspension and add 400  $\mu\text{L}$  1 M  $\text{HClO}_4$ . Stir drastically. Wait for 10 min. To precipitate the protein in the sample 5. Centrifuge the suspension for 10 min at  $1.2 \times 10^4$  g. 6. During this time, prepare 6 sets of 12x75 mm tubes for acid-insoluble CN duplicates. 7. Transfer the supernatant to another tube and neutralize with 100  $\mu\text{L}$  1 M KOH. 8. Centrifuge the suspension for 10 min at  $1.2 \times 10^4$  g. To precipitate  $\text{KClO}_4$  9. Transfer the supernatant to another tube. Measure the final volume of the supernatant. 10. Take 500  $\mu\text{L}$  supernatant for CN assay. X. Preparation of blanks and standards Blanks: add 300  $\mu\text{L}$  D.I. water and run the same as samples but DO NOT ADD ENZYME. Standard curve range: 5 – 15 nmol. Free CN: add 50  $\mu\text{L}$  standard solutions contain 100, and 300  $\mu\text{M}$  L-CN (5, 15 nmol) + 250  $\mu\text{L}$  D.I. water. Total CN: add 50  $\mu\text{L}$  standard solutions contain 100, and 300  $\mu\text{M}$  L-CN (5, 15 nmol) + 450  $\mu\text{L}$

D.I. water. XI. Preparation of adipose tissue total CN 1. Defreeze the tissue overnight. 2. In the morning of the working day, turn on the water bath and centrifuge. 3. Weigh the tube with a Teflon cap. Use spatula to add 200-300 mg minced tissue into the tube. Weigh again. Write down the exact numbers. Use a cotton applicator to clear the sample on the mouth of the tube before weighing. 4. Add 300  $\mu$ L 3.5 M KOH at pH 13. Incubate the solution at 60 oC for 30 min. Vortex the tube drastically. 5. Prepare columns, tubes, racks, and scintillation vials. 6. Incubate the solution at 60 oC for another 30 min. 7. Cool down the solution and add 500  $\mu$ L water. 8. Use dispenser to add 4 mL CM mixture into the tube. Shake upside down 3 times. Wait until there are distinct two layers (or centrifuge for 5 minutes). 9. During this time, prepare 2 sets of 13x100 mm tubes. 10. Transfer the upper layer to another tube. 11. Add 500  $\mu$ L 1 M HClO<sub>4</sub>. Stir drastically. Wait for 10 min. To precipitate the protein 12. Centrifuge the suspension for 10 min at 1.2x10<sup>4</sup> g. 13. Use pasteur pipette to transfer the supernatant to another set of 13x100 mm tubes. 14. Add 175  $\mu$ L 1M KOH. Put the tube into the ice for 10 min. To crystallize KClO<sub>4</sub> 15. Centrifuge the supernatant for 10 min at 1.2x10<sup>4</sup> g. To precipitate KClO<sub>4</sub> 16. Prepare 1 set of 13x100 mm tubes. Weigh the empty tubes. Prepare another set of 12x75 mm tubes. 17. Transfer the supernatant to another tube. 18. Neutralize the supernatant with 50  $\mu$ L 1M HCl. 19. Weigh the tubes again. The net weight is the final volume of the supernatant (assuming 1 g = 1000  $\mu$ L). 20. Take 500  $\mu$ L supernatant for CN assay. XII. Preparation of free CN preparation XIII. Preparation of acid-soluble CN preparation (free + short chain acyl-CN) XIV. Preparation of blanks and standards Blanks: add 500  $\mu$ L D.I. water and run the same as samples but DO NOT ADD ENZYME. Standard range: 1 – 5 nmol. Total CN: add 50  $\mu$ L standard solutions contain 20, 40 and 100  $\mu$ M L-CN (1, 2, 5 nmol) + 225  $\mu$ L D.I. water + 225  $\mu$ L methanol. After washing the CM mixture with 0.2 volume of water, the upper (aqueous) layer contains C:M:W as 3:48:47. Methanol is added to STD tubes to mimic the sample solution.

## Timing

About 2 days for one batch of 20 samples.

## Anticipated Results

Calculation: Subtract the enzyme blank from the zero concentration blank point and use this as a correction factor. The correction factor is added to the counts obtained for each sample. The corrected counts are then applied to the standard curve to determine carnitine content in the sample. Appropriate dilution factors must then be applied to the sample to determine carnitine concentration in the original sample. Liver sample CN range: Free CN: 10 – 15 nmol/sample loading Acid-soluble CN: 10 – 15 nmol/sample loading Acid-insoluble CN: 10 – 15 nmol/sample loading

## References

Zou, W., Noh, S.K., Owen, K.O., & Koo, S.I. (2005) Dietary carnitine enhances the lymphatic absorption of fat and  $\alpha$ -tocopherol in ovariectomized rats. *J. Nutr.* 135: 753-756.  
"http://jn.nutrition.org/content/135/4/753.full":http://jn.nutrition.org/content/135/4/753.full Bhuiyan,

A.K.M.J., Jackson, S., Turnbull, D.M., Aynsley-Green, A., Leonard J.V., and Barlett, K. (1992) The measurement of carnitine and acyl-carnitines: application to the investigation of patients with suspected inherited disorders of mitochondrial fatty acid oxidation. *Clin. Chim. Acta.* 207: 185-204. Cederblad, G, and Lindstedt, S. (1972) A method for the determination of carnitine in picomole range. *Clin. Chim. Acta.* 37: 235-243. MaGarry, J.D., and Forster, D.W. (1976) An improved and simplified radioisotopic assay for the determination of free and esterified carnitine. *J. Lipid Res.* 17: 277-281. Pace, J.A., Wannemacher, R.W., and Neufeld, H.A. (1978) Improved radiochemical assay for carnitine and its derivatives in plasma and tissue extracts. *Clin. Chem.* 24 (1): 32-35.

## **Acknowledgements**

Thanks to Koo, S.I. for mentoring and financial supporting efforts.