

Detection of $O_2^{\cdot-}$ production by chemiluminescence (CL) assay

Naxin Jiang

National University of Singapore

Nguan Soon Tan

National University of Singapore

Bow Ho

National University of Singapore

Jeak Ling Ding

National University of Singapore

Method Article

Keywords: superoxide ion, chemiluminescence, Cypridina luciferin analog

Posted Date: October 31st, 2007

DOI: <https://doi.org/10.1038/nprot.2007.485>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Introduction

Hemoglobin is auto-oxidized from ferrous-hemoglobin to ferric hemoglobin (methemoglobin, metHb) with the production of H_2O_2 . H_2O_2 initiates the catalytic cycle between the ferric (HbFeIII) and ferryl (HbFeIV) hemes, thus activating the pseudoperoxidase activity of metHb, which eliminates the H_2O_2 , producing the superoxide¹. The chemiluminescence (CL) of Cypridina luciferin analog indicates the generation of $O_2^{\cdot -}$ or singlet oxygen, but not that of ozone, hydroxyl radicals or H_2O_2 ². To further confirm that the ROS species produced by metHb and indicated by CLA-CL is superoxide anions but not singlet oxygen, superoxide dismutase (SOD) is applied as a diagnostic confirmation of superoxide anion production.

Procedure

The pseudoperoxidase activity is measured as described², with modifications. 1. Add metHb into a substrate mixture containing 5.3 μM of CLA and 3.3 mM of H_2O_2 in 150 μl of PBS (pH 7.4). 2. Continuously monitor the chemiluminescence, indicated by relative luminescence units (rlu), at one reading per s, for 1 min, with Sirius luminometer (Berthold Detection Systems GmbH, Germany). 3. The pseudoperoxidase activity of the metHb is designated as the relative luminescence units per second (rlu/s). 4. To prove that the ROS species indicated by CLA-CL is superoxide anions, 3 Units/ml of SOD is applied to specifically quench the superoxide anions.

Anticipated Results

The read out of CLA-LA varies from 10^5 - 10^7 rlu/s depending on the concentration of the metHb and the H_2O_2 as the substrate as well as that of CLA as the indicator.

References

1. Alayash, A. I. Hemoglobin-based blood substitutes: oxygen carriers, pressor agents, or oxidants? *Nat Biotechnol.* **17**, 545-549 (1999). 2. Kawano, T., Pinontoan, R., Hosoya, H. & Muto, S. Monoamine-dependent production of reactive oxygen species catalyzed by pseudoperoxidase activity of human hemoglobin. *Biosci Biotechnol Biochem.* **66**, 1224-1232 (2002).

Figures




Figure 1

The pseudoperoxidase activity of methemoglobin (metHb) was demonstrated by CLA-CL. (a) metHb catalyzes the production of $O_2^{\cdot-}$, as shown by the chemiluminescence (CL) assay. (b) SOD significantly reduces metHb-mediated CLA-luminescence, thereby confirming the superoxide anion.