High-throughput Hisx6-tagged protein purification

Chien-Sheng Chen

National Taiwan Ocean University

Heng Zhu

Johns Hopkins University

Method Article

Keywords: high-throughput, protein purification

Posted Date: December 19th, 2007

DOI: <https://doi.org/10.1038/nprot.2007.544>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Introduction

We have developed a high-throughput protein purification protocol that allows us to purify ~4,000 proteins within 10 hours from prepared culture. By combining steps of cell lysis and protein capture on affinity resins in sealed filter plates, we reduced the number of pipetting steps and thus, human errors.

Equipment

Q-Fill2

Procedure

1. Re-suspend cell pellets at 4 °C in 80 µL of lysis buffer, containing 50 mM NaH₂PO₄ at pH 8 with 300 mM NaCl, 20 mM imidazole, CellLytic B (Sigma), Lysozyme (1 mg/mL), Benzonase (50 units/mL), proteinase inhibitor cocktail (Sigma), and PMSF (1 mM/mL). 2. Wash Ni-NTA Superflow (QIAGEN) twice with water and twice with 50 mM NaH₂PO₄ at pH 8 with 300 mM NaCl, 20 mM imidazole using centrifugation. 3. Add 25 µL pre-washed Ni-NTA Superflow (QIAGEN) to each well 4. Transfer the mixtures into bottom-sealed filter plates (Multiscreen Nylon Mesh) 5. Seal the filter plates 6. Incubated for 1.5 hr at 4 °C with vigorously shaking 7. Remove seals 8. Wash the resin-protein complexes 3 times with 250 µL/well of wash buffer I (50 mM NaH₂PO₄ with 300 mM NaCl, 10% glycerol, 20 mM imidazole, 0.01% Triton X-100, at pH 8) and 3 times with Wash buffer II (50 mM NaH₂PO₄ with 150 mM NaCl, 25% glycerol, 20 mM imidazole, 0.01% Triton X-100, at pH 8) using a Q-fill2 (Gentix, UK). 9. Spin the filter plates 10. Add 25 µL of elution buffer (50 mM NaH₂PO₄/150 mM NaCl/25% glycerol/250 mM imidazole/0.01% Triton X-100, pH 7.5) 11. Incubate 15 min at 4 °C 12. Elute proteins into 96-well receiver plates by centrifugation. 13. Repeat 10-12 14. Store the proteins in 96-well plates at -80 °C

Timing

10 hours

Acknowledgements

This work is supported in part by the National Institute of Health (GM071440 to C.H.; U54 RR020839 to H.Z.), W. M. Keck Foundation (to C.H.), the Arnold and Mabel Beckman Foundation (to C.H.), and the Research Corporation (to C.H.).