Appropriateness of Referrals From Primary Care for Lumbar MRI

Susanne Brogaard Krogh (✉ susannebrogaard@gmail.com)

Silkeborg Regional Hospital: Regionshospitalet Silkeborg <https://orcid.org/0000-0002-4563-8548>

Tue Secher Jensen

University of Southern Denmark

Nanna Rolving

DEFACTUM, Central Denmark Region

Malene Laursen

Centre of Elective Surgery, Regional Hospital of Silkeborg

Janus Nikolaj Laust Thomsen

Aalborg University

Casper Brink Hansen

University of Southern Denmark

Christoffer Høj Werenberg

University of Southern Denmark

Erik Rasmussen

University of Southern Denmark

Rune Carlson

University of Southern Denmark

Rikke Krüger Jensen

University of Southern Denmark

Research

Keywords: MRI, Appropriateness of referrals, primary care, lumbar

Posted Date: October 29th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1012188/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Chiropractic & Manual Therapies on February 22nd, 2022. See the published version at <https://doi.org/10.1186/s12998-022-00418-4>.

Abstract

Background:

A number of papers highlight the extent to which low back pain (LBP) is generally mismanaged, especially regarding overuse of magnetic resonance imaging (MRI). International guidelines do not recommend routine imaging, including MRI, and seek to guide clinicians only to refer for imaging based on specific indications. Despite this, several studies show an increase in the use of MRI among patients with LBP and an imbalance between appropriate versus inappropriate use of MRI for LBP. This study aimed to investigate to what extent referrals from general practice for lumbar MRI complied with clinical guideline recommendations in a Danish setting.

Materials and methods:

From 2014-2018, all referrals for lumbar MRI were included from general practitioners in the Central Denmark Region for diagnostic imaging at a public regional hospital. A modified version of the American College of Radiology Imaging Appropriateness Criteria for LBP was used to classify referrals as appropriate or inappropriate, based on the unstructured text in the GPs' referrals. Appropriate referrals included fractures, cancer, symptoms persisting for more than 6 weeks of non-surgical treatment, previous surgery, candidate for surgery or suspicion of cauda equina. Inappropriate referrals were sub-classified as lacking information about previous non-surgical treatment and duration.

Results:

Of the 3,772 retrieved referrals for MRI of the lumbar spine, 55% were selected and a total of 2,051 referrals were categorised. Approximately one quarter (24.5%) were categorised as appropriate, and 75.5% were deemed inappropriate. 51% of the inappropriate referrals lacked information about previous non-surgical treatment, and 49% had no information about the duration of non-surgical treatment. Apart from minor yearly fluctuations, there was no change in the distribution of appropriate and inappropriate MRI referrals from 2014 to 2018.

Conclusion:

The majority of lumbar MRI referrals (75.5%) from general practitioners for lumbar MRI did not fulfil the ACR Imaging Appropriateness Criteria for LBP based on the unstructured text of their referrals. There is a need for referrers to include all guideline-relevant information in referrals for imaging. More research is needed to determine whether this is due to patients not fulfilling guideline recommendations or simply the content of the referrals.

Background

Low back pain (LBP) is one of the most common reasons for patients to seek care in primary care both nationally and internationally [1, 2]. Management strategies for LBP are based on an evaluation of the

individual patient, including case history, physical examination and in some cases imaging. To ensure the best patient care and to optimise management of LBP, several clinical recommendations have been developed, such as those from the American College of Radiology (ACR) Appropriateness Criteria for Low Back Pain [3], the NICE imaging guidelines [4] and the Danish national imaging guidelines [5]. These recommendations are all evidence-based to promote the best clinical practice, including when it is appropriate to use imaging.

Most patients with acute LBP, with or without radiculopathy, have substantial improvements in function and pain within the first 4 weeks, and the guidelines do not recommend routine imaging for those patients [3–5]. In general, imaging is considered for patients with approximately 6 weeks of non-surgical treatment where there has been little or no improvement in their LBP [6]. Furthermore, guidelines recommend that imaging is always considered for patients with suspicion of serious underlying conditions, such as cauda equina syndrome, malignancy, fracture, and infection, i.e. red flag signs and symptoms [6].

Imaging is an important driver of the health care costs associated with LBP, not only because of the direct costs of the test procedures but also because of the downstream effects. Unnecessary imaging can lead to additional tests, follow up on patients, referrals and may even result in invasive procedures of limited or questionable benefit [7].

Despite a clear recommendation against routine imaging of non-specific LBP, this is not well implemented in clinical practice. A systematic review from 2018 showed that 35% of patients referred for lumbar imaging were judged inappropriate due to the absence of red flags for serious pathology, and 32% were judged inappropriate based on the criteria of no clinical suspicion of pathology [8]. Furthermore, a systematic review from 2019 reported that 25% of patients with LBP in primary care underwent imaging and that the use of complex imaging (including MRI) for LBP had increased over the previous two decades [9]. In the Danish national setting, the same pattern of increasing use of MRI was reported, with a 20% increase from 2013 to 2018 (from 70,310 in 2013 to 84,124 in 2018) in patients aged 15 – 85+ years [10].

However, it is unclear to what extent MRI referrals in Denmark are adhering to guideline recommendations. Therefore, this study aimed to investigate how MRI referrals for LBP from general practice clinics in a Danish primary care setting complied with current guideline recommendations.

Objective

To investigate the proportions of appropriate and inappropriate referrals for lumbar spinal MRI from general practice over a period of 5 years.

Method

This study is reported in accordance with the 'Strengthening the Reporting of Observational Studies in Epidemiology' (STROBE) statement [11].

Design

This was a retrospective cross-sectional study.

Setting

In Denmark, general practitioners (GPs) can refer patients with spinal pain for an MRI at a radiology department at a publicly funded hospital. GPs act as gatekeepers for referrals. When referred for an MRI, citizens are free to choose any public hospital, but most patients will choose the local hospital. Data for this study were collected at the Department of Diagnostic Imaging, Silkeborg Regional Hospital in Central Denmark Region from January 1st, 2014 to December 31st, 2018. The Department primarily receives referrals from the municipality of Silkeborg which is the 10th largest of the 98 municipalities in Denmark, with a population of 94,026 (1st of January 2020).

Study population

The study population consisted of referrals from general practice clinics received by the radiology department for patients with LBP referred for a lumbar spinal MRI in the period 2014 - 2018. These referrals were identified by a unique identification number used in the electronic transmission process. This ensured that the sample did not include other primary care providers or hospital departments, such as chiropractors, who have had the right to refer patients for MRI in Denmark since January^{1st} 2014.

The referrals for an MRI of the lumbar spine concerned patients ≥ 18 years of age with LBP, with or without radiating leg pain. The referrals were received by the radiology department and checked for contraindications for MRI. During the data collection period, there were no other criteria for accepting or declining the MRI referral, and according to personal communication with the department, almost all referrals were accepted even if they did not comply with general national guidelines for referral. Referrals were only declined if they did not contain enough information about absolute MRI contraindications, such as non-MRI compliant materials, implants or devices.

Variables

An MRI imaging referral is required to contain information about MRI contraindications and a narrative text, including a short summary of the patient's medical history explaining why the referral is relevant according to current guidelines.

Based on the narrative text, referrals were classified as compliant (appropriate) or non-compliant (inappropriate) according to the 2015 version of the ACR Imaging Appropriateness Criteria for LBP [3]. These criteria (Fig. 1) describe five variants of appropriate MRI referrals ('Variants 2–6'): Variant 2) Suspicion of fracture (e.g. trauma, osteoporosis, chronic steroid use); Variant 3) Suspicion of cancer,

infection, immunosuppression or spondyloarthritis; Variant 4) Candidate for surgery or intervention with persistent or progressive symptoms during or following 6 weeks of conservative management; Variant 5) New or progressing symptoms or clinical findings with a history of prior lumbar surgery; Variant 6) Suspected cauda equina syndrome or rapidly progressive neurological deficit. If the MRI referrals did not include information on any of the above conditions, the referrals were deemed inappropriate (Variant 1). The inappropriate referrals were divided into three subcategories: 1a) no information on previous non-surgical treatment, 1b) no information on the duration of non-surgical treatment, or 1c) other reasons (e.g., errors such as missing narrative text, referrals of cervical / thoracic spine).

In addition to the classification of appropriateness, information on demographic data, including the patient's age and sex, zip code of their residential address, zip code of the referring GP's clinic, MRI referral date and MRI procedure including body region, were collected.

Data collection

MRI referrals were received and stored in the Kodak Carestream RIS (Radiology Information System) version 6.3.0 (Carestream Health, NY, USA). The narrative texts were exported from the RIS archive and were de-identified and uploaded to REDCap electronic data capture tools hosted at Aarhus University REDCap (Research Electronic Data Capture) [12, 13].

Demographic data were extracted from the RIS data. The narrative referral text was used to assess its appropriateness. Five raters participated in the data collection (SBK, CBH, CHW, ER and RC). Each rater entered the classification of the referrals into an Excel spread sheet (Microsoft Excel 2010, Microsoft Corporation, Redmond, WA, USA), developed for the purpose of this study based on Figure 1 and designed to allow only one category per referral to be entered. Details of this process have been described elsewhere[14]. The full dataset was randomly divided into five datasets of equal size by administrative staff in the radiology department who were not part of the study. Due to logistical and time constraints, only a sample representing approximately half of the referrals could be assessed. Prior to data collection, the inter-rater reliability was tested between the five raters. Inter-tester reliability was 'Substantial' to 'Almost perfect' with Kappa values ranging from 0.76 (95% CI: 0.55–0.89) to 0.82 (95% CI: 0.72-0.92) depending on the number of subcategories [14].

Data management and analysis

The appropriate and inappropriate referrals were reported in percentages for the overall categories and for the subcategories. The development over the five-year period was shown graphically by calculating the proportions per year of the two main categories.

Additional variables were presented in tabular form with summarising tables. Baseline characteristics of referred patients were compared between the randomly selected and classified referrals and those not classified. Data management and statistical analysis were performed using Stata version 16 (StataCorp, College Station, Texas, USA).

Results

A total of 4,542 referrals for MRI of the lumbar spine were retrieved. Referrals from general practice clinics accounted for 3,772 (83%) and of these, 2,081 (55%) were allocated for classification. Finally, a total of 2,051 referrals were categorised (Figure 2).

The mean age of the referred categorised patients was 45.1 years and 54.9% were women. There were no differences found between the categorised referrals and those not categorised in relation to the patient's age, sex, year of referral or municipality of residence (Table 1).

Table 1
Baseline characteristics of referred patients by classification status

	Classified	Not classified	Total	P-value	Missing / N (Pct)
n (%)	2,051 (54.8)	1,691 (45.2)	3,742 (100)		0 / 3742 (0.00)
Age, mean (sd)	54.1 (15.8)	54.8 (15.5)	54.4 (15.6)	0.20	0 / 3742 (0.00)
Sex, n (%)					
Men	925 (45.1)	749 (44.3)	1,674 (44.7)		
Women	1,126 (54.9)	942 (55.7)	2,068 (55.3)	0.62	0 / 3742 (0.00)
Year, n (%)					
2014	266 (13.0)	218 (12.9)	484 (12.9)		
2015	220 (10.7)	172 (10.2)	392 (10.5)		
2016	344 (16.8)	271 (16.0)	615 (16.4)		
2017	608 (29.6)	498 (29.5)	1,106 (29.6)		
2018	613 (29.9)	532 (31.5)	1,145 (30.6)	0.85	0 / 3742 (0.00)
Zip code, n (%)					
Outside Eastern Jutland	155 (7.7)	107 (6.4)	262 (7.1)		
Eastern Jutland	1,850 (92.3)	1,554 (93.6)	3,404 (92.9)	0.13	76 / 3742 (2.03)

In the data collection period from 2014 to 2018, a total of 104 general practice clinics referred the 2,051 patients to MRI. The majority of referrals were from general practice clinics in the municipality of 'Silkeborg' (68%) and the rest were from other municipalities in the Central Denmark Region, except for eight referrals (0.39%) received from outside the Region.

Overall, 1,548 (75.5%) of the referrals from general practice clinics did not fulfil the ACR criteria for appropriate referrals, whereas 503 (24.5%) were categorised into one of the five appropriate variants (see Figure 1). Of the five appropriate variants, Variant 5 'New or progressing symptoms or clinical findings with history of prior lumbar surgery' was the most common category (10.5%) followed by Variant 3 'Suspicion of cancer, infection, immunosuppression or spondyloarthritis' (5.8%) and Variant 4 'Candidate for surgery or intervention with persistent or progressive symptoms during or following 6 weeks of conservative management' (5.5%) (Figure 3). Of the 1,548 inappropriate referrals, 782 (50.5%) were subcategorised as Variant 1a because there was no information on previous non-surgical treatment, and 763 (49.3%) as Variant 1b because there was no information on the duration of non-surgical treatment. In three referrals subclassified as 1c 'Other reasons,' the text explained that the patient could not afford or did not wish to engage in conservative care.

The subcategories of appropriate referrals were mostly related to being a candidate for surgery or a worsening of known symptoms (Variants 4 and 5), while suspicion of cancer (2.9%) or spondyloarthritis (2.6%) (Variants 3a and 3b) where the most common pathological reasons for referral (Figure 4).

For referrals categorised as inappropriate, the mean age of patients was 53.3 years (SD 15.65) (range 18-94), and 54.7% were women, and for appropriate referrals, the mean age was 56.4 years (SD 16.0) (range 19-90), and 55.5% were women. The sex of the patient was almost evenly distributed within the appropriate referral categories except for the two smallest categories: Variant 2 'Suspicion of fracture (e.g., trauma, osteoporosis, chronic steroid use)' with 71.0% women (n=22) and Variant 6 'Suspected cauda equina syndrome or rapidly progressive neurological deficit' with 66% women (n=16). Variant 2 had the oldest patients with a mean age of 67.7 years (SD 11.2) and Variant 4 'Candidate for surgery or intervention with persistent or progressive symptoms during or following 6 weeks of conservative management' the youngest with a mean age of 49.3 years (SD 16.3). (Table 2)

Table 2

Characteristics of referred patients by categorisation status (both appropriate and inappropriate referral variants)

	n	Mean age (SD)	Age range	Women (%)	Men (%)
Variant 1	1548	53.4 (15.7)	18-94	54.72	45.28
Variant 2	31	67.7 (11.2)	43-85	70.97	29.03
Variant 3	119	55.9 (17.2)	19-90	52.10	47.90
Variant 4	113	49.4 (16.3)	21-85	51.33	48.67
Variant 5	216	59.0 (14.1)	19-87	56.02	43.98
Variant 6	24	53.3 (16.4)	19-81	66.67	33.33
<i>Inappropriate: Variant 1) Inappropriate referral;</i>					
<i>Appropriate: Variant 2) Suspicion of fracture; Variant 3) Suspicion of cancer, infection, immunosuppression or spondyloarthritis; Variant 4) Candidate for surgery or intervention with persistent or progressive symptoms during or following 6 weeks of conservative management; Variant 5) New or progressing symptoms or clinical findings with a history of prior lumbar surgery; Variant 6) Suspected cauda equina syndrome or rapidly progressive neurological deficit.</i>					

Apart from a statistically insignificant yearly fluctuation, the distribution of appropriate and inappropriate MRI referrals remained stable from 2014 to 2018 (Figure 5). There were no differences in mean age or sex distribution per year.

Discussion

Main findings

This study classified 2,051 lumbar MRI referrals as compliant or non-compliant with the 2015 version of the ACR Imaging Appropriateness Criteria for LBP [3]. The classification was based on the narrative text from the MRI referral, and no patients were clinically evaluated in this study.

Three-quarters (75.5%) of the MRI referrals were deemed inappropriate, and 24.5% were classified as appropriate. The distribution was stable from 2014 to 2018. To our knowledge, no active implementation strategies targeting imaging for spinal pain conditions in general practice were carried out during that period.

Comparison with other studies

Many studies have investigated the appropriateness of lumbar MRI, and a variety of methods have been used [8, 15–18]. Some studies used imaging guidelines to clarify the appropriateness of MRI, and others used clinical symptoms and/or various standards of red flags [19, 20]. The considerable variation in

methods for classification of appropriateness of MRI makes it difficult to make comparisons between studies.

An Australian systematic review from 2018 estimates the overuse and underuse of imaging in the management of LBP [8]. The review included 33 studies and assessed the use of X-ray, CT and/or MRI imaging referrals for patients presenting for care. Inappropriate referrals in people referred for imaging were assessed in 23 studies and showed a pooled effect of 34.8% when 'Absence of red flag clinical features' was used as the inappropriateness criterion and 31.6% when 'No clinical suspicion of pathology' was the criterion. The majority of studies assessed patients referred from general practice. Four studies[20–23] assessed imaging referrals received by radiology departments from primary care physicians, which we considered were comparable to the setting of our study. Rates of inappropriate referrals ranged from 20–47.9% which were considerably lower than the 75.5% found in our study. One study[23] used the same ACR imaging guideline as in our study to assess appropriateness of MRI referrals and reported the highest rate of inappropriate referrals (47.9%) of the four studies.

A scoping review from 2019 included 23 studies describing adult LBP imaging appropriateness in general practice [19]. A range of red flag features was utilised to determine imaging appropriateness. Most studies considered appropriateness in a binary manner by the presence of any red flag feature. Ten different guidelines were referenced in 16 of the 23 studies, while seven studies (30%) used combined methods or modified guidelines. The method for calculating the proportion of inappropriate imaging varied. Ten percent of the studies used the total number of patients presenting with LBP as the denominator, suggesting most studies underestimated the rate of inappropriate imaging and did not capture where imaging was not performed for clinically suspicious LBP.

Both the Australian systematic review from 2018 [8] and the scoping review from 2019 [19] conclude that many different methodologies are used to assess LBP imaging appropriateness.

The current study is therefore not directly comparable to the studies mentioned above. Although they are based on data from MRI referrals, none of them uses the same guideline or checklist to classify referrals, and they do not report the same considerably high number (75.5%) of inappropriate MRI referrals as was found in our study. This large proportion of inappropriate referrals in the current study was probably due to the strict criteria for appropriateness. Only precise information on previous non-surgical treatment and the duration of that treatment in the referral text was used in the evaluation of appropriateness. If information about previous treatment or duration was not provided, the referral was classified as inappropriate, even if text like "*the patients have for some time maintained training*" was available. In cases like this, it was unclear what type or level of training had been performed or what period of time was involved (more or less than 6 weeks), which is essential information in the evaluation of appropriateness when using the ACR imaging guideline. If the criterion of duration was omitted, 38.3% (75.5 minus 37.2%) of the MRI referrals would have been inappropriate which is still higher but closer to the results reported in the previously mentioned literature [8, 19]. It is possible that important clinical symptoms were absent due to oversight in the referrals or that information about non-surgical treatment

or duration was not mentioned because of a lack of knowledge about imaging referral guidelines for LBP. Furthermore, there was a tradition in the department of accepting all referrals despite the lack of information and no feedback procedures existed to inform the GPs that some referrals lacked proper information. This could partly explain why important information was not included in the referrals, which led to a substantially higher number (75.5%) of inappropriate MRI referrals in our study.

Demographic data

More than half of the appropriate referrals were categorised as either Variant 4) 'Candidate for surgery or intervention with persistent or progressive symptoms during or following 6 weeks of conservative management' or Variant 5) 'New or progressing symptoms or clinical findings with a history of prior lumbar surgery'. Both variants are related to clinical management rather than suspicion of serious pathology. In the current study, only 8.5% of all referrals (35% of the appropriate referrals) were referred due to suspicion of serious pathology as in Variants 2, 3 or 6 (fracture, infection, cancer, or cauda equina). In comparison, a study by Gidwani et al. [17] found that 24% of the appropriate MRIs for LBP (n=76.663) had suspicion of red flag conditions as identified by diagnostic codes (IDC-9-CM).

Mean age and sex distribution were relatively comparable between the six variants (Table2). Only Variant 2 (Suspicion of fracture) had a higher percentage of women (70.97%) compared with men (29.03%), and mean age (67.7) was higher when compared to the other variants. This finding is in line with an increasing risk of osteoporotic fractures among women with increasing age.

Perspectives

This study focused on referrals for MRIs of the lumbar spine, in which the narrative text of the referrals was not compared with clinical data. Therefore, it is not possible to determine if a referred patient truly had indications for MRI. Also, it is not possible to measure the proportion of cases where clinical symptoms indicated appropriate MRI but where MRI was not performed (underuse). However, the study reflects what referrals look like in a Danish clinical setting, and it shows that these were not aligned with clinical guideline recommendations. To investigate the 'true' prevalence of guideline-appropriate referrals, future research should contain clinical information at the patient level, including precise information of duration and type of non-surgical treatment. Ideally, appropriate referrals should demonstrate high sensitivity for the detection of serious pathology with a reasonably high specificity to limit unnecessary imaging of patients without serious pathology.

Danish National Clinical Guidelines for non-surgical treatment of patients with recent onset LBP or lumbar radiculopathy [24] was published in 2016. Although it recommends against routine use of MRI for LBP with or without radiculopathy, there was no change in the distribution of appropriate and inappropriate MRI referrals from 2014 to 2018. From a clinical perspective, it seems timely to develop an implementation strategy of imaging guidelines for LBP among GPs to ensure that only patients with a clear indication for MRI are referred and to increase the quality of the referrals.

Methodological considerations

The strengths of this study are the large study sample and the use of a previously tested data collection method that had shown substantial to high inter-rater reliability. The data collection method was based on a well-documented international method for the assessment of imaging guideline appropriateness (ACR)[3]. To ensure a homogeneous practice of referral from general practice clinics and triage by the receiving department, the data were collected from a single imaging department located at a medium-sized hospital with a catchment that included both city and rural areas. However, this could at the same time be a weakness as it is possible that local routines and agreements exist that are not applicable to other parts of the country.

In everyday clinical practice, imaging referrals are not controlled and monitored against clinical findings and symptoms. Although this study cannot tell if the patients had an appropriate indication that was not reflected in the narrative text of the referral, the results do reflect that in everyday practice there is no evidence that guidelines are implemented in either GP referrals or at the receiving imaging department.

Conclusion

The majority of lumbar MRI referrals (75.5%) from general practitioners for lumbar MRI did not fulfil the ACR Imaging Appropriateness Criteria for LBP according to the unstructured text of the referrals. This study cannot determine whether this is due to patients not fulfilling the criteria or to the content of the information in the referrals. There is a need for those referring patients to include all guideline-relevant information in the referrals for imaging. More research is needed to determine whether this is due to patients not fulfilling guideline recommendations or simply to the content of the referrals.

Declarations

Acknowledgements:

We want to thank Hanne la Cour Ginnerup, medical secretary and Andrew Bolas, REDCap manager, from Regional Hospital Silkeborg for their assistance with data collection. We also thank English language editor Asger Torning for proof-reading the paper.

Authors' contributions:

All authors were involved in the design of the study and interpretation of results. SBK, CBH, CHW, ER and RC conducted the data collection, and RKJ did the analyses. SBK wrote the initial draft of the manuscript. All authors have read, improved the draft manuscript, and approved the final manuscript.

Funding:

No funding was received for this study.

Availability of data and materials:

The dataset used in the current study is available from the corresponding author on reasonable request.

Ethics approval and consent to participate:

This study was registered with the Danish Data Protection Agency (j.no. 637374). Danish regulations stipulate that register-based studies do not require research ethics approval.

Competing interests:

RKJ and TSJ are partially employed at the Chiropractic Knowledge Hub funded by the Danish Foundation for Chiropractic Research and Postgraduate Education.

References

1. Hoy D, March L, Brooks P, Blyth F, Woolf A, Bain C, *et al.* The global burden of low back pain: Estimates from the Global Burden of Disease 2010 study. *Annals of the Rheumatic Diseases* 2014; **73**(6): 968–974. DOI: 10.1136/annrheumdis-2013-204428.
2. Flachs E, Eriksen L, Koch M, Ryd J, Dibba E, Skov-Ettrup L, *et al.* *Sygdomsbyrden i Danmark*. Copenhagen: 2015.
3. Patel ND, Broderick DF, Burns J, Deshmukh TK, Fries IB, Harvey HB, *et al.* ACR Appropriateness Criteria Low Back Pain. *Journal of the American College of Radiology* 2016; **13**: 1069–1078. DOI: 10.1016/j.jacr.2016.06.008.
4. National Institute for Health and Care Excellence. *Low back pain and sciatica in over 16s: assessment and management*. 2016.
5. Kallestrup L, Rubak J, Jensen L. *Forløbsprogram for lænderygssmerter*. Region Midtjylland: 2012.
6. Downie A, Williams CM, Henschke N, Hancock M, Ostelo RWJG, De Vet HCW, *et al.* Red flags to screen for malignancy and fracture in patients with low back pain: Systematic review. *BMJ (Online)* 2013; **11**(Dec): 347. DOI: 10.1136/bmj.f7095.
7. Chou R, Fu R, Carrino JA, Deyo RA. Imaging strategies for low-back pain: systematic review and meta-analysis. *The Lancet* 2009; **373**(9662): 463–472. DOI: 10.1016/S0140-6736(09)60172-0.
8. Jenkins HJ, Downie AS, Maher CG, Moloney NA, Magnussen JS, Hancock MJ. Imaging for low back pain: is clinical use consistent with guidelines? A systematic review and meta-analysis. *The Spine Journal* 2018. DOI: 10.1016/j.spinee.2018.05.004.

9. Downie A, Hancock M, Jenkins H, Buchbinder R, Harris I, Underwood M, *et al.* How common is imaging for low back pain in primary and emergency care? Systematic review and meta-analysis of over 4 million imaging requests across 21 years. *British Journal of Sports Medicine* 2019; 1–12. DOI: 10.1136/bjsports-2018-100087.
10. Sundhedsdatastyrelsen. Landspatientsregisteret.
https://www.esundhed.dk/home/registre/landspatientsregisteret/radiologiske_ydelser.
11. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Journal of Clinical Epidemiology* 2008; **61**(4): 344–349. DOI: 10.1016/j.jclinepi.2007.11.008.
12. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)-A metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics* 2009; **42**(2): 377–381. DOI: 10.1016/j.jbi.2008.08.010.
13. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, *et al.* The REDCap consortium: Building an international community of software platform partners. *Journal of Biomedical Informatics* 2019; **95**(December 2018): 103208. DOI: 10.1016/j.jbi.2019.103208.
14. Krogh SB, Jensen TS, Rolving N, Laursen M, Thomsen, Janus Nikolaj Laust Hansen CB, Werenberg CH, *et al.* Categorisation of lumbar spine MRI referrals in Denmark as compliant or non-compliant to international imaging guidelines: an inter-rater reliability study. *Chiropractic and Manual Therapies* 2020; **29**(12): 1–8. DOI: 10.1186/s12998-021-00370-9.
15. Flynn TW, Smith B, Chou R. Appropriate Use of Diagnostic Imaging in Low Back Pain: A Reminder That Unnecessary Imaging May Do as Much Harm as Good. *Journal of Orthopaedic & Sports Physical Therapy* 2011; **41**(11): 838–846. DOI: 10.2519/jospt.2011.3618.
16. Khoury M, Tolentino M, Haj-Ahmad Z, Lilek C, Law MP. Assessing Appropriateness of CT and MRI Referrals for Headache and Lumbar: A Canadian Perspective on Patient-Centered Referrals. *Journal of Medical Imaging and Radiation Sciences* 2019; 1–8. DOI: 10.1016/j.jmir.2019.08.007.
17. Risha Gidwani D, Patricia Sinnott P, Tigran Avoundjian M, Jeanie Lo M, Steven M. Asch MM, and Paul G. Barnett P. Inappropriate Ordering of Lumbar Spine Magnetic Resonance Imaging: Are Providers Choosing Wisely? *American Journal of Managed Care* 2016; **22**(2): 68–76.
18. Wang KY, Yen CJ, Chen M, Variyam D, Acosta TU, Reed B, *et al.* Reducing Inappropriate Lumbar Spine MRI for Low Back Pain: Radiology Support, Communication and Alignment Network. *Journal of the American College of Radiology* 2018; **15**(1): 116–122. DOI: 10.1016/j.jacr.2017.08.005.
19. Yates M, Oliveira CB, Galloway JB, Maher CG. Defining and measuring imaging appropriateness in low back pain studies: a scoping review. *European Spine Journal* 2020; **29**(3): 519–529. DOI: 10.1007/s00586-019-06269-7.
20. Jenkins HJ, Downie AS, Maher CG, Moloney NA, Magnussen JS, Hancock MJ. Imaging for low back pain: is clinical use consistent with guidelines? A systematic review and meta-analysis. *The Spine*

Journal 2018; **18**(12): 2266–2277. DOI: 10.1016/j.spinee.2018.05.004.

21. Kovacs FM, Arana E, Royuela A, Cabrera A, Casillas C, Piñero P, *et al.* Appropriateness of lumbar spine magnetic resonance imaging in Spain. *European Journal of Radiology* 2013; **82**(6): 1008–1014. DOI: 10.1016/j.ejrad.2013.01.017.
22. Emery DJ, Shojania G, Forster AJ, Mojaverian N, Feasby TE. Overuse of Magnetic Resonance Imaging. *JAMA INTERN MED* 2013; **173**(9): 823–825. DOI: 10.1001/jamainternmed.2013.3041.
23. Kennedy SA, Fung W, Malik A, Farrokhyar F, Midia M. Effect of governmental intervention on appropriateness of lumbar MRI referrals: A canadian experience. *Journal of the American College of Radiology* 2014; **11**(8): 802–807. DOI: 10.1016/j.jacr.2013.12.022.
24. Stochkendahl MJ, Kjaer P, Hartvigsen J, Kongsted A, Aaboe J, Andersen M, *et al.* National Clinical Guidelines for non-surgical treatment of patients with recent onset low back pain or lumbar radiculopathy. *European Spine Journal* 2018; **27**(1): 60–75. DOI: 10.1007/s00586-017-5099-2.

Figures

Figure 1

Flow chart for the classification of the referrals according to the modified ACR Imaging Appropriateness Criteria.

Figure 2

Flow of the selection and classification process.

Figure 3

Classification of the referrals according to the modified ACR Imaging Appropriateness Criteria.

Inappropriate referrals: Variant 1) Inappropriate referral; Appropriate referrals: Variant 2) Suspicion of fracture; Variant 3) Suspicion of cancer, infection, immunosuppression or spondyloarthritis; Variant 4) Candidate for surgery or intervention with persistent or progressive symptoms during or following 6 weeks of conservative management; Variant 5) New or progressing symptoms or clinical findings with a history of prior lumbar surgery; Variant 6) Suspected cauda equina syndrome or rapidly progressive neurological deficit.

Figure 4

Classification of the referrals according to subcategorisation of the modified ACR Imaging Appropriateness Criteria. Inappropriate referrals: Variant 1: 1a) no information on previous non-surgical treatment; 1b) no information on the duration of non-surgical treatment; 1c) other reasons. Appropriate referrals: Variant 2) Suspicion of fracture due to a) trauma, b) osteoporosis or c) chronic steroid use; Variant 3a) Suspicion of cancer, 3b) spondyloarthritis, 3c) infection or 3d) immunosuppression; Variant 4) Candidate for surgery or intervention with persistent or progressive symptoms during or following 6 weeks of conservative management; Variant 5) New or progressing symptoms or clinical findings with a history of prior lumbar surgery; Variant 6) Suspected cauda equina syndrome or rapidly progressive neurological deficit.

Figure 5

Trajectory of appropriate and inappropriate MRI referrals from 2014 to 2018.