

# Health-related quality of life among 13–14 year old adolescents with overweight - a mixed methods approach

**Turid Kristin Bigum Sundar**

Universitetet i Oslo

**Kirsti Riiser**

OsloMet - storbyuniversitetet

**Milada Småstuen**

OsloMet - storbyuniversitetet

**Randi Opheim**

Universitetet i Oslo

**Knut Løndal**

OsloMet - storbyuniversitetet

**Kari Glavin**

VID Vitenskapelige Hogskole

**Sølvi Helseth** (✉ [solvi@oslomet.no](mailto:solvi@oslomet.no))

OsloMet - storbyuniversitetet <https://orcid.org/0000-0002-6411-9057>

---

## Research

**Keywords:** Health-related quality of life, adolescents, overweight, obesity, physical activity, mixed methods

**Posted Date:** December 26th, 2019

**DOI:** <https://doi.org/10.21203/rs.2.19545/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Health and Quality of Life Outcomes on May 29th, 2020. See the published version at <https://doi.org/10.1186/s12955-020-01413-0>.

# Abstract

## Background

Overweight and obesity are public concerns with risk of adverse health outcomes. Health-related quality of life (HRQoL) is lower in adolescents than children in general. An increase in body mass index (BMI) is associated with a decrease in HRQoL. The purpose of this study was to measure and explore the HRQoL among adolescents with overweight or obesity who had participated in an intervention study with the aim of increasing PA, reducing BMI and promoting HRQoL.

## Methods

Mixed methods, with a convergent design, were used to investigate how different methodological approaches could expand our understanding of the adolescents' HRQoL. Quantitative post-intervention data on HRQoL were collected among the 84 intervention participants, aged 13–14 years, using the KIDSCREEN 52 questionnaire. The data were compared with a Norwegian reference population of 244 individuals, and analysed using a non-parametric Mann-Whitney test. Qualitative semi-structured interviews were conducted with 21 adolescents from the intervention. A directed approach to content analysis was adopted, using the ten sub-scales from KIDSCREEN 52.

## Results

HRQoL in the intervention sample was significantly reduced on the sub-scale of physical well-being compared to the reference population. The reference population scored significantly lower than the intervention sample on the sub-scale of parent relation and home life. No significant differences were found on the other sub-scales. The qualitative data supported the quantitative findings on the sub-scale of physical well-being, but showed that perceptions of fitness, energy level or health could vary. Regarding parent relations, the interviewees extended this to include relationships to other family members as equally important. Most of the interviewees expressed a negative view of their bodies, but not their clothing or accessories. This may explain why no statistically significant differences were found on these aspects in the results from the KIDSCREEN questionnaire.

## Conclusion

The use of the KIDSCREEN 52 instrument gave important indications about the adolescents' HRQoL. Combining methods enabled a comprehensive approach to research on HRQoL, indicating better ways of providing help. More research using the benefits of mixed methods approaches is needed to further elucidate these findings.

# Introduction

Overweight and obesity are public health concerns globally. Extensive research has investigated the medical challenges associated with these conditions. Growing evidence indicates that overweight or

obesity may impact negatively on well-being and functional health [1, 2]. Adolescence is considered a critical period in life, characterized by physical, psychological and social changes. Most adolescents go through this period without any trauma. However, studies report that in the general population, adolescents rate life satisfaction and health-related quality of life (HRQoL) lower than children, and girls lower than boys [3, 4]. Research shows that this is further aggravated by increased Body Mass Index (BMI) [5, 6]. Adolescents with overweight or obesity are more susceptible to bullying related to weight. This is found to be associated with increased vulnerability, with lower motivation for physical activity (PA), low self-esteem, higher behavioural avoidance and emotional coping strategies [7].

HRQoL is a multidimensional construct for assessing subjective perspectives of psychosocial, physical and functional components of well-being and health as perceived by individuals [8]. The use of such instruments may help identify people at risk of developing health problems, or assist in determining the burden of a health problem, such as overweight or obesity [9]. In the last decade, research reflecting children's views has become increasingly important [10]. From 2001 to 2004, KIDSCREEN, a multidimensional generic questionnaire, was developed and tested in Europe as a collaborative project between 13 countries. The KIDSCREEN instrument comprises three versions for children, KIDSCREEN 52, KIDSCREEN 27 and KIDSCREEN 10, as well as a parent proxy version. The content of the questionnaires is based on literature reviews, focus group interviews with children and parents, and opinions from experts on HRQoL measurements for children. The instruments aim to capture HRQoL from the perspectives of both healthy and chronically ill children by assessing their subjective health and well-being [11].

A survey of HRQoL on 17,159 children across 10 European countries using the generic KIDSCREEN-52 instrument found that overweight or obese children and adolescents had a statistically significant lower mean HRQoL score than children of normal weight. The scores were lowest in the sub-scales of physical well-being, self-perception, social acceptance and bullying [12]. Other studies on overweight or obese children and adolescents reveal similar results [8, 13–16].

Physical activity (PA) during childhood is associated with a multitude of health benefits due to its preventive effects on several physical conditions and its ability to stimulate cognitive performance and mental well-being [17]. A review article and a meta-analysis provide evidence that adolescents who are overweight or obese are less physically active and less physically fit than their leaner counterparts [18, 19]. While physical activity is found to be positively related to HRQoL, sedentary behaviour, on the other hand, is found to be negatively related [20]. A closer look at a combination of Body Mass Index (BMI) and PA among adults indicates that PA is important for HRQoL regardless of weight status [21]. Research shows that the association between PA, body composition and fitness on HRQoL among children has received less attention [21, 22]. Little is known about other factors that may contribute to the results, such as functional difficulties due to overweight, or psychosocial mediators [14]. Recent research shows that self-determination theory (SDT) is useful for understanding perspectives on adolescents' well-being and quality of life [23]. SDT postulates that subjective well-being or feeling positive and being satisfied with life is determined by satisfying the three basic psychological needs; autonomy, competence and relatedness, and that social environments that promote such needs provide optimal conditions for an

individual's development, well-being and motivation [24]. Motivational Interviewing (MI) is a collaborative, person-centred form of counselling approach that can be understood in terms of SDT. The method has shown to be effective in strengthening motivation for change [25].

The main purpose of this study was to measure and explore the HRQoL among adolescents with overweight or obesity who had participated in an intervention study, Young & Active (ClinicalTrials.gov NCT01700309). The aim of the intervention was to increase PA, reduce BMI and promote HRQoL, through the use of principles from SDT and MI [26].

Mixed methods were chosen to investigate how different methodological approaches could expand our understanding of the adolescents' HRQoL.

The research questions were as follows:

1. How do the participants in the Young & Active intervention group respond on the KIDSCREEN questionnaire compared to the Norwegian survey reference population?
2. How do the Young & Active participants describe their health-related quality of life in the interviews?
3. How do the different methodological approaches complement and nuance each other?

## Methods

### Mixed methods

We used a mixed methods approach with a convergent design. The quantitative and qualitative data were collected in parallel during the same time frame [27]. A subsample of interviewees was recruited from the Young & Active population that completed the KIDSCREEN 52 survey [28, 29]. The quantitative and qualitative sets of data were analysed separately, and then brought together for analysis and comparison during the discussion.

### Study context and participants

The 12-week intervention study, Young & Active, was conducted with adolescents, age 13 – 14, from 2012 - 2014. School nurses in three counties from rural and urban districts of south-eastern Norway assisted in the recruitment to Young & Active. Following the recommended height and weight measurement in the 8<sup>th</sup> year [30], adolescents with age and gender adjusted BMI above 25 were invited to participate in the intervention. Adolescents involved in treatment programmes or other interventions were not eligible. A total of 84 adolescents volunteered to participate. As part of the Young & Active study, measurements of height and weight, a standard fitness test, and self-report instruments were individually completed at the adolescents' schools. During the intervention, the adolescents were asked to register all weekly PA. This allowed them to follow their activity level on graphs and figures. They also kept a PA diary and could ask questions about PA if they wished. Once a week, they received online feedback and counselling from one

of the researchers, an experienced physiotherapist. The counselling was based on principles from Motivational Interviewing (MI). Knowing that overweight or obesity, and adolescence in itself, potentially increases adolescents' vulnerability, we considered it relevant to complement the study with qualitative interviews of a sample of the participants, following the post-intervention follow-up in the Young & Active intervention study [12-14].

The qualitative data involved semi-structured interviews with a sample of 21 adolescents from the Young & Active intervention group. All 84 adolescents were asked if they wanted to participate in the interviews. Based on a strategy of convenience sampling (14)[31], we accepted those adolescents who were available in the different schools and municipalities. Seven boys and 14 girls agreed to our request.

In addition, data from a larger Norwegian HRQoL validation study involving 244 individuals, age 13–14, were applied. These data were extracted from a cross-sectional study, conducted across 20 randomly selected schools in Norway, involving 1,123 children, age 8–18 [32].

## **Measurement of health-related quality of life**

HRQoL was measured using the Norwegian version of the KIDSCREEN-52 questionnaire [32]. The questionnaire covers ten subscales of social, mental and physical well-being, with five to seven sub-items rated by each individual using a five-point Likert scale [11, 33]. The scale indicates the frequency of certain behaviours or feelings (1=never to 5=always), or the intensity of an attitude (1=not at all to 5=extremely). The time frame refers to the previous week. The number of items in each dimension is illustrated in Table 1. The dimension scores in KIDSCREEN were converted into a 0–100-point scale, with 100 indicating the best HRQoL and 0 the worst. The KIDSCREEN instrument has been tested on 22,827 children aged 8–18 years across Europe and has shown satisfactory reliability and validity, both on healthy children and children with chronic health conditions [34]. In Norway, the instrument was tested and validated on 1,123 children, aged 8–18 years, showing a Cronbach's alpha value above 0.80 for all KIDSCREEN scales, which suggests good internal consistency reliability for the instrument [11, 12, 32].

Table 1  
Overview of KIDSCREEN-52, a short description [35]

	<b>KIDSCREEN scales</b>	<b>No. of items</b>	<b>Short descriptions</b>
1.	Physical well-being	5	Level of physical activity, energy, fitness and health
2.	Psychological well-being	6	Positive and negative emotions, life satisfaction and optimism
3.	Mood	7	Degree of depressive moods and stressful feelings
4.	Self-perception	5	Positive or negative bodily appearance, through questions about satisfaction with looks, clothes and accessories
5.	Autonomy	5	Freedom of choice and opportunities to create social and leisure time
6.	Parent relation and home life	6	Relationships with parents and home atmosphere
7.	Financial resources	3	Perceived quality of financial resources
8.	Social support and peers	6	Social relations with other adolescents
9.	School environment	6	Perceptions of cognitive capacity, feelings about school
10.	Social acceptance (bullying)	3	Aspects of feeling rejected by peers in school

## Statistical analysis

We used IBM SPSS version 25 to conduct statistical analyses of the KIDSCREEN data. Continuous data were not normally distributed and were therefore described using median and range. Categorical data were presented as counts and percentages.

As the data did not follow normal distribution, non-parametric Mann-Whitney tests were used to examine the differences between the Young & Active sample and the reference sample on the ten dimensions of the KIDSCREEN scales.

In order to achieve an accurate estimate, bootstrapping was applied to the sample. When a sample is relatively small, this is a useful method of increasing the sample size without having to collect repeated samples from the population of interest [36]. P-values < 0.05 were considered statistically significant and

all tests were two-sided. As the study was considered exploratory, no correction for multiple testing was performed.

## **Qualitative interviews**

The research group developed a semi-structured interview guide containing 11 open-ended questions. The interview guide ensured consistency and flexibility in the approach to enable the adolescents' accounts to emerge. Probes were open and specific to the adolescents' comments. The interview guide included questions that focused on the adolescents' everyday lives. Examples of questions included; could you describe yourself to me, your everyday life, health, sleep, self-image, body image, life satisfaction, moods and emotions, family relations, leisure time activities, friendships, school, views on PA and perceptions of participation in Young & Active. The first author conducted all 21 individual interviews following the post-intervention measurements in Young & Active. Interviews took place during school hours at the adolescents' schools with only the adolescent and the interviewer present. Each interview lasted an average of 20 minutes. All interviews were audio-recorded and transcribed verbatim by the first author.

## **Qualitative content analysis**

The interview data were analysed using a qualitative directed content analysis [37]. The ten KIDSCREEN subscales were used as key concepts and theoretical basis (see Table 1).

The first author listened to the recordings and read through the transcripts multiple times, and coded the adolescents' statements according to the ten subscales and definitions in KIDSCREEN 52. All statements were then summarized. The results were discussed and agreed upon in the research group.

## **Ethical considerations**

The study was approved by the Norwegian Regional Committee for Medical and Health Research Ethics (REK no. 2010/2978A). Parents gave their written consent for their children to participate in the Young & Active intervention study and in the qualitative research interviews after completion of the intervention. The adolescents also gave their written assent. Information about the study, confidentiality, potential hazards and benefits, the purpose of the study and the right to withdraw were emphasised, both in the written material and verbally to the adolescents. The anonymity and confidentiality of the participants and the material were ensured by using subject identifier codes instead of names. School nurses were aware of the study and were prepared to help adolescents who needed support in cases of personal difficulties.

## **Results**

## HRQoL data

In total, we analysed KIDSCREEN data on 84 individuals from the intervention sample and 244 from the Norwegian reference sample. The proportion of girls compared to boys was higher in both samples (61% and 58% for Young & Active and the reference population, respectively). Statistically, findings from the Young & Active sample showed a significantly lower HRQoL on the subscale of physical well-being. The Norwegian reference population had a statistically significant lower HRQoL on the parent relations dimension (see Table 2).

Table 2

Comparison of median scores between study sample and reference sample on Kidscreen-52 domains

	Young & Active			Reference			P-value
	N	Median	95% CI	N	Median	95% CI	
Physical well-being	67	55.00	50.00–60.00	244	70.00	67.50–70.00	< 0.01
Psychological well-being	72	77.09	70.83–85.42	244	79.17	75.00–79.17	0.85
Mood	75	82.14	78.57–89.29	244	85.71	82.14–85.71	0.81
Self-perception	72	62.50	47.50–75.00	244	70.00	67.50–75.00	0.46
Autonomy	70	80.00	70.00–87.50	244	70.00	70.00–75.00	0.10
Parent relations	70	83.33	75.00–91.67	244	75.00	75.00–79.17	< 0.05
Financial resources	72	100.00	87.50–100.0	244	83.33	75.00–87.50	0.06
Social support and peers	70	83.33	77.09–87.50	244	70.83	68.75–75.00	0.19
School environment	71	68.75	62.50–75.00	244	62.50	62.50–66.67	0.25
Social acceptance	74	100.00	91.67–100.0	244	91.67	91.67–100.00	0.25

Further, all analyses were stratified by gender (data not shown in table). The subscale of physical well-being showed that both boys and girls from the Young & Active sample reported a statistically significantly lower HRQoL for this subscale compared to the reference population, with values ranging from 45.00–60.00 for girls and from 50.00–70.00 for boys. On the subscale of parent relations, the girls from the reference sample had statistically a significantly lower HRQoL score compared to the girls from

the Young & Active sample, ranging from 70.83–81.20 in the reference sample and from 79.17–91.67 for Young & Active. Among the boys there were no statistically significant differences between the groups on this subscale. No other statistically significant differences were found for any of the remaining subscales.

## Qualitative data

There were qualitative findings on all ten subscales, although the findings on the subscales of Physical Well-being and Self-Perception were the most prominent and will be treated in more detail than the other qualitative data. The findings in the remaining subscales will be presented in a summary. Quotes from the interviewees used in the text have been translated from Norwegian.

## Statements about Physical Well-Being

Most of the adolescents stated that they felt increasingly fit and motivated to engage in PA after participating in Young & Active. However, they still described their health, fitness and energy level as being below average or lower than their peers. For example, one girl said: *I've always struggled with physical activity because I'm not the strongest or the fastest person. I think my fitness level has increased but I haven't noticed that much change (Inf. 10)*. When talking about their health, many adolescents expressed that their health was poor. They attributed this to weighing too much, not being active enough or liking food too much. A few talked about having health problems such as asthma, migraine, or complaints regarding neck, knees or heels. Other adolescents presumed that their health was ok, since they were not ill. Several adolescents emphasized that the extent to which they felt energetic, fit and healthy could vary significantly from week to week or even from day to day, depending on other factors such as amount of sleep, food intake, pain or disease – as one boy said: *My health differs from day to day. Sometimes I feel less energetic, perhaps because I went to bed too late (Inf. 18)*. For some adolescents, feeling energetic was associated with the weather. One girl said: *I feel I can have loads of energy if the weather's nice, then I can go out and run around. But if it's raining, I'll lie on the sofa and play computer games (Inf. 5)*. When enjoying and mastering a physical activity, even pain seemed to be less significant, as one boy stated: *I have some heel problems. I don't notice it when I'm slalom skiing and slalom skiing is my favourite activity (Inf. 1)*. Other adolescents expressed similar opinions and stated that the extent to which they became involved in physical activities or felt energetic depended on their feelings about the physical activity they were thinking about engaging in, as this girl stated: *I guess I have the energy, unless I don't feel like it, and then I can be quite lazy (Inf. 4)*. This girl was regularly active in a sport that she enjoyed, had chosen by herself and which she mastered. When discussing other types of PA, she described herself as a person who didn't like moving. Their feelings about PA also related to exercising together with friends, as this boy stated: *I prefer to exercise with my friends because we push each other. Then I feel I'm getting more out of the exercise and enjoy it (Inf. 6)*.

## Statements about Self-Perception

Most of the adolescents appeared to have a negative view of their bodies, as this girl stated: *I'm not happy with my body. I feel I've grown too fast and that other people look at me. The problem is that all my friends are so thin, and here's me with my large thighs and fat belly. I consider myself fat (Inf. 7)*. Another girl said that she often envied thin girls and wished she could have been like them: *Sometimes I used to think, I want to look like her, and she's so thin and beautiful (Inf. 2)*. For some, thoughts about weight were coupled with their perceptions of self, as this girl explained: *I don't think too highly of myself. It's a bit hard because I'm not used to talking about it, but I'm not happy about myself because I actually think that I weigh too much (Inf. 5)*. Most of the boys appeared to be more preoccupied with building muscles than getting thin, as this boy stated: *I like bigger muscles; they make me feel good (Inf. 1)*. *I'm not that attractive yet, but I will be when I've done some more training (Inf. 17)*. Some of the boys admitted that the main reason they exercised was to lose weight, as this boy stated: *I think a lot about my weight and that's the main reason why I do training (Inf. 3)*. However, like the other boys, he also wanted to achieve weight loss through training and building muscles. A number of the adolescents appeared to distinguish between negative thoughts about their bodies and how they viewed themselves as persons, as one girl said: *I'm sort of satisfied with myself, but not with my body. I want to change my looks and lose weight. But I think my personality is ok (Inf. 11)*. The adolescents' views about themselves were also associated with thoughts about how their friends viewed them, as this boy stated: *I'm a happy person who makes friends easily. I'm satisfied with myself and I think other people like me and regard me as a kind and caring person (Inf. 15)*.

## A summary of findings in the remaining subscales

Within the subscales of psychological well-being and mood, many adolescents stated that they felt happy most of the time but emphasised that this could vary considerably. Some of them said that eating too little could make them feel unhappy and irritable. Several of them stated that if their friends and family were happy, then they were happy, and vice versa. Most of them expressed that spending time with friends made them feel happy and put them in a good mood. They also said that good sleep was important for their moods. Some of them stated that they felt generally insecure about themselves, feeling pressure regarding how they should behave, how they should dress and how they should look.

Within the KIDSCREEN subscales of autonomy, parent relations, financial resources and social acceptance, a few adolescents expressed insufficiency, i.e. feeling a lack of freedom of choice, lack of parental support or too restrictive parents, lack of money to pay for sports activities because of their family's financial situation, or being bullied at school. Most of the adolescents said that they received enough support from their parents and that they had not been bullied. Regarding family relations, they extended this to include siblings and other relatives. Many of them expressed that their feelings were influenced by other family events, such as loss of close family members etc., and not just their direct relationship to their parents.

All the adolescents affirmed that they had friends. Except for a couple of adolescents who told about being bullied at school, the majority said that they were friends with their classmates. All of them stated that they had friends outside school, although for some of them, these were mainly online friends.

Regarding the subscale of school environment, most of the adolescents stated that they liked their schools. The majority said that they felt increased pressure from their parents and teachers about achieving good grades and doing more homework. Several of them expressed that getting bad grades negatively impacted their self-confidence.

## Discussion

We used a mixed methods approach with a convergent design on the post-intervention qualitative and quantitative data [38]. Measuring HRQoL provides important information on how adolescents who are overweight or obese feel about their lives. However, assessing HRQoL through questionnaires is a subjective task, involving cognitive processes that are influenced by biological, psychological and social-environmental factors [39]. Applying both methods provided an opportunity to investigate how the different methodological approaches could complement and nuance each other in understanding the adolescents' HRQoL [37].

PA is associated with many health benefits, including improved HRQoL and self-esteem [40]. Most of the adolescents who had participated in the Young & Active intervention study were still overweight or obese. Post-intervention KIDSCREEN data from the Young & Active study showed that the adolescents had statistically significant lower scores on the subscale of physical well-being compared to the Norwegian reference population. The data was verified as lower among both genders in the intervention group. Other studies on adolescents who are overweight or obese have shown similar results [1, 8, 12, 41, 42].

The qualitative data analysed according to the KIDSCREEN subscale of physical well-being supported the KIDSCREEN results from the Young & Active sample. However, the data highlighted other important factors. Most of the interviewees said that they felt their level of energy, fitness and physical activity was lower than their peers, but still generally better after they had participated in Young & Active. The results from the shuttle run tests in the Young & Active study indicated that the tailored PA intervention had a beneficial short-term effect on cardiorespiratory fitness, HRQoL and BMI [43]. However, the interviewees stated that their physical well-being could vary considerably from day to day, depending on either external or internal factors, such as weather, daily energy level, health conditions, relations to friends and motivation towards an activity. The adolescents' comments demonstrated that the ways in which they might answer questionnaires could vary, depending on their subjective feelings at the time. Also, an analysis of the qualitative data revealed that they had varying understandings of concepts such as health. Some of them described their health as being poor due to being overweight. Others state that they were in good health because they had no diseases. Different ways of understanding concepts could lead to misinterpretation of the same items between subgroups or measurement errors resulting from discrepancies in how the questions were answered in the questionnaires [39].

On the KIDSCREEN subscale of parent relations, the girls in the reference population had statistically significant lower scores than the adolescents in the intervention sample. There was no statistically significant difference between the boys in the two samples. Researchers have suggested that parent-adolescent conflicts peak during early adolescence and that this is a natural part of adolescent development [44, 45]. Research also shows that adolescent girls generally report a lower HRQoL than adolescent boys [46, 47]. This may explain why the girls from the reference population scored lower, although it does not explain the difference between the two groups. Other studies have shown that adolescents with overweight or obesity report a lower HRQoL on their relations to parents and family [48, 49]. The data in this study was post-intervention data. Participation in the intervention may have impacted the adolescents' responses. Getting permission from their parents, and for some being pushed to participate, may have been experienced as supportive. Furthermore, recruitment to studies always entails a selection of participants and the adolescents' parents may have been more initially supportive.

The interview data in the subscale of parent relations supported the findings from the intervention study sample, with a few exceptions. Most of the adolescents expressed positive feelings towards their parents but expanded their descriptions to include relations to other family members as being equally important to their well-being. This has also been reported in another qualitative study [6]. Some of the interviewees appeared to feel that they were more controlled by their parents. These adolescents expressed a generally lower HRQoL. Recent studies have found that satisfaction of the basic psychological needs for autonomy, competence and relatedness significantly predicts positive effects on general well-being and on PA within sport contexts [23, 50]. Studies have shown a correlation between more positive relationships with parents and better weight and health-related behaviours among adolescents, as well as improvements in quality of life among those who experience an increase in parent support [51, 52].

No statistically significant differences were found on the other subscales although the data revealed some dispersion in CI and median between the Young & Active sample and the Norwegian reference sample. The greatest variation was shown on the subscale of self-perception, with the intervention sample showing a CI between 47.50 and 75.00 with a median of 62.50, and the reference sample showing a CI between 67.50 and 75.00 with a median of 70.00. This suggests considerable variation between individuals in the Young & Active sample. KIDSCREEN measurements on this subscale have been found to be lower among adolescents in general compared to children, with decreasing scores among adolescents with overweight and even more so among those who are obese [12, 53]. Research also shows gender differences, with HRQoL among girls declining more than boys during adolescence [47]. In this material, the girls in both groups had lower scores than the boys.

The qualitative data differed from the Young & Active KIDSCREEN results on the subscale of self-perception. Most of the adolescents were concerned about their looks and wanted to change their bodies. However, none of them stated that they were concerned about clothing or other accessories. Recent studies have indicated that adolescents' body image is influenced by the media and current trends in exercise, diet and beauty, which may contribute to unhealthy body perception and beliefs about the ideal body [54]. Body image has always been an important part of societies and cultures, and overweight or

obese individuals are frequently subject to weight-related stigma, either through the media or through bullying [55, 56]. Among adolescents, weight-related stigma is reported to be one of the most prevalent forms of bullying at school [57, 58]. Only a few adolescents in this study stated that they had been bullied. Most of them claimed that they had friends and that they were accepted by their peers. A qualitative study on adolescents' perception of quality of life underlined the importance that adolescents ascribe to friendship [6]. The adolescents in this study described what could be understood as being a coping strategy by separating views of their personalities from views of their bodies, stating that they liked their personality but not their body. Feeling accepted by friends was an important part of a positive view of own personality. The interviewees still expressed weight-related stigma, probably caused by societal influences. If internalized, such sense of stigma could be even more severe and contribute to a reduced HRQoL, as well as interfere with their efforts to improve their health [59, 60]. The general decline in HRQoL among adolescents in this age group regardless of their weight status, as well as not being dissatisfied with clothing and other accessories, may explain why the intervention KIDSCREEN data did not show any statistically significant difference compared to the reference population on this subscale. However, the qualitative data revealed that the adolescents probably struggled more than what emerged from the quantitative data. Furthermore, they experienced increased pressure from parents and teachers regarding succeeding at school. This may have placed a considerable extra burden on the adolescents' self-perception and HRQoL.

## **Strengths and limitations**

The KIDSCREEN data provided important knowledge about HRQoL among adolescents with overweight and obesity that should be taken seriously. Additional qualitative data on the adolescents' experiences helped to extend knowledge about contributors to resilience among these vulnerable groups. Furthermore, the combined data revealed how different methods may engender complementary results and expand our knowledge [28]. The directed content analysis approach to qualitative interviews does present challenges to the naturalistic paradigm. Using a concept such as KIDSCREEN in the processing of data has some inherent limitations in that the data are approached with an informed, but nonetheless, strong bias. Thus it would be more likely to find evidence that is supportive rather than non-supportive [37]. We were aware of this during the process.

The results showed challenges regarding the use of questionnaires, but also some support of the quantitative results. The sample size of the Young & Active is relatively small and may not be representative of a larger population of adolescents with overweight or obesity. Only a sample of 21 participants was interviewed post intervention. If everyone in the intervention group had been interviewed, the qualitative data may have provided different results.

## **Conclusion**

The results of HRQoL surveys may provide important indications of the need for additional exploration and follow-up [8]. The qualitative data showed that perceptions of fitness, energy level or health may vary from day to day, based on both external and internal factors. The results also revealed that the adolescents had different understandings of the concepts and may therefore have answered the questions differently. Combining methods enabled a comprehensive approach to researching health-related quality of life in overweight or obese adolescents. They are vulnerable, specifically in relation to their physical well-being, but also to their self-perception, or at least their body image. Responses regarding self-perception may have provided better results on HRQoL because of the way in which the questions were formulated, integrating clothes and accessories, as well as looks. The qualitative and quantitative data in a mixed methods combination confirm the importance of developing and testing intervention programs that enhance physical well-being and self-perception, using methods that promote resilience and that do not result in adolescents feeling more stress or weight-related stigma. The data underline the importance of supporting the basic psychological needs for autonomy, competence and relatedness for promoting HRQoL. Even though the qualitative findings show that there is a potential risk of misinterpreting the questions, or that other factors may affect the way in which adolescents respond, HRQoL questionnaires such as KIDSCREEN are valuable tools. More research using the benefits of mixed methods approaches may further elucidate our understandings of adolescents as relates to weight and HRQoL.

## **Abbreviations**

SDT: Self-determination theory, HRQoL: Health Related Quality of Life, MI: Motivational Interviewing, PA: Physical activity, BMI: Body Mass Index

## **Declarations**

### **Acknowledgements**

The authors wish to thank study participants for their valuable contribution to this research project. We are also grateful to Kristin Haraldstad for providing basic numerical data on the Norwegian reference group.

### **Funding**

The study was financially supported by the Norwegian Nurses Organisation and the Norwegian Directorate of Health.

### **Availability of data and materials**

The data that support the findings in this study will not be made publicly available, due to participant confidentiality restrictions. Interested investigators may contact the researchers to explore data sharing options.

## Authors' contributions

TS led the design of the study, conducted the qualitative interviews, mixed methods analysis and led the drafting of the manuscript. KR, MS, RO, KL, KG and SH contributed to study design, data analysis and work on the manuscript. All authors read and approved the final manuscript.

## Ethics approval and consent to participate

The study was approved by the Norwegian Regional Committee for Medical and Health Research Ethics (REC no. 2010/2978A, REK sør-øst). The adolescents and their legal parents/guardians gave signed consent for participation in the study as a whole, including the interviews. Information about the study, potential harms and benefits, the purpose of the study and the right to withdraw were emphasized, both in the written material and directly to the adolescents before the interviews started. Anonymity and confidentiality on the audio-recordings and the transcripts were secured by using subject identifier codes instead of names.

## Consent for publication

All participants were notified that their data would be used in reporting finding of this research, and provided assent to use quotes and aggregated data. Consent was provided by the adolescents' parents.

## Competing interests

The authors declare that they have no competing interests.

## References

1. Buttitta M, Iliescu C, Rousseau A, Guerrien A: **Quality of life in overweight and obese children and adolescents: a literature review.** *An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation - Official Journal of the International Society of Quality of Life Research* 2014, **23**:1117-1139.
2. Tsiros MD, Olds T, Buckley JD, Grimshaw P, Brennan L, Walkley J, Hills AP, Howe PRC, Coates AM: **Health-related quality of life in obese children and adolescents.** *Int J Obes* 2009, **33**:387-400.
3. Bonsergent E, Benie-Bi J, Baumann C, Agrinier N, Tessier S, Thilly N, Briançon S: **Effect of gender on the association between weight status and health-related quality of life in adolescents.** *BMC Public Health* 2012, **12**:997.
4. Michel G, Bisegger C, Fuhr D, Abel T: **Age and gender differences in health-related quality of life of children and adolescents in Europe: a multilevel analysis.** *An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation - Official Journal of the International Society of Quality of Life Research* 2009, **18**:1147-1157.
5. Bisegger C, Cloetta B, von Bisegger U, Abel T, Ravens-Sieberer U: **Health-related quality of life: gender differences in childhood and adolescence.** *Soz Präventivmed* 2005, **50**:281-291.

6. Helseth S, Misvær N: **Adolescents' perceptions of quality of life: what it is and what matters.** *J Clin Nurs* 2010, **19**:1454-1461.
7. Thompson I, Hong JS, Lee JM, Prys NA, Morgan JT, Udo-Inyang I: **A review of the empirical research on weight-based bullying and peer victimisation published between 2006 and 2016.** *Educational Review* 2018:1-23.
8. Helseth S, Haraldstad K, Christophersen K-A: **A cross-sectional study of Health Related Quality of Life and body mass index in a Norwegian school sample (8-18 years): A comparison of child and parent perspectives.** 2015.
9. Brundage M, Bass B, Davidson J, Queenan J, Bezjak A, Ringash J, Wilkinson A, Feldman-Stewart D: **Patterns of reporting health-related quality of life outcomes in randomized clinical trials: implications for clinicians and quality of life researchers.** *An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation - Official Journal of the International Society of Quality of Life Research* 2011, **20**:653-664.
10. Ottova V, Erhart M, Rajmil L, Dettenborn-Betz L, Ravens-Sieberer U: **Overweight and its impact on the health-related quality of life in children and adolescents: results from the European KIDSCREEN survey.** *Qual Life Res* 2012, **21**:59.
11. Ravens-Sieberer U, Kidscreen Group E: *The Kidscreen questionnaires : quality of life questionnaires for children and adolescents : handbook.* 3rd edition. edn. Lengerich: Pabst Science Publishers; 2016.
12. Ottova V, Erhart M, Rajmil L, Dettenborn-Betz L, Ravens-Sieberer U: **Overweight and its impact on the health-related quality of life in children and adolescents: results from the European KIDSCREEN survey.** *An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation - Official Journal of the International Society of Quality of Life Research* 2012, **21**:59-69.
13. Pizzi MA, Vroman K: **Childhood Obesity: Effects on Children's Participation, Mental Health, and Psychosocial Development.** *Occupational Therapy In Health Care, 2013, Vol27(2), p99-112* 2013, **27**:99-112.
14. Griffiths LJ, Parsons TJ, Hill AJ: **Self-esteem and quality of life in obese children and adolescents: A systematic review.** vol. 5. pp. 282-304. Oxford, UK; 2010:282-304.
15. Strauss RS, Pollack HA: **Social Marginalization of Overweight Children.** *Arch Pediatr Adolesc Med* 2003, **157**:746-752.
16. Swallen KC, Reither EN, Haas SA, Meier AM, Swallen KC: **Overweight, obesity, and health-related quality of life among adolescents: the National Longitudinal Study of Adolescent Health.** *Pediatrics* 2005, **115**:340-347.
17. Fedewa AL, Ahn S: **The Effects of Physical Activity and Physical Fitness on Children's Achievement and Cognitive Outcomes: A Meta-Analysis.** *Res Q Exerc Sport* 2011, **82**:521-535.
18. Ul-Haq Z, Mackay DF, Fenwick E, Pell JP: **Meta-Analysis of the Association between Body Mass Index and Health-Related Quality of Life among Children and Adolescents, Assessed Using the Pediatric Quality of Life Inventory Index.** *The Journal of Pediatrics* 2013, **162**:280-286.e281.

19. Waters E, de Silva-Sanigorski A, Hall BJ, Brown T, Campbell KJ, Gao Y, Armstrong R, Prosser L, Summerbell CD: **Interventions for preventing obesity in children.** *The Cochrane database of systematic reviews* 2011:CD001871.
20. Gopinath B, Hardy LL, Baur LA, Burlutsky G, Mitchell P: **Physical Activity and Sedentary Behaviors and Health-Related Quality of Life in Adolescents.** *Pediatrics* 2012.
21. Herman MK, Hopman MW, Vandenberg GE, Rosenberg WM: **Physical Activity, Body Mass Index, and Health-Related Quality of Life in Canadian Adults.** *Med Sci Sports Exerc* 2012, **44**:625-636.
22. Gu X, Chang M, Solmon MA: **Physical Activity, Physical Fitness, and Health-Related Quality of Life in School-Aged Children.** *Journal of Teaching in Physical Education* 2016, **35**:117-126.
23. Buttitta M, Rousseau A, Guerrien A, Buttitta M: **A New Understanding of Quality of Life in Children and Adolescents with Obesity: Contribution of the Self-determination Theory.** *Current obesity reports* 2017, **6**:432-437.
24. Ryan RM, Deci EL: **Self-determination theory : basic psychological needs in motivation, development, and wellness.** New York, New York ;London, England: The Guilford Press; 2017.
25. Patrick H, Williams GC: **Self-determination theory: its application to health behavior and complementarity with motivational interviewing.** *International Journal of Behavioral Nutrition and Physical Activity* 2012, **9**:18.
26. Riiser K, Londal K, Ommundsen Y, Sundar T, Helseth SI: **Development and Usability Testing of an Internet Intervention to Increase Physical Activity in Overweight Adolescents.** *JMIR Research Protocols* 2013, **2**.
27. Creswell JW, Plano Clark VL: *Designing and conducting mixed methods research.* 3rd ed. edn. Los Angeles: Sage; 2018.
28. Fetters MD, Curry LA, Creswell JW: **Achieving Integration in Mixed Methods Designs—Principles and Practices.** *Health Serv Res* 2013, **48**:2134-2156.
29. Guetterman TC, Fetters MD, Creswell JW: **Integrating Quantitative and Qualitative Results in Health Science Mixed Methods Research Through Joint Displays.** *Ann Fam Med* 2015, **13**:554-561.
30. The Norwegian Directorate of Health: **Norwegian clinical guideline for for growth measurement of children and adolescents in primary care.** Oslo, Norway: The Norwegian Directorate of Health; 2010.
31. Malterud K, Siersma VD, Guassora AD: **Sample Size in Qualitative Interview Studies.** *Qual Health Res* 2016, **26**:1753-1760.
32. Haraldstad K, Christophersen K-A, Eide H, Nativg GK, Helseth S: **Health related quality of life in children and adolescents: Reliability and validity of the Norwegian version of KIDSCREEN-52 questionnaire, a cross sectional study.** *Int J Nurs Stud* 2011, **48**:573-581.
33. Ravens-Sieberer U, Erhart M, Wille N, Wetzel R, Nickel J, Bullinger M: **Generic Health-Related Quality-of-Life Assessment in Children and Adolescents.** *Pharmacoeconomics* 2006, **24**:1199-1220.
34. Ravens-Sieberer U, Herdman M, Devine J, Otto C, Bullinger M, Rose M, Klasen F: **The European KIDSCREEN approach to measure quality of life and well-being in children: development, current**

- application, and future advances.** *Quality of Life Research* 2014, **23**:791-803.
35. Ravens-Sieberer U, Gosch A, Rajmil L, Erhart M, Bruil J, Duer W, Auquier P, Power M, Abel T, Czemy L, et al: **KIDSCREEN-52 quality-of-life measure for children and adolescents.** *Expert Rev Pharmacoecon Outcomes Res* 2005, **5**:353-364.
36. Singh, Xie KM: **Bootstrap Method.** pp. 46-51; 2010:46-51.
37. Hsieh H-F, Shannon SE: **Three Approaches to Qualitative Content Analysis.** *Qual Health Res* 2005, **15**:1277-1288.
38. Moseholm E, Feters MD: **Conceptual models to guide integration during analysis in convergent mixed methods studies.** *Methodological Innovations* 2017, **10**.
39. Gandhi PK, Revicki DA, Huang IC: **Adolescent body weight and health-related quality of life rated by adolescents and parents: the issue of measurement bias.** *BMC Public Health* 2015, **15**:1192.
40. van Sluijs EMF, Kriemler S: **Reflections on physical activity intervention research in young people – dos, don'ts, and critical thoughts.** *International Journal of Behavioral Nutrition and Physical Activity* 2016, **13**:25.
41. Tsiros MD, Olds T, Buckley JD, Grimshaw P, Brennan L, Walkley J, Hills AP, Howe PRC, Coates AM: **Health-related quality of life in obese children and adolescents.** *Int J Obes (Lond)* 2009, **33**:387.
42. Wille N, Bullinger M, Holl R, Hoffmeister U, Mann R, Goldapp C, Reinehr T, Westenhöfer J, Egmond-Froehlich A, Ravens-Sieberer U: **Health-related quality of life in overweight and obese youths: Results of a multicenter study.** *Health and Quality of Life Outcomes* 2010, **8**:36.
43. Riiser K, Løndal K, Ommundsen Y, Småstuen MC, Misvær N, Helseth S: **The Outcomes of a 12-Week Internet Intervention Aimed at Improving Fitness and Health-Related Quality of Life in Overweight Adolescents: The Young & Active Controlled Trial.** 2015.
44. Laursen B, Coy KC, Collins WA, Laursen B: **Reconsidering changes in parent-child conflict across adolescence: a meta-analysis.** pp. 817-832; 1998:817-832.
45. Smetana JG, Campione-Barr N: **Adolescent development in interpersonal and societal contexts.** *Annu Rev Psychol* 2006, **57**:255.
46. Meade T, Dowswell E, Meade T: **Adolescents' health-related quality of life (HRQoL) changes over time: a three year longitudinal study.** pp. 14-14; 2016:14-14.
47. Michel G, Bisegger C, Fuhr DC, Abel T, The Kg: **Age and gender differences in health-related quality of life of children and adolescents in Europe: a multilevel analysis.** *Qual Life Res* 2009, **18**:1147.
48. de Beer M, Hofsteenge GH, Koot HM, Hirasing R, Delemarre-van de Waal H, Gemke R: **Health-related-quality-of-life in obese adolescents is decreased and inversely related to BMI.** pp. 710-714; 2007:710-714.
49. Kolotkin RL, Zeller M, Modi AC, Samsa GP, Quinlan NP, Yanovski JA, Bell SK, Maahs DM, de Serna DG, Roehrig HR, Kolotkin RL: **Assessing weight-related quality of life in adolescents.** pp. 448-457; 2006:448-457.

50. Bartholomew KJ, Ntoumanis N, Ryan RM, Bosch JA, Thøgersen-Ntoumani C: **Self-Determination Theory and Diminished Functioning: The Role of Interpersonal Control and Psychological Need Thwarting.** *Personality and Social Psychology Bulletin* 2011, **37**:1459-1473.
51. Haines J, Rifas-Shiman SL, Horton NJ, Kleinman K, Bauer KW, Davison KK, Walton K, Austin SB, Field AE, Gillman MW: **Family functioning and quality of parent-adolescent relationship: cross-sectional associations with adolescent weight-related behaviors and weight status.** *International Journal of Behavioral Nutrition and Physical Activity* 2016, **13**:68.
52. Fenner A, Howie E, Straker L, Hagger M: **Exploration of the mechanisms of change in constructs from self-determination theory and quality of life during a multidisciplinary family-based intervention for overweight adolescents.** 2016.
53. Bisegger C, Cloetta B, Von Rueden U, Abel T, Ravens-Sieberer U: *Health-related quality of life: gender differences in childhood and adolescence.* 2005.
54. Voelker DK, Reel JJ, Greenleaf C: **Weight status and body image perceptions in adolescents: current perspectives.** *Adolescent health, medicine and therapeutics* 2015, **6**:149.
55. Pont SJ, Puhl R, Cook SR, Slusser W: **Stigma Experienced by Children and Adolescents With Obesity.** *Pediatrics* 2017, **140**:e20173034.
56. Roberts RE, Duong HT: **Perceived weight, not obesity, increases risk for major depression among adolescents.** *J Psychiatr Res* 2013, **47**:1110-1117.
57. Puhl RM, King KM: **Weight discrimination and bullying.** *Best Practice & Research Clinical Endocrinology & Metabolism* 2013, **27**:117-127.
58. Hand WB, Robinson JC, Stewart MW, Zhang L, Hand SC: **The Identity Threat of Weight Stigma in Adolescents.** *West J Nurs Res* 2017, **39**:991-1007.
59. Lillis J, Thomas JG, Olson K, Wing RR: **Weight self-stigma and weight loss during behavioural weight loss intervention: Self-stigma and weight loss.** 2019.
60. Puhl R, Suh Y: **Health Consequences of Weight Stigma: Implications for Obesity Prevention and Treatment.** *Current Obesity Reports* 2015, **4**:182-190.