

# Polystyrene microplastics do not affect juvenile brown trout (*Salmo trutta f. fario*) or modulate effect of the pesticide methiocarb

Hannah Schmieg (✉ [hannah.schmieg@uni-tuebingen.de](mailto:hannah.schmieg@uni-tuebingen.de))

Eberhard Karls Universität Tübingen <https://orcid.org/0000-0002-3934-9600>

**Sven Huppertsberg**

Hochschule Fresenius gGmbH

**Thomas P. Knepper**

Hochschule Fresenius gGmbH

**Stefanie Kraus**

Eberhard Karls Universität Tübingen

**Katharina Reitter**

Eberhard Karls Universität Tübingen

**Felicitas Rezbach**

Eberhard Karls Universität Tübingen

**Aki S. Ruhl**

Umweltbundesamt

**Heinz R. Köhler**

Eberhard Karls Universität Tübingen

**Rita Triebkorn**

Eberhard Karls Universität Tübingen

---

## Research

**Keywords:** microplastics, pesticide, methiocarb, brown trout, oxidative stress, acetylcholinesterase, histopathology

**Posted Date:** March 9th, 2020

**DOI:** <https://doi.org/10.21203/rs.2.19551/v2>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Environmental Sciences Europe on March 23rd, 2020. See the published version at <https://doi.org/10.1186/s12302-020-00327-4>.

# Abstract

Background There has been a rising interest within the scientific community and the public about the environmental risk related to the abundance of microplastics in aquatic environments. Up to now, however, scientific knowledge in this context has been scarce and insufficient for a reliable risk assessment. To remedy this scarcity of data, we investigated possible adverse effects of polystyrene particles ( $10^4$  particles/L) and the pesticide methiocarb (1 mg/L) in juvenile brown trout (*Salmo trutta* f. *fario*) both by themselves as well as in combination after a 96 h laboratory exposure. PS beads (density 1.05 g/mL) were cryogenically milled and fractionated resulting in irregular shaped particles (<50  $\mu$ m). Besides body weight of the animals, biomarkers for proteotoxicity (stress protein family Hsp70), oxidative stress (superoxide dismutase, lipid peroxidation), and neurotoxicity (acetylcholinesterase, carboxylesterases) were analysed. As an indicator of overall health, histopathological effects were studied in liver and gills of exposed fish. Results Polystyrene particles by themselves did not influence any of the investigated biomarkers. In contrast, the exposure to methiocarb led to a significant reduction of the activity of acetylcholinesterase and the two carboxylesterases. Moreover, the tissue integrity of liver and gills was impaired by the pesticide. Body weight, the oxidative stress and the stress protein levels were not influenced by methiocarb. Effects caused by co-exposure of polystyrene microplastics and methiocarb were the same as those caused by methiocarb alone. Conclusions Overall, methiocarb led to negative effects in juvenile brown trout. In contrast, polystyrene microplastics in the tested concentration did not affect the health of juvenile brown trout and did not modulate the toxicity of methiocarb in this fish species.

## 1. Background

Public and scientific awareness for the problem of environmental pollution with microplastics (MP) has increased considerably over the last years. MP can be found ubiquitously in marine and freshwater ecosystems and even in remote regions (Dris *et al.* 2015, Bergmann *et al.* 2017, Horton *et al.* 2017). While most research focuses on the abundance and possible effects of MP in marine environments, the number of studies in freshwater ecosystems is also rapidly increasing (Horton *et al.* 2017). Nevertheless, many gaps in knowledge about abundance, toxicity, and hazards of MP in freshwater systems still exist.

Possible effects of MP can be related to mechanical injuries caused by the particles (e.g. von Moos *et al.* 2012, Karami *et al.* 2016, Lu *et al.* 2016, Paul-Pont *et al.* 2016, Ding *et al.* 2018 ). Of particular interest are very small particles (mainly nanoplastics) for which uptake in cells were shown (e.g. Mattsson *et al.* 2017, van Pomeroy *et al.* 2017, Ding *et al.* 2018). In addition to this physical damage, MP can also potentially affect organisms due to the leakage of hazardous substances like residual monomers, polymerization solvents, or additives (Lithner *et al.* 2009, Schiavo *et al.* 2018). A third process that should be considered in this context is that MP might adsorb hydrophobic organic pollutants like PAHs, PCBs, or pesticides (reviewed by Wang *et al.* 2018) and transport them into organisms. Sorption of hydrophobic organic pollutants to MP is especially important for freshwater, since the concentrations of these pollutants are generally higher than in marine ecosystems (Dris *et al.* 2015). The sorption of organic

pollutants to MP may alter the bioavailability of the pollutants (e.g. Sleight *et al.* 2017, Kleinteich *et al.* 2018, Rehse *et al.* 2018, Oliveira *et al.* 2013, Nematdoost Haghi and Banaee 2017, Guilhermino *et al.* 2018). Furthermore, Batel *et al.* (2016) showed the possibility of the transfer of organic pollutants adsorbed to MP along an artificial food chain. The biological relevance of MP as vectors for chemicals is still under debate. Several studies suggest the effect of MP to be negligible compared to other exposure pathways and the concentration of organic compounds in most MP to be in equilibrium with the surrounding media (e.g. Bakir *et al.* 2016, Koelmans *et al.* 2016, Beckingham and Ghosh 2017). To assess the risk of MP alone and in combination with organic pollutants, more studies under controlled lab conditions are required.

Residues of pesticides are regularly found in freshwater ecosystems (e.g. Smith *et al.* 2012, Bundschuh *et al.* 2014, Köck-Schulmeyer *et al.* 2014). Methiocarb (or synonym mercaptodimethur) is a carbamate pesticide, which is used as a molluscicide, herbicide, insecticide, acaricide, and as repellent in seed treatment (Worthing *et al.* 1991, European Commission 2014, Loos *et al.* 2018). In the U.S., products containing methiocarb are registered by the EPA for restricted usage (EPA 1994). In September 2019, the European Commission did not renew the approval of methiocarb as an active substance (European Commission 2019b). Prior to this decision, products containing methiocarb were generally approved for usage in the EU, although the application as a molluscicide was already prohibited since 2014 (European Commission 2014, European Commission 2019a). Methiocarb is on the 1<sup>st</sup> surface water Watch List under the Water Framework Directive of the European Union and is recommended for the 2<sup>nd</sup> Watch List (European Commission 2015, Loos *et al.* 2018). The average surface water concentration of methiocarb in the EU-wide monitoring campaign was between 6 - 40 ng/L, and the maximal concentration 109 ng/L.

The purpose of the present study was to investigate whether polystyrene (PS) MP alone and in co-exposure with methiocarb affect juvenile brown trout (*Salmo trutta f. fario*). To determine the amount of methiocarb sorbed on PS particles a separate sorption study would be necessary. Sorption of chemical contaminants to MP involves various mechanisms and depends on physicochemical properties of the sorbate, the sorbent, as well as the medium characteristics (Tourinho *et al.* 2019). The animals were exposed to either PS-MP alone (10<sup>4</sup> particles/L), to methiocarb alone (1 mg/L), or co-exposed to both (10<sup>4</sup> particles/L and 1 mg/L methiocarb) for 96 h. The chosen methiocarb concentration was relatively high to ensure that the effects of methiocarb and hence also a possible modulation could be observed in the experiment. Besides apical endpoints (mortality and body weight), biomarkers for oxidative stress (activity of superoxide dismutase (SOD), and formation of lipid peroxidation (LPO)), proteotoxicity (stress protein family Hsp70) as well as neurotoxicity (acetylcholinesterase (AChE) activity and two protective carboxylesterases (CbE)) were examined. Furthermore, the histopathological status of gills and liver was analysed. Our purpose was to investigate possible mechanistic effects in juvenile brown trout of PS-MP or methiocarb alone at sub-lethal level and whether PS-MP modulate the toxicity of methiocarb at the tested concentrations.

## 2. Materials And Methods

### 2.1. Test organism

In the present study, approximately 11-month-old juvenile brown trout were used. Prior to the experiment, trout were acclimated in a 250 L tank to lab conditions (aerated filtered tap water – iron filter, particle filter, activated charcoal filter) for almost three months. Trout originated from a commercial fish breeder (Forellenzucht Lohmühle, D-72275 Alpirsbach-Ehlenbogen, Germany). In regular controls, the breeding establishment is categorized as category I, disease-free (EU 2006).

### 2.2. Test substances

PS-MP suspensions with a defined number of particles were produced according to the method of Eitzen *et al.* (2019). The polystyrene particles were analysed by different techniques (Raman and FTIR spectroscopy and TED-GC-MS) in scope of the joint project MiWa (for FTIR spectrum see supplement Figure S1). Briefly, transparent polystyrene pellets (Polystyrol 158 K, BASF, Germany) with a density of 1.05 g/mL were cryogenically milled (CryoMill, Retsch, Germany) resulting in irregularly shaped PS-MP particles (Figure 1). The number of PS particles in the pure suspension was determined with a particle counter (SVSS, PAMAS, Germany) by light extinction in a laser-diode sensor (type HCB-LD-50/50). The particle size distribution is provided in Table 1, a figure in the supplement (Figure S2). Subsequently, the particles were fractionated by a micro-sieve (polyamide monofilament) to < 50 µm, and the permeate was used as stock suspension. Particles below 1 µm (e.g. ca. 500 nm) were only rarely found in SEM in samples after cryogenic milling much more excessive than applied for the particles used in this study (compare Eitzen *et al.* 2019).

*Table 1: Size ranges (in µm) and counted particle numbers (per mL) of polystyrene particles in the stock suspension*

size range [ $\mu\text{m}$ ]	Particle number [1/mL ]	size range [ $\mu\text{m}$ ]	particle number [1/mL ]
2-3	9067	30-35	643
3-4	6488	35-40	473
4-5	8872	40-50	511
5-6	4832	50-60	220
6-7	3770	60-70	89
7-8	3484	70-80	43
8-9	3076	80-90	18
9-10	2412	90-100	6
10-12	3259	100-150	5
12-14	2330	150-200	0
14-16	1628	200-250	0
16-18	1195	250-300	0
18-20	1081	300-350	0
20-25	1698	350-400	0
25-30	1041		

Methiocarb was purchased from Sigma-Aldrich (product line: PESTANAL®; CAS number: 2032-65-7; purity 99.8 %; more details in the supplement). The octanol/water partition coefficient of methiocarb is  $\log P_{ow}$  3.08 (ILO 2012). The predicted solubility in water (20 °C) is between 14.23 mg/L and 119.63 mg/L (EPA Chemistry Dashboard). Nevertheless, it was not possible to dissolve 15 mg/L methiocarb without a solvent. Therefore, 0.01% dimethyl sulfoxide (DMSO) was used to dissolve methiocarb in water.

### 2.3. Exposure of juvenile brown trout

Fish were exposed in a static three-block design for 96 h (18.10. - 23.10.2017). Each of the three blocks consisted of five tanks with the different treatment groups: a negative control group (pure water), a solvent control (0.01 % DMSO), PS-MP ( $10^4$  particles/L), methiocarb (1 mg/L), and a co-exposure of  $10^4$  particles/L and 1 mg/L methiocarb (mixture). The test concentration of 1 mg/L methiocarb was selected due to published  $LC_{50}$  values for rainbow trout (*Onchorynchus mykiss*): the reported  $LC_{50}$  (96 h) values vary between  $LC_{50} = 0.198$  mg/L (Johnson and Finley 1980) and  $LC_{50} = 4.82$  mg/L (Pesticide Action Network 2000-2019). However, Altinok *et al.* (2006) found all fish in the experiment exposed to concentrations equal or less 3.75 mg/L survived over two months. The relatively high concentration of 1 mg/L was selected in order to have a high chance for methiocarb to cause effects in brown trout and to be able to observe potential modulation of these effects. Each aquarium contained 10 fish (in 15 L of the corresponding test medium) resulting in 30 fish per treatment (aquaria in triplicates) and 150 fish in total. To be able to consider potential confounding factors, the position of each treatment group within the

blocks was randomized. Exposure took place in a thermo-constant chamber set to 7 °C with a light/dark cycle of 10 h/14 h (tanks were shaded from direct light). Aeration of the tanks was ensured by glass pipettes, which were connected to compressed air via silicone tubes.

In all treatment groups, filtered tap water was used. For the PS-MP treatments, the defined volumes of the stock suspensions (56 240 particles/mL) were added. To avoid a loss of particles in the glass vessels containing the PS-MP suspensions, the vessels were rinsed four times. For each tank containing methiocarb, a stock solution (15 mg/L methiocarb) was prepared as follows: 15 mg methiocarb were dissolved in 1.5 mL DMSO (resulting in a concentration of 0.01 % DMSO in the tanks), and subsequently, 1 L water was added. Finally, the solutions were stirred for two days to ensure a complete dissolution of methiocarb. To provide comparability between the groups, 0.01 % DMSO was also added to the tanks which contained only PS-MP.

Fish were fed daily a defined portion (4 % of body weight) of commercially available fish food (0.8 mm, Biomar, Brande, Denmark). After 48 h and 72 h, 2.5 L of the test medium of each aquarium were removed to get rid of feces and remains of food. Water parameters were measured at the beginning and the end of the experiment (average values: temperature=7.0±0.3 °C, pH=7.3±0.5, oxygen concentration=10.9±0.4 mg/L, oxygen saturation=94.7±2.4 %; conductivity=489.8±15.8 µS/cm; for detailed information see supplement Table S1 and Table S2). Nitrite (NO<sub>2</sub><sup>-</sup>) values did not exceed 0.2 mg/L.

After 24 h, four animals (one fish of block 1 methiocarb, two fish of block 3 methiocarb and one fish of block 2 mixture) were in very poor health conditions and had to be euthanized. Apart from these, no fish died during the experiment.

At the end of the experiment, fish were anesthetized and killed by an overdose of tricaine methanesulfonate (1 g/L MS222, buffered with NaHCO<sub>3</sub>). Subsequently, death of the animals was ensured by severance of the spine. Body weight and length of each animal was recorded prior to dissection (weighted 2.96 ± 1.01 g/6.47 ± 0.73 cm). For histological investigations samples of gills and liver were transferred into fixative (2 % glutardialdehyde diluted in 0.1 M cacodylate buffer; pH 7.6) and stored at 4 °C until further processing. For biochemical analyses, samples of muscle (AChE and CbE activity), brain (LPO analysis), gills (stress protein analysis), and a part containing muscle and kidney (for analysis of SOD activity) were immediately frozen in liquid nitrogen. Subsequently, samples for biochemical analyses were stored at -80 °C.

Additionally, 30 fish (kept in a 250 L tank during the experiment) were sampled in an analogous manner as lab control to be able to recognize possible effects of the exposure procedure (e.g. limited water volume) itself.

## 2.4. Chemical analysis

From each aquarium, 12 mL medium were taken at the beginning and the end of the experiment and frozen at -20 °C until further processing. Samples of the three replicate tanks were analysed separately by

means of HPLC-ESI-MS/MS. Quantification was performed via liquid chromatography coupled to mass spectrometry (LC-MS/MS). As instrumentation, a PerkinElmer Series 200 LC system (PerkinElmer, Waltham, USA) consisting of two Series 200 micro-pumps, a Series 200 vacuum degasser, and a Series 200 autosampler was coupled to a QqLIT mass spectrometer SCIEX QTRAP 3200 (AB SCIEX, Darmstadt, Germany), and ionization was achieved via electrospray ionization (ESI) in positive mode (further information is provided in the supplement).

Separation was performed at 23 °C using a XSelect™ HSS T3 reverse phase column (2.1 mm x 50 mm, 3.5 µm particle size) from Waters (Milford, USA). Injection volume was 10 µL and a gradient was run with a flow rate of 200 µL/min. Eluents used for chromatographic separation were mixtures of methanol and water (Eluent A: MeOH/H<sub>2</sub>O, 5:95, v/v; Eluent B: MeOH/H<sub>2</sub>O, 95:5, v/v) additionally containing 5 mM/L ammonium formate (NH<sub>4</sub>HCO<sub>2</sub>) and adjusted to pH 3 with formic acid (further information are provided in the supplement).

A multiple reaction monitoring (MRM) method was used for quantification. The individual transitions and optimized MS parameters are stated in Table S5 in the supplement.

Unless specified otherwise, all calculations were carried out via Microsoft Excel 2016. The developed method was validated regarding the limit of detection (LoD), the limit of quantification (LoQ), and the method precision and accuracy. LoD and LoQ were determined by application of the signal to noise (S/N) approach. The S/N ratios of two calibration standards ( $\beta$  = 0.25 ng/mL and 0.5 ng/mL) were measured in triplicate, and their S/N ratio was determined. LoD was defined as S/N > 3, and LoQ with S/N > 9. With the determined S/N ratios, the theoretical LoD and LoQ were calculated as LoD=0.04 ng/ml and LoQ=0.13 ng/mL (more details are provided in Table S6 in the supplement). The lowest standard of the calibration (0.25 ng/mL) corresponds to the applied LoQ.

Concentrations of standards used for calibration ranged from 0.25-250 ng/mL, and atrazine was used as internal standard. Measurement precision was determined by multiple (n=5) of three standards and the calculation of their concentration on basis of the obtained calibration. Mean concentrations, relative standard deviation (RSD), and accuracy are stated in Table S7 in the supplement. For quantification, calibration was measured in triplicate (before, in between, and after the samples). Samples were filtered via syringe filters (Carl Roth, SPARTAN® regenerated cellulose, 0.2 µm) after dilution (1:25; 1:100; V.V) with type 1 water and prior to their measurement.

## **2.5. Determination of oxidative stress level**

### **2.5.1. Activity of SOD**

The SOD activity (Cu/Zn SOD, Mn SOD and Fe SOD) in muscle/kidney samples was analysed with the Cayman Chemical superoxide dismutase assay kit (item no. 706002, Cayman Chemical Company, Ann Arbor, USA) in 96-well plates. To remove any blood residues, samples were rinsed in phosphate buffered saline (PBS; pH 7.4) before they were frozen in liquid nitrogen. The dilution of the final samples was

1:150. In the assay, superoxide radicals are generated with xanthine oxidase and hypoxanthine. Subsequently, the superoxide radicals are detected with tetrazolium salt. The absorbance at 450 nm was measured (Bio-Tek Instruments, Winooski VT, USA) after an incubation of 30 min at room temperature. All samples were analysed in duplicates.

### 2.5.2. Level of lipid peroxidases

The level of lipid peroxides was quantified with the ferrous oxidation xylenol orange (FOX) assay. The assay was performed in a slightly modified way for 96-well plates according to Hermes-Lima *et al.* (1995) and Monserrat *et al.* (2003). For homogenization, the brains were 1:2 diluted with HPLC grade methanol and subsequently centrifuged (15 000 rcf, 5 min, 4 °C). Supernatant was stored until the final assay at -80 °C. The assay was performed in 96-well plates. Into each well, 50 µL of 0.75 mM FeSO<sub>4</sub>-solution, 50 µL of 75 mM sulphuric acid, 50 µL of 0.3 mM xylenol orange solution, 40 µL supernatant, and 10 µL bidistilled water were added successively. Triplicates of each sample were tested. Furthermore, a sample blank without FeSO<sub>4</sub> (replaced by bidistilled water) of each sample was measured to correct for potential Fe in the samples. After an incubation of 120 min at room temperature, the absorbance at 570 nm (ABS570) was measured (Bio-Tek Instruments, Winooski VT, USA). Afterwards, in each well 1 µL of 1 mM cumene hydroperoxide solution (CHP) was pipetted. The plates were incubated for another 30 min at room temperature, and a second measurement at 570 nm was conducted. The data of both measurements were related to a master blank (200 µL bidistilled water). Cumene hydroperoxide equivalents (CHPequiv./mg wet weight) were calculated using the following equation: (see Equations in the Supplemental Files)

### 2.6. Stress protein analysis

Hsp70 quantification was performed as described by Dieterich *et al.* (2015). Samples were 1:3 diluted with extraction buffer (80 mM potassium acetate, 5 mM magnesium acetate, 20 mM HEPES, and 2 % protease inhibitor at pH 7.5) and homogenized on ice. The samples were centrifuged (20 000 rcf, 10 min, 4 °C) and the supernatant was divided into two samples. The first was used to quantify the total protein content via Bradford assay (Bradford 1976). Subsequently, the second sample (amount standardized to 40 µg of total protein content) was separated via SDS-PAGE (12 % acrylamide, 0.12 % bisacrylamide, 30 min at 80 V plus 90 min at 120 V). The proteins were blotted on a nitrocellulose membrane and immuno-stained with a monoclonal α-Hsp70 IgG (Dianova, Hamburg, Germany) followed by a secondary peroxidase-coupled α-IgG (Jackson Immunoresearch, West Grove, PA). The optical volume (area x average pixel intensity) of the protein bands was quantified (Image Studio Lite, 4.0.2.1, Li-Cor Biosciences, Lincoln, USA) and put into relation to an internal Hsp70 standard.

### 2.7. Analysis of neurotoxicity

For the analysis of the activity of AChE and two CbE, muscle tissue was diluted 1:5 in tris-LS buffer (20 mM Tris<sub>base</sub>, 20 mM NaCl, inhibitor mix, pH 7.3) and homogenized. After centrifugation (5000 rcf, 10 min, 4 °C), 50 % glycerol was added to the supernatant (1/4 of the amount of the supernatant), and

the mixture was frozen at -20 °C. The protein content in the samples was determined with the Lowry method modified by Markwell *et al.* (1978). The AChE-activity was measured spectrophotometrically at 405 nm (Bio-Tek Instruments, Winooski VT, USA) according to Ellman *et al.* (1961) and modified by Rault *et al.* (2008). CbE activity was determined with the substrates 5 mM 4-nitrophenol acetate (pnpa) and 5 mM 4-nitrophenyl valerate (pnpv) described by Sanchez-Hernandez *et al.* (2009). In all assays, samples were analysed in triplicates. The data were related to the total protein amount (specific activity per milligram total protein content). One unit is described as one micromole substrate hydrolysed per minute.

## 2.8. Histopathology

Liver and gill samples were fixed in 2 % glutardialdehyde diluted in 0.1 M cacodylate buffer (pH 7.6) and stored at 4 °C for at least 4 weeks. Prior to further processing, gill samples were decalcified in a 1:2 mixture of 100 % formic acid and 70 % ethanol. All samples were rinsed three times for 10 min in 0.1 M cacodylate buffer and subsequently three times for 15 min in 70 % ethanol. Dehydration of the samples and embedding into paraffin was achieved in an automated tissue infiltrator (TP 1020, Leica Wetzlar, Germany). With a sledge microtome (SM 2000 R, Leica Wetzlar, Germany), 3 µm thick histological sections were cut. Sections were stained (1) with hematoxylin-eosin (HE) which visualizes nuclei, cytoplasm muscles, and connective tissue and (2) alcian blue - PAS by which glycogen and mucus were stained. In a first step, the slides were evaluated qualitatively to identify pathologies. In a second step, the slides were examined semi-quantitatively in an observer blinded way. The samples were categorized into five classes (1: control; 2: slight reactions; 3: medium reactions; 4: strong reactions; 5: destruction) as suggested by Tribskorn *et al.* (2008). Detailed definitions of the five categorizes are provided in the supplement (Table S8).

## 2.9. Statistical analysis

Data analysis was performed with JMP®14.0.0 (SAS Institute Inc., North Carolina, USA). Standard distribution of the data was checked with the Shapiro Wilk test. Homogeneity of variance was checked with the Levene`s test. If necessary, data were transformed (AChE: root, weight: third root, SOD: fifth root, Hsp70: natural logarithm, Cbe-pnpa: 1/x, Cbe-pnpv: seventh root). Using a t-test for all investigated parameters, we ensured that there were no significant differences between the control and the solvent control. Subsequently, the other treatments were compared to the solvent control. In addition to the comparison to the solvent control, the PS-MP and the methiocarb treatments were compared to the mixture treatment. Comparisons with the lab control were conducted only qualitatively and were not included in the mathematical analyses.

The base  $\alpha$ -level was set to 0.05. Parametric data were analysed with a nested ANOVA including  $\square$ block $\square$  as nesting factor. This allows considering potential confounding factors and avoiding pseudo-replication. The comparison between the single groups was made with Tukey HSD. The data of the activity of CbE with the substrate pnpa could not be transformed to reach homoscedasticity. Thus, a Welch ANOVA was performed to analyse the effect on the activity of CbE-pnpa mathematically. Frequency data were

analysed with likelihood-ratio analysis and a post hoc Bonferroni-Holm correction. Detailed values of the tests are given in the supplement (Table S9).

## 2.10. Credibility of data

The information on the fulfilment of the criteria for reporting and evaluation of ecotoxicity data (CRED) proposed by Moermond *et al.* (2016) is provided in the supplementary information.

# 3. Results

## 3.1. Chemical analyses

At the beginning and the end of the experiment, in both control groups, the methiocarb concentrations in the water phase were below the applied quantification limit (LoQ) of 0.25 µg/L (Table 2). In the PS-MP exposure, no methiocarb in the water phase could be detected (<LoQ). In both exposure groups with methiocarb (methiocarb and mixture), the measured concentrations in the water phase at the beginning of the experiment were 50 % of the nominal concentrations. After 96 h, in the treatment group with solely methiocarb, the concentration was further reduced by approximately 50 %, and in the mixture treatment by 34 %.

*Table 2: Nominal and measured concentrations of methiocarb in the water phase of the different treatment groups. Displayed are the arithmetic means ± standard deviation of the three aquaria. The applied limit of quantification was 0.25 µg/L methiocarb.*

treatment group	nominal concentration	measured concentration at start of experiment	measured concentration at end of experiment
control	0 µg/L	< 0.25 µg/L	< 0.25 µg/L
solvent control	0 µg/L	< 0.25 µg/L	< 0.25 µg/L
PS-MP (10 <sup>4</sup> particles/L)	0 µg/L	< 0.25 µg/L	< 0.25 µg/L
methiocarb	1000 µg/L	463 µg/L ±32 µg/L	237 µg/L ±59 µg/L
mixture PS-MP + methiocarb	1000 µg/L	469 µg/L ±133 µg/L	308 µg/L ±76 µg/L

## 3.2. Mortality and weight

After 48 h, three of the fish exposed to methiocarb and one of the fish exposed to the mixture had to be euthanized due to their poor health conditions. Apart from that, no mortality occurred during the experiment. The total mortality was 2.67 %. Weight (overall mean: 2.96±1.01 g) did not differ between the treatment groups (Table 3; p=0.3384).

### 3.3. Oxidative Stress

To assess the oxidative stress level, SOD activity and the degree of LPO were investigated. No difference ( $p=0.5136$ ) between SOD activity in the different treatment groups was found (Table 3). The degree of LPO of the exposure groups were comparable to the solvent control ( $p=0.5770$ ). For both tested endpoints, neither PS-MP nor methiocarb nor the mixture caused oxidative stress.

Table 3: Summary of data for the investigated endpoints. All data are given as arithmetic means  $\pm$  standard deviation.

	lab control	control	solvent control	PS-MP	methiocarb	mixture
weight [g]	4.02 $\pm 1.09$	2.49 $\pm 0.66$	2.89 $\pm 0.87$	3.21 $\pm 0.91$	2.95 $\pm 1.27$	3.30 $\pm 1.06$
SOD [U/mL]	92.71 $\pm 26.12$	105.29 $\pm 28.81$	106.50 $\pm 24.22$	106.66 $\pm 29.69$	99.88 $\pm 31.59$	99.43 $\pm 35.37$
FOX [CHP-Equiv.]	2.17 $\pm 1.26$	2.56 $\pm 1.09$	1.97 $\pm 1.17$	1.69 $\pm 1.21$	1.83 $\pm 1.52$	1.60 $\pm 1.39$
Hsp70 [relative grey value]	1.71 $\pm 0.29$	1.84 $\pm 0.29$	1.78 $\pm 0.29$	1.72 $\pm 0.26$	1.77 $\pm 0.30$	1.71 $\pm 0.27$
AChE [mU/mg Protein ]	69.32 $\pm 11.55$	88.38 $\pm 24.98$	77.37 $\pm 18.39$	73.93 $\pm 20.28$	31.48 $\pm 11.38$	32.38 $\pm 11.64$
CbE-pnpa [mU/mg Protein ]	82.20 $\pm 9.16$	79.13 $\pm 10.83$	79.10 $\pm 11.38$	76.68 $\pm 7.90$	45.52 $\pm 5.74$	44.48 $\pm 5.01$
CbE-pnpv [mU/mg Protein ]	60.60 $\pm 11.14$	53.89 $\pm 12.38$	54.01 $\pm 12.94$	49.59 $\pm 10.46$	9.10 $\pm 3.24$	9.05 $\pm 1.91$

### 3.4. Proteotoxicity

The analysis of the Hsp70 level did not reveal any differences between control, solvent control, and exposure groups (Table 3;  $p=0.7922$ ).

### 3.5. Neurotoxicity

No difference in the activity of AChE occurred between solvent control and treatment group with PS-MP ( $p=0.8192$ ). However, the methiocarb and the mixture of methiocarb and PS-MP led to a significantly reduced activity of AChE in comparison to the solvent control and PS-MP alone (Figure 2).

The activity of AChE in the methiocarb-exposed fish was 59 % reduced compared to the solvent control ( $p<0.001$ ). In fish exposed to the mixture of PS-MP and methiocarb, the activity of AChE was reduced by 58 % compared to the solvent control ( $p<0.001$ ; Table 3).

Between fish exposed to methiocarb alone and methiocarb plus PS-MP, no differences were found ( $p=0.9951$ ).

The activity of CbE-pnpa was also inhibited by methiocarb and the mixture treatment (Figure 2; DMSO/PS-MP:  $p=0.9357$ ; DMSO/methiocarb, DMSO/mixture and PS-MP/mixture:  $p<0.001$ ;

methiocarb/mixture:  $p=0.8531$ ). The activity of the CbE-pnpa was not altered by PS-MP alone. Methiocarb alone reduced the activity compared to the solvent control by 42 %, the mixture by 44 %.

Even more pronounced was the inhibition of CbE with the substrate pnpv: compared to the solvent control, the activity was reduced by 83 % both in methiocarb-exposed fish and for fish in the mixture treatment (Figure 2; DMSO/mixture, PS-MP/mixture:  $p<0.0001$ ; methiocarb/mixture:  $p=0.7609$ ). Again, PS-MP alone had no influence on the activity of CbE-pnpv. Beside the clear effect of the different treatments, the treatment nested in the block had a significant effect ( $p=0.0322$ ).

## 3.6. Histopathology

### 3.6.1. Liver

Healthy liver tissue consists of large bright cells containing high amounts of glycogen (Figure 3 A and 3 B). Fish from the lab control, control, solvent control, and exposure with PS-MP showed in some cases inflammations and even focal necrosis. In fish exposed to methiocarb alone or the mixture treatment, a reduction of the glycogen stores became obvious (Figure 3 D) resulting in atrophic cells and enlarged intercellular spaces. Furthermore, in both groups containing methiocarb vacuolization and inflammations occurred cumulatively, and in the majority of samples small zones (single cell or few cells) with caryopycnosis and focal necrosis were found (Figure 3 C). In all investigated samples the proportion of necrotic tissue was  $< 10\%$ . Semi-quantitative analyses showed a very similar condition of fish from the control, solvent control, and exposure with PS-MP, whereas livers of fish exposed to methiocarb or the mixture treatment showed significantly stronger reactions (Figure 3 G; DMSO/PS-MP:  $p=0.0577$ , DMSO/methiocarb, DMSO/mixture, PS-MP/mixture:  $p<0.0001$ , methiocarb/mixture:  $p=0.2396$ ).

### 3.6.2. Gills

Observed pathological alterations in gills of brown trout included hyperplasia and hypertrophy of chloride and pillar cells resulting, in some cases, in lamellar fusion (Figure 3 E and F). Furthermore, an increase of mucus secreting cells, lifting of epithelia, oedema, and even necrosis in small areas occurred. Most of these symptoms were observed in the methiocarb and the mixture treatment groups. Lamellar fusion occurred on small parts of the lamellae of circa 20 % in the lab control, the control, and the solvent control samples as well as 10 % of the MP exposure. In the methiocarb and mixture groups it was considerably more common with 47 % (methiocarb) and 42 % (mixture). Oedemas occurred only in 19 % of the methiocarb group and 14 % of the mixture exposed group. They were located at the inner side of the secondary lamellae.

Semi-quantitative analysis revealed a significantly worse condition of the two treatment groups containing methiocarb compared to the solvent control and fish exposed solely to PS-MP (Figure 3 H; DMSO/PS-MP:  $p=0.4562$ , DMSO/methiocarb, DMSO/mixture, PS-MP/mixture:  $p<0.0001$ , methiocarb/mixture:  $p=0.7266$ ). Gill sections were classified in categories 1-4 of 5 categories in total. No difference was found between the control groups and the group containing solely PS-MP.

## 4. Discussion

In the present study, the effects of PS-MP and the pesticide methiocarb either applied alone or in their binary mixture were investigated in juvenile brown trout.

### 4.1. General considerations concerning the conducted experiment

The measured methiocarb concentrations corresponded to approximately half of the nominal concentration in the methiocarb and mixture groups at the beginning of the experiment and 24 % (methiocarb group) and 31 % (mixture group) at the end. Considering the standard deviations, however, there is no relevant difference in the methiocarb concentration of methiocarb and mixture treatment group at the end of the experiment. Methiocarb has a comparable low persistence in laboratory aerobic soil-water systems (Arena *et al.* 2018). The degradation of methiocarb is pH-dependent and higher under alkaline conditions. The main metabolites are methiocarb phenol and methiocarb sulfoxide phenol. The dissipation half-life ( $DT_{50}$ ) of methiocarb in water at pH 7 is 24 days and 0.21 days at pH 9 (EFSA 2006). Thus, a degradation of methiocarb in this experiment was likely, but occurred to a greater extent as it could have been assumed in an environment with a pH of 7.3. During the experiment, beside the abiotic degradation, a metabolization by the fish may have contributed to the decreased methiocarb concentration after 96 h. In humans, mainly cytochrome p450 and flavin-containing monooxygenases in liver and kidney contribute to the metabolism of methiocarb (Usmani *et al.* 2004, Furnes and Schlenk 2005). For longer exposure to methiocarb, regular exchange of the water with freshly-prepared methiocarb solutions or a flow-through exposure is recommended.

### 4.2. Do PS-MP affect juvenile brown trout?

In juvenile brown trout, none of the investigated parameters were affected by  $10^4$  polystyrene particles/L. In the past, MP were found to cause oxidative stress in fish (Lu *et al.* 2016, Paul-Pont *et al.* 2016, Qiao *et al.* 2019). In contrast to these results, other studies found either no effects (Oliveira *et al.* 2013, Luís *et al.* 2015, Fonte *et al.* 2016) or effects in only some of the investigated endpoints for oxidative stress (Avio *et al.* 2015, Ferreira *et al.* 2016, Ding *et al.* 2018, Magni *et al.* 2018). The oxidative defense system is complex and consists of enzymatic (e.g. SOD, catalase (CAT), glutathione reductase) and non-enzymatic compounds (Lushchak 2016). In addition, the types of investigated polymers, their concentrations, their potential additives as well as the size of the used MP highly vary in the different studies. Jeong *et al.* (2016) analyzed the influence of the particle size on oxidative stress responses and showed that there is a clear connection to this parameter: smaller particles cause more oxidative stress. In general, however, the data is still far too limited and experimental designs of the studies are too variable to decide on whether or not MP cause oxidative stress and, in case of any influence, which pathways are affected.

To the best of our knowledge, the present study is the first that investigated potential proteotoxic effects of MP. In the tested concentration, no effect of PS-MP on the stress protein level (Hsp70) was found in brown trout after 96 h exposure.

Also, the activity of AChE and two investigated CbE were not altered after exposure to PS-MP. In the past several studies reported significant reductions about 20 % of AChE activity in common goby (*Pomatoschistus microps*) after exposure to 184 µg/L polyethylene (PE; 1-5 µm) for 96 h (Oliveira *et al.* 2013, Luís *et al.* 2015, Fonte *et al.* 2016). Under the same test conditions, Ferreira *et al.* (2016) only found a reduction of AChE activity by 13 % in common goby. Furthermore, a decrease of AChE activity in Amazonian discus fish *Symphysodon aequifasciatus* was observed after exposure to 200 µg/L fluorescent PE (70–88 µm) for 30 days (Wen *et al.* 2018). In red tilapia (*Oreochromis niloticus*) an even higher inhibition of AChE by 37.7 % was found after exposure to PS nanoplastics (0.1 µm) in concentrations of circa  $1.8 \times 10^6$ ,  $1.8 \times 10^7$ ,  $1.8 \times 10^8$  particles/L for 14 days (Ding *et al.* 2018). Chen *et al.* (2017) did not find any neurotoxic effect of PS-MP (45 µm, 20 particles/mL) in zebrafish (*Danio rerio*) but reported a significant reduction of AChE activity by 40 % after exposure to a high concentration of PS nanoplastics (50 nm,  $1.5 \times 10^{10}$  particles/mL). In general, the mode of action of how MP might cause neurotoxicity remains unclear. In addition to the type of polymer (including different additives), the particle size also seems to be of importance in this context, since neurotoxic effects were mainly found in studies using very small micro- or even nanoplastics. Furthermore, Ding *et al.* (2018) reported accumulation of nanoplastics in the brain of red tilapia. These findings are supported by a study of Mattsson *et al.* (2017) who demonstrated that polystyrene nanoplastics are capable of penetrating the blood-brain barrier of Crucian carp (*Carassius carassius*). The plastic particles we used in the present study were <50 µm with only a low number of particles <1 µm which might explain the lack of influence on neurotoxicity.

In the present study, no alterations in the histopathological status of the liver and gills were found in fish exposed to PS-MP alone when compared to the solvent control group. Also, low-density PE (125-250 µm) caused no alterations in the liver of zebrafish after 3 weeks of exposure (Rainieri *et al.* 2018). Furthermore, besides effects on the intestine Lei *et al.* (2018) did not find any alterations in gills, liver, and kidney of zebrafish exposed to polyamide (PA), polypropylene (PP), polyvinyl chloride (PVC), and PE (each ~70 µm in a concentration up to 10 mg/L) for 10 days. Moreover, In silver barb fry (*Barbodes gonionotus*) no histopathological reactions were caused by PVC (0.1-1000 µm; 0.2, 0.5 and 1.0 mg/L) fragments besides a slight thickening of the intestinal mucosal epithelium (Romano *et al.* 2018). Additionally, Lei *et al.* (2018), did not find any effect of PS-MP (0.1, 1.0 and 5 µm up to 10 mg/L) on tissue integrity of intestine, gills, kidney, and liver of zebrafish after 10 days of exposure. In contrast, Lu *et al.* (2016) observed early inflammation responses as well as lipid droplets in the liver of zebrafish after exposure to 5 µm polystyrene particles in a concentration of  $2.9 \times 10^4$  particles/L for three weeks. In Japanese medaka (*Oryzias latipes*) severe glycogen depletion and fatty vacuolization in the liver occurred, but no alterations were found in gonads after exposure to PE (<0.5mm) for 2 months (Rochman *et al.* 2013, Rochman *et al.* 2014). Karami *et al.* (2016) exposed African catfish (*Clarias gariepinus*) to low-density PE (<60 µm) for 96 h. They observed hyperplasia and sloughing and even necrosis in gills at a concentration of 50 µg/L and even more severe reactions like desquamation of cells at a concentration of 500 µg/L. Additionally, the degree of tissue damage in the liver of the fish was increased after exposure to low-density PE in a concentration of 500 µg/L (Karami *et al.* 2016). The sizes of the

investigated MP is an important parameter since small MP or nanoplastic particles can be taken up by cells (Batel *et al.* 2016, Mattsson *et al.* 2017, Ding *et al.* 2018). In general, there is no evidence for a uniform pattern under which conditions histopathological changes may occur after exposure to MP.

Reports of the environmental concentration of MP in surface waters of a size <50 µm are rather scarce possibly due to difficulties with the sampling and detection methodology (de Sá *et al.* 2018, Triebkorn *et al.* 2019). Nevertheless, reported results of MP indicate higher concentrations of smaller MP compared to larger ones (Triebkorn *et al.* 2019). Measured MP concentrations in the environment varied between 0.5 and 3.1 particles/L (>20 µm) and 1000 and 9000 particles/L (>4 µm) (Su *et al.* 2018, Triebkorn *et al.* 2019). In the present study the concentration of PS particles was 10<sup>4</sup> and therefore higher than the environmental concentration. Our study does not indicate a risk for brown trout at environmental concentrations. Nevertheless, to exclude potential negative effects of MP on brown trout other experiments with longer exposure time and other polymer types are necessary.

### **4.3. Does methiocarb affect juvenile brown trout?**

Three fish exposed to methiocarb as sole pollutant showed strong behavioral reactions after 24 h and had to be euthanized. After 96 h, all other fish exposed to methiocarb also exhibited behavioral abnormalities like slower swimming and reduced escape behavior. To the best of our knowledge, this is the first study that investigated the effects of methiocarb on brown trout. In the present study weight of methiocarb-exposed fish were comparable to the control group. However, 96 h is a relatively short time to observe changes regarding this endpoint in brown trout. To our knowledge no one has analyzed the effect of methiocarb on stress proteins so far, and rarely proteotoxic effects of carbamate pesticides were studied in general. In the present study, methiocarb did not cause any alteration of Hsp70 level of juvenile brown trout. Samples of brown trout showed neither an increase of SOD nor an altered amount of LPO exposure to methiocarb. An increase of LPO and an alteration of reduced glutathione level was found in male Wistar rats fed with 2, 10, and 25 mg/kg methiocarb (Ozden and Alpertunga 2010). Ozden and Alpertunga (2010) found the highest malondialdehyde level in the brain and explained their finding by the comparably large amount of fatty acids in this organ. In another study, Ozden *et al.* (2012) found also an increase of glutathione as well as of the activities of SOD, CAT, and glutathione peroxidase in male Wistar rats after administration of 25 mg/kg methiocarb.

The observed reactions of fish can be seen as a consequence of the AChE inhibition by methiocarb. Acetylcholine (ACh) accumulates in the synaptic cleft leading to a cholinergic crisis (Rosman *et al.* 2009). In the present study, methiocarb led to a reduction of the AChE activity by 59 % in the tested juvenile brown trout. The mode of action of carbamate pesticides is based on carbamylation of AChE and, thereby, inhibition of its ability to hydrolyze acetylcholine (Fukuto 1990). Therefore, it could have been expected that methiocarb reduces the activity of AChE also in brown trout. Comparable effects were observed by Essawy *et al.* (2008) in the land snail *Eobania vermiculata* in which AChE activity was reduced up to 69.3 % by methiocarb. Carboxylesterases play an important role in pesticide detoxification (Wheelock *et al.* 2005). Sanchez-Hernandez *et al.* (2009) suggested that CbE might act as biochemical

barrier for organophosphate pesticides in *Lumbricus terrestris*. Maymó *et al.* (2006) found that esterase activity is higher in western flower thrips (*Frankliniella occidentalis*) with increased resistance against methiocarb. To our knowledge, no studies about the effect of methiocarb on AChE and CbE activities in fish were performed up to date. However, in the present study, CbE as well as AChE was inhibited by methiocarb and no protective effect became obvious. This might possibly be related to the fact that the methiocarb concentration was rather high and all three enzymes were inhibited and the protective effect of CbE ceased.

After exposure to methiocarb, prominent histopathological alterations became evident in livers and gills of the exposed fish. Altinok *et al.* (2006) observed similar effects in gills of rainbow trout exposed to 3.75 and 7.5 mg/L methiocarb. After 96 h, symptoms like lamellar edema, lifting of epithelia, telangiectasis, increased cytoplasmic granularity, and lamellar fusion occurred. Effects were reversible in concentrations below 3.75 mg/L. In fish livers, Altinok *et al.* (2006) found necrosis. Altinok *et al.* (2006) assumed that the alterations are caused by ionic imbalance due to inhibition of AChE activity. In a follow up study, Altinok and Capkin (2007) found no histopathological alterations in liver, kidney, brain, and spleen of rainbow trout after exposure to 2.5 or 3.75 mg/L methiocarb for 21 days. However, in gills of rainbow trout lamellar lifting occurred when fish were exposed to 3.75 mg/L methiocarb for 21 days (Altinok and Capkin 2007). Brown trout seem to be more sensitive to methiocarb than rainbow trout as indicated by the more severe effects in the present study. This was also shown in the past for other environmental stressors (Schmidt-Posthaus *et al.* 2001).

The tested methiocarb concentration (1 mg/L) was considerably higher than the average surface water concentration of 6 - 40 ng/L in an EU wide monitoring campaign (Loos *et al.* 2018). The study was designed to investigate acute effects of methiocarb and a potential modulation of these effects by PS-MP. Nevertheless, the strong acute effects of methiocarb in brown trout elucidate the need for experiments in which brown trout are exposed to methiocarb at environmentally relevant concentrations for a longer exposure time. Such experiments would allow to assess the current risk of methiocarb exposure for brown trout in the environment.

#### **4.4. Do PS-MP modulate the effects of methiocarb?**

When considering all investigated endpoints, fish exposed to the mixture of methiocarb and PS-MP showed the same reactions as fish exposed to methiocarb only. Weight as well as the stress protein level and the level of oxidative stress remained unchanged. However, the activity of AChE and CbE were significantly reduced in the mixture to almost the same extent as caused by methiocarb alone. Furthermore, the observed histopathological alterations in liver and gills in fish of the mixture treatment were comparable to those found in fish exposed only to the pesticide. Thus, the toxicity of methiocarb on brown trout was not modulated by PS-MP. The chemical analytics and the observed effects indicate that no relevant sorption of methiocarb to PS-MP occurred. It is possible that a modulation of the toxicity of methiocarb by PS-MP in the present study was hidden by the comparable strong effect caused by the relatively high methiocarb concentration. To the best of our knowledge, the interaction of carbamate

pesticides and MP have not been investigated before. In their review de Sá *et al.* (2018) identified only one out of 59 studies where no interaction between MP and another tested contaminant was found. Ferreira *et al.* (2016) reported, that the effects of gold nanoparticles were not modulated by PE particles (1-5 µm). Of course, de Sá *et al.* (2018) could not account in their meta-analysis for the probable bias against publishing negative results. Multiple studies show that effects of different pollutants were decreased in combination with MP. For example, PS-MP alleviated the effects of 17 α-ethinylestradiol (EE2) on locomotion in zebrafish (Chen *et al.* 2017). Similar results were found after exposure of zebrafish larvae to EE2, phenanthrene and a mixture of both with PVC particles. In the presence of PVC particles, the expression of cytochrome P4501A and vitellogenin was reduced up to 48 % for EE2 and by 33 % for phenanthrene (Sleight *et al.* 2017). In general, a reduced toxicity of co-contaminants in the presence of MP might be explained by a decreased bioavailability of the chemicals due to sorption to the plastic particles. In the present study, chemical analyses revealed that the concentration of methiocarb was higher in the mixture treatment than in the groups exposed to methiocarb alone after 97 h. Thus, it is probable that methiocarb did not sorb in considerable amounts to the plastic microparticles. In contrast to reports on a reduction of chemical-induced effects by MP, other studies conducted with MP in combination with chemicals found an intensification of such effects. For example, in common carp (*Cyprinus caprio*), MP increased the effects of paraquat on biochemical blood parameters. Thereby, higher concentrations of MP increased the toxicity (Nematdoost Haghi and Banaee 2017). Moreover, Fonte *et al.* (2016) found a significant interaction between PE-MP and cefalexin. In the mixture of both, the effect of cefalexin on the predatory performance of common goby was increased at 20 °C, but reduced at 25 °C. In the present study the toxicity of methiocarb was not enhanced by MP. Compared to the uptake pathway via the water, PS-MP seemed to have a negligible effect on the uptake of methiocarb in juvenile brown trout.

## 5. Conclusion

Based on the results of our study, we conclude that methiocarb heavily impairs the health of brown trout whereas the studied PS-MP in a concentration of 10<sup>4</sup> particles/L do not. It can also be excluded that the studied PS-MP modulate methiocarb-induced effects in brown trout. In general, literature provides a diverse and inconsistent image with respect to the capacity of MP to modulate the toxicity of environmental chemicals. Although our study does not speak for an environmental risk related to the investigated polystyrene particles and their interaction with the pesticide methiocarb, this study provides only a very small piece of knowledge for a defined type and size class of plastics and a single pesticide, and emphasizes the need of further research in this field.

## 6. List Of Abbreviations

AChE	acetylcholinesterase	LC <sub>50</sub>	lethal concentration 50 %
ABS570	absorbance 570 nm	MiWa	microplastics in the water cycle
BPA	bisphenol A	MP	microplastics
CAT	catalase	PA	polyamide
CbE	carboxylesterase	PE	polyethylene
CHP	cumene hydroperoxide	PET	polyethylene terephthalate
EE2	17 $\alpha$ - ethinylestradiol	PS	polystyrene
Equiv	equivalents	PP	polypropylene
LoD	limit of detection	PVC	polyvinyl chloride
LoQ	limit of quantification	SOD	superoxide dismutase
LPO	lipid peroxidation		

## 7. Declarations

### 7.1. Ethics approval and consent to participate

The experiment was approved by the animal welfare committee of the regional council of Tübingen, Germany (authorisation number ZO 2/16).

### 7.2. Consent for publication

Not applicable

### 7.3. Availability of data and material

The datasets used and analysed during the current study are available from the corresponding author on request.

### 7.4. Competing interests

The authors declare that they have no competing interests.

### 7.5. Funding

The experiment was conducted within the joint project MiWa (Microplastics in the water cycle – sampling, sample preparation, analytics, occurrence, removal, and assessment) founded by the German Federal Ministry of Education and Research (support code: 02WRS1378).

### 7.6. Authors' contributions

HS wrote the manuscript except for section 4.2., which was written by SH. She contributed to the design of the experiment, conducted the experiment, analysed (or supervised analyses) of stress-proteins and oxidative stress. SH performed the chemical analysis and revised the whole manuscript. SK and TPK revised the entire manuscript. FR analysed the level of lipid peroxidation as well as the acetylcholinesterase and carboxylesterase activity. KR performed the histopathological analysis and parts of the analysis of the SOD activity; ASR provided the microplastic particles and revised the whole manuscript, HRK and RT designed the experiment, contributed to the interpretation of data and revised the entire manuscript.

## 7.7. Acknowledgements

We are deeply grateful to Prof. Dr. Martin Jekel for the initiation and coordination of the MiWa project. We thank Christoph Fahrenson (ZELMI, TU Berlin) for electron microscopic analyses. For proofreading of the manuscript we thank Sarai Rosenberg and Dr. Sophie Schmieg. Furthermore, the authors wish to thank Dr. Katharina Peschke, Michael Ziegler, Stefanie Jacob, Sabrina Wilhelm, Dr. Simon Schwarz, Dr. Carla Lorenz, Dr. Paul Thellmann, Dr. Andreas Dieterich and Kathrin Leinmüller for discussion, technical assistance and assistance in the lab.

## References

- Altinok I. and Capkin E. (2007). "Histopathology of Rainbow Trout Exposed to Sublethal Concentrations of Methiocarb or Endosulfan." Toxicologic Pathology **35**(3): 405-410.
- Altinok I., Capkin E., Karahan S. and Boran M. (2006). "Effects of water quality and fish size on toxicity of methiocarb, a carbamate pesticide, to rainbow trout." Environmental Toxicology and Pharmacology **22**(1): 20-26.
- Arena M., Auteri D., Barmaz S., Brancato A., Brocca D., Bura L., Cabrera L. C., Chiusolo A., Civitella C., Marques D. C., Crivellente F., Ctverackova L., Lentdecker C. D., Egsmose M., Erdos Z., Fait G., Ferreira L., Greco L., Ippolito A., Istace F., Jarrah S., Kardassi D., Leuschner R., Lostia A., Lythgo C., Magrans J. O., Medina P., Mineo D., Miron I., Molnar T., Padovani L., Morte J. M. P., Pedersen R., Reich H., Sacchi A., Santos M., Serafimova R., Sharp R., Stanek A., Streissl F., Sturma J., Szentes C., Tarazona J., Terron A., Theobald A., Vagenende B., Dijk J. V. and Villamar-Bouza L. (2018). "Peer review of the pesticide risk assessment of the active substance methiocarb." EFSA Journal **16**(10).
- Avio C. G., Gorbi S., Milan M., Benedetti M., Fattorini D., d'Errico G., Pauletto M., Bargelloni L. and Regoli F. (2015). "Pollutants bioavailability and toxicological risk from microplastics to marine mussels." Environmental Pollution **198**: 211-222.
- Bakir A., O'Connor I. A., Rowland S. J., Hendriks A. J. and Thompson R. C. (2016). "Relative importance of microplastics as a pathway for the transfer of hydrophobic organic chemicals to marine life." Environmental Pollution **219**: 56-65.

- Batel A., Linti F., Scherer M., Erdinger L. and Braunbeck T. (2016). "Transfer of benzo[a]pyrene from microplastics to *Artemia nauplii* and further to zebrafish via a trophic food web experiment: CYP1A induction and visual tracking of persistent organic pollutants." Environmental Toxicology and Chemistry **35**(7): 1656-1666.
- Beckingham B. and Ghosh U. (2017). "Differential bioavailability of polychlorinated biphenyls associated with environmental particles: Microplastic in comparison to wood, coal and biochar." Environmental Pollution **220**: 150-158.
- Bergmann M., Wirzberger V., Krumpfen T., Lorenz C., Primpke S., Tekman M. B. and Gerdt G. (2017). "High quantities of microplastic in Arctic deep-sea sediments from the HAUSGARTEN observatory." Environmental Science & Technology **51**(19): 11000-11010.
- Bradford M. M. (1976). "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding." Analytical Biochemistry **72**(1): 248-254.
- Bundschuh M., Goedkoop W. and Kreuger J. (2014). "Evaluation of pesticide monitoring strategies in agricultural streams based on the toxic-unit concept – Experiences from long-term measurements." Science of The Total Environment **484**: 84-91.
- Chen Q., Gundlach M., Yang S., Jiang J., Velki M., Yin D. and Hollert H. (2017). "Quantitative investigation of the mechanisms of microplastics and nanoplastics toward zebrafish larvae locomotor activity." Science of The Total Environment **584-585**: 1022-1031.
- de Sá L. C., Oliveira M., Ribeiro F., Rocha T. L. and Futter M. N. (2018). "Studies of the effects of microplastics on aquatic organisms: What do we know and where should we focus our efforts in the future?" Science of The Total Environment **645**: 1029-1039.
- Dieterich A., Troschinski S., Schwarz S., Di Lellis M. A., Henneberg A., Fischbach U., Ludwig M., Gärtner U., Tribskorn R. and Köhler H.-R. (2015). "Hsp70 and lipid peroxide levels following heat stress in *Xeropicta derbentina* (Krynicky 1836) (Gastropoda, Pulmonata) with regard to different colour morphs." Cell Stress and Chaperones **20**(1): 159-168.
- Ding J., Zhang S., Razanajatovo R. M., Zou H. and Zhu W. (2018). "Accumulation, tissue distribution, and biochemical effects of polystyrene microplastics in the freshwater fish red tilapia (*Oreochromis niloticus*)." Environmental Pollution **238**: 1-9.
- Dris R., Imhof H., Sanchez W., Gasperi J., Galgani F., Tassin B. and Laforsch C. (2015). "Beyond the ocean: contamination of freshwater ecosystems with (micro-) plastic particles." Environmental Chemistry **12**(5): 539-550.
- EFSA (2006). Conclusion on the peer review of methiocarb. Scientific Report EFSA. **79**: 1-82.

- Eitzen L., Paul S., Braun U., Altmann K., Jekel M. and Ruhl A. S. (2019). "The challenge in preparing particle suspensions for aquatic microplastic research." Environmental Research **168**: 490-495.
- Ellman G. L., Courtney K. D., Andres V. and Featherstone R. M. (1961). "A new and rapid colorimetric determination of acetylcholinesterase activity." Biochemical Pharmacology **7**(2): 88-95.
- EPA. "Chemistry Dashboard: Methiocarb." <https://comptox.epa.gov/dashboard/DTXSID3032626> accessed at 26.02.2020
- EPA (1994). R.E.D. facts methiocarb. E. P. Agency. United States.
- Essawy A. E., Abdelmeguid N. E., Radwan M. A., Hamed S. S. and Hegazy A. E. (2008). "Neuropathological effect of carbamate molluscicides on the land snail, *Eobania vermiculata*." Cell Biology and Toxicology **25**(3): 275.
- EU (2006). "Council Directive 2006/88/EC." Official Journal of the European Union **L 328**: 14–56.
- European Commission (2014). "Commission implementing regulation (EU) No 187/2014 amending implementing regulation (EU) No 540/2011 as regards the conditions of approval of the active substance methiocarb." Official Journal of the European Union **L 57**: 24-26.
- European Commission (2015). "Development of the first Watch List under the Environmental Quality Standards Directive."
- European Commission (2019a). "Commission implementing regulation (EU) 2019/707 " Official Journal of the European Union **L 120**: 16-19.
- European Commission (2019b). "Commission implementing regulation (EU) 2019/1606." Official Journal of the European Union **L 250**: 53-55.
- Ferreira P., Fonte E., Soares M. E., Carvalho F. and Guilhermino L. (2016). "Effects of multi-stressors on juveniles of the marine fish *Pomatoschistus microps*: Gold nanoparticles, microplastics and temperature." Aquatic Toxicology **170**: 89-103.
- Fonte E., Ferreira P. and Guilhermino L. (2016). "Temperature rise and microplastics interact with the toxicity of the antibiotic cefalexin to juveniles of the common goby (*Pomatoschistus microps*): Post-exposure predatory behaviour, acetylcholinesterase activity and lipid peroxidation." Aquatic Toxicology **180**: 173-185.
- Fukuto T. R. (1990). "Mechanism of action of organophosphorus and carbamate insecticides." Environmental Health Perspectives **87**: 245-254.
- Furnes B. and Schlenk D. (2005). "Extrahepatic metabolism of carbamate and organophosphate thioether compounds by the flavin-containing monooxygenase and cytochrome p450 systems " Drug Metabolism

and Disposition **33**(2): 214.

Guilhermino L., Vieira L. R., Ribeiro D., Tavares A. S., Cardoso V., Alves A. and Almeida J. M. (2018). "Uptake and effects of the antimicrobial florfenicol, microplastics and their mixtures on freshwater exotic invasive bivalve *Corbicula fluminea*." Science of The Total Environment **622-623**: 1131-1142.

Hermes-Lima M., Willmore W. G. and Storey K. B. (1995). "Quantification of lipid peroxidation in tissue extracts based on Fe(III)xylene orange complex formation." Free Radical Biology and Medicine **19**(3): 271-280.

Horton A. A., Walton A., Spurgeon D. J., Lahive E. and Svendsen C. (2017). "Microplastics in freshwater and terrestrial environments: Evaluating the current understanding to identify the knowledge gaps and future research priorities." Science of The Total Environment **586**: 127-141.

ILO (2012). "International Chemical Safety Cards (ICSC): Methiocarb." ICSC: 1766  
[http://www.ilo.org/dyn/icsc/showcard.display?p\\_version=2&p\\_card\\_id=1766](http://www.ilo.org/dyn/icsc/showcard.display?p_version=2&p_card_id=1766). accessed at 11.12.2019

Jeong C.-B., Won E.-J., Kang H.-M., Lee M.-C., Hwang D.-S., Hwang U.-K., Zhou B., Souissi S., Lee S.-J. and Lee J.-S. (2016). "Microplastic size-dependent toxicity, oxidative stress induction, and p-JNK and p-p38 activation in the monogonont rotifer (*Brachionus koreanus*)." Environmental Science & Technology **50**(16): 8849-8857.

Johnson W. W. and Finley M. T. (1980). Handbook of acute toxicity of chemicals to fish and aquatic invertebrates : summaries of toxicity tests conducted at Columbia National Fisheries Research Laboratory, 1965-78. Resource Publication: 106.

Karami A., Romano N., Galloway T. and Hamzah H. (2016). "Virgin microplastics cause toxicity and modulate the impacts of phenanthrene on biomarker responses in African catfish (*Clarias gariepinus*)."  
Environmental Research **151**: 58-70.

Kleinteich J., Seidensticker S., Marggrander N. and Zarfl C. (2018). "Microplastics reduce short-term effects of environmental contaminants. Part II: Polyethylene particles decrease the effect of polycyclic aromatic hydrocarbons on microorganisms." International journal of environmental research and public health **15**(2): 287.

Köck-Schulmeyer M., Ginebreda A., Postigo C., Garrido T., Fraile J., López de Alda M. and Barceló D. (2014). "Four-year advanced monitoring program of polar pesticides in groundwater of Catalonia (NE-Spain)." Science of The Total Environment **470-471**: 1087-1098.

Koelmans A. A., Bakir A., Burton G. A. and Janssen C. R. (2016). "Microplastic as a vector for chemicals in the aquatic environment: Critical review and model-supported reinterpretation of empirical studies." Environmental Science & Technology **50**(7): 3315-3326.

- Lei L., Wu S., Lu S., Liu M., Song Y., Fu Z., Shi H., Raley-Susman K. M. and He D. (2018). "Microplastic particles cause intestinal damage and other adverse effects in zebrafish *Danio rerio* and nematode *Caenorhabditis elegans*." Science of The Total Environment **619-620**: 1-8.
- Lithner D., Damberg J., Dave G. and Larsson Å. (2009). "Leachates from plastic consumer products – Screening for toxicity with *Daphnia magna*." Chemosphere **74**(9): 1195-1200.
- Loos R., Marinov D., Sanseverino I., Napierska D. and Lettieri T. (2018). Review of the 1st watch list under the Water Framework Directive and recommendations for the 2nd Watch List. Luxembourg, Publications Office of the European Union.
- Lu Y., Zhang Y., Deng Y., Jiang W., Zhao Y., Geng J., Ding L. and Ren H. (2016). "Uptake and Accumulation of Polystyrene Microplastics in Zebrafish (*Danio rerio*) and Toxic Effects in Liver." Environmental Science & Technology **50**(7): 4054-4060.
- Luís L. G., Ferreira P., Fonte E., Oliveira M. and Guilhermino L. (2015). "Does the presence of microplastics influence the acute toxicity of chromium(VI) to early juveniles of the common goby (*Pomatoschistus microps*)? A study with juveniles from two wild estuarine populations." Aquatic Toxicology **164**: 163-174.
- Lushchak V. I. (2016). "Contaminant-induced oxidative stress in fish: a mechanistic approach." Fish Physiology and Biochemistry **42**(2): 711-747.
- Magni S., Gagné F., André C., Della Torre C., Auclair J., Hanana H., Parenti C. C., Bonasoro F. and Binelli A. (2018). "Evaluation of uptake and chronic toxicity of virgin polystyrene microbeads in freshwater zebra mussel *Dreissena polymorpha* (Mollusca: Bivalvia)." Science of The Total Environment **631-632**: 778-788.
- Markwell M. A. K., Haas S. M., Bieber L. L. and Tolbert N. E. (1978). "A modification of the Lowry procedure to simplify protein determination in membrane and lipoprotein samples." Analytical Biochemistry **87**(1): 206-210.
- Mattsson K., Johnson E. V., Malmendal A., Linse S., Hansson L.-A. and Cedervall T. (2017). "Brain damage and behavioural disorders in fish induced by plastic nanoparticles delivered through the food chain." Scientific Reports **7**(1): 11452.
- Maymó A. C., Cervera A., Dolores Garcerá M., Bielza P. and Martínez-Pardo R. (2006). "Relationship between esterase activity and acrinathrin and methiocarb resistance in field populations of western flower thrips, *Frankliniella occidentalis*." Pest Management Science **62**(12): 1129-1137.
- Moermond C. T. A., Kase R., Korkaric M. and Ågerstrand M. (2016). "CRED: Criteria for reporting and evaluating ecotoxicity data." Environmental Toxicology and Chemistry **35**(5): 1297-1309.
- Monserat J. M., Geracitano L. A., Pinho G. L. L., Vinagre T. M., Faleiros M., Alciati J. C. and Bianchini A. (2003). "Determination of Lipid Peroxides in Invertebrates Tissues Using the Fe(III) Xylenol Orange Complex Formation." Archives of Environmental Contamination and Toxicology **45**(2): 177-183.

Nematdoost Haghi B. and Banaee M. (2017). "Effects of micro-plastic particles on paraquat toxicity to common carp (*Cyprinus carpio*): biochemical changes." International Journal of Environmental Science and Technology **14**(3): 521-530.

Oliveira M., Ribeiro A., Hylland K. and Guilhermino L. (2013). "Single and combined effects of microplastics and pyrene on juveniles (0+ group) of the common goby *Pomatoschistus microps* (Teleostei, Gobiidae)." Ecological Indicators **34**: 641-647.

Ozden S. and Alpertunga B. (2010). "Effects of methiocarb on lipid peroxidation and glutathione level in rat tissues." Drug and Chemical Toxicology **33**(1): 50-54.

Ozden S., Catalgol B., Gezginci-Oktayoglu S., Karatug A., Bolkent S. and Alpertunga B. (2012). "Acute effects of methiocarb on oxidative damage and the protective effects of vitamin E and taurine in the liver and kidney of Wistar rats." Toxicology and Industrial Health **29**(1): 60-71.

Paul-Pont I., Lacroix C., González Fernández C., Hégaret H., Lambert C., Le Goïc N., Frère L., Cassone A.-L., Sussarellu R., Fabioux C., Guyomarch J., Albentosa M., Huvet A. and Soudant P. (2016). "Exposure of marine mussels *Mytilus* spp. to polystyrene microplastics: Toxicity and influence on fluoranthene bioaccumulation." Environmental Pollution **216**: 724-737.

Pesticide Action Network (2000-2019). "PAN Pesticide Database, ." <http://www.pesticideinfo.org>. accessed at 1.9.2019

Qiao R., Sheng C., Lu Y., Zhang Y., Ren H. and Lemos B. (2019). "Microplastics induce intestinal inflammation, oxidative stress, and disorders of metabolome and microbiome in zebrafish." Science of The Total Environment **662**: 246-253.

Rainieri S., Conlledo N., Larsen B. K., Granby K. and Barranco A. (2018). "Combined effects of microplastics and chemical contaminants on the organ toxicity of zebrafish (*Danio rerio*)." Environmental Research **162**: 135-143.

Rault M., Collange B., Mazzia C. and Capowiez Y. (2008). "Dynamics of acetylcholinesterase activity recovery in two earthworm species following exposure to ethyl-parathion." Soil Biology and Biochemistry **40**(12): 3086-3091.

Rehse S., Kloas W. and Zarfl C. (2018). "Microplastics reduce short-term effects of environmental contaminants. Part I: Effects of bisphenol A on freshwater zooplankton are lower in presence of polyamide particles." International journal of environmental research and public health **15**(2): 280.

Rochman C. M., Hoh E., Kurobe T. and Teh S. J. (2013). "Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress." Scientific Reports **3**: 3263.

Rochman C. M., Kurobe T., Flores I. and Teh S. J. (2014). "Early warning signs of endocrine disruption in adult fish from the ingestion of polyethylene with and without sorbed chemical pollutants from the

marine environment." Science of The Total Environment **493**: 656-661.

Romano N., Ashikin M., Teh J. C., Syukri F. and Karami A. (2018). "Effects of pristine polyvinyl chloride fragments on whole body histology and protease activity in silver barb *Barbodes gonionotus* fry." Environmental Pollution **237**: 1106-1111.

Rosman Y., Makarovskiy I., Bentur Y., Shrot S., Dushnistky T. and Krivoy A. (2009). "Carbamate poisoning: treatment recommendations in the setting of a mass casualties event." The American Journal of Emergency Medicine **27**(9): 1117-1124.

Sanchez-Hernandez J. C., Mazzia C., Capowiez Y. and Rault M. (2009). "Carboxylesterase activity in earthworm gut contents: Potential (eco)toxicological implications." Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology **150**(4): 503-511.

Schiavo S., Oliviero M., Romano V. and Manzo S. (2018). "Ecotoxicological assessment of virgin plastic pellet leachates in freshwater matrices." Journal of Environmental Accounting and Management **6**(4): 345-353.

Schmidt-Posthaus H., Bernet D., Wahli T. and Burkhardt-Holm P. (2001). "Morphological organ alterations and infectious diseases in brown trout *Salmo trutta* and rainbow trout *Oncorhynchus mykiss* exposed to polluted river water." Diseases of Aquatic Organisms **44**(3): 161-170.

Sleight V. A., Bakir A., Thompson R. C. and Henry T. B. (2017). "Assessment of microplastic-sorbed contaminant bioavailability through analysis of biomarker gene expression in larval zebrafish." Marine Pollution Bulletin **116**(1): 291-297.

Smith R., Middlebrook R., Turner R., Huggins R., Vardy S. and Warne M. (2012). "Large-scale pesticide monitoring across Great Barrier Reef catchments – Paddock to Reef Integrated Monitoring, Modelling and Reporting Program." Marine Pollution Bulletin **65**(4): 117-127.

Su L., Cai H., Kolandhasamy P., Wu C., Rochman C. M. and Shi H. (2018). "Using the Asian clam as an indicator of microplastic pollution in freshwater ecosystems." Environmental Pollution **234**: 347-355.

Tourinho P. S., Kočí V., Loureiro S. and van Gestel C. A. M. (2019). "Partitioning of chemical contaminants to microplastics: Sorption mechanisms, environmental distribution and effects on toxicity and bioaccumulation." Environmental Pollution **252**: 1246-1256.

Triebkorn R., Braunbeck T., Grummt T., Hanslik L., Huppertsberg S., Jekel M., Knepper T. P., Kraus S., Müller Y. K., Pittroff M., Ruhl A. S., Schmiege H., Schür C., Strobel C., Wagner M., Zumbülte N. and Köhler H.-R. (2019). "Relevance of nano- and microplastics for freshwater ecosystems: A critical review." TrAC Trends in Analytical Chemistry **110**: 375-392.

Triebkorn R., Telcean I., Casper H., Farkas A., Sandu C., Stan G., Colărescu O., Dori T. and Köhler H.-R. (2008). "Monitoring pollution in River Mureş, Romania, part II: Metal accumulation and histopathology in

fish." Environmental Monitoring and Assessment **141**(1): 177-188.

Usmani K. A., Karoly E. D., Hodgson E. and Rose R. L. (2004). "In vitro sulfoxidation of thioether compounds by human cytochrome p459 and flavin-containing monooxygenase isoforms with particular preference to the CYP2C subfamily." Drug Metabolism and Disposition **32**(3): 333.

van Pomeran M., Brun N. R., Peijnenburg W. J. G. M. and Vijver M. G. (2017). "Exploring uptake and biodistribution of polystyrene (nano)particles in zebrafish embryos at different developmental stages." Aquatic Toxicology **190**: 40-45.

von Moos N., Burkhardt-Holm P. and Köhler A. (2012). "Uptake and Effects of Microplastics on Cells and Tissue of the Blue Mussel *Mytilus edulis* L. after an Experimental Exposure." Environmental Science & Technology **46**(20): 11327-11335.


Wang F., Wong C. S., Chen D., Lu X., Wang F. and Zeng E. Y. (2018). "Interaction of toxic chemicals with microplastics: A critical review." Water Research **139**: 208-219.

Wen B., Zhang N., Jin S.-R., Chen Z.-Z., Gao J.-Z., Liu Y., Liu H.-P. and Xu Z. (2018). "Microplastics have a more profound impact than elevated temperatures on the predatory performance, digestion and energy metabolism of an Amazonian cichlid." Aquatic Toxicology **195**: 67-76.

Wheelock C. E., Shan G. and Ottea J. (2005). "Overview of carboxylesterases and their role in the metabolism of insecticides." Journal of Pesticide Science **30**(2): 75-83.


Worthing C. R., Hance R. J. and British Crop Protection Council (1991). The Pesticide manual a world compendium, Farnham, Surrey, UK British Crop Protection Council.

## Figures



**Figure 1**

Electron micrograph of the cryo-milled gold sputtered PS particles (Scanning electron microscope (SEM) Zeiss DSM 982 GEMINI; secondary electron detector).



**Figure 2**

Specific activity of acetylcholinesterase and carboxylesterases with substrate pnpa and pnpv in muscular tissue of brown trout. The box plots display the median, the 25th and 75th percentiles as well as minimum and maximum values (whiskers); the dots indicate outliers. Different letters indicate significant differences. Methiocarb and mixture significantly reduced the AChE activity as well as CbE-pnpa and CbE-pnpv activity ( $p < 0.0001$ ).



### Figure 3

A: Control status of liver of juvenile trout with homogenous tissue with large hepatocytes. B: The hepatocytes contain a huge amount of glycogen. C: Occurring reactions in the liver were inter alia inflammations (IN), vascular dilation (VD) and focal necrosis (FN). D: Furthermore, reduced glycogen amount and increased intercellular (IC) spaces in liver samples were found. E: Control status of gills of juvenile trout with regularly shaped secondary lamellae. F: Reaction status of gills showing hypertrophy (HT) and hyperplasia (HP) of cells, lamellar fusion (LF) reactions as well as oedema (OE). A, C, E and F: hematoxylin-eosin staining; B and D: alcian blue-PAS staining. G and H: Semi-quantitative analyses of liver (G) and gills (H). Category 1 represents an excellent health status, 3 a reaction status and 5 a destruction status. 2 and 4 are intermediary classes. No section was assigned to category 5. Statistical comparison showed significantly more reactions in liver and gill samples of methiocarb or methiocarb plus PS-MP exposed fish. Different letters indicate significant differences ( $p < 0.0001$ ).

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Formulas.pdf](#)
- [2SchmiegetalSupplement.pdf](#)