Recurrent disc herniation following percutaneous endoscopic lumbar discectomy preferentially occurs when Modic changes are present

Lu Hao

Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University

Shengwen Li

Haining county people's hospital

Junhui Liu

Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University

Zhi Shan

Zhejiang University School of Medicine Sir Run Run Shaw Hospital

Shunwu Fan

Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University

Fengdong Zhao (✉ zhaofengdong@zju.edu.cn)

Zhejiang University School of Medicine Sir Run Run Shaw Hospital <https://orcid.org/0000-0002-2945-8707>

Research article

Keywords: Modic changes, Recurrent disc herniation, Percutaneous endoscopic lumbar discectomy, Herniated cartilage

Posted Date: December 26th, 2019

DOI: <https://doi.org/10.21203/rs.2.19552/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published at Journal of Orthopaedic Surgery and Research on May 14th, 2020. See the published version at <https://doi.org/10.1186/s13018-020-01695-6>.

Abstract

Objective: To investigate the effect of Modic changes (MCs) and the herniated disc component on recurrent lumbar disc herniation (rLDH) following percutaneous endoscopic lumbar discectomy (PELD).

Methods: We included 102 (65 males, 37 females, aged 20–66 yr) inpatients who underwent PELD from August 2013 to August 2016. All patients underwent CT and MRI preoperative. The presence and type of Modic changes were assessed. During surgery the herniated disc component of each patient was classified into two groups: Nucleus pulposus group, Hyaline cartilage group. The association of herniated disc component with Modic changes was investigated. The incidence of recurrent disc herniation was assessed on more than 2-year follow-up.

Results: In total, 11 patients were lost to follow up; the other 91 were followed up for 24–60 months. Of the 91 patients, 99 discs underwent PELD; 28/99 (28.3%) had MCs. Type I and II MCs were seen in 9 (9.1%) and 19 (19.2%), respectively; no type III MCs were found. Among 28 endplates with MCs, according to the herniated disc component, 18/28 (64.3%) showed evidence of hyaline cartilage in the intraoperative specimens, including 6/9 and 12/19 endplates with type I and II MCs, respectively. Among 71 endplates without MCs, 14/71 (19.7%) showed evidence of hyaline cartilage in the intraoperative specimens. Hyaline cartilage was more common in patients with MCs ($P < 0.05$). We found 2 cases of rLDH in the non-MC group ($n = 71$); 6 cases rLDH were found in the MC group ($n = 28$), including 2 and 4 cases for types I and II, respectively. There was no significant difference between types I and II ($P > 0.05$). rLDH was more common in patients with MCs ($P < 0.05$). We found 5 rLDH cases in the hyaline cartilage group ($n = 32$); 3 rLDH cases were found in the nucleus pulposus group ($n = 67$). rLDH was more common in the hyaline cartilage group ($P < 0.05$).

Conclusions: MCs were associated with the herniated cartilage disc component. rLDH following PELD preferentially occurs when MCs or the herniated cartilage are present.

Introduction

Percutaneous endoscopic lumbar discectomy (PELD), a minimally invasive spinal procedure, has become increasingly well accepted by both surgeons and patients who suffer from lumbar disc herniation (LDH). Compared with conventional open surgery, this technique allows the removal of the herniated disc through a very small skin incision.^[1] However, along with the widespread use of PELD, many researchers have stated concerns regarding recurrence.^[2, 3]

Recurrent lumbar disc herniation (rLDH) has been defined as disc herniation at the same level with the reappearance of the same pain after a pain-free interval and magnetic resonance imaging (MRI) confirmation. The true incidence of rLDH after lumbar discectomy is unclear. The recurrence rate of LDH has been reported to be 5–15%.^[3–8] There have been many studies designed to determine the recurrence of LDH, and various risk factors were suggested including disc degeneration, trauma, age, smoking,

gender, and obesity.^[3, 4] Radiologically identifiable factors, such as disc degeneration, disc height, and sagittal range of motion have been shown to be related to spinal instability and consequently to rLDH.^[8–10]

Modic changes (MCs), which are present as signal alterations in the vertebral endplate and adjacent bone marrow, are found on T1- and T2-weighted MRI. These changes are associated with vertebral endplate fissures and disc herniation.^[11–13] However, little is known about the effect of MCs on rLDH.

MCs are known to be associated with LDH-containing cartilaginous fragments.^[11–13] Schmid et al. reported the presence of a cartilaginous endplate in the extruded disc material in 63% of patients with MCs^[11] and the association between the cartilage endplate and spontaneous resorption has been specifically investigated.^[12] However, the effect of LDH-containing cartilaginous fragments on rLDH has received little attention.

The purpose of this study was to investigate the effect of MCs and the herniated disc component on rLDH following PELD.

Materials And Methods

Inclusion criteria

Patients were included if they had radicular pain for at least 3 months that was refractory to 6 weeks of conservative treatment with or without neurological deficit, numbness in the lumbar spine, buttock, and/or lower extremity; and MRI and computed tomography (CT) demonstrated anatomical LDH correlating with symptoms and no disc calcification.

Exclusion criteria

Patients were excluded if they had any of the following: prior lumbar surgery at another institution, disc calcification, segmental instability, vertebral fractures, spinal infection, other types of degenerative disc disease, tumors, or pregnancy.

Recurrent disc herniation

Recurrent herniation was defined as follows: (1) patients showed a successful PELD operation as confirmed by a pain-free interval of at least one month and (2) reappearance of the same pain as presentation and MRI confirmation of the recurrent herniation on the same level.

CT and MRI

CT scan of the lumbar spine was performed using a 16-slice CT scanner (GE LightSpeed Pro 16; GE Healthcare) with a detector configuration of 16×1.25 mm. A standard lumbar spine protocol with a tube voltage of 120 kV, tube current of 100–650 mA, and a rotation time of 0.8 seconds was used. Automatic

tube current modulation based on the patient's size and X-ray attenuation was used. The slice thickness and reconstruction interval were 1.25 mm and 0.625 mm, respectively.

MRI images of the lumbar spine were obtained using a General Electric 1.5-T magnet with a T1-weighted sequence (repetition time/echo time, 560 ms/12 ms; field of view, 320 × 256; receiver bandwidth, variable; 4.0-mm slice with a gap of 1.0 mm; number of excitations, 3) and a T2-weighted sequence (repetition time/echo time, 3000 ms/100 ms; field of view, 320 × 256; receiver bandwidth, variable; 4.0-mm slice with a gap of 1.0 mm; number of excitations, 3).

Patients

According to the inclusion and exclusion criteria, we included 102 (65 males, 37 females; aged 20–66 years) inpatients who underwent PELD from August 2013 to August 2016. All patients had undergone CT and MRI examinations that were evaluated retrospectively. The presence, location, and type of MCs were assessed from the MRI scans. During surgery, the herniated disc component of each patient was classified into two groups: the nucleus pulposus group and hyaline cartilage group. The association of herniated disc component with MCs was investigated. The incidence of recurrent disc herniation was assessed during a follow-up period of at least 2 years (Fig. 1).

Patients were divided into the nucleus pulposus group and hyaline cartilage group, and the MC group and non-MC group. The incidence of rLDH was compared between the nucleus pulposus and hyaline cartilage groups, and between the MCs and non-MC groups.

Ethical approval was obtained from the Medical Ethics Committee of the hospital. Additionally, all patients provided written informed consent for their information to be stored in the hospital's database and used for research.

Data analysis

Data were inputted into Excel (Microsoft Corp., Redmond, Washington) and quantitative results were expressed in terms of a mean and standard deviation. Data were transferred to SPSS version 20.0 (IBM Corp., Armonk, NY, USA). Group t-tests (two-sided), and chi-square tests, were used to compare mean values, and proportions, respectively. A significance level of 0.05 was used, and $p < 0.05$ was considered significant without multiple test adjustment.

Results

In total, 11 patients were lost to follow up; the other 91 patients were followed up for more than two years (24–60 months). Of the 91 patients, 99 discs underwent PELD, and 28/99 (28.3%) had MCs; type I and II MCs were seen in 9 (9.1%), 19 (19.2%), respectively. No type III MCs was found.

Among 28 endplates with MCs, according to the herniated disc component, 18/28 (64.3%) showed evidence of hyaline cartilage in the intraoperative specimens, including 6/9 endplates with type I MCs and

12/19 endplates with type II. There was no significant difference between types I and II ($P > 0.05$). Among 71 endplates without MCs, 14/71 (19.7%) showed evidence of hyaline cartilage in the intraoperative specimens. Hyaline cartilage was more common in patients with MCs ($P < 0.05$).

There were 2 cases of recurrent disc herniation in the non-MC group ($n = 71$); 6 cases of recurrent disc herniation were found in the MC group ($n = 28$), including 2 and 4 cases of types I and II, respectively. There was no significant difference between types I and II ($P > 0.05$). Recurrent disc herniation was more common in patients with MCs ($P < 0.05$).

There were 5 cases of recurrent disc herniation in the hyaline cartilage group ($n = 32$) and 3 cases of recurrent disc herniation in the nucleus pulposus group ($n = 67$). Recurrent disc herniation was more common in the hyaline cartilage group ($P < 0.05$).

Discussion

Summary of findings

To our knowledge, this study is the first to investigate the effect of MCs and the herniated disc component on rLDH following PELD. We found that hyaline cartilage is more common in patients with MCs in the intraoperative specimens, and recurrent disc herniation is more common in patients with MCs. Recurrent disc herniation was more common in the hyaline cartilage group. rLDH following PELD preferentially occurs when MCs or the herniated cartilage are present.

Modic changes and herniated cartilage

The variable composition of herniated tissue has been described previously, with various proportions of nucleus pulposus, annulus, hyaline cartilage, and bone being reported.[14, 15] Tanaka et al.^[16] suggested that the cartilaginous endplate-osteochondral junction is weak, whereas the cartilaginous endplate-inner annulus fibrosus connection is strong, so herniated tissue usually contains a cartilaginous endplate. Rajasekaran et al.^[15] reported that a high proportion of Indian patients had herniations containing cartilage and bone.

MCs, which present as signal alterations in the vertebral endplate and adjacent bone marrow, are found on T1- and T2-weighted MRI. LDH is known to be a strong risk factor for MCs, especially type I.[17, 18] MCs are known to be associated with LDH-containing cartilaginous fragments.^[11–13] Schmid et al. reported the presence of a cartilaginous endplate in the extruded disc material in 63% of patients with MCs. Our study confirmed the results; 18/28 (64.3%) of our patients showed evidence of hyaline cartilage in the intraoperative specimens, including 6/9 endplates with type I MCs, and 12/19 endplates with type II. Among 71 endplates without MCs, 14/71 (19.7%) showed evidence of hyaline cartilage in the intraoperative specimens. Hyaline cartilage was more common in patients with MCs.

Effect of Modic changes and herniated cartilage on rLDH

MCs are associated with the herniated cartilage disc component; however, the effect of MCs and the herniated disc component on the rLDH following PELD has received little attention. Our study showed that rLDH preferentially occurs when MCs or the herniated cartilage are present. For the endplates with MCs, the connection between the cartilage endplate and the vertebral body is relatively weak. The cartilage endplate may be separated from the vertebral body and herniated with nucleus pulposus, so the herniated tissue usually contains a cartilage endplate.

For patients with MCs, these conditions usually imply that the endplate structure has been damaged. The herniated cartilaginous endplate is only one part of the damaged endplate, because the connection between the cartilage endplate and the vertebral body is relatively weak, so other parts of the endplate easily separate from the vertebral body and herniate with the nucleus pulposus, and rLHD occurs. Similar to the MCs endplate, if the herniated cartilage is present in the intraoperative specimen, which also reflects damage to the endplate structure, other parts of the endplate easily separate from the vertebral body and herniate with nucleus pulposus, and rLHD preferentially occurs.

Our study has some limitations. Firstly, it was a retrospective study with a small sample size. The relatively small sample size limits the accuracy of correlation between the rate of rLDH and the type of MCs. Secondly, the study focused on the phenotypic association between rLDH and MCs or the herniated disc component, while no correlative mechanism was studied. In particular, we did not confirm the results using biomechanics and histomorphologic methods. Evidently, further study is needed in the future.

Nevertheless, this study suggests that rLDH preferentially occurs after PELD when MCs are present. If we choose PLED for patients with MCs, we need to inform these patients that they have a higher incidence rate of rLHD; fusion may be a better choice, especially for old patients. Meanwhile, if the herniated cartilage is present in the intraoperative specimen, we also should inform these patients that they have a higher incidence rate of rLHD. Such patients may require a second operation.

Conclusions

MCs are associated with the herniated cartilage disc component. rLHD following PELD preferentially occurs when MCs or the herniated cartilage are present.

Declarations

Ethics approval and consent to participate

The study was reviewed and approved by the institutional review board and the ethics committee of our institution. Patients or their family members agreed to our study, and signed the informed consents.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

All authors state that we have no competing of interest.

Fundings

This study was partially supported by National Natural Science Foundation of China (project number 81871796 and 81802192).

Authors' contributions

All authors were involved in conception and design. Study design: LH, JHL, SWL, SWF and FDZ. Study conduct: LH and ZS. Data collection and analysis: ZS, SWL. Data interpretation: SWL, and FDZ. Drafting manuscript: JHL, LH, and FDZ. All authors take responsibility for the integrity of the data analysis.

Acknowledgements

NO

Authors' information

1 Department of Orthopaedics, Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University.

2 Department of Orthopaedics, Haining county people's hospital.

Abbreviations

Modic changes (MCs)

recurrent lumbar disc herniation (rLDH)

percutaneous endoscopic lumbar discectomy (PELD)

References

1.

- Ruetten S, Komp M, Merk H, Godolias G. Full-endoscopic interlaminar and transforaminal lumbar discectomy versus conventional microsurgical technique: a prospective, randomized, controlled study. *Spine*. 2008;33:931–9. doi:10.1097/BRS.0b013e31816c8af7.
- 2.
- Cheng J, Wang H, Zheng W, Li C, Wang J, Zhang Z, Huang B, Zhou Y. Reoperation after lumbar disc surgery in two hundred and seven patients. *International orthopaedics*. 2013;37:1511–7. doi:10.1007/s00264-013-1925-2.
- 3.
- Kim JM, Lee SH, Ahn Y, Yoon DH, Lee CD, Lim ST. Recurrence after successful percutaneous endoscopic lumbar discectomy. *Minimally invasive neurosurgery: MIN*. 2007;50:82–5. doi:10.1055/s-2007-982504.
- 4.
- Suk KS, Lee HM, Moon SH, Kim NH. Recurrent lumbar disc herniation: results of operative management. *Spine*. 2001;26:672–6.
- 5.
- Connolly ES. Surgery for recurrent lumbar disc herniation. *Clin Neurosurg*. 1992;39:211–6.
- 6.
- Fandino J, Botana C, Viladrich A, Gomez-Bueno J. Reoperation after lumbar disc surgery: results in 130 cases. *Acta Neurochir*. 1993;122:102–4.
- 7.
- Mobbs RJ, Newcombe RL, Chandran KN. Lumbar discectomy and the diabetic patient: incidence and outcome. *Journal of clinical neuroscience: official journal of the Neurosurgical Society of Australasia*. 2001;8:10–3. doi:10.1054/jocn.2000.0682.
- 8.
- Axelsson P, Karlsson BS. Intervertebral mobility in the progressive degenerative process. A radiostereometric analysis. *European spine journal: official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical. Spine Research Society*. 2004;13:567–72. doi:10.1007/s00586-004-0713-5.
- 9.
- Hasegawa K, Kitahara K, Hara T, Takano K, Shimoda H, Homma T. Evaluation of lumbar segmental instability in degenerative diseases by using a new intraoperative measurement system. *Journal of neurosurgery Spine*. 2008;8:255–62. doi:10.3171/spi/2008/8/3/255.
- 10.
- Zhao F, Pollintine P, Hole BD, Dolan P, Adams MA. Discogenic origins of spinal instability. *Spine*. 2005;30:2621–30.
- 11.
- Schmid G, Witteler A, Willburger R, Kuhnen C, Jergas M, Koester O. Lumbar disk herniation: correlation of histologic findings with marrow signal intensity changes in vertebral endplates at MR imaging. *Radiology*. 2004;231:352–8. doi:10.1148/radiol.2312021708.
- 12.

Shan Z, Fan S, Xie Q, Suyou L, Liu J, Wang C, Zhao F. Spontaneous resorption of lumbar disc herniation is less likely when modic changes are present. *Spine*. 2014;39:736–44.

doi:10.1097/brs.0000000000000259.

13.

Joe E, Lee JW, Park KW, Yeom JS, Lee E, Lee GY, Kang HS. Herniation of cartilaginous endplates in the lumbar spine: MRI findings. *AJR American journal of roentgenology*. 2015;204:1075–81.

doi:10.2214/ajr.14.13319.

14.

Moore RJ, Vernon-Roberts B, Fraser RD, Osti OL, Schembri M. The origin and fate of herniated lumbar intervertebral disc tissue. *Spine*. 1996;21:2149–55.

15.

Rajasekaran S, Bajaj N, Tubaki V, Kanna RM, Shetty AP. ISSLS Prize winner: The anatomy of failure in lumbar disc herniation: an in vivo, multimodal, prospective study of 181 subjects. *Spine*. 2013;38:1491–500. doi:10.1097/BRS.0b013e31829a6fa6.

16.

Tanaka M, Nakahara S, Inoue H. A pathologic study of discs in the elderly. Separation between the cartilaginous endplate and the vertebral body. *Spine*. 1993;18:1456–62.

17.

Zhang YH, Zhao CQ, Jiang LS, Chen XD, Dai LY. Modic changes: a systematic review of the literature. *European spine journal: official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical. Spine Research Society*. 2008;17:1289–99.

doi:10.1007/s00586-008-0758-y.

18.

Becker GT, Willburger RE, Liphofor J, Koester O, Schmid G. [Distribution of MRI signal alterations of the cartilage endplate in pre-operated patients with special focus on recurrent lumbar disc herniation]. *RoFo: Fortschritte auf dem Gebiete der Rontgenstrahlen der Nuklearmedizin*. 2006;178:46–54. doi:10.1055/s-2005-858919.

Legends

Legends.

Figures

Figure 1

T1- weighted (a) and T2-weighted (b) MR images of lumbar endplates showed the presence of type II Modic changes (indicated by high signal on T1 and T2) on L4/5 level. MRI showed disc herniation on the same level (c and d), the herniated cartilage was present in the intraoperative specimen (e).

Reappearance of the same pain as presentation and MRI confirmation of the recurrent herniation on the same level after 6months follow up (f and g), then fusion was performed (h and i).