Merkel Cell Carcinoma With Gastric Metastasis, A Rare Presentation: Case Report and Literature Review

Vittorio Durastante (✉ vittorio.durastante@asufc.sanita.fvg.it)

Tolmezzo Hospital Unit: Presidio Ospedaliero di Tolmezzo

Antonello Conte

Tolmezzo Hospital Unit: Presidio Ospedaliero di Tolmezzo

Pier Paolo Brollo

Università degli Studi di Udine <https://orcid.org/0000-0001-5943-9737>

Carlo Biddau

Ospedale Santa Maria degli Angeli di Pordenone

Michele Graziano

Ospedale di Latisana: Presidio Ospedaliero di Latisana

Vittorio Bresadola

Università degli Studi di Udine: Università degli Studi di Udine

Research Article

Keywords: Merkel cell carcinoma, neuroendocrine tumors, gastric surgery, gastric metastasis, immunotherapy for cancer

Posted Date: October 28th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1019469/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Journal of Gastrointestinal Cancer on March 7th, 2022. See the published version at <https://doi.org/10.1007/s12029-022-00817-z>.

Abstract

Introduction.

Merkel cell carcinoma (MCC) is an infrequent, but highly aggressive neuroendocrine neoplasm of the skin with a propensity for recurrence and metastasis. We report a rare case of gastric metastatic localization of this cancer by focusing on the diagnostic, clinical and surgical approach to the patient.

Case report.

Clinical presentation begins with a peripheral lymphadenopathy whose immunohistochemical characterization identifies the lymphatic dissemination of the disease. Gradually the patient develops a severe anaemic state which requires several blood transfusions and surgical gastric resection to remove a large bleeding lesion of the antral region. The histopathological analysis of the specimen confirms the metastatic origin from MCC, but the primitive lesion remains unknown.

Discussion.

Since this clinical situation is very rare, we conducted a review of the literature selecting the few cases reported, in order to evaluate the current knowledge on this topic. Metastatic involvement of the stomach from Merkel cell carcinoma is a rare presentation of this disease progression with a frequent delay in formulating the correct diagnosis and in further treatment which may be life-threatening for the patient. As regards the local treatment, there is no specific guideline, and the therapeutic indication should be tailored on the specific case.

Introduction

Merkel cell carcinoma (MCC) is an infrequent, but highly aggressive neuroendocrine neoplasm of the skin with a propensity for recurrence and metastasis. There are very few epidemiological information regarding MCC. A recent analysis in the US population reports that the MCC incidence rate in 2013 was 0.7 cases/100.000 person-years, which increases exponentially with age from 0.1 to 1.0 to 9.8 (per 100.000 person-years) among age groups 40-44 years, 60-64 years, and ≥ 85 years, respectively [1]. Another epidemiological study conducted in Finland reports an incidence in 1989-2008 of 0.11/100.000 for men and 0.12/100.000 for women, with a mean age at diagnosis of 76 years (range 27-100) and 69% of the patients were women [2]. The 5-year survival ratio varies from up to 60-70% (stage I disease) to less than 15% (stage IV disease) depending on the considered population and it is strictly correlated with the stage at the diagnosis [3–5]. Ultraviolet (UV) radiation exposure, reduced skin pigmentation, history of other sun exposure linked skin cancer (melanoma in particular), Merkel cell polyomavirus infection (MCPyV+), immunosuppression (leukemia or lymphoma, HIV infection, transplanted patient) and chronic inflammatory disorders are known risk factors for MCC development [6–12]. In up to 10% of patients, MCC is diagnosed after histopathological analysis of enlarged lymph nodes specimen with no evident primary cutaneous tumour and an improved prognosis than in cases of evident cutaneous MCC with lymph node metastasis. Besides regional lymph nodes, metastases are commonly found in the skin, distant lymph nodes, lungs, adrenal glands, liver, brain and bones [13–15]. Metastatic spreading to the stomach is a very rare evolution of the disease [16]. Due to this infrequent presentation, there is often a delayed diagnosis and no unified consent about its treatment which, in the majority of the cases, aim to control the symptoms of the metastatic lesion as a bridge to systemic chemo-/immunotherapeutic treatment. In this context, we report our experience of a case of Merkel cell carcinoma with symptomatic gastric metastasis and a contextual review of the current literature.

Methods

After reporting our clinical case, we performed a review of the literature by searching the following terms on PubMed and Scopus databases: “Merkel cell carcinoma” AND “metastasis” AND (“gastric” OR “stomach”). Subsequently, a cross-check of the reference lists of selected articles was performed to find any missed papers. The research has been performed by three independent reviewers (P.R.B., C.B. and M.G.) and closed on the 30th September 2021. Only cases of gastric metastasis from MCC, in the absence of further confounding conditions, were selected. At the end, 20 publications have been considered relevant for the research query (Figure 1) [16–35]. The information available for each clinical case (including the case report described herein) regarding the age, gender, ethnicity, primary site, therapy for primary site, time between initial diagnosis/therapy and metastasis presentation, management of metastasis, outcome at time of report, time between metastatic diagnosis and death have been collected and compared.

Case Report

A 61-year-old Caucasian man with no previous medical history presented to surgical evaluation with rapidly enlarging right inguinal lymph nodes. Surgical excisional biopsy of an inguinal lymph node confirmed the diagnosis of lymphatic metastasis from Merkel cell carcinoma by immunohistochemical assessment (IHC); no primitive skin lesion was found during dermatological evaluation. PET-CT scan (with ¹⁸F-deoxy-glucose) showed suspicious tracer accumulations in the left colonic tract (benign tubular adenomas and a hyperplastic polyp confirmed at histological examination after endoscopic excision) and in some mediastinal lymph nodes. After the initial diagnosis and initial staging, a complete right inguino-crural lymphadenectomy was performed, followed by a cumulative 50 Gy radiant treatment of the region (2 Gy daily, five days a week, 25 sessions). After two months, the follow up abdominal-MRI showed a highly vascularized lesion of the antral wall of the stomach (30 mm in diameter) which aroused the suspicion of a gastrointestinal stromal tumor (GIST). After one month, a total body CT scan showed no possible metastatic nodularity and confirmed the presence of the gastric lesion (47 mm in diameter) with perigastric lymph nodes enlargement (Figure 2). Four months after inguinal radiotherapy, the patient started to develop increasing fatigue followed by emission of melena. The subsequent severe anaemia (Hb 7.6 g/dL) required urgent hospital admission and several blood transfusions were performed; the

patient always remained hemodynamically stable with no cardiorespiratory symptoms. During gastric endoscopy a bioptic specimen of the lesion was collected and the IHC analysis revealed a metastatic MCC localization. In order to control the chronic bleeding, the patient underwent an open gastric antral resection with Roux-en-Y gastro-jejunal anastomosis (Figure 3). Following the surgical procedure, no other blood transfusions were necessary with a rapid and spontaneous increasing in hemoglobin level (10.8 g/dL at hospital discharge); there were no post-operative complications and the patient was discharged on seventh post-operative day. IHC analysis of the surgical specimen confirmed the presence of a MCC metastatic localization on the antral wall of the stomach (positivity for CK20, chromogranin A and synaptophysin; negativity for TTF1 and CD20) with multiple lymph nodal involvements. The patient has been selected, after oncological evaluation, for a first line immunotherapy with avelumab (Bavencio®, 800 mg iv every two weeks), an anti-PDL1 monoclonal antibody. Radiotherapy will be considered depending on the grade of disease control during follow-up.

Discussion

As emerged from our literature review, gastric metastasis from MCC are a very rare clinical presentation with only 21 cases reported so far, including this one (Table 1). With the limits of these small numbers, we noticed that the average age of presentation of the metastatic localization is 70.6-year-old (SD 11.7-year-old), ranging from 46 to 92-year-old; 5 cases are women (24%), while 16 are men (76%). With the same limitations, the average time from the initial diagnosis to gastric metastatic presentation is 22.3 months (SD 16.7 months), ranging from 2 to 72 months. The primary tumor site was unknown in only 2 cases (10%). In the other 19 cases with known primitive lesion: 12 (63%) were located to the lower part of the body (below the umbilical line); 4 (21%) were located to the head and neck region; 3 (16%) were located to the upper extremities. These lesions have been mainly treated at their primary site with surgical resection in variable combinations with chemotherapy and/or local radiotherapy; lymph nodes dissection has been reported in only 4 cases (19%). The signs and symptoms related to gastric metastasis were aspecific and mainly related to gastrointestinal bleeding (melena, hematemesis, fatigue, anaemia, abdominal pain) and metastatic disease (fatigue, weight loss, persistent fever, abdominal pain), only one patient had an asymptomatic presentation. The gastric metastasis has been surgically treated in 8 cases (38%) in order to control the gastrointestinal bleeding and reduce the mass effect as a bridge to systemic therapy; in 2 cases a perforation has occurred, one at presentation and one during chemotherapy. In 13 cases (62%) adjuvant chemotherapeutic treatment has been performed with various combinations of platinum-based (cisplatin, CDDP; carboplatin, CBDCA) regimens and the possible addition of etoposide (EPEG), taxanes and anthracyclines. Chemo-radiotherapy has been reported in one case (5%), while radiotherapy after chemotherapy has been adopted in 2 cases (10%). Of the 8 reported cases from 2016, only 2 (25%) have received immunotherapy for metastatic disease. At the time of case reporting, 8 patients were alive (38%), while 12 were deceased (57%), one was not declared. Of the 12 deceased patient, 10 had this negative outcome due to metastatic progression and in 9 cases the time between metastasis development and death has been reported. With the limits of these small numbers, the average time between gastric metastatic involvement and death is 3.1 months (SD 2.8 months), ranging from 3 weeks to 9 months. It is important to underline that each one of these 9 patients has been treated with standard chemotherapy and none of them has received immunotherapeutic drugs.

According to the most recent guidelines (National Comprehensive Cancer Network, NCCN [36] and European Dermatology Forum - European Association of Dermato-Oncology - European Organization for Research and Treatment of Cancer, EDF-EADO-EORTC [37]) on the management of Merkel cell carcinoma, a wide surgical resection (with 1 to 2 cm margins down to the muscle fascia or the pericranium, regardless of tumour size) of the primary lesion, if present, is the first step of treatment. Surgical biopsy of enlarged peripheral lymph nodes is always recommended. In patients with clinically negative lymph nodes, sentinel lymph node biopsy (SLNB) must be performed whenever possible, regardless of the size of the primary tumor. If lymphatic spreading is confirmed, a complete lymph nodal dissection and/or elective regional radiotherapy to the draining lymph node basin is the current standard of therapy, but none of these have been compared in a randomized setting. When the primary tumor site is known, an adjuvant wide-field radiotherapy might be considered to reduce the risk of nodal spread or recurrence [38,39]. The most common chemotherapeutic treatments (adjuvant or for systemic disease) for MCC are platinum-based (cisplatin, CDDP; carboplatin, CBDCA) regimens, with the possible addition of etoposide (EPEG), taxanes and anthracyclines, but no standardized protocols have been released and their efficacy in advanced stage disease is poor [40]. After the introduction of immunotherapy in 2016, the PD1-PDL1 (programmed cell death-1 and programmed cell death ligand-1) immune-checkpoint pathway has become a key therapeutic target in reactivating immune responses against MCC. Today pembrolizumab [41] and avelumab [42] are indicated as first choice systemic therapy option for metastatic disease (for avelumab, regardless of the PDL1 expression status of the tumour).

Metastatic involvement of the stomach from Merkel cell carcinoma is a rare presentation of this disease progression. Due to its infrequency and later presentation after the primary site/lymph nodal involvement, there is a delay in correctly diagnosing the metastatic disease. This feature often causes a subsequent delay in further systemic treatment and may be life-threatening for the patient. As regards the local treatment of MCC-related gastric metastasis, there is no specific guideline, and the therapeutic indication should be tailored on the specific case.

Declarations

Conflict of interest. The authors declare that they have no conflict of interest.

Human/animal rights. All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2008.

Informed consent. Informed consent was obtained from the patient to be included in the study.

References

1. Paulson KG, Park SY, Vandeven NA, et al. Merkel cell carcinoma: Current US incidence and projected increases based on changing demographics. *J Am Acad Dermatol.* 2018;78(3):457-463.e2. doi:10.1016/j.jaad.2017.10.028

2. Kukko H, Böhling T, Koljonen V, et al. Merkel cell carcinoma – A population-based epidemiological study in Finland with a clinical series of 181 cases. *Eur J Cancer*. 2012;48(5):737–42. doi:10.1016/j.ejca.2011.06.001
3. Agelli M, Clegg LX. Epidemiology of primary Merkel cell carcinoma in the United States. *J Am Acad Dermatol*. 2003;49(5):832–41. doi:10.1016/S0190-9622(03)02108-X
4. Fields RC, Busam KJ, Chou JF, et al. Five Hundred Patients With Merkel Cell Carcinoma Evaluated at a Single Institution. *Ann Sur*. 2011;254(3):465–73; discussion 473-5. doi:10.1097/SLA.0b013e31822c5fc1
5. Schadendorf D, Lebbé C, Zur Hausen A, et al. Merkel cell carcinoma: Epidemiology, prognosis, therapy and unmet medical needs. *Eur J Cancer*. 2017;71:53–69. doi:10.1016/j.ejca.2016.10.022
6. Miller RW, Rabkin CS. Merkel cell carcinoma and melanoma: etiological similarities and differences. *Cancer Epidemiol Biomarkers Prev*. 1999;8(2):153–158.
7. Howard RA, Dores GM, Curtis RE, et al. Merkel Cell Carcinoma and Multiple Primary Cancers. *Cancer Epidemiol Biomarkers Prev*. 2006;15(8):1545–9. doi:10.1158/1055-9965.EPI-05-0895
8. Starrett GJ, Marcelus C, Cantalupo PG, et al. Merkel Cell Polyomavirus Exhibits Dominant Control of the Tumor Genome and Transcriptome in Virus-Associated Merkel Cell Carcinoma. *mBio*. 2017;8(1):e02079-16. doi:10.1128/mBio.02079-16
9. Ziprin P, Smith S, Salerno G, et al. Two cases of Merkel cell tumour arising in patients with chronic lymphocytic leukaemia. *Br J Dermatol*. 2000;142(3):525–8. doi:10.1046/j.1365-2133.2000.03370.x
10. Engels EA, Frisch M, Goedert JJ, et al. Merkel cell carcinoma and HIV infection. *Lancet*. 2002;359(9305):497–8. doi:10.1016/S0140-6736(02)07668-7
11. Clarke CA, Robbins HA, Tatalovich Z, et al. Risk of Merkel Cell Carcinoma After Solid Organ Transplantation. *J Natl Cancer Inst*. 2015;107(2):dju382. doi:10.1093/jnci/dju382
12. Sahi H, Sihto H, Artama M, et al. History of chronic inflammatory disorders increases the risk of Merkel cell carcinoma, but does not correlate with Merkel cell polyomavirus infection. *Br J Cancer*. 2017;116(2):260–264. doi:10.1038/bjc.2016.391
13. Chen KT, Papavasiliou P, Edwards K, et al. A better prognosis for Merkel cell carcinoma of unknown primary origin. *Am J Surg*. 2013;206(5):752–7. doi:10.1016/j.amjsurg.2013.02.005
14. Becker JC, Stang A, DeCaprio JA, et al. Merkel cell carcinoma. *Nat Rev Dis Primers*. 2017;3:17077. doi:10.1038/nrdp.2017.77
15. Xue Y, Thakuria M. Merkel Cell Carcinoma Review. *Hematol Oncol Clin North Am*. 2019;33(1):39–52. doi:10.1016/j.hoc.2018.08.002
16. Hu ZI, Schuster JA, Kudelka AP, et al. Merkel Cell Carcinoma With Gastric Metastasis and Review of Literature. *J Cutan Med Surg*. 2016;20(3):255–8. doi:10.1177/1203475415623509
17. Canales LI, Parker A, Kadakia S. Upper gastrointestinal bleeding from Merkel cell carcinoma. *Am J Gastroenterol*. 1992;87(10):1464–1466.
18. Krasagakis K, Almond-Roesler B, Zouboulis CC, et al. Merkel cell carcinoma: Report of ten cases with emphasis on clinical course, treatment, and in vitro drug sensitivity. *J Am Acad Dermatol*. 1997;36(5 Pt 1):727–32. doi:10.1016/S0190-9622(97)80325-8
19. Idowu MO, Contos M, Gill S, et al. Merkel cell carcinoma: a report of gastrointestinal metastasis and review of the literature. *Arch Pathol Lab Med*. 2003;127(3):367–369.
20. Cubiella J, Salgado M, Riú M, et al. Metástasis gástrica por tumor de Merkel: una causa inusual de hemorragia digestiva alta. *Rev Esp Enferm Dig*. 2004;96(2):150–1. doi:10.4321/S1130-01082004000200011
21. Shalhub S, Clarke L, Morgan MB. Metastatic Merkel cell carcinoma masquerading as colon cancer. *Gastrointest Endosc*. 2004;60(5):856–8. doi:10.1016/S0016-5107(04)02173-X
22. Li M, Liu C. Cytokeratin 20 Confirms Merkel Cell Metastasis to Stomach. *Applied Appl Immunohistochem Mol Morphol*. 2004;12(4):346–9. doi:10.1097/00129039-200412000-00010
23. Hizawa K, Kurihara S, Nakamori M, et al. An autopsy case of Merkel cell carcinoma presenting aggressive intraabdominal metastasis and duodenal obstruction. *Nihon Shokakibyo Gakkai Zasshi*. 2007;104(9):1383–1386.
24. Wolov K, Tully O, Mercogliano G. Gastric Metastasis of Merkel Cell Carcinoma. *Clin Gastroenterol Hepatol*. 2009;7(3):A26. doi:10.1016/j.cgh.2008.07.024
25. Rosa F, Pacelli F, Papa V, et al. Merkel cell carcinoma metastatic to the stomach. *ANZ J Surg*. 2010;80(1-2):100–1. doi:10.1111/j.1445-2197.2009.05181.x
26. Temiz P, Semin A, Lerzan A, et al. Gastric metastasis of Merkel cell carcinoma: case report. *Med J Trakya Univ*. 2012;27(3):323–326.
27. Syal NG, Dang S, Rose J, et al. Gastric metastasis of merkel cell cancer–uncommon complication of a rare neoplasm. *J Ark Med Soc*. 2012;109(7):134–136.
28. Parikh MP, Samo S, Ganipiseti V, et al. Gastric metastasis of Merkel cell carcinoma, a rare cause of gastrointestinal bleeding: case report and review of the literature. *J Gastrointest Oncol*. 2014;5(4):E68-E72. doi:10.3978/J.ISSN.2078-6891.2014.029
29. Capella C, Marando A, Longhi E, et al. Primary gastric Merkel cell carcinoma harboring DNA polyomavirus: first description of an unusual high-grade neuroendocrine carcinoma. *Hum Pathol*. 2014;45(6):1310–4. doi:10.1016/j.humpath.2014.01.019
30. Santos-Juanes J, Palomo C, Rodrigo L, et al. Anaemia due to gastric haemorrhage by a metastatic Merkel cell carcinoma. *Prz Gastroenterol*. 2015;10(4):244–6. doi:10.5114/pg.2015.51183
31. Pedrazzani C, Vitali M, Guglielmi A. A Case of Unexpected Gastric Mass. *JAMA Surg*. 2015;150(12):1187–8. doi:10.1001/jamasurg.2015.0981
32. Hulstaert E, Smith V, Mielants H, et al. Rapidly growing gastric metastasis of Merkel cell carcinoma, an unusual cause of melena. *Acta Clin Belg*. 2016;71(4):263–6. doi:10.1080/17843286.2015.1105612

33. Trivedi D, Collins V, Roberts E, et al. Perforated gastric metastasis of Merkel cell carcinoma: Case report and review of the literature. *Human Pathology: Case Reports*. 2017;8:20–3. doi:10.1016/j.ehpc.2017.01.001
34. Ha JY, Park SE, Kim HS, et al. A case report of recurrent Merkel cell carcinoma with synchronous metastases to the heart and stomach. *Medicine (Baltimore)*. 2018;97(44):e13032. doi:10.1097/MD.00000000000013032
35. Kini JR, Tapadia R, Shenoy S. Merkel Cell Carcinoma Metastasis to Stomach: an Infrequent Culmination of a Rare Neoplasm. *J Gastrointest Cancer*. 2018;49(1):78–80. doi:10.1007/s12029-016-9852-2
36. Bichakjian CK, Olencki T, Aasi SZ, et al. Merkel Cell Carcinoma, Version 1.2018, NCCN Clinical Practice Guidelines in Oncology. *J Natl Compr Canc Netw*. 2018;16(6):742–774. doi:10.6004/jnccn.2018.0055
37. Lebbé C, Becker JC, Grob JJ, et al. Diagnosis and treatment of Merkel Cell Carcinoma. European consensus-based interdisciplinary guideline. *Eur J Cancer*. 2015;51(16):2396–403. doi:10.1016/j.ejca.2015.06.131
38. Servy A, Maubec E, Sugier PE, et al. Merkel cell carcinoma: value of sentinel lymph-node status and adjuvant radiation therapy. *Ann Oncol*. 2016;27(5):914–9. doi:10.1093/annonc/mdw035
39. Bhatia S, Storer BE, Iyer JG, et al. Adjuvant Radiation Therapy and Chemotherapy in Merkel Cell Carcinoma: Survival Analyses of 6908 Cases From the National Cancer Data Base. *J Natl Cancer Inst*. 2016;108(9):djw042. doi:10.1093/jnci/djw042
40. Nghiem P, Kaufman HL, Bharmal M, et al. Systematic literature review of efficacy, safety and tolerability outcomes of chemotherapy regimens in patients with metastatic Merkel cell carcinoma. *Future Oncol*. 2017;13(14):1263–1279. doi:10.2217/fon-2017-0072
41. Nghiem PT, Bhatia S, Lipson EJ, et al. PD-1 Blockade with Pembrolizumab in Advanced Merkel-Cell Carcinoma. *N Engl J Med*. 2016;374(26):2542–52. doi:10.1056/NEJMoa1603702
42. D'Angelo SP, Russell J, Lebbé C, et al. Efficacy and Safety of First-line Avelumab Treatment in Patients With Stage IV Metastatic Merkel Cell Carcinoma. *JAMA Oncol*. 2018;4(9):e180077. doi:10.1001/jamaoncol.2018.0077

Tables

Table 1. Literature review of 21 cases (this article included) of gastric metastasis from Merkel cell carcinoma (MCC).

Case report	Age (years)	Gender	Ethnicity	Site of primary MCC	Treatment of primary site lesion	Time between initial diagnosis or therapy and metastasis (months)	Clinical presentation	Treatment of metastasis	Outcome (at the time of report)	Time between metastatic disease and death
Canales et al. (1992) [17]	67	M	Unknown	Right gluteal region	Surgical resection and local RT	11	Generalized fatigue, orthostatic symptoms, melena	Total gastrectomy, CP, VCR, DOX	Receiving CT	NA
Krasagakis et al. (1997) [18]	81	M	Unknown	Right hand (5th finger)	Surgical resection, axillary lymph node dissection, RT, CT	12	GI bleeding	Surgical resection of gastric mass	Deceased for metastatic disease	3 weeks
Idowu et al. (2003) [19]	79	F	Unknown	Right inguinal	Surgical resection, RT, CT	48	Light-headedness, melena, 27.2-kg weight loss within 6 months	Not described	Deceased for metastatic disease	4 months
Cubiella et al. (2004)* [20]	74	M	Unknown	Right retroauricular	Surgical resection, CT	23	Hematemesis, melena	Surgical resection of gastric mass	Deceased for metastatic disease	2 months
Shalhub et al. (2004) [21]	62	M	Unknown	Right neck	Surgical resection	72	Persistent fevers, jaundice	CBDCA, EPEG	Deceased for metastatic disease	1 month
Li and Liu (2004) [22]	46	M	Caucasian	Left preauricular	Surgical resection, RT, CT	8	Melena	CDDP, DOX	Receiving CT	NA
Hizawa et al. (2007)* [23]	85	F	Japanese	Right eyelid	Surgical resection, RT	20	Vomit	Palliative care	Deceased for metastatic disease	2 months
Wolov et al. (2009) [24]	80	M	Unknown	Right upper extremity	Surgical resection, RT	24	Progressive fatigue, syncope, melena	Combined CT-RT with 4 cycles of CBDCA	Deceased	Unknown
Rosa et al. (2010) [25]	72	F	Unknown	Right inguinal	Surgical resection, RT, CT	24	Light-headedness, epigastric pain, hematemesis, weight loss	Billroth II resection, 2 cycles of adjuvant CT	Alive after 24 months of follow-up	NA
Temiz et al. (2010) [26]	75	M	Unknown	Left thigh	Surgical resection	10	Perforated duodenal ulcer	Incidental resection of gastric mass	Deceased	17 months
Syal et al. (2012) [27]	68	M	Unknown	Left leg	Surgical resection, RT, CT	Unknown	Melena, fatigue, exertional dyspnea, epigastric pain	CT, RT	Receiving RT	NA
Parikh et	60	M	Hispanic	Right inguinal	CT, RT	4	Fatigue,	3 cycles of	Unknown	Unknown

al. (2014) [28]								weakness, black-colored stools	CT		
Capella et al. (2014) [29]	72	M	Unknown	Unknown	NA	2	Epigastric pain and 9-kg weight loss within 1 month.	Total gastrectomy and splenectomy, EPEG, CDDP	Deceased for metastatic disease	2 months	
Santos-Juanes et al. (2016) [30]	52	M	Unknown	Right thigh	Surgical resection, right inguinal lymph nodes dissection, RT, CBDCA, EPEG	24	Anaemia	CT	Deceased for metastatic disease	6 months	
Pedrazzani et al. (2016) [31]	70	F	Unknown	Left knee	Surgical resection	19	Asymptomatic	No adjuvant therapy	Alive after 12 months follow-up	NA	
Hu et al. (2016) [16]	92	M	Caucasian	Left upper extremity	Surgical resection, RT	15	Fatigue, early satiety, syncope, melena, 9-kg weight loss within 1 month	Palliative care	Deceased for metastatic disease	5 weeks	
Hulstaert et al. (2016) [32]	77	M	Caucasian	Abdominal wall (left sided paraumbilical)	Surgical resection	36	Melena	CBDCA, EPEG (refused surgery)	Receiving CT	NA	
Trivedi et al. (2017) [33]	52	M	Vietnamese	Left gluteal region	Biopsy, RT	36	Ulcerated skin lesion, gastric perforation during CT	CDDP, EPEG, emergency wedge gastrectomy	Alive after 18 months follow-up	NA	
Ha et al. (2018) [34]	82	M	Unknown	Left gluteal region	Surgical resection, RT	36	Nausea, vomiting, 10-kg weight loss within 1 month	CT, RT, immuno-therapy	Deceased for metastatic disease	Unknown	
Kini et al. (2018) [35]	75	F	Unknown	Left gluteal region	Surgical resection, left pelvic and inguinal lymph nodes dissection	12	Recurrent vomiting and jaundice	CT	Deceased for metastatic disease	9 months	
Durastante et al. (2021)	61	M	Caucasian	Unknown (right inguinal lymph nodes enlargement)	Right inguinal lymph nodes dissection, RT	9	Fatigue, anaemia and melena	Partial gastrectomy, immuno-therapy (anti-PDL1)	Receiving immuno-therapy (RT under evaluation)	NA	

M: male; F: female; RT: radiotherapy; CT: chemotherapy; CT-RT: chemo-radiotherapy; GI: gastro-intestinal; CP: cyclophosphamide; VCR: vincristine; DOX: doxorubicin; CBDCA: carboplatin; EPEG: etoposide; CDDP: cisplatin; NA: not applicable.

* publication not in English language

Figures

Figure 1

Study attrition diagram

Figure 2

Diagnostic images of our case (a) MRI slice showing the gastric metastatic lesion of the antral region of the stomach (arrow) (b) CT scan slice showing the same lesion (47 mm in diameter) and a nearby enlarged perigastric lymph node

Figure 3

Post-resection view of the gastric metastasis from Merkel cell carcinoma after the opening of the surgical specimen