

# Parental, Gestational, and Early-life Exposure to Indoor Environmental Hazardous Factors on Allergic Rhinitis Among Preschool Children in Urumqi City: A Case-control Study

**Haonan Shi**

School of public health, Xinjiang Medical University

**Huizhen Qi**

Department of Neurology, The Fifth Affiliated Hospital of Xinjiang Medical University

**Lan Jiang**

Department of Laboratory Medicine, Xinjiang Uyghur Autonomous Region Maternal and Child Health Hospital

**Yuting Lin**

Department of Laboratory Medicine, Xinjiang Uyghur Autonomous Region Maternal and Child Health Hospital

**Tingting Wang**

School of Nursing & Health Management, Shanghai University of Medicine & Health Sciences

**Jian Yao**

School of public health, Xinjiang Medical University

**Huijuan Dong**

School of public health, Xinjiang Medical University

**Junwen Lu**

School of public health, Xinjiang Medical University

**Yue Wang**

School of public health, Xinjiang Medical University

**Zhe Yin**

School of public health, Xinjiang Medical University

**Xinyu Qi**

Department of Health Surveillance, Customs Health and Quarantine Service of Urumqi

**Qi Yan** (✉ [13579939479@163.com](mailto:13579939479@163.com))

School of public health, Xinjiang Medical University

**Keywords:** Parental, Gestational, Early-life Exposure, Preschool Children

**Posted Date:** November 8th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-1020475/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

## Background

Allergic rhinitis is a disease associated with impaired quality of life and heredity. This study aimed to investigate the association of allergic rhinitis (AR) in preschool children with exposure to indoor environment-related factors early in life.

## Methods

In August 2019, we implemented a study among 2020 preschool children in Urumqi City using a case-control design. The study included parental reports for the occurrence of AR in children, parental history of respiratory disease, and indoor environmental correlates of maternal exposure from 1 year prior to pregnancy until the child's age of 0-1 years.

## Results

Mode of birth (caesarean section) ( $OR=1.31$ ,  $95\%CI=1.02\sim1.67$ ), father with AR ( $OR=2.67$ ,  $95\%CI=2.08\sim3.44$ ), mother with AR ( $OR=3.70$ ,  $95\%CI=2.88\sim4.74$ ), mother with asthma ( $OR=3.11$ ,  $95\%CI=1.18\sim8.20$ ), and mother with newly purchased furniture in the parents' residence during pregnancy ( $OR=1.49$ ,  $95\%CI=1.03\sim2.14$ ) were risk factors for AR in children.

## Conclusions

The focus of allergic rhinitis should be on children with a family history of AR and asthma and caesarean delivery. Primary prevention efforts for AR in preschool children are avoiding exposure of children to indoor environmental hazardous factors early in life.

## Background

Allergic rhinitis (AR) refers to a noninfectious inflammatory disease of the nasal mucosa involving multiple immunocompetent cells and cytokines mediated by IgE after allergen exposure in atopic individuals[1]. Most people ignore the harms posed by AR because of its milder symptoms, but several studies have confirmed that AR causes a range of serious harms. An important study from the Chinese lung health study group published in *The Lancet* in 2019[2] showed that AR is an important risk factor for asthma in adults. In addition, symptoms of AR adversely affect patients' work and quality of life, and in severe cases, sleep-disordered breathing can occur[3]. AR also causes symptoms such as reduced learning ability and poor concentration in children and adolescents[4]. A 2012 study from 10 cities in China involving an indoor environment and child health (CCHH) showed a prevalence of physician-diagnosed AR ranging from 2.2% to 23.9% in Chinese preschool children, including 9.8% in Urumqi City[5]. Since Urumqi City is a multiethnic aggregation site, there are large differences in lifestyle habits among ethnic groups, and the climate is dry, which may affect the transmission of allergens within the environment. It is therefore necessary to carry out a large epidemiological survey study of AR in preschool

children as soon as possible. A 2019 study suggested that exposure to environmental risk factors early in life may increase the incidence of AR in children[6].

For the reasons set out above, the prevalence of AR among preschool-aged children and mothers' residential environmental factors from preconception to the age of 1 year in 60 kindergartens in Xinshi District, Shayibake District, Tianshan District, Shuimogou District, Toutunhe District, and Midong District were investigated in August 2019. Exploring the association between exposure to indoor environment-related factors and AR in children during this period provides a scientific and theoretical basis for the prevention of AR in children.

## **Methods**

### **Investigated subjects**

In August 2019, according to the administrative division of Urumqi, 8-12 kindergartens were randomly selected from six districts, namely, Xinshi District, Shayibake District, Tianshan District, Shuimogou District, Toutunhe District, and Midong District. A total of 60 kindergartens with AR diagnosed by doctors were investigated. In the same period, children who were not diagnosed with AR in the same kindergarten were used as controls. This study was approved by the ethics committee of the First Affiliated Hospital of Xinjiang Medical University, and the parents of the investigated children provided written informed consent.

### **Investigation contents and methods**

Questionnaires were developed for all children in the case group and control group; the questionnaire contents were adapted from the questionnaires used in the CCHH survey study and were slightly revised according to the specific context of Urumqi City.

The specific survey included:

- (1) Demographic characteristics, including sex, ethnicity, and age at home address.
- (2) Feeding status, including whether they were only children, whether they were breastfed, and duration of breastfeeding.
- (3) Prevalence of AR in children and family members, including AR in children and parents and AR-related symptoms.
- (4) Dwelling environment, including housing type, renovation situation, and furniture purchase situation.
- (5) Living habits, including indoor pets and house plants, smoking, air conditioning, and the use of a new air filtering system.

(6) Dietary habits, including meal types and times.

Before the survey began, our subject group first had contact with the Education Bureau of Urumqi City and the kindergarten teachers, and uniform professional training was provided to the responsible teachers involved in the survey class before the survey. Consent for the questionnaire to be administered to the parents of the children was given in their kindergartens by the preschool teachers, and the questionnaires were completed by the parents or other guardians of the children at home. They were asked to complete the form within 1 week and submit it to the responsible teacher at the kindergarten, who then returned the questionnaire and uniformly sent it to the Education Bureau of Urumqi City. All questionnaires were reviewed by more than 2 trained subject team members, and ineligible questionnaires were excluded.

## Determination of outcomes

(1) AR: parent reports that the child has had at least 1 occasion in which AR has been definitively diagnosed by a physician, with the age of onset determined by the time of the first diagnosis of AR.

(2) Passive smoking: refers to the exposure of a nonsmoker to the smoke emitted from the ignition cigarette or to the smoke exhaled by the smoker for at least 1 day per week[7].

## Statistical analysis

Epi Data 3.1 was used to establish the database, and SPSS 26.0 software was used for data analysis and processing. Univariate analysis used the  $\chi^2$  test. Factors associated with the onset of AR in preschool children were analysed using multivariate logistic regression, reporting odds ratios (odds ratios, ORs) and 95% confidence intervals (95% confidence intervals, 95% CIs). All statistical analyses were performed using  $\alpha=0.05$  two-sided test criterion, and  $P<0.05$  was considered statistically significant.

## Results

### General condition of the investigated subjects

A total of 404 cases were collected in this study, including 225 boys and 179 girls with a mean age of  $5.18 \pm 0.99$  years and a mean age at onset of  $3.29 \pm 1.11$  years. A total of 1616 individuals were matched in control group in a 1:4 ratio of cases to controls. The control group used age and sex as matching specifications. There were 900 boys and 716 girls with a mean age  $5.13 \pm 1.02$  years in the control group. The distribution of mode of birth, father with AR, mother with AR, and mother with asthma showed significant differences between the two groups (all  $P<0.05$ ). There were no significant differences in the distribution of ethnicity, gestational cycle, birth weight, or duration of breastfeeding between the two groups (all  $P>0.05$ ) (Table 1).

## Univariate analysis of AR and indoor environmental factors

The results of univariate analysis suggested that renovation or appearance of dampness or mould in the parents' residence before the mother's pregnancy, newly purchased furniture or passive smoking in the parents' residence during the mother's pregnancy, renovation or passive smoking in the children's residence during the child's 0-1 years of age, and father smoking were the main environmental factors contributing to the development of AR in children (all  $P < 0.05$ ) (Table 2).

## Multivariate logistic regression analysis of AR onset

General condition survey analysis was introduced into the logistic regression model along with variables that were meaningful in the results of univariate analysis; suffering from AR was the dependent variable (0 = no, 1 = yes). Five variables of the final residency main effect model were associated with the onset of AR in preschool children. Mode of birth (caesarean section), father with AR, mother with AR, mother with asthma, and newly purchased furniture in the parents' residence during pregnancy were risk factors for childhood AR (all  $P < 0.05$ ) (Table 3).

## Discussion

Previous investigators have suggested that the prevalence of AR is influenced by both genetic[8] and environmental[9] factors. This study found that the parental origin of AR contributed significantly to the pathogenesis of AR in children. Concomitant with the ongoing development of genome-wide association studies (GWAS), multiple genes implicated in the pathogenesis of AR have been identified thus far. Within the past 10 years alone, there have been more than 20 investigative studies on genetic susceptibility genes for AR. A 2018 study by Waage et al.[10] also identified 41 risk loci for AR, which also suggested that AR has high genetic susceptibility. In addition to AR in parents, this investigation identified maternal asthma as a risk factor for AR in children, the mechanism of which is not fully defined. However, this conclusion was similarly reached in a 2007-2008 survey by Wu et al.[11].

Asthma and AR are both more common allergic diseases with similar aetiologies, often coexisting in the same family or patient[12]. In fact, AR as a significant risk factor for asthma in adults[2] has been previously reported. As early as 2008, a study from Western Europe[13] was published in *The Lancet* proposing that either allergic or nonallergic rhinitis could predict the onset of asthma in adults. Regarding how maternal asthma contributes to the onset of AR in children, we speculate that on the one hand, there may be an overlap with susceptibility genes. In 2007, the first GWAS on asthma in *Nature* identified a locus (ORMDL3, GSDML, ZPBP2, and IKZF3) on chromosome 17q21 that confers an increased risk of childhood asthma[14]. A subsequent study from Japan[15] also confirmed that variants in this locus were involved in the pathogenesis of AR. On the other hand, AR shares the same environmental influences as asthma. A study by Zhao and colleagues[16] found that passive smoking and petkeeping could increase the risk of AR and asthma in children. In addition, there are many genes associated with asthma and AR

pathogenesis and environmental influencing factors that may be responsible for the increased risk of developing AR in children with maternal asthma.

The present investigation found that caesarean section is a risk factor for AR in children. Previous studies have also suggested that children born by caesarean section have a higher risk of respiratory and allergic diseases than those born vaginally[17,18]. In particular, elective caesarean section is more likely to cause respiratory morbidity in children[19]. Many studies have confirmed that caesarean section has an important effect on the development and maturation of the neonatal immune system. This may be because children delivered via the birth canal can be exposed to normal beneficial gut microbes from the maternal vagina during delivery. Caesarean section children are not exposed to these microorganisms, which leads to an imbalance of intestinal flora and the occurrence of diseases such as allergies and autoimmunity[20,21]. The study by Azad et al.[22] also confirmed that infants born by caesarean section had significantly lower richness and diversity of gut flora than those born vaginally, which may be one of the reasons why children born by caesarean section are more susceptible to AR.

For indoor environmental factors, this study only found an association between newly purchased furniture in the parents' residence during the mother's pregnancy and the onset of AR in children. Newly produced furniture, renovation materials, etc., contain large amounts of volatile organic compounds, of which formaldehyde, benzene, toluene, and xylene are the most common substances[23]. Inhalation of such substances at high concentrations irritate the respiratory tract, inducing respiratory disease in children, and formaldehyde is the most representative of these. Many current studies[24] have confirmed that formaldehyde can cause respiratory injury. In addition, substances such as formaldehyde can also cross the placental barrier and act on the embryo, causing abnormal foetal growth and development.

Several studies have previously confirmed that indoor passive smoking[16], parental smoking[25], keeping houseplants[26], petkeeping[16], and indoor dampness phenomena[27] are closely related to the development of respiratory diseases in children, although this study did not draw the above conclusions. This may be because this study employed a case-control design, and for the case group, subjects were selected strictly according to the physician diagnosed AR. Some children who presented with AR symptoms but were not definitively diagnosed with AR were not included in the case group. This may have been overly conservative, resulting in the above factors not being found to be associated with AR pathogenesis in children. Additionally, it has been suggested that outdoor air pollution may negatively affect respiratory health but may also enhance sensitization to some plants and fungi[28]. In 2012, a "coal to gas" project was launched in Urumqi City, Xinjiang, which greatly improved the outdoor air quality and may have impacted some factors in the analysis. Therefore, based on the results of the present investigation, it is not yet possible to demonstrate, that the abovementioned factors are not associated with the onset of AR in children. Further confirmation through population-based cohort studies is required to further clarify the contributing factors to the onset of AR in children.

## Conclusion

Caesarean section, parents with AR, mother with asthma, and buying new furniture in the mother's home during pregnancy are risk factors for AR in preschool children. We should actively carry out AR prevention for cesarean section preschool children with a family history of AR and asthma, prevent children from contacting harmful factors in the indoor environment early in life.

## Abbreviations

AR: allergic rhinitis; *CI*: confidence interval; *OR*: odds ratio

## Declarations

### Ethics approval and consent to participate

The study was conducted strictly in accordance with the Declaration of Helsinki and approved by the research ethics committee of Fudan University [protocol no. IRB00002408 & FWA00002399], all parents and class teachers of the children under investigation have signed written informed consent.

### Consent for publish

Not applicable

### Availability of data and materials

The datasets generated and/or analysed during the current study are not publicly available due the data belongs to the School of Public Health of Fudan University but are available from the corresponding author on reasonable request.

### Competing interests

The authors declare that they have no competing interests.

### Funding

All phases of this study were supported by National Natural Science Foundation of China (grant no. 81860179 and 81861138005), Thirteenth Five Year Key Disciplines (Public Health and Preventive Medicine) and Special Environment and Health Research Key Laboratory Project in Xinjiang for initiating the original research project.

### Authors Contributions

HS designed the study, participated in the field investigation phase, and wrote the airicle.

HQ, TW and QY onceptualized and designed the study, and investigated and revised the manuscript.

LJ, YL, JY, HD, JL, YW, ZY and XQ participated in the field investigation phase.

All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

## Acknowledgements

Acknowledgement is given to National Natural Science Foundation of China (grant no. 81860179, 81861138005), Thirteenth Five Year Key Disciplines (Public Health and Preventive Medicine) and Special Environment and Health Research Key Laboratory Project in Xinjiang for initiating the original research project. Researchers within the School of Public health of Xinjiang Medical University and Urumqi Education Bureau must also be acknowledged for assisting with data collection.

## References

- [1]Seidman M D, Gurgel R K, Lin S Y, et al. Clinical Practice Guideline: Allergic Rhinitis Executive Summary. *Otolaryngology Head and Neck Surgery*, 2015;152(2):197-206. DOI: 10.1177/0194599814562166.
- [2]Huang K W, Yang T, Xu J Y, et al. Prevalence, risk factors, and management of asthma in China: a national cross-sectional study. *The Lancet*, 2019;394(10196):407-418. DOI: 10.1016/S0140-6736(19)31147-X.
- [3]Nathan R A. The burden of allergic rhinitis. *Allergy and Asthma Proceedings*, 2007;28(1):3-9. DOI: 10.2500/aap.2007.28.2934.
- [4]Blaiss M S. Pediatric allergic rhinitis: physical and mental complications. *Allergy & Asthma Proceedings the Official Journal of Regional & State Allergy Societies*, 2008;29(1):1-6. DOI: 10.2500/aap2008.29.3072.
- [5]Zhang Y P, Li B Z, Huang C, et al. Ten cities cross-sectional questionnaire survey of children asthma and other allergies in China. *Chinese Science Bulletin*, 2013;58(34):4182-4189. DOI: 10.1007/s11434-013-5914-z.
- [6]To T, Zhu J, Stieb D, et al. Early Life Exposure to Air Pollution and Incidence of Childhood Asthma, Allergic Rhinitis and Eczema. *European Respiratory Journal*, 2019;55(2):1900913. DOI: 10.1183/13993003.00913-2019.
- [7]Luo L F, Chen J W, Luo M H, et al. Exposure to second-hand smoke and its influencing factors among 15-69 years old residents in Guangzhou city. *Chinese Journal of Public Health*, 2019;35(8):973-976. DOI: 10.11847/zgggws1120915.
- [8]Li J, Zhang Y, Zhang L. Discovering susceptibility genes for allergic rhinitis and allergy using a genome-wide association study strategy. *Current Opinion Allergy Clinical Immunology*, 2015;15(1):33-40. DOI: 10.1097/ACI.000000000000124.

- [9]Dunlop J, Matsui E, Sharma H P. Allergic Rhinitis: Environmental Determinants. *Immunology Allergy Clinics*, 2016;36 (2):367-377. DOI: 10.1016/j.iac.2015.12.012.
- [10]Waage J, Standi M, Curtin J A, et al. Genome-wide association and HLA fine-mapping studies identify risk loci and genetic pathways underlying allergic rhinitis. *Nat Genet*, 2018;50(8):1072-1080. DOI: 10.1038/s41588-018-0157-1.
- [11]Wu K Y, Li Z Q, Xue F, et al. Allergic rhinitis in women and the impact on incidence of allergic diseases in their offspring. *Chinese Journal of Otorhinolaryngology-skull Base Surgery*;2010, 16(4):261-265. DOI: CNKI:SUN:ZEBY.0.2010-04-008.
- [12]Rhodes H L, Thomas P, Sporik R, et al. A birth cohort study of subjects at risk of atopy: twenty-two-year followup of wheeze and atopic status. *American Journal of Respiratory and Critical Care Medicine*, 2002;165(2): 176-180. DOI: 10.1164/ajrccm.165.2.2104032.
- [13]Shaaban R, Zureik M, Soussan D, et al. Rhinitis and onset of asthma: a longitudinal population-based study. *The Lancet*, 2008;372(9643):1049-1057. DOI: 10.1016/S0140-6736(08)61446-4.
- [14]Moffatt M F, Kabesch M, Liang L M, et al. Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature*, 2007;448(7152):470-473. DOI: 10.1038/nature06014.
- [15]Tomita K, Sakashita M, Hirota T, et al. Variants in the 17q21 asthma susceptibility locus are associated with allergic rhinitis in the Japanese population. *Allergy*, 2013;68(1): 92-100. DOI: 10.1111/all.12066.
- [16]Zhao Y, Liu Y Q, Liu M M, et al. Interactive effects of environmental tobacco smoke and pets ownership on respiratory diseases and symptoms in children. *Chinese Journal of Pediatrics*, 2013;51(2):96-100. DOI: 10.3760/cma.j.issn.0578-1310.2013.02.004.
- [17]Hansen A K, Wisborg K, Ulbjerg N, et al. Elective caesarean section and respiratory morbidity in the term and near-term neonate. *Acta Obstetrica et Gynecologica Scandinavica*, 2007;86(4):389-394. DOI: 10.1080/00016340601159256.
- [18]Li S Y, Song Y J, Dong J Y, et al. Influence of delivery mode on cognition and body development of infants. *Chinese Journal of Woman and Child Health Research*, 2014;25(6):922-924. DOI: CNKI:SUN:SANE.0.2014-06-004.
- [19]Hansen A K, Wisborg K, Ulbjerg N, et al. Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *BMJ*, 2008;336(7635):85-87. DOI: 10.1136/bmj.39405.539282.BE.
- [20]Biasucci G, Benenati B, Morelli L, et al. Cesarean delivery may affect the early biodiversity of intestinal bacteria. *Journal of Nutrition*, 2008;138(9):1796s-1800s. DOI: 10.1089/jmf.2007.0500.

- [21]Renz-Polster H, David M R, Buist A S, et al. Caesarean section delivery and the risk of allergic disorders in childhood. *Clinical & Experimental Allergy*, 2010;35(11):1466-1472. DOI: 10.1111/j.1365-2222.2005.02356.x.
- [22]Azad M B, Konya T, Maughan H, et al. Gut microbiota of healthy Canadian infants: profiles by mode of delivery and infant diet at 4 months. *CMAJ*, 2013;185(5): 385-394. DOI: 10.1503/cmaj.121189.
- [23]Tang X, Bai Y, Duong A, et al. Formaldehyde in China: Production,consumption, exposure levels,and health effects. *Environment International*, 2009;35(8):1210-1224. DOI: 10.1016/j.envint.2009.06.002.
- [24]Yan C H, Yi J H. Study progress on mechanism of formaldehyde induced asthma. *China Occupational Medicine*, 2015;42(6):682-684. DOI: 10.11763/j.issn.2095-2619.2015.06.020.
- [25]Liu W, Huang C, Hu Y, et al. Associations between indoor environmental smoke and respiratory symptoms among preschool children in Shanghai, China. *Chinese Science Bulletin*, 2013;58(34):4211-4216. DOI: 10.1007/s11434-013-5728-z.
- [26]Xie L F, Zhu L, Zhang F Z, et al. Relationship between full flowering stages of anemophilous and ligneous plants and consultation rate of allergic rhinitis. *Journal of Clinical Otorhinolaryngology Head and Neck Surgery*, 2016;30(15):1206-1209. DOI: 10.13201/j.issn.1001-1781.2016.15.007.
- [27]Zhao Z H, Zhang X, Liu R R, et al. Prenatal and early life home environment exposure in relation to preschool children’s asthma, allergic rhinitis and eczema in Taiyuan, China. *Chinese Science Bulletin*, 2013;58(34):4245-4251. DOI: 10.1007/s11434-013-5705-6.
- [28]Eguiluz-Gracia I, Mathioudakis A G, Bartel S, et al. The need for clean air: the way air pollution and climate change affect allergic rhinitis and asthma. *Allergy*, 2020; 75(9): 2170-2184. DOI: 10.1111/all.14177.

## Tables

**Table 1. Demographics and characteristics of the AR and controls groups.**

Factors	AR group (%)	Control group (%)	$\chi^2$	<i>P</i> value
Nation			1.22	0.27
Han	336 (90.6)	1433 (88.7)		
Other ethnic groups	38 (9.4)	183 (11.3)		
Mode of birth			11.45	<0.01
Antepartum	178 (44.1)	864 (53.5)		
Caesarean section	226 (55.9)	752 (46.5)		
Gestational cycles			1.35	0.25
Premature delivery	3 (0.7)	24 (1.5)		
Term birth	401 (99.3)	1592 (98.5)		
Birth weight (g)			1.81	0.41
<2500	15 (3.7)	81 (5.0)		
2500~8000	348 (86.1)	1393 (86.2)		
≥8000	41 (10.1)	142 (8.8)		
Duration of breastfeeding			5.35	0.38
No breastfeeding	23 (5.7)	108 (6.7)		
<1 month	29 (7.2)	85 (5.3)		
1 month - 4 months	48 (11.9)	167 (10.3)		
4 month - 6 months	47 (11.6)	172 (10.6)		
6 months - 1 year	107 (26.5)	492 (30.4)		
≥1 year	150 (37.1)	592 (36.6)		
Father suffering from AR			95.71	<0.01
No	214 (53.0)	1249 (77.3)		
Yes	190 (47.0)	367 (22.7)		
Mother suffering from AR			162.24	<0.01
No	197 (48.8)	1292 (80.0)		
Yes	207 (51.2)	324 (20.0)		
Asthma in father			2.84	0.09
No	397 (98.3)	1603 (99.2)		

Yes	7 (1.7)	13 (0.8)		
Asthma in mother			13.91	<0.01
No	393 (97.3)	1606 (99.4)		
Yes	11 (2.7)	10 (0.6)		

**Table 2. Univariate analysis of AR and indoor environmental factors in preschool children between AR and control groups.**

Factors	AR group (%)	Control group (%)	$\chi^2$	P value
Furniture was newly purchased in parents' residence before pregnancy			1.85	0.17
No	298 (73.8)	1244 (77.0)		
Yes	106 (26.2)	372 (23.0)		
Renovation in parents' residence before pregnancy			10.02	<0.01
No	319 (79.0)	1380 (85.4)		
Yes	85 (21.0)	236 (14.6)		
Dampness or mould was noted in the parents' residence before pregnancy			5.31	0.02
No	348 (86.1)	1456 (90.1)		
Yes	56 (13.9)	160 (9.9)		
Furniture was newly purchased in the parents' residence during pregnancy			12.86	<0.01
No	338 (83.7)	1454 (90.0)		
Yes	66 (16.3)	162 (10.0)		
Renovation in parents' residence during pregnancy			4.73	0.30
No	367 (90.8)	1517 (93.9)		
Yes	37 (9.2)	99 (6.1)		
Dampness or mould was noted in the parents' residence during pregnancy			3.79	0.05
No	363 (89.9)	1499 (92.8)		
Yes	41 (10.1)	117 (7.2)		
Furniture was newly purchased in children's homes between 0 and 1 year of age			2.13	0.15

No	359 (88.9)	1474 (91.2)		
Yes	45 (11.1)	142 (8.8)		
Residence renovation in children's homes between 0 and 1 year of age			5.23	0.02
No	371 (91.8)	1532 (94.8)		
Yes	33 (8.2)	84 (5.2)		
Dampness or mould was noted in the parents' residence between 0 and 1 year of age			3.82	0.05
No	368 (91.1)	1516 (93.8)		
Yes	36 (8.9)	100 (6.2)		
A hairy pet was kept in children's homes between 0 and 1 year of age			0.00	0.95
No	392 (97.0)	1569 (97.1)		
Yes	12 (3.0)	47 (2.9)		
Flowering plants were grown in children's homes between 0 and 1 year of age			1.04	0.31
No	300 (74.3)	1239 (76.7)		
Yes	104 (25.7)	377 (23.3)		
Father smoking			7.53	<0.01
No	284 (70.3)	1242 (76.9)		
Yes	120 (29.7)	374 (23.1)		
Mother smoking			0.63	0.43
No	403 (99.8)	1607 (99.4)		
Yes	1 (0.2)	9 (0.6)		
Passive smoking in the mother's residence during pregnancy			16.09	<0.01
No	272 (67.3)	1244 (77.0)		

Yes	132 (32.7)	372 (23.0)		
Passive smoking in children's homes between 0 and 1 year of age			15.17	<0.01
No	252 (62.4)	1168 (72.3)		
Yes	152 (37.6)	448 (27.7)		

**Table 3. Multivariate logistic regression analysis of the onset of AR in preschool children between AR and control groups.**

Factors	Reference group	$\beta$	<i>S-x</i>	Wald $\chi^2$	<i>P</i> value	<i>OR</i>	95% <i>CI</i>
Mode of birth	Antepartum	0.26	0.12	4.49	0.03	1.30	1.02~1.66
Father suffering from AR	No	0.99	0.13	60.74	<0.01	2.68	2.09~3.44
Mother suffering from AR	No	1.32	0.13	108.87	<0.01	3.74	0.92~4.79
Mother suffering from asthma	No	1.09	0.49	4.96	0.03	2.97	1.14~7.76
Furniture was newly purchased in the parents' residence during pregnancy	No	0.58	0.22	6.99	<0.01	1.79	1.16~2.76