De-implementation of Urinary Catheters in Neurosurgical Patients During the Operation and on the Ward: a Mixed-methods Study Protocol

Jeanne-Marie Nollen (✉ J.M.Nollen@lumc.nl)

Leiden University Medical Center: Leids Universitair Medisch Centrum <https://orcid.org/0000-0002-1241-5380>

Anja H. Brunsveld-Reinders

Leiden University Medical Center: Leids Universitair Medisch Centrum

Wilco Peul

Leiden University Medical Center: Leids Universitair Medisch Centrum

Wouter van Furth

Leiden University Medical Center: Leids Universitair Medisch Centrum

Study protocol

Keywords: De-implementation, urinary catheters, mixed-methods, healthcare professionals, patients

Posted Date: November 17th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1021538/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Indwelling urinary catheters (IDUCS) are frequently routinely inserted during transsphenoidal pituitary gland tumour surgery or spinal fusion surgery despite literature stating that there are no indications for using IDUCS during or after these surgeries. The aim of the study is to reduce the number of inappropriately inserted IDUCs in during or after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery with an operation time of <4 hours.

Methods

A pragmatic, before-and-after mixed-methods observational study including medical chart analysis, satisfaction surveys with patients and healthcare professionals and multidisciplinary group interviews to assess the effectiveness of and experiences with various non-invasive de-implementation strategies aimed to decrease the number of inappropriate IDUCS inserted during and after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery in a multicentre context.

Discussion

This paper presents the study protocol of a multi-centred before and after trial that aims to reduce inappropriate IDUC use after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery, and thereby reducing UTIs, shortening hospital stay and increasing patient comfort. The results can be used to de-implement IDUCs after a broad range of surgeries on several wards.

Trial registration

The study has been submitted to the Dutch Trial Register (NTR).

Contributions To The Literature

- The implementation of a variety of de-implementation strategies focussed on the healthcare professional as well as patients that focus on reducing indwelling urinary catheter use and it's complications;
- A greater understanding of patients' experiences with urinating after transsphenoidal resection of pituitary gland tumours and spinal fusion operations;
- Facilitates multidisciplinary discussion on the use of IDUCs in the postoperative phase.

Background

Indwelling urinary catheter (IDUC) placement in instances of neurosurgical interventions such as anterior skull base operations (e.g. transsphenoidal resection of pituitary gland tumours) and spinal fusion

operations (spondylodesis) has become routine for various reasons (1, 2). In patients where an IDUC was not placed during surgery, these will frequently be inserted upon their return at the neurosurgical ward.

Current literature highlights a distinction between appropriate and inappropriate IDUC use in daily practice. The following reasons are generally viewed as appropriate:

- Urinary retention and bstruction of the bladder (3);
- Surgery time > 4 hours (4);
- Mobility restriction $\geq$ 24 hours postoperative (3);
- Administration of large contents of infusion fluid and/or diuretics during operation (3);
- The need to measure the urine production every hour postoperative (5).

Despite abovementioned appropriate reasons for IDUC placement, there are a number of arguments to be made against IDUC use including prolonged recovery time and increased health risks of a different nature. It is commonly known that IDUCs are associated with urinary tract infections (UTIs), non-infection complications (e.g. pain, discomfort, haematuria, mobility restriction and the feeling the need to urinate) and delayed mobilization (6). UTIs need to be treated with antibiotics which can lead to antibiotic resistance (7, 8), and cases of hospital acquired UTIs are associated with longer hospital stay and additional costs (9). The restriction on patient's ability to mobilize due to the IDUC prolongs their recovery time as research shows that early mobilization postoperatively decreases the risk complications and morbidity (e.g. respiratory decompensation/pneumonias, deep venous thrombosis/pulmonary embolism) (10).

The decision to insert an IDUC per- and postoperative transsphenoidal resection of pituitary gland tumours and spinal fusion operations, is based on a number of considerations including but not limited to: the detection of the post-surgical complications diabetes insipidus (DI), post-operative mobility restriction (maximum 24-hours), urinary retention, and convenience for nurses when a patient has an IDUC.

Diabetes insipidus

The key argument in favour of IDUC placement is that it helps ensure close monitoring of fluid balance which is key in early detection and diagnosis of the most common postoperative complication after pituitary surgery, DI, which is characterized by polydipsia and polyuria and can lead to dehydration when left undetected (11). Despite IDUCs are known to contribute to the accuracy of fluid output measures following pituitary surgery, the fluid balance does not need to be measured hourly postoperative (12). Monitoring the fluid balance closely every 6-12 hours following transsphenoidal pituitary surgery sufficient for ensuring early detection and diagnosis of diabetes insipidus (DI) (2, 13, 14).. After spondylodesis operations, there is no additional risk of fluid disturbances, therefore monitoring a fluid balance postoperatively is not obligated (15).

Mobility restriction

In general, after transsphenoidal resection of pituitary gland tumours and spondylodesis operations, postoperative mobilisation restriction occurs only in rare cases, and the duration of the surgery generally does not exceed the twenty-four hour limit (3, 4, 14, 16-19).

Urinary retention

Another common reason IDUCs are inserted postoperatively is post-operative urinary retention (POUR). Urinary retention is the inability to empty the bladder while it being full (20). After anesthesia and surgery without IDUC placement, postoperative urinary retention (POUR) is common, with reported incidence of 5% – 70% after general surgery and up to 50% after spinal surgery (21, 22). Despite POUR being indicated as an appropriate reason for IDUC insertion, intermittent catheterization has been named in literature as preferred intervention due to a lower risk of UTIs (3)

Convenience

IDUCs are frequently inserted after surgery due to convenience of care for nurses, especially after pituitary surgery where one of the nurses' main tasks is to monitor the fluid balance (23). IDUCs reduce nurses' workload as there is no need to mobilize patient to the restroom and collect the urine in bedpans (24).

Alternatives

Instead of inserting an IDUC without an appropriate reason, the urinary output can be measured with the aid of non-invasive, lower risk tools including an urinal or bedpan (19). When a patient is unable to urinate postoperatively, bladder scanners can help assess the urinary retention after which intermittent catheterization can be executed.

If an IDUC is inserted per- or postoperative, literature, regardless of the surgical diagnosis, indicates that IDUCs should be removed promptly, preferably within 24-hours postoperatively, since patients who have an IDUC have an increased risk of 3-7% of developing a urinary tract infection, per extra day the IDUC remains in place (3, 25). However, it is unknown to what extent IDUCs are removed within this 24-hour timeframe after transsphenoidal resection of pituitary gland tumours and spinal fusion operations.

Since IDUCs might be used to a greater extent and possibly for a longer period than is deemed appropriate by literature after these surgeries, with all health consequences this entails, this protocol describes a study to evaluate the effectiveness of multiple de-implementation strategies to reduce the inappropriate use of IDUCs during the operation and in the postoperative phase on the ward. Therefore, the goal of this study is: *“no IDUC, unless...”*

Methods

Design

This pragmatic, mixed-methods observational study collects medical chart data, satisfaction survey data and multidisciplinary group interviews data to assess the effectiveness of and experiences with various non-invasive de-implementation strategies aimed to decrease the number of inappropriate IDUCS inserted during and after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery in a multicentre context. The study has a before-and-after design and a timespan of 2.5 years starting in 2020. The medical chart assessment continues throughout the entire duration of the study whereas the satisfaction surveys and group interviews take place both before and after the de-implementation. The surveys will be held with both patients and healthcare professionals whereas the group interviews will only be held with healthcare professionals. Quantitative methods are used to assess the effect of the de-implementation strategies on IDUC related outcomes including IDUC placement, complications and patients' and healthcare professionals' experiences and the group interviews are used to get insight in the role of each specific professional regarding IDUC use in the patients' journey from pre-operative consult to being discharged.

We have six specific aims:

1. To reduce the number of inappropriate inserted IDUCs in the hospital during and after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery with an operation time of <4 hours;
2. To assess the frequency of intermitted urinary catheterization after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery;
3. To reduce the number of UTIs after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery;
4. To assess the number of urinary retention bladders in relation to the number of IDUCs placed during and after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery;
5. To better understand patients' experiences and to provide a broad understanding of potential factors contributing to patient satisfaction in relation to urinating in the postoperative phase;
6. To investigate healthcare professionals' experiences with IDUCs and the experienced consequences after IDUC de-implementation.

Setting

The study will take place in one university hospital in which both transsphenoidal pituitary gland tumour surgery and spinal fusion surgery is executed and four general hospitals where only spinal fusion surgery is performed. The de-implementation strategies will be implemented in four intervention hospitals: one university hospital and three general hospitals. One general hospital is designated for the control group since, according to the hospitals' neurosurgeons, IDUCs are not routinely placed in this hospital. All hospitals are located in the Randstad, which is the most densely populated area in the Netherlands, and selected based on the following criteria: 1. transsphenoidal pituitary gland tumour surgery and spinal fusion surgery is executed and 2. IDUC use is routinely reported in the medical chart.

Study population

The study population consists of two groups: 1. patients who underwent/will undergo transsphenoidal pituitary gland tumour surgery or spinal fusion surgery and 2. healthcare professionals (e.g. neurosurgeons, neurosurgical residents, operation assistants, recovery nurses, neurosurgical ward nurses).

All patients who underwent transsphenoidal pituitary gland tumour surgery or spinal fusion surgery in 2019 and 2020, and are aged 18 and older, are included in the medical chart assessment. Patients who will undergo transsphenoidal pituitary gland tumour surgery or spinal fusion surgery in 2021 or 2022, and are aged 18 and older, are eligible for the study and have to give consent for the medical chart assessment and survey. Patients who meets any of the following criteria will be excluded from participation: an operation time > 4 hours; having a mobility restriction $\geq$ 24 hours postoperative, having pre-existing bladder complications for which an IDUC is used pre-operatively; peri- or postoperative neurological deficit (e.g. paresis); having pre-existing psychological problems; being unable to understand and/or execute instructions from healthcare professionals and not speaking fluent Dutch or English fluently. If patients are operated in 2021 or 2022 and informed consent is not obtained for the medical chart assessment or the survey, they will be excluded from the study.

Healthcare professionals working as a neurosurgeon, neurosurgical resident, operation assistant, recovery nurse or neurosurgical ward nurse are eligible to participate in the survey and group interviews. All participants must be aged 18 or over and give consent to participate. Healthcare professionals who do not give consent for the survey and/or the group interviews are excluded from the study.

Main outcome

The primary study parameter is the number of IDUCs that are placed during and after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery (spondylodesis). Data corresponding to the IDUC placement will also be collected. Data will be obtained from patients' medical charts.

Secondary outcomes

Secondary outcomes linked to the medical chart assessment are: the incidence of intermittent urinary catheterization, the incidence and volume of urinary retention bladders, incidence of urinary tract infections. Secondary outcomes from the patients' surveys are the postoperative experiences with and without IDUC use and the implications for the recovery process. Outcomes related to healthcare surveys are experiences with postoperative IDUC use and the consequences of de-implementing IDUCs. Secondary outcomes related to the group interviews are perceptions on the role of each specific professional regarding IDUC use in the patients' journey from pre-operative consult to being discharged.

Medical chart assessment

During the pre-operative consult, patients will be asked to participate in the study, thereby participating in the medical chart assessment, by the nurse or resident who attends the consult. The following items will be systematically collected from each medical record:

- The incidence of IDUC placement, including date of insertion, time of placement, location of insertion, reason of insertion and which discipline inserted the IDUC;
- The incidence of intermittent urinary catheterization, including: date and time of insertion, location of insertion, reason of insertion and which discipline inserted the catheter;
- The incidence and volume of urinary retention bladders, including: date of urinary retention and where the urinary retention was noticed

A retention bladder has a urine volume of more than 500 ml (26);

- The incidence of urinary tract infections

The diagnosis of a symptomatic urinary tract infection, with or without an IDUC, is the detection of bacteria and leukocytes in the presence of clinical symptoms (27). This definition is chosen since asymptomatic urinary tract infections can be expected when testing urine from an IDUC without the presence of symptoms and do not require antibiotic therapy (28). Clinical symptoms include painful and frequent urination, fever, flank pain and general malaise (29). The pathogen and leukocytes are detected and identified by using midstream urine for a urine sediment. The sediment must contain $>10^3$ cfu/mL bacteria and >5 leukocytes (27, 30);

- The operation time in minutes;
- The duration of stay in recovery room in minutes;
- The date of operation;
- Age;
- Gender;
- Length of hospital stay in days'

Satisfaction surveys

The patient satisfaction survey will be designed to get insight in patients' experiences postoperatively and to provide a broad understanding of potential factors contributing to patient satisfaction in relation to urinating in the postoperative phase. The healthcare professional satisfaction survey will be designed to acquire a greater understanding of healthcare professionals' experiences with IDUCs and the experienced consequences after IDUC de-implementation. Both surveys will be tested by pilot participants selected from the neurosurgical ward and the operation room. After piloting and revision, the surveys will be sent to all eligible healthcare professionals via their work-email in the before and after measurement phase. During the pre-operative consult, patients will be asked to participate in the survey by the nurse or resident

who attends the consult. Patients will receive a hardcopy of the survey if they are admitted to the hospital.

Group interviews

A purposive sampling method will be used to create a diverse and representative sample of at least one professional from each profession. Healthcare professionals will be asked to participate in the focus group via their work e-mail. The participants are all working in one of the four participating hospitals and there will be no mixing between hospitals as policies and procedures can differ per site. The group interviews will be held at a date and place most suitable for the participants in a meeting room in the specific hospital. The group interview will focus on the following topics: 1. participants' experiences with the current IDUC policy pre- and postoperatively, 2. perceptions and experiences with intercollegiate collaboration and communications regarding IDUC use and 3. perceptions regarding the patients' role. In addition, demographics including information on age, working experience and gender will be collected at the beginning of the group interviews. The interviews will be led by an independent moderator. At least one of the researchers will also attend the group interviews to answer specific questions related to the topics. The expected duration of the interviews is 60 – 80 minutes. The interviews will be taped and transcribed verbatim.

De-implementation strategies

In this study, multiple de-implementation strategies will be used to decrease the number of inserted IDUCS in pituitary and spinal fusion patients. The de-implementation strategies focus primarily on healthcare professionals. The rationale behind using multiple strategies is that the components positively influence one another and add to acquiring the wanted effect (31). The de-implementation strategies will take place in the four intervention hospitals. There will be no strategies implemented in the control hospital.

Flowcharts

Three flowcharts (figure 1, 2 and 3) were created based on the indications for appropriate IDUC use in combination with the treatment of POUR. The flowcharts advocate intermittent catheterization over inserting an IDUC as this intervention has a lower risk of UTIs (3). Flowchart 1 is designed to use during prior to the operation when deciding on IDUC placement.

Education

Neurosurgeons, neurosurgical residents, operation assistants, recovery nurses and neurosurgical ward nurses will receive education regarding appropriate and inappropriate IDUC use during and after pituitary and spondylodesis surgery. The information sessions consists of a presentation presented by the researcher will be held once per intervention hospital. Figures 1, 2 and 3 will be used as basis for the educational programme. Additionally, the importance of reducing IDUC use as well as possible complications will be discussed. Healthcare professionals will receive information on how to document IDUC use comprehensively and thorough (e.g. date and time of insertion, location of insertion, reason of insertion,

which discipline inserted the IDUC, date and time of urinary retention and volume of urinary retention). Due to the COVID-19 restrictions, the educational sessions will be organised in online events.

Information

Information regarding the existence of the study will be distributed among the healthcare professionals in the hospitals to create awareness. Intranet, social media and hospital newsletters will be used for dissemination.

Reminders

Informational posters regarding (in)appropriate IDUC use will be placed in the breakrooms of the operation theatre, neurosurgical residents and on the neurosurgical ward .

Organizational strategies

To ensure a structural change in IDUC use, the new policy will be established in the formal and informal rules of each hospital. This means that procedures and protocols regarding inserting an IDUC will be changed.

Feedback

The outcome measures of the collected data regarding IDUC at baseline and after measurements will be communicated to each hospital during the study once per month by sending newsletters to participating healthcare professionals.

Patient information

During the pre-operative consult with the neurosurgeon, all patients will receive information regarding the use of an IDUC during and after the surgery. Patients will receive an infographic explaining the reason for IDUC reduction including alternatives use.

Sample size

In the academic hospital, approximately 150 patients undergo a transsphenoidal pituitary gland tumour surgery per year. In all five hospitals combined, approximately 657 patients undergo a spinal fusion surgery (spondylodesis) per year. The duration of the study is 2,5 years which means that the medical charts of a total of 375 pituitary patients and 1643 spondylodesis patients can be included in the study.

Patients will be asked to fill in the patient satisfaction survey for a period of two months during the basement measurement period as well as in the after measurement period. Per month, 12-13 patients will undergo pituitary surgery. Therefore, a total of 48 – 52 pituitary patients will be asked to participate in the survey. In the hospitals combined, 55 patients will have a spondylodesis operation every month, which means that over a period of four months 219 patients will be asked to fill in the survey.

For all five hospitals combined, there are 650 healthcare professionals who are involved in the care for pituitary and/or spondylodesis patients. These participants will be asked to participate in a satisfaction survey at baseline measurement as well as the after measurement. Group interviews will be held in each intervention hospital in the baseline and after measurement phase. The group interviews will consist of six to eight participants as literature indicates that this number is sufficient (32). Per hospital 12-16 healthcare professionals will be asked to participate. In total, 60 to 80 healthcare professionals will be asked to participate in the group interviews.

Analysis

We used a combination of qualitative and quantitative data to answer the primary and secondary outcomes. The medical chart research and the satisfaction surveys will be analysed with quantitative techniques while the group interviews will be analysed with the aid of qualitative methods. A deletion method will be used to eliminate missing data.

Primary outcome

The primary study parameter will be the number of IDUCS that are placed during and/or after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery (spondylodesis). The number of inappropriately/appropriately placed IDUCS will be determined with the aid of figures 1, 2 and 3. The software programme SPSS is used during the analysis. The data will be analysed for all hospitals combined with a logistic regression with corrections for several baseline characteristics of the population (e.g. age, sex, type of operation, hospital and COVID-19 period). Data corresponding with the IDUC placement (e.g. date of insertion, time of placement, location of insertion, reason of insertion and which discipline inserted the IDUC) will be analysed with descriptive statistics. The data will be analysed per hospital with the aid of descriptive statistics. Continuous data will be presented as median (interquartile range) or mean (standard deviation) and where appropriate categorical variables as number (percentages). Graphic data displays may also be used to summarize the data. Descriptive statistics will also be presented for the baseline measurement and the after measurement separately.

Since the control hospital states that there are no IDUCS inserted during/after spondylodesis operations prior to the study, it will be determined to what degree the data from this hospital can be incorporated in the analysis after the baseline measurement. If there are (almost) no IDUCS inappropriately placed, the data will only be analysed with descriptive statistics. The data will be incorporated in the logistic regression if IDUCS are frequently inappropriately inserted.

Secondary outcome

Medical chart

The incidence of intermittent urinary catheterization, the incidence of urinary retention and the incidence of urinary tract infections will be analyzed equally to the primary study parameter.

Surveys

The surveys from the healthcare professionals will be analyzed with the aid of a paired non-parametric T-test. The surveys from the patients will be analyzed with a non-paired non-parametric test. Demographics will be analyzed with descriptive statistics.

Group interviews

After the interviews are transcribed, the software program Atlas.ti will be used to analyze the data. The grounded theory will be used as a framework for the analysis (33, 34). This analysis involves three sequential phases of coding: open, axial and selective coding (35). An iterative approach was used which implies that data collection and analysis occurred simultaneously (36, 37). Two researchers will independently code the transcripts and afterwards discuss the findings to reach consensus about the interpretation.

Ethics and funding

Approval from the Ethical Committee was obtained for all five hospitals either at site level or where this did not exist, from a scientific committee at the site. The researchers will adhere to ethical standard for research involving people. In addition, all researchers will follow their institutional ethical requirements. Funding sources did not have a role in the writing of this manuscript or the decision to submit the publication. Before the operations, the surveys and the group interviews, patients and healthcare professionals will be given an informed consent form as well as information on the study and the participants' rights. It will be specifically stated that participation is voluntary, that participants can withdraw at any time and that confidentiality is guaranteed through anonymization. Per request, the results of the study will be communicated to the participants by email.

Discussion

This paper presents the study protocol of a multi-centred before and after trial that aims to reduce inappropriate IDUC use after transsphenoidal pituitary gland tumour surgery and spinal fusion surgery, and thereby reducing UTIs, shortening hospital stay and increasing patient comfort. Besides developing and executing de-implementation strategies to accomplish a reduction of used IDUCs, the study focusses on patients' and healthcare professionals' experiences with IDUCs in daily practice and the consequences for the care system. Several challenges are anticipated while executing the study. Since there are multiple study sites, frequent and clear communication between the researchers and the different departments in each hospital is needed. Additionally, due to professionals' busy schedules, planning the group interviews ahead is necessary to ensure a sufficient number of participants. The results from this study can be used to de-implement IDUCs after a broad range of surgeries on several wards.

Abbreviations

DI = Diabetes Insipidus

DURIN-study = De-implementation of urinary catheters in neurosurgical patients

IDUC = Indwelling Urinary Catheter

POUR = Post-Operative Urinary Retention

UTI = Urinary Tract Infection

Declarations

Ethics approval and consent to participate

The Medical Ethics Committee of the university hospital and the four general hospital approved the study (N20. 152).

Consent for publication

Patients operated in 2021 or 2022 will sign informed consent regarding publishing their data. All healthcare professionals will sign informed consent regarding publishing their data.

Availability of data and materials

Data will be available on Zenodo.

Competing interests

The authors have no conflicts of interest to report.

Funding

Funding was received for this study Radboudumc. Specifically from the department IQ Healthcare program "Doen of laten? Het bevorderen van gepaste zorg".

Authors' contributions

All authors contributed to the study conception and design. Material preparation and data collection and analysis will be performed by JMN. Data analysis will be performed by JMN and AHBR. The first draft of this manuscript was written by JMN. AHBR, WvF and WP commented on previous versions of the manuscript. JMN, AHBR, WvF and WP read and approved the final manuscript.

Acknowledgements

Not applicable.

Author's information

Jeanne-Marie Nollen, RN, MSc¹, Anja H. Brunsveld-Reinders, RN, PhD², Wouter R. van Furth, MD, PhD¹, Wilco C. Peul, MD, PhD, MPH, MBa¹

¹Department of neurosurgery, Leiden University Medical Center, the Netherlands, ² Department of quality and patient safety, Leiden University Medical Center, the Netherlands

Email addresses:

J.M.Nollen@lumc.nl

A.H.Brunsveld-Reinders@lumc.nl

W.C.Peul@lumc.nl

W.R.van_Furth@lumc.nl

References

1. Strickland AR, Usmani MF, Camacho JE, Sahai A, Bruckner JJ, Buraimoh K, et al. Evaluation of risk factors for postoperative urinary retention in elective thoracolumbar spinal fusion patients. *Global Spine Journal*. 2021;11(3):338-44.
2. Edate S, Albanese A. Management of electrolyte and fluid disorders after brain surgery for pituitary/suprasellar tumours. *Hormone research in paediatrics*. 2015;83(5):293-301.
3. Gould CV, Umscheid CA, Agarwal RK, Kuntz G, Pegues DA, Control HICPACJI, et al. Guideline for prevention of catheter-associated urinary tract infections 2009. 2010;31(4):319-26.
4. Sadeghi M, Leis JA, Laflamme C, Sparkes D, Ditrani W, Watamaniuk A, et al. Standardisation of perioperative urinary catheter use to reduce postsurgical urinary tract infection: an interrupted time series study. *BMJ Quality & Safety*. 2019;28(1):32.
5. Nicolle LE. Urinary Catheter-Associated Infections. *Infectious Disease Clinics of North America*. 2012;26(1):13-27.
6. Saint S, Trautner BW, Fowler KE, Colozzi J, Ratz D, Lescinskas E, et al. A Multicenter Study of Patient-Reported Infectious and Noninfectious Complications Associated With Indwelling Urethral Catheters. *JAMA Internal Medicine*. 2018;178(8):1078-85.
7. Chardavoyne PC, Kasmire KE. Appropriateness of Antibiotic Prescriptions for Urinary Tract Infections. *West J Emerg Med*. 2020;21(3):633-9.
8. Pujades-Rodriguez M, West RM, Wilcox MH, Sandoe JJE. Lower Urinary Tract Infections: Management, Outcomes and Risk Factors for Antibiotic Re-prescription in Primary Care. 2019;14:23-31.

9. Mitchell BG, Ferguson JK, Anderson M, Sear J, Barnett AJJoHI. Length of stay and mortality associated with healthcare-associated urinary tract infections: a multi-state model. 2016;93(1):92-9.
10. Epstein NE. A review article on the benefits of early mobilization following spinal surgery and other medical/surgical procedures. *Surg Neurol Int.* 2014;5(Suppl 3):S66-S73.
11. Lamas C, del Pozo C, Villabona C. Clinical guidelines for management of diabetes insipidus and syndrome of inappropriate antidiuretic hormone secretion after pituitary surgery. *Endocrinol Nutr.* 2014;61(4):e15-24.
12. Garrahy A, Moran C, Thompson CJ. Diagnosis and management of central diabetes insipidus in adults. *Clinical endocrinology.* 2019;90(1):23-30.
13. Prete A, Corsello SM, Salvatori R. Current best practice in the management of patients after pituitary surgery. *Ther Adv Endocrinol Metab.* 2017;8(3):33-48.
14. Zada G, Woodmansee W, Iuliano S, Laws E. Perioperative Management of Patients Undergoing Transsphenoidal Pituitary Surgery. *Asian journal of neurosurgery.* 2010;5:1-6.
15. Lall M. Nursing care of the patient undergoing lumbar spinal fusion. *Journal of Nursing Education and Practice.* 2017;8:44.
16. Shimanskaya V, Wagenmakers M, Bartels R, Boogaarts J, Grotenhuis J, Hermus A, et al. Toward Shorter Hospitalization After Endoscopic Transsphenoidal Pituitary Surgery: Day-by-Day Analysis of Early Postoperative Complications and Interventions. *World Neurosurgery.* 2018;111.
17. Burgess LC, Wainwright TW, editors. What is the evidence for early mobilisation in elective spine surgery? A narrative review. *Healthcare*; 2019: Multidisciplinary Digital Publishing Institute.
18. Gould C, Umscheid C, Agarwal R. Toolkit for Reducing Catheter-Associated Urinary Tract Infections in Hospital Units: Implementation Guide.
19. Christian E, Harris B, Wrobel B, Zada G. Endoscopic endonasal transsphenoidal surgery: Implementation of an operative and perioperative checklist. *Neurosurgical focus.* 2014;37:E1.
20. Palese A, Buchini S, Deroma L, Barbone F. The effectiveness of the ultrasound bladder scanner in reducing urinary tract infections: a meta-analysis. *Journal of clinical nursing.* 2010;19(21-22):2970-9.
21. Boulis NM, Mian FS, Rodriguez D, Cho E, Hoff JT. Urinary retention following routine neurosurgical spine procedures. *Surgical Neurology.* 2001;55(1):23-7.
22. Baldini G, Bagry H, Aprikian A, Carli F. Postoperative urinary retention: anesthetic and perioperative considerations. *Anesthesiology.* 2009;110(5):1139-57.
23. Hu FW, Yang DC, Huang CC, Chen CH, Chang CM. Inappropriate use of urinary catheters among hospitalized elderly patients: Clinician awareness is key. *Geriatr Gerontol Int.* 2015;15(12):1235-41.
24. Quinn M, Ameling JM, Forman J, Krein SL, Manojlovich M, Fowler KE, et al. Persistent Barriers to Timely Catheter Removal Identified from Clinical Observations and Interviews. *The Joint Commission Journal on Quality and Patient Safety.* 2020;46(2):99-108.
25. Nicolle LE. Catheter associated urinary tract infections. *Antimicrob Resist Infect Control.* 2014;3:23-.

26. Lee YY, Tsay WL, Lou MF, Dai YT. The effectiveness of implementing a bladder ultrasound programme in neurosurgical units. *J Adv Nurs*. 2007;57(2):192-200.
27. Schmiemann G, Kniehl E, Gebhardt K, Matejczyk MM, Hummers-Pradier E. The diagnosis of urinary tract infection: a systematic review. *Dtsch Arztebl Int*. 2010;107(21):361-7.
28. Trautner BW, Cope M, Cevallos ME, Cadle RM, Darouiche RO, Musher DM. Inappropriate treatment of catheter-associated asymptomatic bacteriuria in a tertiary care hospital. *Clinical Infectious Diseases*. 2009;48(9):1182-8.
29. Vasudevan R. Urinary tract infection: an overview of the infection and the associated risk factors. *J Microbiol Exp*. 2014;1(2):00008.
30. Van Pinxteren B, Knottnerus B, Geerlings S, Visser H, Klinkhamer S, Van der Weele G. NHG-Standaard Urineweginfecties (derde herziening). *Huisarts Wet*. 2013;56(6):270-80.
31. Colla CH, Mainor AJ, Hargreaves C, Sequist T, Morden N. Interventions Aimed at Reducing Use of Low-Value Health Services: A Systematic Review. *Medical Care Research and Review*. 2016;74(5):507-50.
32. Nyumba T, Wilson, Derrick C, Mukherjee N. The use of focus group discussion methodology: Insights from two decades of application in conservation. *Methods in Ecology and Evolution*. 2018;9:20-32.
33. Van Houdt S, Sermeus W, Vanhaecht K, De Lepeleire J. Focus groups to explore healthcare professionals' experiences of care coordination: towards a theoretical framework for the study of care coordination. *BMC Fam Pract*. 2014;15:177-.
34. Nili A, Tate M, Johnstone D, Gable G. A Framework for Qualitative Analysis of Focus Group Data in Information Systems 2014.
35. Walker D, Myrick F. Grounded theory: An exploration of process and procedure. *Qualitative health research*. 2006;16(4):547-59.
36. Chapman A, Hadfield M, Chapman C, JotRCoPoE. Qualitative research in healthcare: an introduction to grounded theory using thematic analysis. 2015;45(3):201-5.
37. Mayer I, JJoS, Retailing, Marketing. Qualitative research with a focus on qualitative data analysis. 2015;4(9):53-67.

Figures

Figure 1

Flowchart IDUC placement during surgery

Figure 2

Flowchart IDUC placement in recovery room

Figure 3

Flowchart IDUC placement in neurosurgical ward