

Unravelling the genetic architecture of grain quality traits under optimum and low-nitrogen stress conditions using tropical maize (*Zea mays* L.) germplasm

Noel Ndlovu

National University of Ireland, Galway

Charles Spillane

NUIG: National University of Ireland Galway

Peter C. McKeown

NUIG: National University of Ireland, Galway

Jill E. Cairns

International Maize and Wheat Improvement Centre: Centro Internacional de Mejoramiento de Maiz y Trigo

Biswanath Das

International Maize and Wheat Improvement Centre

Manje Gowda (✉ pgrgowda@gmail.com)

International Maize and Wheat Improvement Centre: Centro Internacional de Mejoramiento de Maiz y Trigo <https://orcid.org/0000-0003-4434-6364>

Research Article

Keywords: Grain, quality traits, nitrogen, tropical maize, GWAS, genomic prediction

Posted Date: November 10th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1021932/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Soils in sub-Saharan Africa are nitrogen deficient due to low fertilizer use and inadequate soil fertility management practices. This has resulted in a significant yield gap for the major staple crop maize, which is undermining nutritional security and livelihood sustainability across the region. Dissecting the genetic basis of grain protein, starch and oil content under nitrogen-starved soils can increase our understanding of the governing genetic systems and improve the efficacy of future breeding schemes. An association mapping panel of 410 inbred lines and four bi-parental populations were evaluated in field trials in Kenya and South Africa under optimum and low nitrogen conditions and genotyped with 259,798 SNP markers. Genetic correlations demonstrated that these populations may be utilized to select higher performing lines under low nitrogen stress. Furthermore, genotypic, environmental and GxE variations in nitrogen-starved soils were found to be significant for oil content. Broad sense heritabilities ranged from moderate (0.18) to high (0.86). Under low nitrogen stress, GWAS identified 42 SNPs linked to grain quality traits. These significant SNPs were associated with 51 putative candidate genes. Linkage mapping identified multiple QTLs for the grain quality traits. Under low nitrogen conditions, average prediction accuracies across the studied genotypes were higher for oil content (0.78) and lower for grain yield (0.08). Our findings indicate that grain quality traits are polygenic and that using genomic selection in maize breeding can improve genetic gain. Furthermore, the identified genomic regions and SNP markers can be utilized for selection to improve maize grain quality traits.

Key Message

Genome-wide association study (GWAS) demonstrated that multiple genomic regions influence grain quality traits under nitrogen-starved soils. Using genomic prediction, genetic gains can be improved through selection for grain quality traits.

Introduction

Maize (*Zea mays* L.) yields in sub-Saharan Africa (SSA) are amongst the lowest in the world (FAO 2021). Average yields in this region range from 1-3 t ha⁻¹, well below the global average of around 5 t ha⁻¹ (Prasanna et al. 2020, 2021). Over a quarter of households in SSA are deemed persistently food insecure, with that figure climbing to 40% during the dry season (Fraval et al. 2019). Furthermore, demand for maize in this region is expected to triple by the year 2050 as a result of rapid population growth (Ekpa et al. 2018). While drought stress and increased climate variability are linked to low yields, low fertilizer use is also a key driver of the maize yield gap in this region (Tittonell and Giller 2013). While average N fertiliser use in SSA by smallholder farmers has increased over the past decade, however it remains very low at 17.9 kg N ha⁻¹ (Jayne and Sanchez 2021). This difference is particularly pronounced on female managed plots which tend to receive less nitrogen inputs than male management plots within a farm (Cairns et al. 2021; Farnworth et al. 2017).

For vulnerable populations, increased dietary diversity and consumption of nutrient-rich food is essential for increased nutrition (Poole et al. 2021). However, in more than 20 countries in SSA, maize accounts for more than 30% of calories consumed (Goredema-Matongera et al. 2021). In Lesotho, Malawi, Zambia, and Zimbabwe, the average per capita consumption of maize is more than 100 kg per person per year (i.e., 300 g per day), roughly half of daily calorie intake (Cairns et al. 2021; Prasanna et al. 2021). High-quality protein sources include eggs, meat, and dairy products, but vulnerable populations in SSA have limited access to these foods and rely heavily on maize as a primary source of protein (Nuss and Tanumihardjo 2011). Average protein supply from maize is 10 g per capita per day in SSA, with up to 35 g per capita per day in southern Africa (FAO 2021). In Burkina Faso, Eswatini, Ethiopia, Lesotho, Malawi, Mozambique, Nigeria, Tanzania, Togo, Zambia and Zimbabwe, maize provides a greater source of protein than protein derived from animal sources (FAO 2021).

Generally, maize grain has low oil (4.4%) and protein (9.1%) contents but a relatively high starch content (73.4%) basing on dry matter measurements (Dei 2017). In temperate maize, breeding has led to significant increase in grain yield. However, grain protein content is estimated to have decreased 0.3% per decade and grain starch content has increased at 0.3% per decade (Duvick 2005). Grain oil content has also reduced over time in temperate maize (Scott et al. 2006). Grain quality is linked closely to soil quality (Wood et al. 2018). Maize protein content increases as the level of N applied increases (Zhang et al. 2020). To date, studies looking at the effect of reduced N level on grain quality have used significantly higher levels of N than is relevant to smallholder farmers in SSA. Under low nitrogen stress in SSA environments, previous maize grain quality assessment studies, such as Abu et al. (2021), Ngaboyisonga and Njoroge (2014), and (Oikeh et al. 1998), used N rates ranging from 30 - 120 kg ha⁻¹. Such application rates are higher than the average N application rates (which is between 12 - 16 kg ha⁻¹) (Heffer and Prud'homme 2015; Sheahan et al. 2014) in smallholder agriculture in SSA. Incorporating grain quality traits into breeding can involve significant costs associated with grain nutrient analyses, morphological characterizations, and associated trait complexities stymie progress in improving maize grain quality. The high-cost requirement of measuring grain quality traits stems from the wet chemistry procedures required to create near-infrared spectroscopy (NIRS) calibration curves, which are relatively expensive. There is a significant opportunity to assess and employ the potential of genomic selection in the improvement of grain quality

traits in maize. This reinforces the need for more rapid progress to improve maize grain quality components (particularly, protein, starch, and oil content) through accelerated and more efficient breeding enabled by molecular markers.

The integration of genomic tools such as genomic-wide association studies (GWAS), linkage mapping and genomic selection (GS) with traditional breeding approaches have increased the efficiency of grain quality and Low N tolerance selection. These techniques have been used to identify causal variants for Low N tolerance. Linkage mapping is a common method for locating quantitative trait loci (QTL) based on a segregating population derived from a cross of two parental lines with significantly divergent phenotypes (Xiao et al. 2017). Linkage mapping can be combined with GWAS to provide a more comprehensive strategy for identifying markers linked with a trait of interest, by the discovery of trait linked markers by GWAS and their validation through linkage mapping. The most robust markers can then be employed for marker-assisted selection (MAS). However, the accuracy with which genetic maps are constructed is dependent on the mapping population since doubled haploid (DH) lines, near-isogenic lines (NILs), recombinant inbred lines (RILs), and backcross lines are extremely effective yet to date labour-intensive and time-consuming to generate (Rao et al. 2021). GS has been highlighted as a powerful option for selecting polygenic traits that are difficult to select through MAS. GS can accomplish this by employing genome-wide dense markers for predictions, and therefore can support association analyses to determine the genetic basis of key traits (Bentley et al. 2014). Galli et al. (2020) reported that GWAS could identify both additive and dominant genetic effects that influence Low N tolerance in maize.

To understand how low N stress affects key grain quality traits such as grain protein, starch, and oil content, this study was performed using a tropical maize population under low N and optimum conditions across multi-location field trials in Kenya and South Africa. The study had the objectives of (i) assessing the genetic architecture of low soil-N tolerant maize test crosses using their responses to grain quality and yield traits under two management conditions (optimum and low soil N); (ii) identifying the significant quantitative trait nucleotides (QTNs) and putative candidate genes and QTLs for quality traits in tropical maize germplasm tested in multiple locations; and (iii) assessing the potential of utilizing GS in the improvement of grain quality traits.

Materials And Methods

Germplasm, experiment design and management

An association mapping panel of 410 tropical maize lines developed under CIMMYT's Improved Maize for African Soils (IMAS) project (Ertiro et al. 2020a) in collaboration with the Kenya Agricultural and Livestock Research Organisation (KALRO) and Agricultural Research Council (ARC, South Africa) was used. The study also evaluated two DH populations from CIMMYT's Heterotic group B (221 lines of CML550/CML504, 115 lines of CML550/CML511), and two DH population from CIMMYT's Heterotic group A (175 lines of CML505/LaPostaSeqC7-F64-2-6-2-2 and 131 lines of CML536/LaPostaSeqC7-F64-2-6-2-2) (Ertiro et al. 2020b). Test cross hybrids were generated by crossing all inbred lines with a broadly adapted CIMMYT maize inbred line tester from the opposite heterotic group.

The field trials were designed and conducted by Das et al. (2019) and Ertiro et al. (2020a) who previously studied and published on different objectives using data from these same trials. Three optimal and six low N-stressed sites were used to screen the testcross progenies (Table 1). Experiments conducted in the same site over several years were classified as separate environments. The study employed an alpha-lattice design with two replications. The sites for low N trials were depleted for soil N content by growing sorghum for several years without applying any external N fertilizer. At planting, triple phosphate (46% P_2O_5) was applied to the low N trials at the rate of 50 kg P_2O_5 /ha. On optimum trials, diammonium phosphate (DAP) fertilizer was used at the rate of 54 kg N per hectare. Optimum trials were top-dressed with urea fertilizer at the rate of 138 kg N per hectare three weeks after planting. All trials under both optimum and low-N were irrigated as required to avoid any moisture stress. Trials under both conditions were kept weed-free and other standard agronomic practices were conducted.

Measurements of grain yield and quality traits

Data were recorded for grain yield and quality traits (i.e., protein, oil, and starch contents). Shelled grain yield was measured in kilograms (kg) and converted to tons per hectare and reported at 12.5% moisture. Protein, starch, and oil content were measured using a FOSS Infratec TM 1241 from 500-gram samples of grain taken from each plot and are reported as a percentage of whole grain. The FOSS Infratec is a non-destructive whole-grain analyzer that uses near-infrared reflectance (NIR) to estimate quality parameters. Five 100-gram subsamples were assayed and the mean reading for each parameter was reported per plot.

Table 1
Description of the field trials used in the study.

Country	Site name	Management options	Germplasm used	Season
Kenya	Embu	Low N	IMAS	2011B, 2012A
		Low N	CML550/CML504	2014A
		Low N	CML550/CML511	2014A
	Kiboko	Optimum	IMAS	2011A, 2012A
		Low N	IMAS	2011A, 2011B, 2012A
		Optimum	CML505/LaPostaSeqC7-F64-2-6-2-2	2015B
		Low N	CML505/LaPostaSeqC7-F64-2-6-2-2	2015A, 2015B
		Optimum	CML550/CML511	2014A
		Low N	CML550/CML504	2015A, 2015B
		Low N	CML550/CML511	2015A
South Africa	Cedara	Optimum	IMAS	2011B
		Low N	IMAS	2011B

***A = main season; B =off-season**

Phenotypic data analysis

The restricted maximum likelihood (REML) approach was used to conduct analyses of variance (ANOVA) using the META-R program for each and across environments under optimum and Low-N conditions (Alvarado et al. 2020). The linear mixed model was used to calculate all variance components. The study treated replication as fixed effect and all other treatment effects as random. On an entry-mean basis, the broad-sense heritability was estimated using the genotypic to phenotypic variance ratio from the derived variance components. Furthermore, to determine the genotypic effects of the investigated lines for each and across environments, best linear unbiased estimation (BLUE) and best linear unbiased prediction (BLUP) were obtained. For GWAS and linkage mapping, BLUPs were used. On the other hand, BLUEs were used for GS analyses. The classification of the genotypic correlation coefficients followed the guidelines provided by Profillidis and Botzoris (2019). To determine the impact of low N stress on the aforementioned traits, we used a t-test to compare the mean values of the two management conditions. We also determined the percentage change (i.e., decrease or increase) in grain quality trait performance.

Genotyping-by-sequencing (GBS)

DNA was extracted according to the CIMMYT high-throughput mini-prep Cetyl Trimethyl Ammonium Bromide (CTAB) method (Semagn (2014). Following the protocol presented in Elshire et al. (2011), maize DNA samples were genotyped using a restriction enzyme (*ApeKI*) and 96-plex multiplexing at the Institute of Biotechnology at Cornell University, USA. The Institute of Genomic Diversity (IGD) at Cornell provided raw GBS data for a maximum of 955,120 SNP loci spread throughout the 10 maize chromosomes (Ertiro et al. 2020a). Raw data was filtered for linkage mapping according to the criteria used in Ertiro et al. (2020a) of >10 percent minor allele frequency (MAF) and no missing data. Furthermore, the genotype data were filtered for GWAS using the Trait Analysis by Association, Evolution, and Linkage (TASSEL v.5.2.7.2) software, with a baseline count of SNPs on 90% and a MAF of >5% of the sample size as presented in Ertiro et al. (2020a). Principal Component Analysis (PCA) was carried out in TASSEL (v.5.2.7.3), as were genetic distances and kinship.

Genome-wide association study analysis

The R package 'FarmCPU-Fixed and random model Circulating Probability Unification' (Liu et al. 2016) was used for GWAS analysis. FarmCPU utilized the first three PCs derived by TASSEL as input for GWAS. The kinship was computed using FarmCPU's default kinship algorithm as presented in Ertiro et al. (2020a). The Manhattan and quantile-quantile (QQ) plots, GWAS findings, and a table of marker effects of user-provided variables were all produced by the FarmCPU using the "GAPIT" function. The putative genes were checked on the: (i) maize gdb website (<http://www.maizegdb.org/>) and (ii) ensemble website

(<http://plants.ensembl.org/biomart/martview/ce35c2dc12e78418361fb4cffa43bdbe>) using the BLAST tool against the representative genome B73 version 2.

QTL mapping and genomic prediction

The four DH populations were genotyped with GBS and data was further filtered to a manageable size using TASSEL software with >0.10 MAF, $<5\%$ heterozygosity, and 90% the minimum count of the total size (Bradbury et al., 2007; Sironik et al., 2019). In all the populations, homozygous marker loci for both parents and uniformly distributed polymorphic markers between parents were retained. Linkage maps were constructed by using QTL IciMapping version 4.1 in all four DH populations. BIN, is an inbuilt tool implemented in QTL IciMapping was used to remove the highly correlated SNPs. This resulted into retain 2,699, 1,962, 1,985 and 2,086 high-quality SNPs in CML550/CML504, CML550/CML511, CML505/LaPostaSeqC7-F64-2-6-2-2 and CML536/LaPostaSeqC7-F64-2-6-2-2, respectively. These SNPs were used to construct linkage maps using the MAP function. IciMapping used the grouping, ordering, and rippling steps to construct a linkage map. The Kosambi genetic distance mapping function which assumes that recombination events influence the occurrence of adjacent recombination was used. BLUP values across environments for the DH populations were used in QTL detection analysis (Meng et al., 2015). The mapping populations were grouped by the SNPs and the significant difference between the means (P -value <0.0001) was detected based on the markers that were linked to a QTL controlling the selected target trait (Collard et al., 2005). The highest peak of one LOD that supports the confidence interval was used to declare the significance of the QTL map position on both sides of the QTL (Hackett, 2002). The phenotypic variation explained (PVE) by each QTL and together for all QTLs for each trait was estimated. The origin of the favourable allele for each trait was identified based on the sign of the additive effects of each QTL.

BLUEs across environments for each trait in each population were used in the GS analysis. The Ridge-regression BLUP (RR-BLUP, Zhao et al. 2012) with fivefold cross-validation for each trait was used for the analysis. A sample of 4,000 SNPs with all data values, equally distributed throughout the genome, and MAF >0.05 was chosen from the GBS data for the IMAS panel and all four DH populations. Individual DH population and the IMAS set were sampled to form a training and prediction set. The prediction accuracy was calculated as the correlation between the observed phenotypes and genomic estimated breeding values (GEBVs) divided by the square root of heritability (Dekkers 2007). In each population, 100 iterations were done for the sampling of the training and validation sets.

Results

Effect of low N stress on grain yield and quality traits

There was significant variation in protein, starch and oil content, and grain yield within all four biparental DH populations and the IMAS panel under optimum and low N stress conditions (Figure 1 and Table 2). In the IMAS panel and CML505/LaPostaSeqC7-F64-2-6-2-2 DH pop, yield under low N stress was reduced by 59% and 48%, respectively. In DH pop CML550/CML511, the mean yield under low N stress was 5.45 t ha^{-1} , however, this was a reduction of 47% relative to optimal conditions. Low N stress significantly ($p < 0.01$) reduced protein and oil content (except in DH pop CML505/LaPostaSeqC7-F64-2-6-2-2) but had no significant effect on starch content. Although the level of N stress and therefore the reduction in grain yield was the lowest in DH pop CML550/CML511, both protein and oil content had the largest reduction in this population under low N stress.

The genotypic, environmental, and genotype-environment interaction effects (G, E and G x E, respectively) were significant at $p \leq 0.05$ for yield and quality traits (Table 2). For protein, starch, and oil content under optimal conditions, the magnitude of genotypic variance was greater compared to low N stress conditions. Under low N conditions, the effect of genotype, environment, and G x E interactions on oil content was significant across all genotypes tested. Interestingly, under the same conditions, the genotypic effects on protein content were only significant in DH pops CML505/LaPostaSeqC7-F64-2-6-2-2 and CML536/LaPostaSeqC7-F64-2-6-2-2. The G x E effects on protein content and starch content in DH pop CML550/CML511 were significant. The zero estimates of G x E interactions for starch and protein content (observed on DH pop CML550/CML511 under low N stress) indicate that genotypic performance for these traits was stable across the tested environments. H^2 values of each trait under both optimal and low N stress are presented in Table 2. In general, H^2 of all traits was lower under low N stress than optimal, with the exception of starch content which increased under low N stress across all populations.

Grain yield was negatively correlated with protein content across populations, regardless of N stress level (Table 3). Similarly, starch content showed a negative correlation with protein and oil content across genotypes and management options. A weak positive correlation was reported between protein and oil content across populations and N levels. The only exceptions were in DH pop CML550/CML504 under optimum conditions where oil content was significantly (at $p < 0.01$) and negatively correlated to protein content ($r = -0.92^{**}$).

Table 2
Genetic parameters for IMAS panel and four DH Populations evaluated under optimum and low N stress conditions in multiple environments.

	Optimum					Low N				
	Mean	σ^2_G	σ^2_{GE}	σ^2_e	h^2	Mean	σ^2_G	σ^2_{GE}	σ^2_e	h^2
IMAS Panel										
Grain yield (t/ha)	8.47	0.59**	0.38**	1.37	0.62	3.49	0.08**	0.13**	0.67	0.50
Protein content (%)	10.53	0.15**	0.02**	0.33	0.71	8.07	0.06**	0.08**	0.33	0.59
Starch content (%)	69.17	0.10**	0.02**	0.28	0.55	70.95	0.11**	0.07**	0.53	0.62
Oil content (%)	5.24	0.05**	0.01**	0.03	0.86	5.01	0.04**	0.03**	0.06	0.72
DH pop1 (CML550/CML504)										
Grain yield (t/ha)	-	-	-	-	-	4.12	0.17**	0.00	0.98	0.41
Protein content (%)	-	-	-	-	-	8.10	0.06**	0.03**	0.19	0.47
Starch content (%)	-	-	-	-	-	70.97	0.08**	0.00	0.51	0.38
Oil content (%)	-	-	-	-	-	5.36	0.03**	0.00	0.03	0.79
DH pop2 (CML550/CML511)										
Grain yield (t/ha)	10.40	0.77**	-	2.10	0.42	5.45	0.27**	0.11**	0.77	0.52
Protein content (%)	10.96	0.12**	-	0.10	0.72	7.98	0.12**	0.00	0.23	0.67
Starch content (%)	68.12	0.08**	-	0.15	0.52	69.65	0.12**	0.00	0.30	0.62
Oil content (%)	6.24	0.03**	-	0.02	0.79	5.81	0.01*	0.04**	0.05	0.20
DH pop3 (CML505/LaPostaSeqC7-F64-2-6-2-2)										
Grain yield (t/ha)	5.08	0.07*	-	0.17	0.45	2.70	0.03*	0.09**	0.30	0.21
Protein content (%)	8.21	0.17**	-	0.21	0.61	7.52	0.05*	0.07**	0.26	0.33
Starch content (%)	70.96	0.07*	-	0.61	0.18	71.68	0.09**	0.07**	0.51	0.34
Oil content (%)	5.05	0.02*	-	0.04	0.48	5.25	0.03*	0.00	0.05	0.66
DH pop4 (CML536/LaPostaSeqC7-F64-2-6-2-2)										
Grain yield (t/ha)	-	-	-	-	-	3.78	0.21**	0.12**	0.81	0.54
Protein content (%)	-	-	-	-	-	7.61	0.05*	0.06**	0.17	0.48
Starch content (%)	-	-	-	-	-	71.58	0.02*	0.09**	0.40	0.18
Oil content (%)	-	-	-	-	-	5.06	0.02*	0.01*	0.04	0.66
*, ** Significant at $p < 0.05$ and $p < 0.01$ level, respectively										

Protein content had a negative correlation with grain yield ($r = -0.41^{**}$) and starch content ($r = -0.54^{**}$) in the IMAS panel under optimum conditions. Similarly, a weak positive correlation was observed in the IMAS panel, DH pops CML550/CML504 and CML505/LaPostaSeqC7-F64-2-6-2-2 between protein content and oil content under low N stress. Starch and oil contents were shown to be significantly ($p < 0.05$) and negatively correlated under optimum ($r = -0.65$) and low N stress ($r = -0.18$) in the IMAS panel. The genotypic correlation coefficients for DH pops CML505/LaPostaSeqC7-F64-2-6-2-2 and CML536/LaPostaSeqC7-F64-2-6-2-2 showed that oil content under optimum conditions had no correlation with protein content ($r = 0.00$). As demonstrated in Table 3, further significant ($p < 0.05$ and $p < 0.01$) trait correlations were established among the phenotypic parameters measured across the management conditions for each set of genotypes. The observed correlations between grain quality and yield can be useful for selection decisions or trade-offs in genotypic selection.

Table 3

Genotypic correlations between grain yield and grain quality traits evaluated under optimum and low N stress management. Values in shaded cells represent genetic correlations between traits under low N stress, values in unshaded cells represent genetic correlations between traits under optimum conditions.

	Grain yield	Protein content	Starch content	Oil content
IMAS Panel				
Grain yield	-	-0.03	0.28*	0.01
Protein content	-0.41**	-	-0.30**	0.10*
Starch content	0.01	-0.54**	-	-0.18**
Oil content	-0.17*	0.07	-0.65**	-
DH pop1 (CML550/CML504)				
Grain yield	-			
Protein content	-0.10*	-		
Starch content	-0.39**	-0.37**	-	
Oil content	0.21**	0.03	-0.84**	-
DH pop2 (CML550/CML511)				
Grain yield	-	-0.21*	0.17*	0.18*
Protein content	-0.24**	-	-0.78**	0.09
Starch content	-0.12	-0.61**	-	-0.72**
Oil content	0.25**	-0.92**	-0.95**	-
DH pop3 (CML505/LaPostaSeqC7-F64-2-6-2-2)				
Grain yield	-	-0.43**	0.52**	-0.36**
Protein content	-0.24**	-	-0.53**	0.46**
Starch content	0.30**	-0.23*	-	-0.97**
Oil content	0.02	0.00	-0.84**	-
DH pop4 (CML536/LaPostaSeqC7-F64-2-6-2-2)				
Grain yield	-			
Protein content	-0.29**	-		
Starch content	0.26**	-0.13*	-	
Oil content	-0.20*	0.00	-0.95**	-
*, ** Significant at p<0.05 and p<0.01 level, respectively.				

GWAS analyses

Of the 337,113 GBS SNPs used to genotype 410 genotypes, 77.1% (259,798) remained after filtering using the >5% MAF and 10% missing per marker criteria (Supplementary Figure S1). The kinship relations among the IMAS panel were determined using the filtered 259,798 SNP markers and depicted as a genetic cluster, indicating that the panel of genotypes are split into four potential genetically differentiated subgroups. The heatmap of the panel's kinships was used to predict the magnitude of the existing relationships in the genotypes: this established that the genotypes were not closely related and that there is no strong population structure (Supplementary Figure S2). Further partition of the population structure of the IMAS panel using STRUCTURE 2.3.4 is presented in an earlier study by Kibe et al. (2020a) and Gowda et al. (2015). PCA was carried out using 259,798 high-quality SNPs (Supplementary Figure S3). The first principal component (PC1) accounted for roughly 4.5% of the overall variation, whereas the second principal component (PC2) explained 2.5% (Supplementary Figure S3). Calculation of genome-wide LD using 259,798 SNPs showed a significant decline in LD as genetic distance rose, with different rates of

attenuation for each of the ten chromosomes (Supplementary Figure S4). Figures 2 and 3 depict the GWAS findings for protein, starch, and oil content, and yield across the two N managements as Manhattan and Q-Q plots of p -values evaluating the anticipated and observed $-\log_{10} p$ -values. Sixty-one SNPs were significantly ($P=2 \times 10^{-5}$, $p=0.1$ False Discovery Rate (FDR)) associated with the protein, starch, and oil content, and yield under optimal conditions and were spread across 10 chromosomes (Table 4). Under low N conditions, 42 SNPs are linked to the aforementioned traits.

Table 4

Chromosomal positions and SNPs significantly associated with grain yield, protein, starch, and oil content detected by SNP-based GWAS in the IMAS association mapping panel under optimum and low N management conditions.

SNP	Chr	Position	MLM-P value	MAF	Allele effect	Putative candidate genes	Predicted function of candidate gene
Grain yield under optimum N							
S6_114082980	6	114082980	1.51E-08	0.24	-0.08	GRMZM2G165969	Serine-type endopeptidase activity
S9_20607559	9	20607559	7.62E-08	0.23	-0.08	GRMZM2G391473	Protein serine/threonine kinase activity
S10_138021889	10	138021889	1.42E-07	0.07	0.11	GRMZM2G061287	Regulation of DNA-templated transcription
S3_2218349	3	2218349	2.01E-07	0.09	0.10	GRMZM2G159307	ATP binding and magnesium chelatase activity
S4_240579103	4	240579103	2.41E-07	0.23	0.07	GRMZM2G491366	Unknown
S5_200110933	5	200110933	4.55E-07	0.33	-0.06	GRMZM2G144898	Sexual reproduction
S5_171518940	5	171518940	6.87E-07	0.24	0.06	GRMZM2G043250	Zinc ion binding
S1_279913054	1	279913054	1.46E-06	0.06	0.12	GRMZM2G300990	Protein serine/threonine kinase activity
S7_115546635	7	115546635	1.98E-06	0.39	-0.05	AC189790.2_FG003	Unknown
S3_220130640	3	220130640	4.03E-06	0.20	0.06	GRMZM2G100911	Protein serine/threonine kinase activity
S8_164576912	8	164576912	8.04E-06	0.19	-0.06	GRMZM2G121440	Shoot apex development
S10_132510876	10	132510876	1.39E-05	0.39	-0.05	AC194695.3_FG003	Unknown
Grain yield under low N							
S7_124403566	7	124403566	1.04E-05	0.34	0.14	GRMZM2G007249	1-aminocyclopropane-1-carboxylate oxidase activity
S7_174752436	7	174752436	1.05E-05	0.08	-0.24	GRMZM2G108166	DNA binding and helicase activity
S2_30999187	2	30999187	1.85E-05	0.21	-0.15	GRMZM2G052078	Shoot apex and pollen development
S7_170251419	7	170251419	2.39E-05	0.23	0.14	GRMZM2G024484	NADH dehydrogenase (ubiquinone) activity
S5_11883140	5	11883140	2.72E-05	0.35	0.13	GRMZM2G090435	Homiothermy
Protein content under optimum N							
S1_17679954	1	17679954	9.57E-06	0.14	-0.13	GRMZM2G410623	Shoot apex and pollen development
S1_214242607	1	214242607	1.90E-05	0.14	-0.15	GRMZM2G127429	Aconitate hydratase activity
S10_114836465	10	114836465	2.69E-05	0.28	0.10	GRMZM2G049681	Shoot apex and pollen development
Protein content under low N							

*MAF - Minor allele frequency; MAE – minor allele effect; ^a The exact physical position of the SNP can be inferred from marker's name, for example, S1_82702920: chromosome 1; 82,702,920 bp (Ref Gen_v2 of B73)

SNP	Chr	Position	MLM-P value	MAF	Allele effect	Putative candidate genes	Predicted function of candidate gene
S4_120988951	4	120988951	2.36E-05	0.27	0.11	GRMZM2G032314	Shoot apex and pollen development
S3_198394847	3	198394847	3.47E-05	0.17	0.13	GRMZM2G037102	Unknown
Starch content under optimum N							
S2_174345463	2	174345463	6.92E-06	0.21	-0.16	GRMZM2G033694	Histone-lysine N-methyltransferase activity
S2_174345465	2	174345465	6.92E-06	0.21	-0.16	GRMZM2G033694	Histone-lysine N-methyltransferase activity
S6_5158703	6	5158703	6.97E-06	0.30	-0.15	GRMZM2G125138	Shoot apex and pollen development
S8_3430590	8	3430590	1.06E-05	0.13	-0.18	GRMZM2G013461	Cysteine-type endopeptidase inhibitor activity
S3_180044790	3	180044790	1.13E-05	0.15	-0.17	GRMZM2G493214	Unknown
S5_10542862	5	10542862	1.25E-05	0.18	0.15	GRMZM2G114789	Nucleic acid-binding
S6_60978968	6	60978968	1.32E-05	0.09	-0.22	GRMZM2G171810	Unknown
S7_14465153	7	14465153	2.14E-05	0.21	0.17	GRMZM2G104325	ATP biosynthesis process
Starch content under low N							
S5_10542862	5	10542862	1.15E-05	0.18	0.17	GRMZM2G114789	Nucleic acid-binding
S6_5158703	6	5158703	1.56E-05	0.30	-0.16	GRMZM2G125138	Shoot apex and pollen development
S2_174345463	2	174345463	1.99E-05	0.21	-0.17	GRMZM2G033694	Histone-lysine N-methyltransferase activity
S2_174345465	2	174345465	1.99E-05	0.21	-0.17	GRMZM2G033694	Histone-lysine N-methyltransferase activity
S6_60978968	6	60978968	2.02E-05	0.09	-0.24	GRMZM2G171810	Unknown
Oil content under optimum N							
S6_150886775	6	150886775	8.06E-14	0.11	0.12	GRMZM2G077789	DNA binding and pollen development
S6_60978968	6	60978968	6.93E-12	0.09	0.11	GRMZM2G171810	Unknown
S6_117596170	6	117596170	5.58E-10	0.30	0.06	GRMZM2G505444	Unknown
S2_148879075	2	148879075	2.63E-09	0.14	-0.08	AC211891.4_FG001	Zinc ion binding
S5_50087909	5	50087909	1.09E-08	0.22	0.06	GRMZM2G132464	Hydrolase activity, hydrolyzing O-glycosyl compounds
S5_61427444	5	61427444	1.65E-08	0.17	-0.07	GRMZM2G417435	Shoot apex and pollen development
S3_169317292	3	169317292	3.50E-08	0.19	-0.07	GRMZM2G034423	Unknown
*MAF - Minor allele frequency; MAE – minor allele effect; ^a The exact physical position of the SNP can be inferred from marker's name, for example, S1_82702920: chromosome 1; 82,702,920 bp (Ref Gen_v2 of B73)							

SNP	Chr	Position	MLM-P value	MAF	Allele effect	Putative candidate genes	Predicted function of candidate gene
S1_191845162	1	191845162	9.38E-08	0.15	-0.07	GRMZM2G026696	Unknown
S2_20538422	2	20538422	5.52E-07	0.11	0.07	GRMZM2G702954	Unknown
S6_36950637	6	36950637	5.58E-07	0.09	-0.08	GRMZM2G571740	Unknown
S1_209355590	1	209355590	2.08E-06	0.41	0.04	GRMZM2G059138	Pollen and leaf development
S9_1260192	9	1260192	3.91E-06	0.50	-0.04	GRMZM2G500782	Unknown
Oil content under low N							
S6_60978968	6	60978968	2.15E-12	0.09	0.12	GRMZM2G171810	Unknown
S5_20201057	5	20201057	9.72E-12	0.43	0.06	GRMZM2G070523	DNA binding and shoot apex development
S8_19834231	8	19834231	5.96E-09	0.15	0.07	AC218957.3_FG003	Unknown
S2_214956407	2	214956407	2.83E-08	0.24	-0.07	GRMZM2G153877	Shoot apex and pollen development
S3_221547612	3	221547612	1.03E-07	0.45	-0.05	GRMZM2G028037	Protein serine/threonine kinase activity
S2_196110550	2	196110550	1.82E-07	0.37	-0.05	GRMZM2G113618	Monoxygenase and indoleacetaldoxime dehydratase activity
S9_2472485	9	2472485	6.15E-07	0.30	-0.05	GRMZM5G833563	DNA polymerase and nucleotidylexotransferase activity
S6_150841558	6	150841558	8.01E-07	0.11	0.07	GRMZM2G378586	Shoot apex and pollen development
S10_139315324	10	139315324	1.32E-06	0.19	0.05	GRMZM2G080516	DNA-binding transcription factor activity
S5_214518378	5	214518378	1.60E-06	0.48	-0.04	GRMZM2G008834	Protein binding
S2_20859604	2	20859604	9.48E-06	0.20	0.05	GRMZM2G014400	Nucleic acid-binding and shoot apex development
S1_269023923	1	269023923	2.31E-05	0.10	-0.06	GRMZM2G155767	Phosphorelay sensor kinase activity
*MAF - Minor allele frequency; MAE – minor allele effect; ^a The exact physical position of the SNP can be inferred from marker's name, for example, S1_82702920: chromosome 1; 82,702,920 bp (Ref Gen_v2 of B73)							

Under optimal conditions, three SNPs linked with protein content were significant on chromosomes 1 (*S1_17679954* and *S1_214242607*) and 10 (*S10_114836465*). Under low N stress, however, two different SNPs *S3_198394847* and *S4_120988951* were associated with protein content. Starch content (low N) was associated with five SNPs, the most significant of which was *S5_10542862*. Eight SNPs on chromosomes 2 (*S2_174345463* and *S2_174345465*), 3 (*S3_180044790*), 5 (*S5_10542862*), 6 (*S6_5158703* and *S6_60978968*), 7 (*S8_3430590*), and 8 (*S7_14465153*) were linked with the starch content under optimum conditions. Under optimal conditions, twelve SNPs were significantly associated with oil content, with one-third of these loci located on chromosome 6. Twelve SNPs, with loci on all chromosomes except 4 and 7, were significantly linked with oil content under low N. For starch and oil content at low N conditions, only the significant SNP on chromosome 6 (*S6_60978968*) was co-detected. The proportion of detected SNPs for the other traits are presented in Table 4. The Q-Q plot for grain yield and oil content under optimum conditions and oil content under low N stress revealed that some observed *P*-values were more significant as the marker points migrated from the dotted red line towards the y-axis.

To elucidate the molecular and physiological mechanisms controlling grain quality traits under optimum and low N conditions, candidate genes were identified (Harper et al. 2016). On all chromosomes, a total of 51 candidate genes were discovered (Table 4). The lowest number of candidates (2) and the highest number (12) were related to protein content under low N and oil content under both optimum and low N, respectively. From these candidates, 80.39% (41 genes) were functionally annotated, whereas 19.61% (10 genes) were classified as unknown proteins. The study revealed four candidate genes with protein serine/threonine kinase activity that play a role in soil N response. Under optimum conditions, GRMZM2G159307 and GRMZM2G104325 were encoded as ATP binding proteins for grain yield and starch content, respectively. GRMZM2G10816 (yield), GRMZM2G070523 and GRMZM2G080516 (oil content) were associated with DNA biosynthesis under low N stress conditions. Under both optimal and low N circumstances, GRMZM2G033694 was annotated in the Histone-lysine N-methyltransferase family. Genes coding for shoot apex development were discovered to be associated with grain yield, protein, starch, and oil content under low N stress.

QTLs associated with grain yield and quality traits

The four populations used in this study for linkage mapping were also used in our earlier study (Ertiro et al., 2020a) which includes detailed information about genetic maps. Table 5 shows the detected QTLs and their positions and genetic effects. In DH pop CML550/CML504, two QTLs each were detected for grain yield and starch content, three QTL for protein content and five QTL for oil content under low N stress. The PVE by these QTL was varied from 4.67 to 22.19% and together the total PVE was varied from 12.5% for grain yield to 47.9% for oil content. In DH pop CML550/CML511, one QTL each were detected for grain yield, starch content and oil content under low N stress. In DH pop CML505x LaPostaSeqC7-F64-2-6-2-2, five QTL were detected for grain yield with one QTL on chromosome 3 having a major effect with 12.17% of PVE. For protein content nine QTL were detected with all individually having minor effects except a QTL on chromosome 3 with 11.78% of variance explained. For starch content, three QTL each were detected under optimum and low N conditions with two major effects QTL on chromosome 8. For oil content three QTL were detected under optimum and six QTL were identified under low N conditions with one common QTL on chromosome 2 across management conditions. Four major effect QTL were identified for oil content on chromosomes 2, 4 and 5 which explained >10% of the phenotypic variation (Table 5). In DH pop CML536xLapostaSeqiaF64, one QTL each were detected for protein and oil content and three QTL were detected for starch content, with one major effect QTL at chromosome 4 which contributes 20.3% of phenotypic variation for oil content.

Table 5

Number of QTL detected for grain yield, grain protein, starch and oil content under optimum (opt) and low-nitrogen (LN) stress across environments in four populations.

Trait Name	QTL Name	Chr	Position (cM)	Left Marker	Right Marker	LOD	PVE(%)	TPVE (%)	Add	Fav allele
DH pop1 (CML550xCML504)										
Grain yield_LN	<i>qGY7_10</i>	7	629	S7_7419076	S7_10288357	3.31	6.60	12.51	0.22	CML550
	<i>qGY8_15</i>	8	578	S8_14728619	S8_15994213	3.54	6.85		0.22	CML550
Protein content LN	<i>qPC1_115</i>	1	525	S1_112639745	S1_162463938	6.60	11.04	30.51	0.13	CML550
	<i>qPC4_185</i>	4	115	S4_184892495	S4_185566354	3.38	5.39		0.09	CML550
	<i>qPC8_10</i>	8	128	S8_7675588	S8_12273955	4.71	8.30		0.11	CML550
Starch content LN	<i>qSC1_24</i>	1	47	S1_24583363	S1_24676752	3.44	6.94	19.24	-0.13	CML504
	<i>qSC4_174</i>	4	95	S4_173800701	S4_174954893	4.56	9.06		-0.15	CML504
Oil content LN	<i>qOC1_268</i>	1	313	S1_268152564	S1_268431295	3.28	3.49	47.92	-0.04	CML504
	<i>qOC3_03</i>	3	294	S3_3559699	S3_3912927	7.98	8.84		-0.07	CML504
	<i>qOC6_60</i>	6	35	S6_56452467	S6_63537451	4.30	4.67		-0.05	CML504
	<i>qOC6_133</i>	6	103	S6_132237821	S6_133515728	17.69	22.19		-0.11	CML504
	<i>qOC8_165</i>	8	503	S8_164378522	S8_166561727	5.97	6.95		-0.06	CML504
DH pop2 (CML550xCML511)										
Grain yield_LN	<i>qGY5_15</i>	5	610	S5_4278140	S5_202395053	2.75	8.79	8.62	-0.74	CML511
Starch content LN	<i>qSC1_180</i>	1	288	S1_175655223	S1_187977553	2.75	11.52	11.45	0.22	CML550
Oil content LN	<i>qOC3_60</i>	3	206	S3_58480638	S3_156799630	2.65	10.47	9.78	-0.09	CML511
DH pop3 (CML505xLapostaSequiaF64)										
Grain yield Opt	<i>qGY3_187</i>	3	207	S3_186485761	S3_189293786	3.40	9.47	8.86	-0.037	LPSF64
Grain yield LN	<i>qGY3_196</i>	3	175	S3_195927184	S3_202609108	9.91	12.17	26.28	-0.093	LPSF64
	<i>qGY3_207</i>	3	954	S3_206952639	S3_208331031	3.92	4.41		0.063	CML505
	<i>qGY4_160</i>	4	28	S4_159978522	S4_160277631	3.99	4.59		-0.063	LPSF64
	<i>qGY4_155</i>	4	524	S4_154382211	S4_155910715	6.79	8.06		0.076	CML505
Protein content_Opt	<i>qPC4_140</i>	4	162	S4_125492338	S4_141413020	3.04	8.46	14.15	-0.002	LPSF64
	<i>qPC4_180</i>	4	577	S4_179648189	S4_180236732	10.68	4.23		-0.003	LPSF64
	<i>qPC5_70</i>	5	291	S5_69971990	S5_73044866	4.16	8.46		-0.002	LPSF64
	<i>qPC6_157</i>	6	57	S6_156035854	S6_157412461	3.95	4.23		-0.002	LPSF64
	<i>qPC6_40</i>	6	652	S6_36947497	S6_45922975	3.04	4.23		-0.002	LPSF64
	<i>qPC7_126</i>	7	166	S7_125835683	S7_127144774	4.59	8.46		0.003	CML505
	<i>qPC10_142</i>	10	102	S10_141963062	S10_142398096	3.92	8.46		0.002	CML505
Protein content_LN	<i>qPC3_187</i>	3	208	S3_180516308	S3_189293786	5.27	11.78	24.15	0.076	CML505

* Chr = Chromosome, LOD = Logarithm of Odds; add = additive effect; PVE = phenotypic variance explained; fav allele = parental line contributing the favorable allele for trait, QTL name composed by the trait code followed by the chromosome number in which the QTL was mapped and a physical position of the QTL

Trait Name	QTL Name	Chr	Position (cM)	Left Marker	Right Marker	LOD	PVE(%)	TPVE (%)	Add	Fav allele
	<i>qPC4_146</i>	4	193	S4_145319458	S4_147746777	3.63	7.73		-0.062	LPSF64
Starch content_Opt	<i>qSC2_14</i>	2	47	S2_13290784	S2_18626409	3.93	8.63	23.99	0.056	CML505
	<i>qSC4_205</i>	4	677	S4_204860243	S4_206663242	3.42	3.70		0.049	CML505
	<i>qSC5_14</i>	5	380	S5_13609325	S5_14471659	3.77	8.63		0.051	CML505
Starch content_LN	<i>qSC8_123</i>	8	294	S8_122410298	S8_123512730	11.35	7.00	19.95	-0.313	LPSF64
	<i>qSC8_124</i>	8	298	S8_123512730	S8_124357554	18.85	12.86		0.427	CML505
	<i>qSC9_24</i>	9	224	S9_23280315	S9_30187968	3.32	1.73		0.176	CML505
Oil content_Opt	<i>qOC2_186</i>	2	480	S2_185692862	S2_186464021	4.89	10.53	35.22	0.030	CML505
	<i>qOC4_70</i>	4	506	S4_68323590	S4_75504829	6.01	12.56		-0.032	LPSF64
	<i>qOC5_182</i>	5	158	S5_182450446	S5_181213050	4.35	8.94		-0.028	LPSF64
Oil content_LN	<i>qOC2_186</i>	2	482	S2_185692862	S2_186464021	6.12	7.87	54.86	0.042	CML505
	<i>qOC3_223</i>	3	87	S3_224567900	S3_222406989	3.35	4.14		-0.031	LPSF64
	<i>qOC4_60</i>	4	188	S4_57947900	S4_75164940	3.42	4.32		-0.031	LPSF64
	<i>qOC5_183</i>	5	148	S5_183857938	S5_182051683	11.99	17.31		-0.063	LPSF64
	<i>qOC7_08</i>	7	470	S7_8930324	S7_7716737	7.89	10.71		0.052	CML505
	<i>qOC9_80</i>	9	263	S9_79412157	S9_83242552	7.44	9.87		-0.048	LPSF64
DH pop4_CML536xLaPostaSequiaF64										
Protein content_LN	<i>qPC5_67</i>	5	197	S5_68309122	S5_66022006	3.12	16.04	15.90	0.051	CML536
Starch content_LN	<i>qPC2_08</i>	2	255	S2_8288623	S2_10208	3.09	6.16	37.02	0.105	CML536
	<i>qSC4_32</i>	4	245	S4_36777242	S4_31674554	9.80	20.36		0.200	CML536
	<i>qSC9_135</i>	9	127	S9_135814387	S9_134561093	5.20	9.65		0.138	CML536
Oil content_LN	<i>qOC5_21</i>	5	229	S5_21876475	S5_20198056	3.40	8.25	8.19	-0.049	LPSF64
* Chr = Chromosome, LOD = Logarithm of Odds; add = additive effect; PVE = phenotypic variance explained; fav allele = parental line contributing the favorable allele for trait, QTL name composed by the trait code followed by the chromosome number in which the QTL was mapped and a physical position of the QTL										

The RR-BLUP model (Endelman 2011) was used to estimate the performance of maize genotypes for grain quality traits for each population (Figure 4 and Supplementary Table S2). Under low nitrogen conditions, average prediction accuracies across the studied genotypes were higher for oil content (0.78) and lower for grain yield (0.08). In IMAS panel we observed the prediction accuracy of 0.41, 0.38, 0.39 and 0.44 under optimum and 0.35, 0.35, 0.41 and 0.56 under low N conditions, respectively. Interestingly, in DH pop CML550/CML504 outperformed other DH populations in terms of genomic prediction accuracy. Under low N, CML550/CML504 had the best prediction accuracy for protein (0.66), oil (0.73), and starch (0.7) content. The prediction accuracy for protein content was highest in DH pop CML550/CML504 both under optimum ($r = 0.66$) as well as under low N stress ($r = 0.69$). For starch content under low N, prediction correlation was highest for CML550/CML504 ($r = 0.70$) followed by the CML536x LaPostaSeqC7-F64-2-6-2-2 ($r = 0.56$), IMAS panel ($r = 0.41$), CML550x CML511 ($r = 0.26$) and CML505x LaPostaSeqC7-F64-2-6-2-2 ($r = 0.23$). CML536x LaPostaSeqC7-F64-2-6-2-2 had the highest prediction correlation for oil content under low N ($r = 0.78$), followed by CML550/CML504 ($r = 0.73$) and CML505x LaPostaSeqC7-F64-2-6-2-2 ($r = 0.71$). CML550x CML511 had the lowest prediction for grain yield ($r = 0.08$), protein ($r = 0.17$) starch ($r = 0.16$), and oil content ($r = 0.11$).

Discussion

Significant levels of malnutrition (Christian and Dake 2021) and food insecurity (Giller 2020) continue to be experienced by maize-dependent smallholder farming populations in SSA that cultivate in nitrogen-depleted soils. Unravelling the genetic architecture of grain yield and quality traits through GWAS and GS is critical for the development of superior genotypes conferring high expression of grain

quality traits both under optimum and low N stress. This study aimed to understand the underlying genetics of low N stress on grain quality traits by combining GWAS with the IMAS panel, QTL detection, and GS in four bi-parental populations. Grain quality traits, notably protein, starch, and oil content, are critical for reducing the incidence of undernutrition in SSA. Understanding the performance of grain quality traits and associated genetic markers under low N stress can aid in the development of maize lines with high protein, starch, and oil content.

Phenotypic analyses showed that protein, starch, and oil content were significantly decreased under low N stress compared to optimal conditions across all tested genotypes. This is consistent with the findings of Liu et al. (2008), who found that in lower N conditions, protein and oil content are considerably reduced. However, the same research reported the opposite for starch content increased under lower levels of N stress. According to Jahangirloo et al. (2021) and Simić et al. (2020), high N conditions are associated with higher protein content and yield. On the other hand, low N substantially decreases protein concentration and all zein fractions apart from β -zeins, according to research conducted in Zimbabwe by Shawa et al. (2021). The impact of soil nutrient management on oil content was significant in the study of Ray et al. (2019). That study reported that using N as a component in NPK blends increased the quantity of saturated fatty acids while decreasing the percentage of unsaturated fatty acids in grain maize oil. Kaplan et al. (2017) suggested that N fertilizer application in combination with adequate irrigation has a favourable effect on the oil content. However, in SSA due to the impoverished economic situation of smallholder farmers, N fertilization is not currently an accessible solution to combat endemic undernutrition. Therefore, maize lines showing high protein, starch and oil content under low N stress should be considered for incorporation into maize breeding programs targeting the SSA region.

In low N stress environments, genetic variability is crucial for the effective selection of enhanced grain quality traits in maize. Ertiro et al. (2020a) asserted that, due to the intrinsic unpredictability of various traits of interest, phenotypic data for trials conducted under low N conditions typically show poor heritability. However, under both optimum and low N environments, our study estimated wide genetic variances and moderate to high broad sense heritabilities. Estimates of heritability ranging from moderate to high imply that the traits have the potential to be enhanced by recurrent selection (Gowda et al. 2021). The influence of G, E, and G x E interactions on oil content was significant under low N conditions across all genotypes examined. The genotypic effects on protein content were significant in some of the genotypes tested (CML505/LaPostaSeqC7-F64-2-6-2-2 and CML536/LaPostaSeqC7-F64-2-6-2-2) under low N conditions. The G x E effects of CML550/CML511 on protein and starch content were significant. The detected significant genotypic variation for the assessed traits in this study indicated the possibility of selecting for improved protein, starch, and oil content under low N stress. Among the three grain quality traits investigated, starch content had the lowest H^2 estimate, whereas oil content had the highest H^2 estimate under low N conditions. Oil content's high broad-sense heritability suggests that its narrow-sense heritability may be even greater, implying that significant genetic gain for this trait is attainable.

Grain yield had a negative genotypic correlation with protein content in all populations, regardless of N level. This supports Arisede et al. (2020)'s findings that increased grain yield was associated with decreased grain protein content in both susceptible and tolerant maize hybrids when residual soil N was low, despite the fact that tolerant hybrids showed a substantially smaller loss in grain protein content. It is well known that choosing between yield and quality is hard in breeding. The observed relationship between grain yield and quality traits in this research under both optimal and low N conditions imply that selecting for grain yield alone will not increase protein, starch, or oil content. On the other hand, protein content had a significant negative correlation with starch content in all populations, which is consistent with previous results (Liu et al. 2008; Zheng et al. 2021). Thus, to increase grain quality, particularly under low N stress, there is a need to select for both grain yield and grain quality. Obviously, this would be very expensive, and a negative relationship makes breeders balance in the selection of these traits hence the need to investigate the potential of molecular breeding in the improvement of these traits.

Association studies targeting protein, oil, and starch content in maize have been conducted utilizing a range of genotypes and marker sets (Alves et al. 2019; Cook et al. 2012; Zheng et al. 2021). The use of GWAS in maize genetics has been highly effective in discovering causal genes for grain quality traits (Zheng et al. 2021). In particular, GWAS is an effective technique for mapping loci linked with complex plant traits in genetically heterogeneous populations (Deng et al. 2021). The power of detection of GWAS is dependent on the LD between the markers and QTL. In outcrossing plant species such as maize, LD declines at a short distance and rapidly (Dinesh et al. 2016). In this study, the LD declined rapidly across physical distance (Kibe et al., 2020a), showing that the IMAS panel has significant genetic diversity and was, therefore, suitable for GWAS.

Candidate genes and SNPs discovered by GWAS for maize grain nutrient content can provide critical information for maize breeding efforts focusing on developing high-quality varieties (Zheng et al. 2021). In this study, GWAS identified 42 SNPs linked to the grain quality traits studied under low N conditions. However, there were no overlapping SNPs for grain quality traits under low N. Under low N stress, two SNPs on chromosomes 3 (*S3_198394847*) and 4 (*S4_120988951*) were discovered to be linked to protein content. Moreover, 12 SNPs with loci on all chromosomes except 4 and 7 were also shown to be substantially associated with oil content under low N stress. The genetic regions

identified in this work through GWAS will be increasingly relevant in future breeding approaches for accurate selection of high grain quality and to increase tolerance of maize lines to low N stress.

Comparison of SNPs identified in this study under low N and optimum conditions revealed no overlapping of SNPs for grain yield and protein content possibly we were not able to detect the common variants responsible for these traits in different management conditions. Nonetheless, for starch content, all the SNPs detected under low N were also detected under optimum conditions. Whereas for the oil content SNP *S6_60978968* on chromosome 6 was consistently detected in both the conditions and also found common SNPs in bin 2.01, 5.03, 6.05 and 9.01 on chromosome 2, 5, 6 and 9, respectively (Table 4). Further comparison of the detected SNPs with the previous studies revealed some overlapping with earlier reported QTL (Wang et al. 2016; Zheng et al. 2021). For instance, SNP *S1_214242607* associated with grain protein content under optimum was closely located with marker detected through GWAS (Zheng et al. 2021) and co-located with QTL detected in two populations (Wang et al. 2016). SNP *S1_191845162* detected for oil content was co-located within the QTL (bnlg2086-umc1122 interval) reported by Zhang et al. (2008) and Wang et al. (2010, umc1395-umc2237 interval). Marker *1_190758142* detected through GWAS for oil content by Zheng et al. (2021) was also located within the same QTL region pointing to the importance of the region for improving oil content in maize. Another SNP *S2_148879075* detected for oil content was located within the QTL region (bnlg108-phi092 interval) reported by Zhang et al. (2008). SNP *S1_17679954* for protein content was co-located within the QTL detected for oil content on chromosome 1 (umc1685-umc1044 interval) in F₃ population (Wang et al. 2010). Nevertheless, some SNPs did not coincide with earlier reports in terms of their physical location. This possibly due to several reasons like these SNPs might be specific to the population in this study, the variation for quality traits in these populations is different, and different methods used to estimate quality traits in different studies also contribute to variation. However, new specific SNPs detected in this study need further validation, nevertheless, these results can serve as a reference for future studies.

We identified 51 candidate genes potentially underlying the molecular and physiological processes governing grain quality traits under optimum and low N environments. The identification of candidate genes based on associated SNPs can aid with the identification of genes important in grain quality performance under optimal and low N environments. Under low N stress, genes coding for shoot apex growth were revealed to be linked with grain yield, protein, starch, and oil content. Peng et al. (2010) asserted that shoot growth, rather than root size, is a good indicator of N sufficiency in maize. The research also identified four candidate genes with protein serine/threonine kinase activity that play a role in soil N response. Protein kinases are well-known regulators of the response of plants to abiotic stresses (Diédhiou et al. 2008; Kulik et al. 2011; Mao et al. 2010). GRMZM2G159307 and GRMZM2G104325 encode ATP binding proteins for grain yield and starch content, respectively. ATP binding proteins are essential for cellular motility, membrane transport and the control of different metabolic activities (Chauhan et al. 2009). ATP-binding has also been reported in several studies to influence the maintenance of homeostasis in plants under both abiotic and biotic stresses (Dahuja et al. 2021; Franz et al. 2011; Jarzyniak and Jasiński 2014). GRMZM2G033694 was assigned to the Histone-lysine N-methyltransferase family at both optimum and low N conditions. It is important to note, however, that these candidate genes should be further validated before being used in breeding schemes. Further functional research on the candidate genes discovered in this study is necessary to validate their possible utility in high grain quality breeding under low N conditions.

Linkage mapping in four populations found multiple QTLs for the studied grain quality traits. Zheng et al. (2021) alluded that, numerous grain nutritional quality QTLs in maize have been identified by genetic dissection of nutrient quality over the last two decades using traditional QTL mapping. Despite the discovery of QTLs and genes that confer superior maize grain quality in some studies, further sources of genetic variation are likely to exist among currently unexplored populations. QTL analyses in four DH populations revealed 8, 13, 12 and 15 potential QTLs associated with grain yield, protein, starch, and oil content, respectively. One QTL on chromosome 3 (*qGY3_187*) for grain yield is overlapped with major effect QTL (*qPC3_187*) for protein content and located between 180 and 189 Mb, which might be an interesting region to improve both protein and grain yield by considering their negative relationship. Zhang et al. (2015) also identified a consistent QTL (umc1644-phi102228 interval) in the same genomic region for protein content. Another QTL *qPC1_115* in CML550/CML504 which explained 11% of the phenotypic variance was consistent with earlier reported QTL (phi001-umc1988 interval) by Zhang et al. (2015) and *qPC10_142* detected on chromosome 10 was consistent with QTL (SYN37373 - PZE110095199 interval) reported by Wang et al. (2016) in recombinant inbred line population. There was one major effects QTL (>10% phenotypic variance explained) for grain yield (*qGY3_196*), three QTL each for protein content (*qPC1_115*, *qPC3_187*, *qPC5_67*) and starch content (*qSC1_180*, *qPC4_32*, *qPC8_124*) and six QTLs for oil content (*qOC2_186*, *qOC3_60*, *qOC4_70*, *qOC5_183*, *qOC6_133* and *qOC7_08*) were detected in four biparental populations. A major QTL (*qSC1_180*) identified in DH pop CML550x CML511, explaining about 11.5% of total phenotypic variance and located between 175 and 188 Mb, was consistent with a QTL (SYN367-PZE101031077 interval) observed in a RIL population by Wang et al. (2016). Similarly, another QTL for starch content (*qSC8_124*) located between 123 and 124 Mb also coincided with earlier reported QTL on chromosome 8 (PZE108069534-SYN19928 interval; Wang et al. (2016)). Similarly, major QTL for oil content *qOC2_186* was also overlapped with earlier detected QTL, indicating several consistent regions for quality traits across genetic back grounds which supports their stable nature and is

amenable for MAS-based improvement. Overall, several QTLs were consistent with the previous studies indicating their reliability to be used in applied breeding.

The promise of GS in tropical maize for various traits of interest has been evaluated in a range of studies (Azmach et al. 2018; Beyene et al. 2019; Crossa et al. 2014; Gowda et al. 2021). The relative merits of GS over phenotypic selection influence its widespread application in breeding programs (Kibe et al. 2020a). Moderate to high accuracies observed in this study for the bi-parental populations and IMAS panel offer promise in breeding for quality traits in tropical maize. Under N-starved soils, average prediction accuracies across the studied genotypes (Figure 4) were higher for oil content (0.78) and lower for grain yield (0.08) which ascribed to their differences in their genetic architecture. CML550/CML504 exhibited the best protein (0.69), oil (0.73), and starch (0.70) content prediction accuracies under low N stress. GS levels have a direct influence on the degree of trait genotypic diversity and heritability in each population (Kibe et al. 2020a). This is confirmed by this study, especially for oil content which had the highest genetic prediction accuracy and H^2 estimates. Prediction accuracy of the IMAS panel was in agreement with various studies on moderately complex traits like resistance for grey leaf spot (Kibe et al. 2020a), common rust (Kibe et al. 2020b), *Striga* (Gowda et al. 2021), maize lethal necrosis and maize chlorotic mottle virus (Sitonik et al. 2019). In the IMAS panel, the observed moderate prediction accuracy can be attributed to its genetic structure and high LD between adjacent markers, which could also be credited to its moderate heritability. Overall, this study indicates that utilizing a common training population to predict grain quality trait performance under low N stress in many linked but separate populations can be beneficial.

Conclusions

To investigate the genetic basis of protein, starch, and oil content performance under low N stress, we employed a single panel consisting of 410 tropical maize lines for GWAS and genomic prediction. QTL mapping was also used to investigate the underlying genetic architecture in four bi-parental populations to better understand the grain quality traits. The genotypic correlations of the grain quality traits investigated indicated that these populations can be used to select better-performing lines under low N stress. GWAS identified 42 SNPs associated with grain quality traits. In addition, several QTLs for the examined grain quality traits were identified by linkage mapping across populations. The genomic regions identified can be used for selection efforts to enhance grain quality trait performance in low-nitrogen soils. Furthermore, the findings showed that including GS in maize breeding can successfully support phenotypic selection to improve grain quality trait performance under low N stress. Future work should, therefore, focus on validating the identified QTLs to enhance the efficacy of maize breeding in SSA.

Declarations

Acknowledgements

The authors are grateful to the International Maize and Wheat Improvement Center (CIMMYT) and partner scientists and technicians who generated the germplasm, designed and conducted the experiments which we used to explore our objectives. Gratitude is also extended to the Cornell University Genotyping Facility and Prof. Ed Buckler's group for providing genotyping services. Technical support received from the Plant & AgriBiosciences Research Centre, Ryan Institute, National University of Ireland Galway (NUI Galway) in drafting the manuscript is acknowledged. The authors would also want to express their gratitude to Kevin Vail Pixley for his edits and reviews.

Funding

The research was supported by the Bill and Melinda Gates Foundation (B&MGF), and the United States Agency for International Development (USAID) through the Stress Tolerant Maize for Africa (STMA, B&MGF Grant # OPP1134248) Project, AG2MW (Accelerating Genetic Gains in Maize and Wheat for Improved Livelihoods, B&MGF Investment ID INV-003439) project, Improved Maize for African Soils (IMAS) Project (&MGF Grant # OPPGD1017), and the CGIAR Research Program on Maize (MAIZE). MAIZE receives W1&W2 support from the Governments of Australia, Belgium, Canada, China, France, India, Japan, Korea, Mexico, Netherlands, New Zealand, Norway, Sweden, Switzerland, U.K., U.S., and the World Bank. This study was made possible by a grant from the Bill & Melinda Gates Foundation to CIMMYT and a Fellowship award from Irish Aid to NN.

Author Contributions

NN, CS, PCM, JC, BD and MG conceived the experiment; BD, MG, and JC conducted the field evaluations and phenotyping; MG, and JC coordinated the GBS experiments; NN, CS, PCM and MG carried out the GWAS and GS analyses; NN, CS, PCM, JC, BD and MG interpreted the results and drafted the manuscript.

Conflict of Interest

The authors declare no conflict of interest.

References

- Abu P, Badu-Apraku B, Ifie BE, Tongoona P, Melomey LD, Offei SK (2021) Genetic diversity and inter-trait relationship of tropical extra-early maturing quality protein maize inbred lines under low soil nitrogen stress. *Plos one* 16:e0252506.
- Alvarado G, Rodríguez FM, Pacheco A, Burgueño J, Crossa J, Vargas M, Pérez-Rodríguez P, Lopez-Cruz MA (2020) META-R: A software to analyze data from multi-environment plant breeding trials. *Crop J* 8:745-756, 10.1016/j.cj.2020.03.010
- Alves ML, Carbas B, Gaspar D, Paulo M, Brites C, Mendes-Moreira P, Brites CM, Malosetti M, Van Eeuwijk F, Patto MCV (2019) Genome-wide association study for kernel composition and flour pasting behavior in wholemeal maize flour. *BMC Plant Biol* 19:1-17.
- Arisede C, Mainassara Z-A, Jill C, Amsal T, Cosmos M, Bish D, Benhildah M, Mike O, Maruthi PB (2020) Low-N stress tolerant maize hybrids have higher fertilizer N recovery efficiency and reduced N-dilution in the grain compared to susceptible hybrids under low N conditions. *Plant Prod Sci* 23:417-426.
- Azmach G, Menkir A, Spillane C, Gedil M (2018) Genetic Loci Controlling Carotenoid Biosynthesis in Diverse Tropical Maize Lines. *G3 (Bethesda)* 8:1049-1065. 10.1534/g3.117.300511
- Bentley AR, Scutari M, Gosman N, Faure S, Bedford F, Howell P, Cockram J, Rose GA, Barber T, Irigoyen J (2014) Applying association mapping and genomic selection to the dissection of key traits in elite European wheat. *Theor Appl Genet* 127:2619-2633.
- Beyene Y, Gowda M, Olsen M, Robbins KR, Pérez-Rodríguez P, Alvarado G, Dreher K, Gao SY, Mugo S, Prasanna BM (2019) Empirical comparison of tropical maize hybrids selected through genomic and phenotypic selections. *Front plant sci* 10:1502.
- Cairns JE, Chamberlin J, Rutsaert P, Voss RC, Ndhlela T, Magorokosho C (2021) Challenges for sustainable maize production of smallholder farmers in sub-Saharan Africa. *J Cereal Sci*:103274.
- Chauhan JS, Mishra NK, Raghava GP (2009) Identification of ATP binding residues of a protein from its primary sequence. *BMC Bioinform* 10:1-9.
- Christian AK, Dake FA (2021) Profiling household double and triple burden of malnutrition in sub-Saharan Africa: prevalence and influencing household factors. *Public Health Nutrition*:1-14.
- Cook JP, McMullen MD, Holland JB, Tian F, Bradbury P, Ross-Ibarra J, Buckler ES, Flint-Garcia SA (2012) Genetic architecture of maize kernel composition in the nested association mapping and inbred association panels. *Plant Physiol* 158:824-834.
- Crossa J, Perez P, Hickey J, Burgueno J, Ornella L, Cerón-Rojas J, Zhang X, Dreisigacker S, Babu R, Li Y (2014) Genomic prediction in CIMMYT maize and wheat breeding programs. *Heredity* 112:48-60.
- Dahuja A, Kumar RR, Sakhare A, Watts A, Singh B, Goswami S, Sachdev A, Praveen S (2021) Role of ATP-binding cassette transporters in maintaining plant homeostasis under abiotic and biotic stresses. *Physiologia Plantarum* 171:785-801.
- Das B, Atlin GN, Olsen M, Burgueño J, Tarekegne A, Babu R, Ndou EN, Mashingaidze K, Moremoholo L, Ligeyo D (2019) Identification of donors for low-nitrogen stress with maize lethal necrosis (MLN) tolerance for maize breeding in sub-Saharan Africa. *Euphytica* 215:1-15.
- Dei HK (2017) Assessment of maize (*Zea mays*) as feed resource for poultry. *Poultry science*:1-32.
- Dekkers J (2007) Prediction of response to marker-assisted and genomic selection using selection index theory. *J Anim Breed Gen* 124:331-341.
- Deng S, Caddell DF, Xu G, Dahlen L, Washington L, Yang J, Coleman-Derr D (2021) Genome wide association study reveals plant loci controlling heritability of the rhizosphere microbiome. *The ISME Journal*:1-14.
- Diédhiou CJ, Popova OV, Dietz K-J, Gollmack D (2008) The SNF1-type serine-threonine protein kinase SAPK4 regulates stress-responsive gene expression in rice. *BMC Plant Biol* 8:1-13.

- Dinesh A, Patil A, Zaidi P, Kuchanur P, Vinayan M, Seetharam K (2016) Genetic diversity, linkage disequilibrium and population structure among CIMMYT maize inbred lines, selected for heat tolerance study. *Maydica*,61(M29).
- Duvick DN (2005) The contribution of breeding to yield advances in maize (*Zea mays* L.). *Advances in agronomy* 86:83-145,
- Ekpa O, Palacios-Rojas N, Kruseman G, Fogliano V, Linnemann AR (2018) Sub-Saharan African maize-based foods: technological perspectives to increase the food and nutrition security impacts of maize breeding programmes. *Global Food Secur* 17:48-56.
- Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE (2011) A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS one* 6:e19379.
- Endelman JB (2011) Ridge regression and other kernels for genomic selection with R package rrBLUP. *The plant genome* 4.
- Ertiro BT, Labuschagne M, Olsen M, Das B, Prasanna BM, Gowda M (2020a) Genetic dissection of nitrogen use efficiency in tropical maize through genome-wide association and genomic prediction. *Frontiers in plant science* 11:474.
- Ertiro TB, Olsen M, Das B, Gowda M, Labuschagne M (2020b) Genetic dissection of grain yield and agronomic traits in maize under optimum and low-nitrogen stressed environments. *Int J Mol Sci* 21:543.
- FAO (2021) FAOStat. Food and Agriculture Organization of the United Nations, Rome. <https://www.fao.org/faostat/en/>
- Farnworth CR, Stirling C, B. Sapkota T, Jat ML, Misiko M, Attwood S (2017) Gender and inorganic nitrogen: what are the implications of moving towards a more balanced use of nitrogen fertilizer in the tropics? *International Journal of Agricultural Sustainability* 15:136-152.
- Franz S, Ehlert B, Liese A, Kurth J, Cazalé A-C, Romeis T (2011) Calcium-dependent protein kinase CPK21 functions in abiotic stress response in *Arabidopsis thaliana*. *Mol Plant* 4:83-96.
- Fraeval S, Hammond J, Bogard JR, Ng'endo M, van Etten J, Herrero M, Oosting SJ, de Boer IJ, Lannerstad M, Teufel N (2019) Food access deficiencies in sub-Saharan Africa: prevalence and implications for agricultural interventions. *Frontiers in Sustainable Food Systems* 3:104.
- Galli G, Alves FC, Morosini JS, Fritsche-Neto R (2020) On the usefulness of parental lines GWAS for predicting low heritability traits in tropical maize hybrids. *PLoS one* 15:e0228724.
- Giller KE (2020) The food security conundrum of sub-Saharan Africa. *Global Food Secur* 26:100431.
- Goredema-Matongera N, Ndhlela T, Magorokosho C, Kamutando CN, van Biljon A, Labuschagne M (2021) Multinutrient biofortification of maize (*Zea mays* L.) in Africa: current status, opportunities and limitations. *Nutrients* 13:1039.
- Gowda M, Das B, Makumbi D, Babu R, Semagn K, Mahuku G, Olsen MS, Bright JM, Beyene Y, Prasanna BM (2015) Genome-wide association and genomic prediction of resistance to maize lethal necrosis disease in tropical maize germplasm. *Theor Appl Genet* 128:1957-1968.
- Gowda M, Makumbi D, Das B, Nyaga C, Kosgei T, Crossa J, Beyene Y, Montesinos-López OA, Olsen MS, Prasanna BM (2021) Genetic dissection of *Striga hermonthica* (Del.) Benth. resistance via genome-wide association and genomic prediction in tropical maize germplasm. *Theor Appl Genet* 134:941-958.
- Harper L, Gardiner J, Andorf C, Lawrence CJ (2016) MaizeGDB: the maize genetics and genomics database. *Plant Bioinformatics*. Springer, pp 187-202.
- Heffer P, Prud'homme M (2015) Fertilizer outlook 2015–2019. 83rd IFA annual conference, Istanbul (Turkey), pp 25-27.
- Illés Á, Mousavi SN, Bojtor C, Nagy J (2020) The plant nutrition impact on the quality and quantity parameters of maize hybrids grain yield based on different statistical methods. *Cereal Res Commun* 48:565-573.
- Jahangirlou MR, Akbari GA, Alahdadi I, Soufizadeh S, Parsons D (2021) Grain Quality of Maize Cultivars as a Function of Planting Dates, Irrigation and Nitrogen Stress: A Case Study from Semiarid Conditions of Iran. *Agriculture* 11:11.
- Jarzyniak KM, Jasiński M (2014) Membrane transporters and drought resistance—a complex issue. *Frontiers in plant science* 5:687.
- Jayne TS, Sanchez PA (2021) Agricultural productivity must improve in sub-Saharan Africa. *Science* 372:1045-1047.

Kaplan M, Kale H, Karaman K, Unlukara A (2017) Influence of different irrigation and nitrogen levels on crude oil and fatty acid composition of maize (*Zea mays* L.). *Grasas y Aceites* 68:e207-e207.

Kibe M, Nair SK, Das B, Bright JM, Makumbi D, Kinyua J, Suresh LM, Beyene Y, Olsen MS, Prasanna BM, Gowda M (2020a) Genetic Dissection of Resistance to Gray Leaf Spot by Combining Genome-Wide Association, Linkage Mapping, and Genomic Prediction in Tropical Maize Germplasm. *Front Plant Sci* 11, 10.3389/fpls.2020.572027.

Kibe M, Nyaga C, Nair SK, Beyene Y, Das B, Bright JM, Makumbi D, Kinyua J, Olsen MS, Prasanna BM (2020b) Combination of linkage mapping, GWAS, and GP to dissect the genetic basis of common rust resistance in tropical maize germplasm. *Int J Mol Sci* 21:6518.

Kulik A, Wawer I, Krzywińska E, Bucholc M, Dobrowolska G (2011) SnRK2 protein kinases—key regulators of plant response to abiotic stresses. *Omics: a journal of integrative biology* 15:859-872.

Liu X, Huang M, Fan B, Buckler ES, Zhang Z (2016) Iterative usage of fixed and random effect models for powerful and efficient genome-wide association studies. *PLoS Genet* 12:e1005767.

Liu Z, Xie H, Tian G, Chen S, Wang C, Hu Y, Tang J (2008) QTL mapping of nutrient components in maize kernels under low nitrogen conditions. *Plant Breed* 127:279-285,

Mao X, Zhang H, Tian S, Chang X, Jing R (2010) TaSnRK2. 4, an SNF1-type serine/threonine protein kinase of wheat (*Triticum aestivum* L.), confers enhanced multistress tolerance in Arabidopsis. *Journal of experimental botany* 61:683-696.

Ngaboyisonga C, Njoroge K (2014) Quality protein maize under low-nitrogen and drought: genotype by environment interaction for grain and protein qualities. *Agricultural Journal* 9:68-76.

Nuss ET, Tanumihardjo SA (2011) Quality protein maize for Africa: closing the protein inadequacy gap in vulnerable populations. *Advances in Nutrition* 2:217-224.

Oikeh S, Kling J, Okoruwa A (1998) Nitrogen fertilizer management effects on maize grain quality in the West African moist savanna. *Crop Sci* 38:1056-1161.

Peng Y, Niu J, Peng Z, Zhang F, Li C (2010) Shoot growth potential drives N uptake in maize plants and correlates with root growth in the soil. *Field Crops Research* 115:85-93,

Poole N, Donovan J, Erenstein O (2021) Agri-nutrition research: revisiting the contribution of maize and wheat to human nutrition and health. *Food Policy* 100:101976.

Prasanna B, Nair SK, Babu R, Gowda M, Zhang X, Xu Y, Olsen M, Chaikam V, Cairns JE, Zaman-Allah M (2020) Increasing genetic gains in maize in stress-prone environments of the tropics. *Genomic Designing of Climate-Smart Cereal Crops*. Springer, pp 97-132.

Prasanna BM, Cairns JE, Zaidi PH et al. (2021) Beat the stress: breeding for climate resilience in maize for the tropical rainfed environments. *Theor Appl Genet* 134, 1729–1752. <https://doi.org/10.1007/s00122-021-03773-7>.

Profillidis V, Botzoris G (2019) Chapter 5—Statistical Methods for Transport Demand Modeling. *Modeling of Transport Demand*; Profillidis, VA, Botzoris, GN, Eds.,

Rao PG, Behera T, Gaikwad AB, Munshi A, Srivastava A, Boopalakrishnan G (2021) Genetic analysis and QTL mapping of yield and fruit traits in bitter melon (*Momordica charantia* L.). *Scientific reports* 11:1-10.

Ray K, Banerjee H, Dutta S, Hazra AK, Majumdar K (2019) Macronutrients influence yield and oil quality of hybrid maize (*Zea mays* L.). *PLoS one* 14:e0216939,

Scott MP, Edwards JW, Bell C, Schussler J, Smith J (2006) Grain composition and amino acid content in maize cultivars representing 80 years of commercial maize varieties. *Maydica* 51:417.

Semagn K (2014) Leaf tissue sampling and DNA extraction protocols. *Molecular Plant Taxonomy*. Springer, pp 53-67.

Shawa H, van Biljon A, Labuschagne MT (2021) Protein quality and quantity of quality protein maize (QPM) and non-QPM hybrids under optimal and low nitrogen conditions. *Cereal Chemistry* 98:507-516.

- Sheahan M, Barrett CB, Sheahan MB (2014) Understanding the agricultural input landscape in sub-Saharan Africa: Recent plot, household, and community-level evidence. World Bank Policy Research Working Paper.
- Simić M, Dragičević V, Mladenović Drinić S, Vukadinović J, Kresović B, Tabaković M, Brankov M (2020) The contribution of soil tillage and nitrogen rate to the quality of maize grain. *Agronomy* 10:976.
- Sitonik C, Suresh LM, Beyene Y et al (2019) Genetic architecture of maize chlorotic mottle virus and maize lethal necrosis through GWAS, linkage analysis and genomic prediction in tropical maize germplasm. *Theor Appl Genet.* 132:2381-2399 <https://doi.org/10.1007/s00122-019-03360-x>
- Tittonell P, Giller KE (2013) When yield gaps are poverty traps: The paradigm of ecological intensification in African smallholder agriculture. *Field Crops Research* 143:76-90.
- Wang Y, Li J, Li Y, Wei M, Li X, Fu J (2010) QTL detection for grain oil and starch content and their associations in two connected F2:3 populations in high-oil maize. *Euphytica* 174:239–252.
- Wang Z, Liu N, Ku L, Tian Z, Shi Y, Guo S, Su H, Zhang L, Ren Z, Li G (2016) Dissection of the genetic architecture for grain quality-related traits in three RIL populations of maize (*Zea mays* L.). *Plant Breed* 135:38-46.
- Wood SA, Tirfessa D, Baudron F (2018) Soil organic matter underlies crop nutritional quality and productivity in smallholder agriculture. *Agriculture, ecosystems & environment* 266:100-108.
- Xiao Y, Liu H, Wu L, Warburton M, Yan J (2017) Genome-wide association studies in maize: praise and stargaze. *Mol Plant* 10:359-374.
- Zhang H, Jin T, Huang Y, Chen J, Zhu L, Zhao Y, Guo J (2015) Identification of quantitative trait loci underlying the protein, oil and starch contents of maize in multiple environments. *Euphytica* 205:169-183.
- Zhang J, Lu X, Song X, Yan J, Song T, Dai J, Rocheford T, Li J (2008) Mapping quantitative trait loci for oil, starch, and protein concentrations in grain with high-oil maize by SSR markers. *Euphytica* 162:335-344.
- Zhang L, Liang Z-y, He X-m, Meng Q-f, Hu Y, Schmidhalter U, Zhang W, Zou C-q, Chen X-p (2020) Improving grain yield and protein concentration of maize (*Zea mays* L.) simultaneously by appropriate hybrid selection and nitrogen management. *Field Crops Research* 249:107754.
- Zhao Y, Gowda M, Liu W, Würschum T, Maurer HP, Longin FH, Ranc N, Reif JC (2012) Accuracy of genomic selection in European maize elite breeding populations. *Theor Appl Genet* 124:769-776.
- Zheng Y, Yuan F, Huang Y, Zhao Y, Jia X, Zhu L, Guo J (2021) Genome-wide association studies of grain quality traits in maize. *Scientific Reports* 11:1-12.

Figures




Figure 1

Phenotypic distribution for grain yield and quality traits evaluated under optimum and Low N stress conditions. The sky blue and red color plots represent the trials conducted under optimum and low N stress conditions, respectively. DH pop1= CML550/CML504; DH pop2= CML550/CML511; DH pop3= CML505/LaPostaSeqC7-F64-2-6-2-2; and DH pop4= CML536/LaPostaSeqC7-F64-2-6-2-2.




Figure 2

Manhattan and quantile-quantile plots generated using a mixed linear model for grain yield (A), Protein content (B), starch content (C) and Oil content (D) under optimum management. The significance level ($P = 2 \times 10^{-5}$ at 0.1 False Discovery Rate (FDR)) is represented by the dashed horizontal line. The X-axis shows the position of SNPs along the 10 maize chromosomes, with various colours indicating distinct chromosomes. The Y-axis shows the $-\log_{10}(P \text{ observed})$ in each analysis.




Figure 3

Manhattan and quantile-quantile plots generated using a mixed linear model for grain yield (A), protein content (B), starch content (C) and oil content (D) under low N stress management. The significance level ($P = 2 \times 10^{-5}$ at 0.1 False Discovery Rate (FDR)) is represented by the dashed horizontal line. The X-axis shows the position of SNPs along the 10 maize chromosomes, with various colours indicating distinct chromosomes. The Y-axis shows the $-\log_{10}(P \text{ observed})$ in each analysis.




Figure 4

Genome-wide prediction accuracies for grain yield and quality traits in IMAS panel (A), DH pop2= CML550/CML511 (B), DH pop3= CML505/LaPostaSeqC7-F64-2-6-2-2 (C), DH pop1= CML550/CML504 (D) and DH pop4= CML536/LaPostaSeqC7-F64-2-6-2-2 (E). Blue and red colour box plots indicate traits were evaluated under optimum and low N stress conditions, respectively.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterials.docx](#)