

Does the Medical Insurance System Play a Real Role in Reducing Catastrophic Economic Burden in Elderly Patients With Cardiovascular Disease in China? Implication for Accurately Targeting Vulnerable Characteristics.

Meiyan Ma

Research Center of Public Policy and Management, School of Health Management, Harbin Medical University, Harbin, Heilongjiang, China

Li Ye (✉ liye8459@163.com)

Harbin Medical University

Yuze Li

Department of Medicine, Jiamusi University

Nianshi Wang

Research Center of Public Policy and Management, School of Health Management, Harbin Medical University, Harbin, Heilongjiang, China

Qunhong Wu

Research Center of Public Policy and Management, School of Health Management, Harbin Medical University, Harbin, Heilongjiang, China

Linghan Shan

Research Center of Public Policy and Management, School of Health Management, Harbin Medical University, Harbin, Heilongjiang, China

Qi Xia

Research Center of Public Policy and Management, School of Health Management, Harbin Medical University, Harbin, Heilongjiang, China

Xiyu Zhang

Research Center of Public Policy and Management, School of Health Management, Harbin Medical University, Harbin, Heilongjiang, China

Wenqing Miao

Research Center of Public Policy and Management, School of Health Management, Harbin Medical University, Harbin, Heilongjiang, China

Wanxin Tian

Research Center of Public Policy and Management, School of Health Management, Harbin Medical University, Harbin, Heilongjiang, China

Dingyun You

School of public health, Kunming Medical University, Kunming, Yunnan, China

Baoguo Shi

Department of Economics, School of Economics, Minzu University of China, Beijing, China

Heng Li

Hospital Development institute of Shanghai Jiao Tong University, Shanghai, China

Tao Sun

Department of Health Service Management, School of Medicine, Hang Zhou Normal University, Hangzhou, Zhejiang, China

Xuelian Fu

The Second Affiliated Hospital of Harbin Medical University, Harbin, Heilongjiang, China

Yanhua Hao

Research Center of Public Policy and Management, School of Health Management, Harbin Medical University, Harbin, Heilongjiang, China

Han Gao

The Affiliated Tumor Hospital of Harbin Medical University, Harbin, Heilongjiang, China

Research

Keywords: cardiovascular disease, catastrophic health expenditure, impoverishment by medical expense, medical insurance, elderly households, China

Posted Date: November 11th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-102328/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Globalization and Health on March 29th, 2021. See the published version at <https://doi.org/10.1186/s12992-021-00683-7>.

Does the Medical Insurance System Play a Real Role in Reducing Catastrophic Economic Burden in Elderly Patients with Cardiovascular Disease in China? Implication for Accurately Targeting Vulnerable Characteristics.

Economic Burden in Elderly Patients with Cardiovascular Disease in China.

Meiyan Ma¹, Ye Li^{1*}, Nianshi Wang¹, Qi Xia¹, Mingli Jiao¹, Qunhong Wu¹, Linghan Shan¹, Xiyu Zhang¹, Dingyun You², Wanxin Tian¹, Heng Li³, Tao Sun⁴, Xuelian Fu⁵, Baoguo Shi⁶, Yanhua Hao¹, Han Gao⁷

¹Department of Social Medicine, School of Health Management, Harbin Medical University, Harbin, Heilongjiang, China;

²School of public health, Kunming Medical University, Kunming, Yunnan, China;

³Hospital Development institute of Shanghai Jiao Tong University, Shanghai, China;

⁴Department of Health Service Management, School of Medicine, Hangzhou Normal University, Hangzhou, Zhejiang, China;

⁵The Second Affiliated Hospital of Harbin Medical University, Harbin, Heilongjiang, China;

⁶Department of Economics, School of Economics, Minzu University of China, Beijing, China;

⁷The Affiliated Tumor Hospital of Harbin Medical University, Harbin, Heilongjiang, China.

Corresponding author: Ye Li, Harbin Medical University, Harbin, Heilongjiang, 150086, China. E-mail addresses: liye8459@163.com

Qunhong Wu, Harbin Medical University, Harbin, Heilongjiang, 150086, China. E-mail addresses.

● Abstract

- **Background:** The vulnerability of cardiovascular disease (CVD) patients' health abilities, combined with the severity of the disease and the overlapping risk factors, can cause such people to bear the economic burden of the disease due to the medical services. We estimated the economic burden of CVD, and identified the weak link in the design of the medical insurance.
- **Methods:** Data from 5,610 middle-aged and elderly with CVD were drawn from the 2015 wave of China Health and Retirement Longitudinal Study (CHARLS). The recommended method of the World Health Organization (WHO) was adopted to calculate catastrophic health expenditure (CHE), impoverishment by medical expenses (IME), and applied the treatment-effect model to analyze the determinants of CHE.
- **Results:** The incidence of CHE was 19.9% of the CVD, compared to 7.6% of the sample households trapped in IME. The incidence of CHE in CVD participating in medical insurance was 2.6% higher than for uninsured families (16.3%). Family size, health satisfaction, combine with other chronic diseases, having hospitalization and disabled members, and participating in insurance were found to be significantly associated with the CHE.
- **Conclusions:** Elderly with physical vulnerabilities were more prone to CHE. The medical insurance only reduced barriers to accessing health resources for elderly with CVD, but lacked policy inclination for high-utilization populations, and had poorly accurate identification of vulnerable characteristics of CVD, which in turn affects the economic protection ability of the medical insurance. The dispersion between the multiple medical

security systems leads to the existence of blind spots in the economic risk protection of individuals and families.

- **Keywords:**cardiovascular disease, catastrophic health expenditure, impoverishment by medical expense, medical insurance, elderly households,China

Introduction

Aging, rapid urbanization, and shifting disease patterns are causing cardiovascular disease (CVD) to become established as one of the primary diseases in the world.[1.2] Worldwide, roughly 17.9 million people die of CVD each year, and the absolute number of deaths from chronic CVD worldwide increased by 42.4% between 1990–2015 (2017).[3.4]The incidence and mortality of CVD are also serious in China. As of 2015, there were 290 million people suffering from CVD in China, including more than 9.5 million cases of heart disease.[5] In addition, between 1990–2013, the total number of CVD deaths increased by 46% in China, mainly due to population aging.[6] The increasing trend of CVD has brought a heavy economic burden to the world at the population and household levels.[7] It is estimated that as of 2015, the estimated annual global cost of CVD is expected to increase by 16%, from 906 billion to more than 104.4 billion by 2030.[8] In India, 25% of families with a member with CVD experience catastrophic expenditure, and 10% are driven into poverty.[9]

The particularity of the internal disease structure and the characteristics of the elderly have made China one of the countries with a high incidence of CVD. In China, the Disability Adjusted of Life Years(DALY) caused by cardiovascular disease has reached 582.255 million, accounting for roughly 20% of the world's cardiovascular disease, which is nearly 9% higher than the international average.[10] Annual inpatient care cost has risen at a steady rate of 30%

since 2004.[11] Compared to Western Europe, China suffers a 50% higher mortality rate from CVD and faces a larger economic burden.[12] Annual direct medical expenses for cardiovascular diseases in China are more than 130 billion yuan, accounting for more than 22% of the total medical expenses in China during the same period, which is 10% higher than Australia.[13] Some research indicates that during the 30 years from 2010–2040, the annual economic benefits of reducing cardiovascular mortality by 1% in China is equivalent to 68% of real GDP in 2010, exceeding US \$10.7 trillion.[14]

China's basic medical insurance system consists of three types of medical insurance scheme : urban employees' insurance medical(MIUE), urban resident insurance medical(MIUR), and the new rural cooperative medical system(NCMS).[15] The basic medical insurance for economic protection with two basic goals: first, to ensure that all people have access to high-quality care; and second, that everyone has the ability to pay for these services to maintain and improve their health.[16]

China has basically achieved 95% coverage of medical and health services. However, the inefficiency of the health care system and the uneven distribution of medical resources make the cost-effectiveness of continuous medical reforms lower, and the coverage and depth of the medical insurance system needs to be improved. Studies have demonstrated that the actual effect of this coverage has been offset by the rapid escalation of medical expenses, especially for the elderly group that combines age and disease.[17] Some scholars measured the disease burden of CVD patients aged 60 years and older in 2011–2013 and found that the direct and indirect medical costs of residents rose from 4,938 yuan to 5,717 yuan, and the OOP (Out-of-pocket) increased by roughly 12 percentage points (from 58.1% to 70%).[18] The

resident's medical insurance reimbursement ratio is much lower than the European Union's (EU) major disease reimbursement ratio of not less than 60%.[19]The World Health Organization (WHO) reports that a reasonable OOP is 15–20%.[20] Once the literature exceeds the prescribed warning value, it will often cause vulnerable groups, such as rural residents and the elderly with low economic incomes, to fall into excessive cash health expenditures, resulting in “ impoverishment by medical expenses” and “returning from poverty.”

The above data indicate that the vulnerability of CVD patients' health abilities, combined with the severity and recurrence of the disease itself and the overlapping risk factors, caused such people to bear the heavy economic burden of the disease due to the use of medical services. Therefore, in the study we adopt the WHO recommended method to verify the health-reducing effect of the health care system, that is, whether the medical insurance system reduces the risk of CHE in cardiovascular patients over the age of 45. And it is important to understand whether medical insurance system increases the accessibility and affordability of health care, which provides good evidence for further ways to alleviate health poverty.

Methods

Data source and sampling method

The data used to calculate the rates of catastrophic health expenditure (CHE) and impoverishment by medical expenses (IME) were obtained from the 2015 China Health and Pension Tracking Survey (CHARLS) database, which is designed to collect microscopic information from >45-year-olds. CHARLS covers 450 communities in 150 counties from 28

of the 32 provinces in mainland China. Face-to-face household interviews were conducted by qualified investigators. Households were randomly selected from maps and listings within each rural or urban community by four-stage stratified cluster sampling to select eligible individuals. Data were collected through questionnaires, and quality control was implemented by supervisors and included GPS comparison, data verification, recording verification, and telephone verification. A total of 5,610 samples (2,507 households) were finally obtained after cleaning the incomplete data and missing values.

Statistical analysis

Calculation method for CHE and IME

We adopted the WHO's recommended method to calculate CHE and IME.[21] CHE was defined as an OOP payment for health care equaling or exceeding 40% of a household's capacity to pay. IME was defined as consumption expenditure equal to or higher than household subsistence expenditure but lower than the subsistence expenditure (SE) net of OOP health payments. The key expenditure indicators involved in the calculation process were as follows:

Equivalent family size: $eqsize_h = hsize_h^{0.56}$

Equivalent food expenditure: total household food expenditure divided by equivalent family size

$$eqfood_h = \frac{food_h}{eqsize_h}$$

Poverty Line (pl): the weighted average food expenditure of a household, whose food expenditure as a share of household consumption expenditure fell between the 45th and 55th percentiles of the entire sample.

$$pl = \frac{\sum w_h * eqfood_h}{\sum w_h}$$

Household subsistence expenditure (SE): was calculated using food expenditure as a share of total household consumption expenditure.

$$se_h = pl * eqsize_h$$

A household's capacity to pay (CTP): was defined as non-subsistence spending of a household as a share of total household consumption expenditure.

$$ctp_h = exp_h - se_h \quad \text{if } se_h \leq food_h$$

$$ctp_h = exp_h - food_h \quad \text{if } se_h > food_h$$

Out-of-pocket health expenditure (OOP): the payments made by households for their health services without third-party compensation.

$$oopctp_h = \frac{oop_h}{ctp_h}$$

Household consumption expenditure (exp): comprises both monetary and in-kind payments on all goods and services and the money value of the consumption of home-made products.

The consumption quintile: ranked by equalized per capital household consumption expenditure weighted with the standard household size rather than the actual household scale.

$$eq \exp_h = exp_h / eqsize_h$$

Treatment effect model

The relationship between participating in the medical insurance system and CHE is characterized by a joint causal relationship. There are implicit selection biases in participating in the medical insurance system, such as the average community participation rate,

self-assessment of health status, and other unobserved characteristics associated with initial selection. To address the bias caused by hidden selectivity bias and joint causality, we applied a instrumental variable (IV) method called the therapeutic effect model.[22,23]

Instrument indicators

We assume that indicators of medical insurance are endogenous. In order to address this endogeneity, we needed instrumental variables, which are related to endogenous predictors (medical insurance) but are not related to the error term of outcome variables (CHE)[24]The premise of using the instrumental variable method is that there are endogenous variables in the regression equation. So, it is necessary to identify endogenous variables (whether participation in medical insurance) in the regression equation. In our paper, we verify this problem by DWH test. In other words, the DWH test checks whether the endogenous predictor is truly endogenous. (The heterogeneity test result shows that, $P = 0.000 < 0.05$, so we use the DWH test)

By reviewing existing literature, we initially identified two possible instruments: self-assessment of health status and average community participation rate. Generally speaking self-reported poorly-healthy groups are more likely to choose to take insurance medical to maintain their health.[25] In addition, the average community participation rate of residents' basic medical insurance will have an impact on the willingness of families to participate in insurance. People living in the same community have certain common characteristics, such as family economic level and health management awareness, which will make them participate in medical insurance. Moreover, the basic medical insurance plan in China is a community or company-based unit. For example, the MIUE is driven by the company, and the NCMS is that

the village cadres mobilize the villagers to participate in the medical insurance scheme, so that the community participation rate is highly correlated with the willingness to participate in the family.[26]

Good instrument should satisfy two main criteria known as relevance and validity criteria, that is to say, good instruments would be correlated with the endogenous variable (relevance criteria) but not to be correlated with the error terms in the model of the outcome variable (validity criteria).

We use the Over-identification Test and GMM regression to check the validity and relevance.[27,28] In addition, in order to further investigate the weak instrumental variable problem, we also perform redundancy-test. The results show that community participation rate becomes our ultimate effective instrumental variable. The results regarding the validity and relevance are **presented in Table1.**

Table 1. Validity test and correlation test

The treatment effect model includes a two-stage regression. In the first phase, we regression the outcome variables of the covariates (the heterogeneity test results show that $P = 0.000 < 0.05$, so we use GMM as our regression equation); based on the results of the first phase, we add instrument variables and the outcome variable performs a quadratic regression.

Outcome variable

We created a binary indicator for CHE as the outcome variable (1=occurrence, 0=no occurrence).

Covariates

In our study, we used three sets of information as covariates for this paper: Social

demographic characteristics, specifically age, gender, marital status, education level of household head, family size, and having a member over 65 years old. The health status of family members, including whether the household has at least one member with a chronic disease, who has been admitted to the hospital, or has a disability. The demand and utilization of health services for household heads, mainly including hospitalization rate, outpatient rate, and non-admission rate (defined as the percentage of patients who need to be hospitalized but are not hospitalized).

Results

Sample characteristics

Table 2. Patients' demographic characteristics

Table 2 summarizes the characteristics of CVD in 2015. The total sample comprised 2,507 households and 5,610 individuals, of which 50.2% of household heads with CVD were male, and around 65.7% of the elderly were aged <64. Over 59.4% had an elementary to junior high education level, only 23.3% of the CVD patients had a health satisfaction status of "good," with >60.4% having one or more other chronic diseases. Furthermore, 97.1% of the CVD patients had medical insurance.

Health-care needs and service utilization

Table 3. Health care demand, service utilization, and reimbursement of household head

Regarding health service utilization for the high-risk group of elderly CVD patients, the rate of monthly prevalence was 11.4% overall. A higher proportion of patients with CVD used inpatient services (14.8%), and more than 19.8% of respondents reported the utilization of

outpatient services over the past month. The above indicators demonstrate different trends for different populations with cardiovascular diseases as follows:

Cardiovascular patients with a high level of health service demand but low utilization of health services and low reimbursement rate. For example, families with cardiovascular disease patients and >3 other chronic diseases had a high demand for health services. Its prevalence rate and hospitalization rate were 13.8% and 23.5%, respectively. However, the health service utilization (non-admission rate 9.8%) and the reimbursement level (48.5%) were lower than cardiovascular disease patients without other chronic diseases.

In terms of family economic level and type of medical insurance, cardiovascular patients with a high level of health service demand and high health utilization but low reimbursement. Household heads of cardiovascular diseases with NCMS had a higher prevalence (14.3%) and outpatient rate (20.5%), but the hospitalization reimbursement ratio (39.7%) was lower by nearly 30% than MIUE (69.8%). Patients who participated in MIUR of cardiovascular diseases demonstrated a similar pattern. It is worth noting that among the wealthiest households with cardiovascular diseases, the hospitalization rate and visit rate were 16.4% and 20.9%, respectively, but the hospitalization reimbursement ratio was only 49.2%, less than 50%.

CHE, and IME in different households

Table 4. CHE, and IME in different households

We further measured the incidence of CHE and IME in elderly with CVD, and 19.9% of all interviewed CVD patients encountered CHE, and the IME occurred in 7.6% of the overall population.

We also found that as the family size increased for CVD patients, the risk of households being trapped in CHE and IME was reduced.. For example, the incidence of CHE among the household with one people(8.1%) was twice that of having no other diseases (14.7%). Furthermore, families with inpatients and disability members of CVD increased the risk of CHE and had a greater financial burden.,and the incidence of CHE were 29.3% and 31.1%, which is more than 10% higher than that of those without inpatients and disability members of CVD (18.6% and 19.5%).

It is worth noting that income level is not the main reason for the high incidence of CHE in patients with CVD. Among them, sub-poor families with CVD had the highest incidence rate of CHE (22.5%) and health expenditure burden(22.1%),,respectively,higher than the richest households with CVD .As far as the type of insurance is concerned, among the three basic medical insurances, cardiovascular families with NCMS had the greatest economic burden of disease, and the risk of CHE was also the highest (21.6%), nearly 8.3% higher than that of NIUE (13.3%).

The treatment-effect model for patients with CVD

Table 5. Results of treatment-effect model for patients with CVD

Community participation rate was added as an effective instrumental variable to the treatment-effect model. The results were as follows: family size, health satisfaction, combination with other chronic diseases, having hospitalization members, having disabled members, and participating in insurance were all found to be significantly associated with the odds of encountering CHE with CVD ($P<0.05$).

Social demographic perspective: The incidence of CHE and family size with CVD have

developed in reverse—that is, as the cardiovascular family grows in size, the risk of CHE is reduced by 4.99 percentage points.

Health needs and utilization perspective: Combination with other chronic diseases, health satisfaction, having hospitalization members, and having disabled members had a positive correlation with the risk of CHE in cardiovascular patients, and with an increase of combining other chronic diseases or hospitalization members increased, the risk of a family's CHE increased by 4.68 percentage points and 10.01 percentage points, respectively.

Medical insurance perspective: It is worth noting that cardiovascular patients participating in medical insurance increased, the risk of CHE and increased the risk of being trapped in CHE by 28.9 percentage points.

Medical insurance level

As shown in the above figure, among patients with cardiovascular disease, rural population accounts for 75.1% of the sample size, and the CHE of NCMS was the riskiest, at 21.6%. In addition, the treatment-effect model shows that the probability of CHE participating in health insurance families with cardiovascular disease has increased by 29.8 percentage points. Therefore, to identify the NCMS of key bottlenecks, we conducted a specific influencing factor analysis and found that families with cardiovascular disease combined with >3 other chronic diseases (38.88%) were the primary cause of high CHE in NCMS. Next, in turn, being more than 75 years old (33.33%), having disabled members (33.33%), having hospitalization members (32.41%), and combining two chronic diseases (30.74%) were the top five factors influencing high CHE of families with cardiovascular disease in NCMS. The results are **presented in Fig.1.**

Discussion

The proportion of households associated with CHE and IME in elderly patients with CVD were 19.9% and 7.6%, respectively. These rates were higher than those of developed countries, but were lower than those of low-income countries.[29,30]The incidences of CHE participating in medical insurance were 3.6% higher than those of uninsured households (16.3%). In addition, families with cardiovascular disease have a much higher risk of IME than families without cardiovascular disease (7.0%) and the overall population (7.2%).It can be seen that cardiovascular disease patients' risk tolerance for health care payments is actually lower than the average in China. At the same time, our study also found that cardiovascular families with chronic patients, inpatients, and disabled members, are at a higher risk of falling into poverty because of the cost of health care, and they are becoming a stubborn group with a high burden of CVD disease under the medical insurance system.

As a kind of vulnerable group, patients with cardiovascular disease are mainly characterized as a high-risk group integrating physiological, social, and health factors. We need to identify risk factors for patients with CVD and identify vulnerable groups and then better play the economic protection role of the medical insurance system. As mentioned earlier, despite these governmental efforts, many disadvantaged groups are not considered target populations for benefit enhancement. Through comprehensive analysis, we found that the disadvantaged elderly population with mental health has the following characteristics:

Older groups with physical vulnerability are more prone to CHE.

In the first place, age growth, loss of healthy capital, and decline in physiology are inevitable for the elderly. However, elderly people are at a disadvantage in accessing resources and fail

to enjoy social welfare policies fairly.[31,32]For example, the incidence of CHE in elderly people over 75 years old in the NCMS was 33.33%, second only to the merger of three chronic diseases (33.88%).The WHO reported that 23% of the world's disease burden is on older people, and chronic non-communicable diseases have a major impact on this burden.[33]Furthermore, the morbidity and concurrency of elderly patients with CVD may eventually lead to premature death and disability, while long-term health care costs, drug costs, and rehabilitation costs greatly increase the risk of CHE.[34] Our results show that the hospitalization rate (23.5%), rate of visits (24.5%), and incidence of CHE (31.8%) of patients with more than three chronic diseases are far higher than those without chronic diseases (13.3%, 17.7%, 14.7%, respectively), and the treatment-effect model shows that the combination with other chronic diseases increases the risk of 4.68 percentage points of CHE. Past evidence has demonstrated that the likelihood of using health care (e.g, hospitalization) increases in the presence of chronic or multiple conditions.[35]It can be seen that with the increase of the number of chronic diseases, the addition of other chronic diseases will prolong the hospitalization time on the basis of the original single disease, which will cause CVD patients to superimpose the cost of other diseases when they bear the economic pressure from their cardiovascular disease burden. Studies have found that one-third of adults have multiple chronic diseases, equivalent to 3/4 of the elderly in developed countries.[36]Therefore, suffering from a variety of chronic diseases has become a major health problem for the elderly in the future, greatly increasing the risk of CHE.

The medical insurance system only guarantees the basic health utilization threshold for elderly patients with CVD and lacks policy inclination for high-utilization populations.

The existing medical insurance system reduces the threshold of health service utilization for vulnerable groups, such as those with cardiovascular diseases, but only achieves the first goal of medical insurance—that is, to ensure that all people have access to high-quality care. Our results show that the incidence of CHE for high-demand people with inpatients and disabled patients is much higher than that of the normal population. In our paper, families with disabled patients had higher prevalence (21.7%) and outpatient rates (21.8%) than those without disabilities (13.4% and 19.6%, respectively). However, the hospitalization reimbursement ratio was only 40.6%, far below the overall level (44.5%), and OOP accounted for 35.6% of total household health expenses, which is much higher than that of OECD countries.[37] A study in South Korea showed that families with disabilities face higher CHE than those without people with disabilities, and annual living expenses for OOP medical expenses are roughly 1.2 to 1.4 times greater.[38] This may be due to physical or mental disability in disabled patients leading to job loss or reduced earnings, while higher medical care needs due to disability increase the burden of high medical costs.[39] Even relatively small expenditures are catastrophic for poor families, and excessive out-of-pocket health care spending can lead to poverty.[40]

The medical insurance system has poorly accurate identification of vulnerable characteristics, which in turn affects the economic protection ability of the medical insurance system for patients with cardiovascular diseases.

China's medical insurance policy aims to solve the problem of “people falling into poverty due to illnesses” and to ensure that most people are not reduced to poverty because of health-related issues. However, our results indicate that the risk of CHE for CVD patients

participating in medical insurance schemes has increased by 28.9%, which had the highest incidence of CHE for NCMS. It can be seen that the medical insurance system has the disadvantages of insufficient protection of policies in reducing the economic burden of residents' medical care and maintaining residents' health rights and health. China's health sector reform has achieved unprecedented progress, but protecting vulnerable groups from health care-related impoverishment remains a challenge. Specifically, we need to reconsider benefit packages and redesign social health care insurance programs in order to further protect the elderly population with cardiovascular disease.

Inequality between types of health insurance systems: binary urban and rural structure divided by place of residence. The household registration system in China directly affects the ability to obtain various medical benefits.[41] In our paper, the hospitalized reimbursement ratio of MIUE with cardiovascular disease patients was 69.8%, while that of the NCMS was only 39.7%, accounting for only half of the MIUE. Urban residents are expected to have a greater awareness about their health and better access to health insurance (especially private health insurance) and hence are more likely to obtain health insurance. The health needs of people in rural areas are not able to be converted into effective medical needs in time due to lower income levels, high medical prices, and inadequate medical care, thus bearing the risk of greater CHE. A Chinese study shows that NCMS cannot prevent CHE from happening in poor families but only reduces the incidence of CHE in wealthy families.[42] This differentiated design of health care benefits can sometimes lead to social inequalities, often with the same disease, at different costs, and patients with higher socioeconomic status usually enjoy better health insurance and higher service utilization. In

rural areas, this should further increase the reimbursement ratio of outpatient and inpatient expenditures to the elderly with cardiovascular diseases who receive treatment at every kind of hospital.[43]

Inequality of different income groups under the same medical insurance system. Our research indicated that the incidence of CHE in sub-poor households with CVD (22.5%) was higher than 5% of households with the highest income. Its OOP accounts for 22.1% of CTP, and the economic burden of disease was higher than that of high-income families by nearly 10%. A similar phenomenon occurred in India. According to the data, India's medical expenditure accounts for only 0.9% of the GDP, which is lower than the 2.8% of the GDP of less developed countries. However, with only government spending, among the five economic subgroups, the poorest households with less income receive only 10% of medical care, while 20% of the wealthiest households receive up to 33% of social subsidies, three times as much as the poorest households.[44] The heavy burden of disease can lead to a difficult family life in low-income groups, which thus fall into the evil cycle of “poor illness due to illness and illness due to poverty.” Therefore, the internal system design of medical insurance should strengthen the economic support and protection for poor families and achieve certain policy inclinations.

The inconsistencies between multiple health care systems lead to blind spots in the economic risk protection of individuals and families.

China has established multiple medical security systems, including basic medical insurance, major illness insurance, commercial medical insurance, social medical assistance, and charity assistance. However, the complexity of cardiovascular disease and the difference in the

income of the population mean that various systems have not woven a standard, unified safety net to prevent impoverishment by medical expense, which is mainly reflected in the following two points.

Firstly, the particularity of cardiovascular disease can lead to the loss of labor, the indirect cost of nursing expenses, and transportation expenses, and long-term drug maintenance will also increase the economic burden of patients. These neglected indirect costs and drug costs have become blind spots in the economic protection of basic health care systems. In our study, elderly cardiovascular patients still have to pay more than half (55.5%) of their medical expenses after reimbursement by basic medical insurance schemes. Moreover, there is no effective mechanism to connect basic insurance schemes, major disease schemes, and medical assistance insurance systems, resulting in this group of patients falling into poverty after paying for the high medical expense.

Secondly, our result also shows that sub-poor households have the highest incidence of CHE at 22.5%, and the OOP accounts for 22.1% of CTP, much higher than the poorest households. The medical security system has implemented certain health expense reduction policies for these “extremely poor families” without economic sources or micro-income, including lowering the deductible line, increasing the proportion of reimbursement, and improving or eliminating top-up measures. The high-risk marginalized population with an economic income slightly higher than that of the poorest households, which is prone to CHE, has not reached the standard of assistance for supplementary medical assistance and has failed to be covered by a multilevel medical security system.

It can be seen that exploring the integration of multilevel medical insurance systems,

promoting the complementary functions and overlapping effects of multilevel medical insurance systems, and jointly solving the poverty problems caused by cardiovascular patients have become the target task of the current prevention and treatment process of chronic diseases.

Conclusions

The findings revealed that the incidence of CHE and IME were relatively high among elderly households with CVD patients. The elderly who had poor health satisfaction, combination with other chronic diseases, inpatients, and disabled people were the high-risk groups of CVD. Medical insurance programs have not reduced the risk of CHE or relieved the financial burden of CVD patients in China, especially the elderly who participate in the NCMS. Therefore, there is an urgent need for Chinese policymakers to improve the accuracy of medical health insurance schemes to target the vulnerable characteristics of elderly cardiovascular patients. While improving the accessibility of health services for cardiovascular patients, the next step should be to focus on achieving another function of the medical health insurance system, which is to maximize the economic protection for the vulnerable en-rollers and reducing their risk of being trapped in the poverty, rather than increasing the risk. It is worth noting that our research has its limitations. First, the data were self-reported surveys that may be affected by measurement errors. Second, the study used cross-sectional samples rather than panel data, making it difficult to test the long-term effects of health insurance and other control variables on CHE in cardiovascular patients.

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and materials: The datasets generated during and/or analyzed during the current study are available in the CHARLS repository, [<http://charls.pku.edu.cn/pages/data/111/zh-cn.html>]

Competing interests: The authors declare that they have no competing interests.

Funding: This work was supported by the National Natural Science Foundation (71874045, 71403073, 71333003, 71804036) 、 China Postdoctoral Science Foundation (2016M590296)、 and Heilongjiang Postdoctoral Foundation (LBH-Z14166).

Authors' Contributions: Conceptualization, Han Gao, Xiyu Zhang and Yuze Li; Formal analysis, Meiyang Ma, Heng Li and Wenqing Miao; Funding acquisition, Ye Li and Linghan Shan; Investigation, Ye Li; Methodology, Meiyang Ma; Project administration, Nianshi Wang, Linghan Shan and Baoguo Shi; Software, Nianshi Wang, Xuelian Fu and Qi Xia; Validation, Qunhong Wu and Dingyun You; Visualization, Wanxin Tian and Xiyu Zhang; Writing – original draft, Meiyang Ma; Writing – review & editing, Ye Li, Tao Sun and Yanhua Hao.

Acknowledgement: Each author contributed to the concept, design, research, data analysis, drafting of the article, and we acknowledge the outstanding contributions of Yuze Li, Nianshi Wang who contributed equally to the first author of this article. I have obtained written permission from all persons named in the Acknowledgement. And this article has 4564 words in total.

Authors' information: Meiyang Ma¹, Ye Li^{1,*}, Yuze Li², Nianshi Wang¹ , Qunhong

Wu¹, Linghan Shan¹, Qi Xia, Xiyu Zhang¹, Wenqing Miao, Wanxin Tian, Dingyun You³,
Baoguo Shi⁴, Heng Li⁵, Tao Sun⁶, Xuelian Fu⁷, Yanhua Hao¹, Han Gao⁸

¹Research Center of Public Policy and Management, , School of Health Management, Harbin
Medical University, Harbin, Heilongjiang, China;

²Department of Medicine, Jiamusi University, Jiamusi Heilongjiang 154007, China

³School of public health, Kunming Medical University, Kunming, Yunnan, China;

⁴Department of Economics, School of Economics, Minzu University of China, Beijing,
China;

⁵Hospital Development institute of Shanghai Jiao Tong University, Shanghai, China;

⁶Department of Health Service Management, School of Medicine, Hang Zhou Normal
University, Hangzhou, Zhejiang, China;

⁷The Second Affiliated Hospital of Harbin Medical University, Harbin, Heilongjiang, China;

⁸The Affiliated Tumor Hospital of Harbin Medical University, Harbin, Heilongjiang, China.

References

[1] Li Y , Wang D D , Ley S H , et al. Potential Impact of Time Trend of Life-Style Factors on
Cardiovascular Disease Burden in China. *J Am Coll Cardiol*, 2016, 68(8):818-833.

[2] Tuppin P , Rivière, Sébastien, Rigault A , et al. Prevalence and economic burden of
cardiovascular diseases in France in 2013 according to the national health insurance scheme
database. *Arch Cardiovasc Dis*, 2016:S187521361630047X.

[3] Hsu W Y . A decision-making mechanism for assessing risk factor significance in
cardiovascular diseases. *Decis Support Syst*, 2018, 115:64-77.

[4] GBD 2015 Eastern Mediterranean Region Cardiovascular Disease Collaborators. Burden

of cardiovascular diseases in the Eastern Mediterranean Region, 1990-2015: findings from the Global Burden of Disease 2015 study. *Int J Public Health*, 2018, 63(Suppl 1):1-13.

[5] China Cardiovascular Disease Report 2018. Edited by the National Cardiovascular Center. Beijing: China Encyclopedia Publishing House, 2019.1.

[6] Zhou M, Wang H, Zhu J, et al. Cause-specific mortality for 240 causes in China during 1990-2013: a systematic subnational analysis for the Global Burden of Disease Study 2013. *Lancet*. 2016; 387:251-72.

[7] Al-Mallah M H , Farah I , Al-Madani W , et al. The Impact of Nurse-Led Clinics on the Mortality and Morbidity of Patients with Cardiovascular Diseases: A Systematic Review and Meta-analysis. *J Cardiovasc Nurs*, 2016, 31(1):89.

[8] National Academies of Sciences, Engineering, and Medicine. Global Health and the Future Role of the United States: A Report of the National Academies of Sciences, Engineering, and Medicine by the Committee on Global Health and the Future of the United States, Board on Global Health, Health and Medicine Division. Washington, DC: The National Academies Press, 2017.

[9] Engelgau MM, Saharty S, Kudesia P, et al. Capitalizing on the demographic transition: Tackling noncommunicable diseases in South Asia. World Bank: Washington, DC; 2011.

[10] Murray CJ, Vos T, Lozano R, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012; 380 (9859) :2197-2223.

[11] National Center for CVDs of China. Report on CVDs in China (2016). Beijing, China: Encyclopedia of China Publishing House, 2017.

[12] Yusuf S, Reddy S, Anand S. Global Burden of Cardiovascular Diseases Part II: Variations in Cardiovascular Disease by Specific Ethnic Groups and Geographic Regions and Prevention Strategies. *Circulation*, 2001, 104(23):2855-2864.

[13] .Australian Institute of Health and Welfare (AIHW). Australia' s Health 2014; Australia' s Health Series No. 14; Cat. No. AUS 178; AIHW: Canberra, Australia, 2014. Published online on February 1 2017. <https://www.aihw.gov.au/reports/australias-health/australias-health-2014/contents/table-of-contents> .

[14] .World Bank: Toward a Healthy and Harmonious Life in China: Stemming the Rising Tide of Non-Communicable Diseases. 2011. Published online on July 26 2011. <https://www.worldbank.org/en/news/feature/2011/07/26/toward-health-harmonious-life-china-stemming-rising-tide-of-non-communicable-diseases>.

[15] Li Y , Wu QH , Li CJ, et al. Catastrophic Health Expenditure and Rural Household Impoverishment in China: What Role Does the New Cooperative Health Insurance Scheme Play? *Plos one*, 2014, 9(4):e93253~e93253.

[16] Fanourgiakis J, Kanoupakis E. Catastrophic healthcare expenditure during economic recession in the field of cardiovascular disease. *Expert Rev Pharm Out*, 2014, 14(1):5-8.

[17] Liu X , Sun X , Zhao Y , et al. Financial protection of rural health insurance for patients with hypertension and diabetes: repeated cross-sectional surveys in rural China. *BMC Health Serv Res*, 2016, 16(1):481.

[18] Li C, Jian W. Economic burden of cardiovascular disease in middle-aged and elderly people-An empirical analysis based on China's health and pension tracking survey. *Chinese*

Journal of Health Policy Research, 2017, 10(5): 75-80.

[19] European Commission. Published online on March 26, 2014. <https://ec.europa.eu/eurostat/en/web/products-statistical-working-papers>.

[20] World Health Organization. Health Financing Strategy for the Asia Pacific Region (2010-2015). Switzerland: World Health Organization, 2009.

[21] Xu K. Distribution of health payments and catastrophic expenditures methodology. Geneva: Department of Health System Financing, World Health Organization, 2005.

[22] Wooldridge J M . Further results on instrumental variables estimation of average treatment effects in the correlated random coefficient model. *Econ Lett*, 2003, 79(2):185-191.

[23] Jelena A, Milena P, Bernd R, Wim G. Catastrophic Health Care Expenditure among Older People with Chronic Diseases in 15 European Countries. *Plos One*. 2016;11(7):e0157765.

[24] Lewbel A . Endogenous selection or treatment model estimation. *J Econometrics*, 2007, 141(2):777-806.

[25] Wang XW. A Study on the Impact of Health on the Labor Supply Behavior of the Mid-aged and Elderly of China. Anhui University of Finance and Economics, 2016.

[26] Hu HW, Zhang L, Li JY, Du YX, Wang JR. Does Urban Residents Basic Medical Insurance Worsen the Medical Burden to the Elderly?-System Evaluation with Household Medical Burden. *Chinese Scientific Research on Aging*, 2015;(4):51-61.

[27] Gu NY, Gai Y, Hay JW. The effect of patient satisfaction with pharmacist consultation on medication adherence: An instrumental variable approach. *Pharm Pract*, 2008;6(4).

[28] Hall AR., Rudebusch GD, Wilcox DW. Judging Instrument Relevance in Instrumental

Variables Estimation. *Int Econ Rev*, 1996; 37(2): 283-298.

[29] .Jelena Arsenijevic .Catastrophic Health Care Expenditure among Older People with Chronic Diseases in 15 European Countries. *PLoSOne*. 2016; 11(7): e0157765.

[30] .Tolla M T , Norheim O F , Verguet, Stéphane, et al. Out-of-pocket expenditures for prevention and treatment of CVD in general and specialised cardiac hospitals in Addis Ababa, Ethiopia: a cross-sectional cohort study. *BMJ Glob Health*, 2017, 2(2):e000280.

[31] .Sónia Ribeiro, Cláudia Furtado, João Pereira. Association between cardiovascular disease and socioeconomic level in Portugal ☆ *Revista Portuguesa de Cardiologia*, 2013, 32(11):847-854.

[32] .Ngalesoni F N , Ruhago G M , Mori A T , et al. Equity impact analysis of medical approaches to cardiovascular diseases prevention in Tanzania. *Soc Sci Med*, 2016:S0277953616304695.

[33] .WHO. Health in 2015: from MDGs to SDGs. Geneva: World Health Organization, 2015.

[34] .Dittrich L , Stara D . The Impact of Aging Population on the Rise of the Health Care Cost in the Czech Republic. *Int Adva Econo Res*, 2013, 19(1):11-17.

[35] .Hopman P , Heins M J , Rijken M , et al. Health care utilization of patients with multiple chronic diseases in The Netherlands: Differences and underlying factors. *Eur J Intern Med*, 2015, 26(3):190-196.

[36] .Cotter H, Emma S. The global burden of multiple chronic conditions: A narrative review. *Prev Med Rep*, 2018, 12(284-293).

[37] .Organization for Economic Cooperation and Development. Health at a glance 2013:

2015 .http://dx.doi.org/10.1787/health_glance-2013-en.

[38] .Lee J E , Shin H I , Do Y K , et al. Catastrophic Health Expenditures for Households with Disabled Members: Evidence from the Korean Health Panel.J Korean Med Sci, 2016, 31(3).

[39] .Zuvekas S H , Selden T M . Mental Health and Family Out-of-Pocket Expenditure Burdens. Med Care Res Rev, 2010, 67(2):194-212.

[40] .Gotsadze G , Zoidze A , Rukhadze N . Household catastrophic health expenditure: evidence from Georgia and its policy implications. BMC Health Serv Res, 2009, 9(1):69-0.

[41] .Li C, Young B R, Jian W. Association of socioeconomic status with financial burden of disease among elderly patients with cardiovascular disease: evidence from the China Health and Retirement Longitudinal Survey.Bmj Open,2018, 8(3):e018703.

[42] .Guo N , Iversen T , Lu M , et al. Does the new cooperative medical scheme reduce inequality in catastrophic health expenditure in rural China? BMC Health Serv Res, 2016, 16(1):653.

[43] .Wang Z , Li X , Chen M . Catastrophic health expenditures and its inequality in elderly households with chronic disease patients in China.Int J Equity Health, 2015, 14(1):8.

[44]. Meng Q, Xu L, Zhang Y, et al. Trends in access to health services and financial protection in China between 2003 and 2011: a cross-sectional study. Lancet, 2012; 379: 805-14.

Figures




Figure 1

Different vulnerable groups of CHE by NCMS

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Highlights.docx](#)