Caputo SIR Model for COVID-19 under Optimized Fractional Order

Ali S. Alshomrani

King Abdul Aziz University

Malik Z. Ullah

King Abdul Aziz University

Dumitru Baleanu (✉ dumitru@cankaya.edu.tr)

Cankaya University <https://orcid.org/0000-0002-0286-7244>

Research Article

Keywords: Ulam-Hyers stability, Least-squares, Caputo, Pandemic, SIR model

Posted Date: November 5th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-102574/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Caputo SIR Model for COVID-19 under Optimized Fractional Order

Ali S. Alshomrani¹, Malik Z. Ullah¹, Dumitru Baleanu^{2*,3}

¹Department of Mathematics, King Abdul Aziz University, Jeddah, Saudi Arabia.

^{2*}Department of Mathematics, Cankaya University, Ankara, Turkey

³Institute of Space Sciences, Magurele- Bucharest, Romania

* Correspondence to: dumitru@cankaya.edu.tr (D. Baleanu)

October 23, 2020

Abstract

Everyone is talking about coronavirus from last couple of months due to its exponential spread throughout the globe. Lives have become paralyzed and as many as 180 countries are so far affected with 928,287 (14 September, 2020) deaths within couple of months. Ironically, 29,185,779 are still active cases. Having seen such drastic situation, a relatively simple epidemiological SIR model with Caputo derivative is suggested unlike more sophisticated models being proposed nowadays in the current literature. Major aim of the present research study is to look for possibilities and extents to which the SIR model fits the real data for the cases chosen from 01 April to 15 March, 2020, Pakistan. To further analyze qualitative behavior of the Caputo SIR model, uniqueness conditions under the Banach contraction principle are discussed and stability analysis with basic reproduction number is investigated using Ulam-Hyers and its generalized version. Best parameters have been obtained via nonlinear least-squares curve fitting technique. The infectious compartment of the Caputo SIR model fits the real data better than classical version of the SIR model [4]. Average absolute relative error under the Caputo operator is about 48% smaller than the one obtained in the classical case ($\nu = 1$). Time series and 3D contour plots offer social distancing to be the most effective measure to control the epidemic.

Keywords: Ulam-Hyers stability; Least-squares; Caputo; Pandemic; SIR model.

1 Introduction

Literature includes mathematical models for the pathways for the transmission of infectious ailments. These models play an important role in quantifying and assessing the efficient control and preventive measures of infectious ailments [1, 2, 3]. It has been proved in multifarious ways that mathematical modeling is a very flexible and efficient way of researching the dynamics of transmission of infectious ailments. Mathematical analysis and numerical simulations can be used to create and evaluate control measures that are convincing. As for the model of compartmental ailments, there are several infectious ailments, beginning with the very classic SIR model to the more complex ones [4].

In December 2019, there was an emergence of a novel ailment in China which was later declared as pandemic named COVID-19 by world health organization [5]. It is well known that the COVID-19 pandemic has given rise to fearful living and coexistence around the globe, with millions of people worldwide infected. The mortality-recovery ratio appears to be in a positive proportion. Nevertheless, due to the sensitivity of Polymerase Chain Reaction, the absence or presence of the previously infected host is observed and the recovery rate appears to be promising in the absence of any curative vaccine. The challenge facing health care professionals, the World Health Organization and the Center for Disease Control in each quarter was whether reinfection could occur after a COVID-19 patient had been clinically treated. The subtle nature of the disease has led many scientists and medical practitioners to massively embark on multiple studies to combat and avoid the spread of the disease [6, 7, 8, 9, 10, 11, 12].

The seven known human COVID-19, four are the known human influenza pathogens. SARS-CoV, MERS-CoV and 2019-nCoV are responsible for severe respiratory diseases [13]. While the COVID-19 has long been developed and studied by researchers and medical professionals, many people are

lacking knowledge of the disease and vaccines and antiviral drugs are still not available to directly prevent or treat the infection. The last major outbreak of SARS-CoV in China was in 2003. It was an acute respiratory infectious ailments with a large risk of demise. China managed the SARS outbreak through numerous inspections and effective preventive measures. The time of incubation for COVID-19 is substantial and very long compared to SARS. Many studies computed different periods of incubation for the COVID-19, as an example, 5.2 days [14], 3.0 days [15] and 4.75 days [16]. In [15], an incubation period of up to 24 days is reported, this extends to 38 days as reported in Enshi Tujia and Miao autonomous prefecture in Hubei province of China. Notice that people afflicted with asymptomatics are very numerous [19] and in comparison to SARS-CoV and MERS-CoV the death rate is much inferior [15]. Genetic virus studies show that SARS-CoV and 2019-nCoV are 85% homologous [17]. But 2019-nCoV binds ACE2 to an affinity higher than SARS-CoV [18]. In the positive reported cases induced at COVID-19, SARS will be surpassed by the end of 29 January 2020. The variability of the asymptomatic cases, the incubation time and the super transmissibility of the virus bring considerable difficulties in controlling the epidemics.

Over the last few decades, many scientists have shown that the fractional models can more accurately explain natural phenomena than the differential equations of the integer-order. Because of this advantage, the fractional calculus has taken on the importance and popularity of modeling realistic cases, especially those with memory effects [20, 21, 22, 23]. Furthermore, the application of the fractional calculus is used in many fields of social sciences, engineering, and biology [24, 25, 26, 27]. Given this importance, we feel inspired to investigate and examine a new fractional version of the model involving Caputo operator. To the best of our knowledge this is the first time Caputo operator has been employed for the model being considered.

2 Formulation of Caputo SIR Model

Although, there are various complex models proposed recently for transmission dynamics of COVID-19 as discussed in the section 1 but we have used relatively simple and most popular model used in the field of mathematical epidemiology. The SIR model [4] is the stepping stone of all the models proposed after it. This is the reason to choose the SIR model to investigate dynamics of the COVID-19 pandemic while using the Caputo differential operator. The SIR models consists of three mutually distinct categories. The first are those people who could potentially catch the disease $S(t)$, second are those who currently have the disease and can infect others $I(t)$ (infectives), and then $R(t)$ stands for removed and this is the group of people who have already caught to the disease and have now either recovered from the disease or have died with the disease. With all mathematical models, we have to make various assumptions to simplify the real world phenomena because things are just too complicated to express everything in a set of simple equations so here the first assumption that we make is that the epidemic is sufficiently short so it does not last that long so that we can assume that the total population remains constant. The second assumption in the model relates to the way in which the disease is transmitted and we assume that the rate of increase in the infectives is proportional to the contact between susceptibles and infectives and we assume that this occurs at a constant rate. Finally, our third assumption relates to the removal rate and this is the category $R(t)$ so there is a constant rate this could be a death rate or a recovery rate. Having made the assumptions, a set of equations (1–3) can be written down that are going to govern the model.

$$\mathbb{D}\left(S(t)\right) = \beta \frac{SI}{N}, \quad (1)$$

$$\mathbb{D}\left(I(t)\right) = \beta \frac{SI}{N} - \gamma I, \quad (2)$$

$$\mathbb{D}\left(R(t)\right) = \gamma I, \quad (3)$$

where $N = S + I + R$ and $\mathbb{D}$ stands for the classical integer-order differential operator. We now have three differential equations for three categories people within the population. So, the number of susceptibles is going to decrease according to the number of contacts (contact rate = β) between infectives and susceptibles. Similarly, the number of infectives will increase due to contact between people and decrease from people either recovering or dying as a result of the disease. Finally, the removed category people that no longer can catch the disease either because they have recovered or died and this is going to increase at the constant rate (recovery rate = γ) depending on how many infectives there are. Next, we need some initial data before we can solve the system of differential equations. Therefore,

$$S(0) = S_0 > 0, I(0) = I_0 > 0, R(0) = R_0 \geq 0. \quad (4)$$

There are many instances wherein classical epidemiological models have been re-investigated using operators of non-local nature. It is because of their memory property which makes them most suitable tool to capture dynamics for the spread of a disease since epidemics are known to have memory retaining characteristics.

Having being inspired from plethora of research works carried out in fractional mathematical epidemiology (see, for example [28, 29, 30, 31, 32, 33, 34, 35] and most of the references cited therein), we have introduced the following Caputo type SIR model wherein its distinguished feature is that the dimensional inconsistency of the model has been removed by carrying the fractional order ν in the power of biological parameters β and γ .

$$\begin{aligned} {}^C\mathbb{D}_{0,t}^\nu(S(t)) &= \frac{1}{\Gamma(1-\nu)} \int_0^t (t-\psi)^{-\nu} \frac{d}{dt} S(\psi) d\psi = -\beta^\nu \frac{SI}{N}, \\ {}^C\mathbb{D}_{0,t}^\nu(E(t)) &= \frac{1}{\Gamma(1-\nu)} \int_0^t (t-\psi)^{-\nu} \frac{d}{dt} E(\psi) d\psi = \beta^\nu \frac{SI}{N} - \gamma^\nu I, \\ {}^C\mathbb{D}_{0,t}^\nu(R(t)) &= \frac{1}{\Gamma(1-\nu)} \int_0^t (t-\psi)^{-\nu} \frac{d}{dt} R(\psi) d\psi = \gamma^\nu I, \end{aligned} \quad (5)$$

where ${}^C\mathbb{D}_{0,t}^\nu$ stands for the Caputo (non-integer order) differential operator with order of the system (5) to be $\nu \in (0, 1)$.

3 Existence and Uniqueness Results

Among the vital areas in the concept of noninteger-order differential equations is the concept of existence and the uniqueness of solutions in a dynamical system. Recently, the theory has attracted many researchers' attention [36]. By means of fixed point theorem, we report the existence and uniqueness of (5). Take into account the proposed model (5) in the form as comes next:

$$\begin{cases} {}^C\mathcal{D}_{0+}^\nu S(t) = \Lambda_1(t, S, E, R), \\ {}^C\mathcal{D}_{0+}^\nu E(t) = \Lambda_2(t, S, E, R), \\ {}^C\mathcal{D}_{0+}^\nu R(t) = \Lambda_3(t, S, E, R), \end{cases} \quad (6)$$

with

$$\begin{cases} \Lambda_1(t, S, E, R) = -\beta^\nu \frac{SE}{N}, \\ \Lambda_2(t, S, E, R) = -\beta^\nu \frac{SE}{N} - \gamma^\nu E, \\ \Lambda_3(t, S, E, R) = \gamma^\nu E. \end{cases} \quad (7)$$

Therefore, (5) can accommodate the form

$$\begin{cases} {}^C\mathcal{D}_0^\alpha \Upsilon(t) = \mathcal{R}(t, \Upsilon(t)); & t \in J = [0, b], \quad 0 < \alpha \leq 1, \\ \Upsilon(0) = \Upsilon_0 \geq 0, \end{cases} \quad (8)$$

only if

$$\begin{cases} \Upsilon(t) = (S, E, R)^T, \\ \Upsilon(0) = (S_0, E_0, R_0)^T, \\ \mathcal{R}(t, \Upsilon(t)) = (\Theta_i(t, S, E, R))^T, \quad i = 1, 2, 3, \end{cases} \quad (9)$$

with T representing the transpose. Now, we can write (8) as

$$\begin{aligned} \Upsilon(t) &= \Upsilon_0 + \mathcal{J}_{0+}^{\nu} \mathcal{R}(t, \Upsilon(t)) \\ &= \Upsilon_0 + \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} \mathcal{R}(\kappa, \Upsilon(\kappa)) d\kappa. \end{aligned} \quad (10)$$

Consider the Banach space on $[0, b]$, with $[0, b]$ denoting a set of continuous functions from on $\mathbb{R}$ with the associated norm given by $\|\Upsilon\| = \sup_{t \in \mathcal{J}} |\Upsilon|$ be $\mathbb{E} = C([0, b]; \mathbb{R})$. Also, take into account the following theorem: .

Theorem 3.1. *Surmise that $\mathcal{R} \in C([\mathcal{J}, \mathbb{R}])$ and also maps $\mathcal{J} \times \mathbb{R}^5$ (bounded-subset) to relatively $\mathbb{R}$ (compact subsets). Additionally, $\exists$ a constant $L_{\mathcal{R}} > 0$ so that*

(A₁) $|\mathcal{R}(t, \Upsilon_1(t)) - \mathcal{R}(t, \Upsilon_2(t))| \leq L_{\mathcal{R}} |\Upsilon_1(t) - \Upsilon_2(t)|; \forall t \in \mathcal{J}$ and each $\Upsilon_1, \Upsilon_2 \in C([\mathcal{J}, \mathbb{R}])$. (10) which is an equivalence of (5) accomodates a unique solution only if $\Omega L_{\mathcal{R}} < 1$, and

$$\Omega = \frac{b^{\nu}}{\Gamma(\nu + 1)}.$$

Proof. Take into account $P : \mathbb{E} \rightarrow \mathbb{E}$ expressed as

$$(P\Upsilon)(t) = \Upsilon_0 + \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} \mathcal{R}(\kappa, \Upsilon(\kappa)) d\kappa. \quad (11)$$

(11) shows that the unique solution for (5) represents the fixed point of P . Moreover, take $\sup_{t \in \mathcal{J}} \|\mathcal{R}(t, 0)\| = M_1$ and $\kappa \geq \|\Upsilon_0\| + \Omega M_1$. Therefore, it will be suffices to verify $P\mathbb{H}_{\kappa} \subset \mathbb{H}_{\kappa}$, and the set is given by

$\mathbb{H}_\kappa = \{\Upsilon \in \mathbb{E} : \|\Upsilon\| \leq \kappa\}$, is convex and closed. Now, for any $\Upsilon \in \mathbb{H}_\kappa$, we have

$$\begin{aligned}
|(P\Upsilon)(t)| &\leq |\Upsilon_0| + \frac{1}{\Gamma(\nu)} \int_0^t (t-\kappa)^{\nu-1} |\mathcal{R}(\kappa, \Upsilon(\kappa))| d\kappa \\
&\leq \Upsilon_0 + \frac{1}{\Gamma(\nu)} \int_0^t (t-\kappa)^{\nu-1} [|\mathcal{R}(\kappa, \Upsilon(\kappa)) - \mathcal{R}(\kappa, 0)| + |\mathcal{R}(\kappa, 0)|] d\kappa \\
&\leq \Upsilon_0 + \frac{(\mathcal{L}_\mathcal{R}\kappa + M_1)}{\Gamma(\nu)} \int_0^t (t-\kappa)^{\nu-1} d\kappa \\
&\leq \Upsilon_0 + \frac{(\mathcal{L}_\mathcal{R}\kappa + M_1)}{\Gamma(\nu+1)} b^\nu \\
&\leq \Upsilon_0 + \Omega(\mathcal{L}_\mathcal{R}\kappa + M_1) \\
&\leq \kappa,
\end{aligned} \tag{12}$$

which affirms the result. Further, for a given $\Upsilon_1, \Upsilon_2 \in \mathbb{E}$, one reaches

$$\begin{aligned}
|(P\Upsilon_1)(t) - (P\Upsilon_2)(t)| &\leq \frac{1}{\Gamma(\nu)} \int_0^t (t-\kappa)^{\nu-1} |\mathcal{R}(\kappa, \Upsilon_1(\kappa)) - \mathcal{R}(\kappa, \Upsilon_2(\kappa))| d\kappa \\
&\leq \frac{\mathcal{L}_\mathcal{R}}{\Gamma(\nu)} \int_0^t (t-\kappa)^{\nu-1} |\Upsilon_1(\kappa) - \Upsilon_2(\kappa)| d\kappa \\
&\leq \Omega \mathcal{L}_\mathcal{R} |\Upsilon_1(t) - \Upsilon_2(t)|,
\end{aligned} \tag{13}$$

indicating that $\|(P\Upsilon_1) - (P\Upsilon_2)\| \leq \Omega \mathcal{L}_\mathcal{R} \|\Upsilon_1 - \Upsilon_2\|$. Thus, as a consequence of Banach contraction rule, (5) possess a unique solution on $\mathcal{J}$.

Now, we want show the existence of solutions of (5) by means Schauder fixed point principle.

Lemma 3.2. *Take $M \neq \emptyset$ as a bounded, convex and closed subset of a Banach Space $\mathbb{E}$. Consider P_1, P_2 as an operators in obideince of the following:*

- $P_1\Upsilon_1 + P_2\Upsilon_2 \in M$, when $\Upsilon_1, \Upsilon_2 \in M$;
- P_1 is continuous and compact;
- P_2 is a contraction mapping.

Then $\exists u \in M$ so that $u = P_1 u + P_2 u$.

Theorem 3.3. *Surmise that $\mathcal{R} : \mathcal{J} \times \mathbb{R}^5 \rightarrow \mathbb{R}$ is continuous and satisfies condition (A_1) . Additionally, take into account that*

$$(A_2) \quad |\mathcal{R}(t, \Upsilon)| \leq \mathcal{K}(t), \text{ for all } (t, \Upsilon) \in \mathcal{J} \times \mathbb{R}^5 \text{ and } \mathcal{K} \in C([0, b], \mathbb{R}_+).$$

Then (5) possesses at least one solution on $\mathcal{J}$ only if

$$\mathcal{L}_K \|\Upsilon_1(t_0) - \Upsilon_2(t_0)\| < 1.$$

Proof. Considering $\sup_{t \in \mathcal{J}} |\mathcal{K}(t)| = \|\mathcal{K}\|$ and $\zeta \geq \|\Upsilon_0\| + \Omega \|\mathcal{K}\|$, with $\mathbf{B}_\zeta = \{\Upsilon \in \mathbb{E} : \|\Upsilon\| \leq \zeta\}$.

Take into account P_1, P_2 on $\mathbf{B}_\zeta$ given by

$$(P_1 \Upsilon)(t) = \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} \mathcal{R}(\kappa, \Upsilon(\kappa)) d\kappa \quad t \in \mathcal{J},$$

and

$$(P_2 \Upsilon)(t) = \Upsilon(t_0), \quad t \in \mathcal{J}.$$

Therefore, $\forall \Upsilon_1, \Upsilon_2 \in \mathbf{B}_\zeta$, we have

$$\begin{aligned} \|(P_1 \Upsilon_1)(t) + (P_2 \Upsilon_2)(t)\| &\leq \|\Upsilon_0\| + \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} \|\mathcal{R}(\kappa, \Upsilon_1(\kappa))\| d\kappa \\ &\leq \|\Upsilon_0\| + \Omega \|\mathcal{K}\| \\ &\leq \zeta < \infty. \end{aligned} \tag{14}$$

Thus, $P_1 \Upsilon_1 + P_2 \Upsilon_2 \in \mathbf{B}_\zeta$.

Now, the contraction of P_2 will be proved.

Given any $t \in \mathcal{J}$ and $\Upsilon_1, \Upsilon_2 \in \mathbf{B}_\zeta$, yields

$$\|(P_2 \Upsilon_1)(t) - (P_2 \Upsilon_2)(t)\| \leq \|\Upsilon_1(t_0) - \Upsilon_2(t_0)\|. \tag{15}$$

Having $\mathcal{R}$ as a continuous function, P_1 is continuous. Furthermore, $\forall t \in \mathcal{J}$ and $\Upsilon_1 \in \mathbf{B}_\zeta$,

$$\|P_1 \Upsilon\| \leq \Omega \|\mathcal{K}\| < +\infty,$$

P_1 is uniformly bounded. At last, we prove that P_1 is compact. Start with $\sup_{(t, \Upsilon) \in \mathcal{J} \times \mathbf{B}_\zeta} |\mathcal{R}(\kappa, \Upsilon(\kappa))| = \mathcal{R}^*$, yields

$$\begin{aligned} |(P_1 \Upsilon)(t_2) - (P_1 \Upsilon)(t_1)| &= \frac{1}{\Gamma(\nu)} \left| \int_0^{t_1} [(t_2 - \kappa)^{\nu-1} - (t_1 - \kappa)^{\nu-1}] \mathcal{R}(\kappa, \Upsilon(\kappa)) d\kappa \right. \\ &\quad \left. + \int_{t_1}^{t_2} (t_2 - \kappa)^{\nu-1} \mathcal{R}(\kappa, \Upsilon(\kappa)) d\kappa \right| \\ &\leq \frac{\mathcal{R}^*}{\Gamma(\nu)} [2(t_2 - t_1)^\nu + (t_2^\nu - t_1^\nu)] \\ &\rightarrow 0, \text{ as } t_2 \rightarrow t_1. \end{aligned} \tag{16}$$

Hence, P_1 is equicontinuous and so is relatively compact on $\mathbf{B}_\zeta$. In accordance with Arzelá Ascoli principle, P_1 is compact on $\mathbf{B}_\zeta$. Having satisfied all the claims of Lemma 3.2, (5) possess at least on solution on $\mathcal{J}$.

4 Stability Analysis

Here, the stability of (5) will be scrutinized the aspect of Ulam-Hyers and generalized Ulam-Hyers [37, 38]. It has been proved that, stability analysis is important for an approximate solution. Assume $\varepsilon > 0$ and take into account the inequality

$$|^C \mathcal{D}_{0+}^\nu \bar{\Upsilon}(t) - \mathcal{R}(t, \bar{\Upsilon}(t))| \leq \varepsilon, \quad t \in J. \tag{17}$$

and $\varepsilon = \max(\varepsilon_j)^T$, $j = 1, \dots, 5$.

Definition 4.1. (8) which is the equivalence of (5), is Ulam-Hyers stable if $\exists \mathcal{X}_\mathcal{R} > 0$, so that $\forall$

$\varepsilon > 0$ and a solution $\bar{\Upsilon} \in \mathbb{E}$ holding for (4.1), there is the unique solution $\Upsilon \in \mathbb{E}$ for (8), with

$$|\bar{\Upsilon}(t) - \Upsilon(t)| \leq \mathcal{X}_{\mathcal{R}}\varepsilon, \quad t \in J,$$

where $\mathcal{X}_{\mathcal{R}} = \max(\mathcal{X}_{\mathcal{R}_j})^T$.

Definition 4.2. (8), which is the equivalence with (5), is referred to generalized Ulam-Hyers stable if $\exists$ a continuous function $\vartheta_{\mathcal{R}} : \mathbb{R}_+ \rightarrow \mathbb{R}_+$, with $\vartheta_{\mathcal{R}}(0) = 0$, so that $\forall$ solution $\bar{\Upsilon} \in \mathbb{E}$ of (17), there is the unique solution $\Upsilon \in \mathbb{E}$ for (8), so that

$$|\bar{\Upsilon}(t) - \Upsilon(t)| \leq \vartheta_{\mathcal{R}}\varepsilon, \quad t \in J,$$

where $\vartheta_{\mathcal{R}} = \max(\vartheta_{\mathcal{R}_j})^T$.

Remark 4.3. A function $\bar{\Upsilon} \in \mathbb{E}$ satisfies (17), if and only if $\exists$ a function $h \in \mathbb{E}$ with the property as comes next:

$$(i) \quad |h(t)| \leq \varepsilon, \quad h = \max(h_j)^T, \quad t \in J;$$

$$(ii) \quad {}^C\mathcal{D}_{0+}^{\nu} \bar{\Upsilon}(t) = \mathcal{R}(t, \bar{\Upsilon}(t)) + h(t), \quad t \in J.$$

Lemma 4.4. Surmise that $\bar{\Upsilon} \in \mathbb{E}$ holds for (17), then $\bar{\Upsilon}$ holds for the following:

$$\left| \bar{\Upsilon}(t) - \bar{\Upsilon}_0 - \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} \mathcal{R}(\kappa, \bar{\Upsilon}(\kappa)) d\kappa \right| \leq \Omega\varepsilon. \quad (18)$$

Proof. By utilizing (ii) of 4.3,

$${}^C\mathcal{D}_{0+}^{\nu} \bar{\Upsilon}(t) = \mathcal{R}(t, \bar{\Upsilon}(t)) + h(t),$$

and Lemma 4.4, we have that

$$\bar{\Upsilon}(t) = \bar{\Upsilon}_0 + \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} \mathcal{R}(\kappa, \bar{\Upsilon}(\kappa)) d\kappa + \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} h(\kappa) d\kappa. \quad (19)$$

Considering (i) of 4.3, yields

$$\begin{aligned} \left| \bar{\Upsilon}(t) - \bar{\Upsilon}_0 - \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} \mathcal{R}(\kappa, \bar{\Upsilon}(\kappa)) d\kappa \right| &\leq \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} |h(\kappa)| d\kappa \\ &\leq \Omega \varepsilon. \end{aligned} \quad (20)$$

Hence, we complete the result.

Theorem 4.5. *Surmise that $\mathcal{R} : J \times \mathbb{R}^5 \rightarrow \mathbb{R}$ is continuous $\forall \Upsilon \in \mathbb{E}$ and assumption (A_1) holds with $1 - \Omega \mathcal{L}_{\mathcal{R}} > 0$. Then, (8) which is the equivalence of (5) is Ulam-Hyers and consequently, generalized Ulam-Hyers stable.*

Proof. Surmise that $\bar{\Upsilon} \in \mathbb{E}$ holds for (17) and $\Upsilon \in \mathbb{E}$ be a unique solution (8). Therefore, $\forall \varepsilon > 0, t \in J$ and Lemma 4.4, yields

$$\begin{aligned} |\bar{\Upsilon}(t) - \Upsilon(t)| &= \max_{t \in \mathcal{J}} \left| \bar{\Upsilon}(t) - \Upsilon_0 - \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} \mathcal{R}(\kappa, \Upsilon(\kappa)) d\kappa \right| \\ &\leq \max_{t \in \mathcal{J}} \left| \bar{\Upsilon}(t) - \bar{\Upsilon}_0 - \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} \mathcal{R}(\kappa, \bar{\Upsilon}(\kappa)) d\kappa \right| \\ &\quad + \max_{t \in \mathcal{J}} \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} |\mathcal{R}(\kappa, \bar{\Upsilon}(\kappa)) - \mathcal{R}(\kappa, \Upsilon(\kappa))| d\kappa \\ &\leq \left| \Upsilon(t) - \bar{\Upsilon}_0 - \frac{1}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} \mathcal{R}(\kappa, \bar{\Upsilon}(\kappa)) d\kappa \right| \\ &\quad + \frac{\mathcal{L}_{\mathcal{R}}}{\Gamma(\nu)} \int_0^t (t - \kappa)^{\nu-1} |\bar{\Upsilon}(\kappa) - \Upsilon(\kappa)| d\kappa \\ &\leq \Omega \varepsilon + \Omega \mathcal{L}_{\mathcal{R}} |\bar{\Upsilon}(t) - \Upsilon(t)|. \end{aligned}$$

So,

$$\|\bar{\Upsilon} - \Upsilon\| \leq \mathcal{X}_{\mathcal{R}} \varepsilon,$$

where

$$\mathcal{X}_{\mathcal{R}} = \frac{\Omega}{1 - \Omega \mathcal{L}_{\mathcal{R}}}.$$

Equating $\vartheta_{\mathcal{R}}(\varepsilon) = \mathcal{X}_{\mathcal{R}}\varepsilon$ so that $\vartheta_{\mathcal{R}}(0) = 0$. One concludes that the (5) is stable for both Ulam-Hyers and generalized Ulam-Hyers.

5 Estimation of Parameters

A disease model like the one being investigated in the present research study, is widely accepted and validated if its simulations for infectious compartment agree well with those of available real cases of the disease. In order to do so, one needs to obtain values of the parameters of the model in a best possible way so that they cause the model to be in good agreement with the available data. This brings into attention a few existing methods for getting best values of such parameters. Bayesian technique, probability plotting, maximum likelihood estimation and least squares method are among the methods that exist for parameter estimation. In this study, we have used least squares method to compute the parameters (β = transmission rate and γ = recovery rate) of the model (5). Since, there are only two non-demo-graphical parameters for the proposed model, they are best fitted using the real cases of COVID-19 pandemic throughout Pakistan (source <http://covid.gov.pk/stats/pakistan>). Daily cases of the disease are taken from April 01 - - May 15, 2020 when the present research study was being conceived and prepared.

In the least-squares approach, the objective function requires to be minimized. The objective is based upon adjusting the parameters for a model function to fit a data set in a best possible way. A simple data set consists of n points (data pairs) $(x_k, y_k), k = 1, \dots, n$, where x_k shows an independent variable and y_k stands for dependent variable whose value is computed using observation. The model function possesses the form $g(x, p)$, where m adjustable parameters are kept in the vector p . The goal is to notice the parameters for the model that fits the data in best possible way. Such a fit of a

model to a data point is measured by its residual, defined as the difference between the real available value of the dependent variable and the value predicted by the model:

$$r_k = y_k - g(x_k, p) \quad (21)$$

The least-squares method obtains the optimal parameter values by minimizing the sum of squared residuals as shown below:

$$S = \sum_{k=1}^n (r_k)^2 = \sum_{k=1}^n (y_k - g(x_k, p))^2 \quad (22)$$

For initial conditions, the overall population of Pakistan is found as $N(0) = 212.2\text{M}$, the initial susceptible population is estimated to be $S(0) = 212197711$, the initial exposed population is estimated to be $E(0) = 2289$ and the initial recovered population is estimated to be $R(0) = 0$. There are only two biological parameters best fitted via least square fitting method thereby yielded best fit of the model's solution to the real cases chosen from Pakistan as depicted by the Figure 1 under the classical situation, that is, when $\nu = 1$ wherein the best parameters are as follows: $\beta = 3.0918$ and $\gamma = 3.0190$ for the transmission and recovery rate, respectively. The average absolute relative error between the real cases and the model's simulations for the infectious compartment is decreased. Such a value for the error is approximately $3.3603e - 02$. The Figure 1 shows the real cases by black circles while the best fitted curve is shown by the blue solid line and the residuals are shown by side. The biological parameters included in the model are listed in the Table 1 along with their best estimated values obtained via least-squares technique. These parameters have finally produced the value of the basic reproduction number equal to $\mathcal{R}_0 = 1.0241$ for the real COVID-19 cases under the classical situation in Pakistan from April 01 - - May 15, 2020. Moreover, similar analysis for the Caputo fractional order model is carried out wherein the best fitted parameters are found as: $\beta = 3.9405$ and $\gamma = 3.8010$ for the transmission and recovery rate, respectively. One distinguished feature in the present research study is the computation of optimum value of the fractional order parameter ν which is $8.561591e-01$. Hence, the basic reproduction number is obtained as $\mathcal{R}_0 = 1.0313$ and the corresponding average

absolute relative error is $1.7360e-02$ for the Caputo model whereas $3.3603e-02$ is the average absolute relative error when $\nu = 1$. This clearly reveals the superiority of the fractional operator over the classical one. The Figure 2 shows the real cases by black circles while the best fitted curve obtained under the Caputo fractional case is shown by the blue solid line and the residuals are shown by side.

Table 1: Best fitted parameters employed in the model (5) for its classical $\nu = 1$ and fractional $\nu = 8.561591e - 01$ version.

Parameter	Description	Value		Sources
β	Transmission rate	3.0918^C	3.9405^F	Fitted
γ	Recovery rate	3.0190^C	3.8010^F	Fitted

Figure 1: (a) Best data fitting for the COVID-19 daily cases (black circles) in Pakistan from April 01 to May 15, 2020 with the infectious class $I(t)$ (solid blue line) of the model under the classical case ($\nu = 1$) and the (b) respective residuals.

Figure 2: Best data fitting for the COVID-19 daily cases (black circles) in Pakistan from April 01 to May 15, 2020 with the infectious class $I(t)$ (solid blue line) of the model under the Caputo operator ($\nu = 8.561591e - 01$) and the (b) respective residuals.

6 Sensitivity Analysis

In this portion, the concept of sensitivity analysis is employed to discover the robust significance of the generic parameters that are present in the basic reproduction number $\mathcal{R}_0$. Further, with the aid of parameter values from reliable assumptions, both analytic and numerical values of the parameters in $\mathcal{R}_0$ are obtained. The analytic expressions obtained can be used to shed some light on how to control the onset of the model in variant localities, if and only if the dynamics follow model (1). The threshold value $\mathcal{R}_0$ is a quantity of which lowering the number to less than unity is considered as the major way of curtailing and aborting the spread of the ailment. The sensitivity index technique is used to measure the most sensitive parameters in the model, those with positive sign are considered as highly and proportionally sensitive for increasing the value of $\mathcal{R}_0$ while those with negative sign are less sensitive for the decrease of $\mathcal{R}_0$ and the other category are neutrally sensitive (with zero relative sensitivity). It is popularly known that, the cause of rapid transmission is associated directly with

the basic reproduction number $\mathcal{R}_0$. The elasticity indices of $\mathcal{R}_0$ to the associated parameters in the model are defined as [39]:

$$\Upsilon_{\mathbf{P}_i}^{\mathcal{R}_0} = \frac{\partial \mathcal{R}_0}{\partial \mathbf{P}_i} \times \frac{\mathbf{P}_i}{\mathcal{R}_0}, \quad (23)$$

where $\mathcal{R}_0$ denotes the basic reproduction ratio and $\mathbf{P}_i$ is as stated above. $\mathcal{R}_0$ for the model (5) under consideration is defined by the following expression:

$$\mathcal{R}_0 = \left(\frac{\beta}{\gamma} \right)^\nu. \quad (24)$$

Thus, after some computation we reach:

$$\begin{aligned} \Upsilon_\beta &= \nu, \\ \Upsilon_\gamma &= -\nu. \end{aligned} \quad (25)$$

Table 2: The elasticity indices for $\mathcal{R}_0 = 1.0313$ to the parameters of the model (5).

Parameter	Baseline value	Elasticity index
β	3.9405 ^F	8.561591e − 01
γ	3.8010 ^F	−8.561591e − 01

The numerical values showing the relative significance of the R_0 parameters are given in table 2. Positive relationship parameter is β , while negative-relations is γ . A negative relationship indicates that an increase in this parameters' value would help to reduce the brutality of the disease. While a positive relationship indicates that an increase in the values of that parameter would have a substantial effect on the frequency of the spread of the ailment. Figure 3(a) and 3(b) display the numerical signs stated in the Table 2.

(a)

(b)

Figure 3: Normalized Local Sensitivity Indices of $\mathcal{R}_0$ for each parameter

7 Numerical Simulations

This is the place wherein we get deep insights into the model's dynamical behavior. The present section offers the numerical simulations of the Caputo model (5) while using the biological parameters as listed in the Table 1. We have employed the FDE12 technique in order to get the solution of the nonlinear model shown in the Equations (5) and achieved the graphical results based upon parameters which are taken to be as variables. Susceptible $S(t)$, infectious $I(t)$ and recovered $R(t)$ populations are investigated with different values of the parameters. It may also be noted that the optimized value of the fractional order $\nu = 8.561591e - 01$ is used during each simulation. As shown in (a) plot of the Figure 4 that the susceptible individuals decrease as the transmission rate increases, on the contrast, the susceptible individuals decrease as the recovery rate decreases as shown by (b) plot of the Figure 4. Similarly, the slightly increased value of the transmission rate β is responsible for spread of the virus (Figure 5 (a)) and the number of infectious individuals decrease when there is an improvement in their recovery which is observed in (b) plot of the Figure 5. Surprisingly, increasing value of β causes an increase in the recovered population after a certain level of time interval as shown in (a) plot of the Figure 6 and similar surprising behavior is noted in (b) plot of the Figure 6. However, this should be obvious since with rapid transmission rate there will be more effective strategies used by public health sectors and concerned government to overcome the situation leading to an increase in the recovered population with similar sort of explanation for (b) plot of the Figure 6.

We have also investigated effects of fractional order ν on the dynamical behavior of the Caputo model (5) for all three compartments. As can be seen in the Figure 7 that the susceptible population decreases if ν increases whereas the Figure 8 shows an increase in the infectious population for increasing values of ν and this makes sense because susceptible population, after being infected, is shifted to I compartment thereby leading to an increase therein. It is observed in the Figure 9 that the recovered population propels forward if ν gets large. Finally, we have also shown dynamics of the basic reproduction number $\mathcal{R}_0$ under different effects of the transmission rate β and recovery rate γ

in the Figure 10 wherein even a slightly large value of β brings the reproduction number near to 1 which is clearly an alarming situation for policy makers to devise an effective approach to prevent $\mathcal{R}_0$ to be greater than 1.

Figure 4: Behavior of susceptible population under varying values of β and γ .

Figure 5: Behavior of infectious population under varying values of β and γ .

Figure 6: Behavior of recovered population under varying values of β and γ .

Figure 7: Behavior of susceptible population for different values of the fractional order parameter ν .

Figure 8: Behavior of infectious population for different values of the fractional order parameter ν .

Figure 9: Behavior of recovered population for different values of the fractional order parameter ν .

Figure 10: Behavior of basic reproductive number.

8 Conclusion

Using Caputo differential operator famous to have non-local nature which is perfectly suitable to investigate transmission dynamics of a disease, we have fractionalized SIR epidemic model having order ν without violating dimensional consistency between parameters and the operator itself. Proposed Caputo SIR model is used to comprehend behavior of devastating disease of COVID-19 recently emerged in the entire community of humankind. The model has shown promising results for it is proved to have a unique solution on the basis of Banach contraction principle. Using Ulam-Hyers and its generalized version, the Caputo model is shown to be stable. Fitted parameters of the model, using real incidence cases of the virus from April 01 to May 15, 2020, are obtained under least-squares approach wherein the fractional order ν is also optimized to be $8.561591e - 01$: one of the major contributions in the present work. Using these values, it has been proved that the Caputo model outperforms its classical version by 48% while the basic reproductive number is $\mathcal{R}_0 = 1.0313$. Sensitivity of the parameters β and γ relating with $\mathcal{R}_0$ is also investigated. Various numerical simulations carried out suggest that the epidemic can effectively be controlled only if its contact rate is lowered and this is possible when people observe social distancing, and wear protective masks. On the other hand, an under-developed country like Pakistan will be in huge trouble if such measures are not strictly followed upon. Our future study will explore the effects of non-singular differential operators on the standard SIR model.

Availability of data and materials

We used the real data for the cases chosen from 01 April to 15 March, 2020, Pakistan which is freely available under <http://covid.gov.pk/stats/pakistan>.

Acknowledgements

Not applicable.

Funding

Not applicable.

Competing interests

The authors declare that they have no competing interests regarding the publication of this paper.

Authors' contribution

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript

Authors' details

¹Department of Mathematics, King Abdul Aziz University, Jeddah, Saudi Arabia. ^{2*}Department of Mathematics, Cankaya University, Ankara, Turkey. ³Institute of Space Sciences, Magurele-Bucharest, Romania.

References

- [1] Djordjevic, J.; Silva, C.J.; Torres, D.F.M. A stochastic SICA epidemic model for HIV transmission. *Applied Mathematics Letters* 2018, 84, 168–175.

- [2] Ndairou, F.; Area, I.; Nieto, J.J.; Silva, C.J.; Torres, D.F.M. Mathematical modeling of Zika disease in pregnant women and newborns with microcephaly in Brazil. *Mathematical Methods in the Applied Sciences* 2018, 41, 8929–8941. doi:10.1002/mma.4702.
- [3] Rachah, A.; Torres, D.F.M. Dynamics and optimal control of Ebola transmission. *Mathematics in Computer Science* 2016, 10, 331–342. doi:10.1007/s11786-016-0268-y.
- [4] Brauer, F.; Castillo-Chavez, C.; Feng, Z. *Mathematical Models in Epidemiology*; Springer- Verlag New York, 2019.
- [5] Coronavirus disease (COVID-2019) situation reports, World Health Organization. <https://www.who.int/317emergencies/diseases/novel-coronavirus-2019/situation-reports/>. Assessed 2 August 2020
- [6] He, S., Peng, Y. and Kehui, K. SEIR modeling of the COVID-19 and its dynamics. *Nonlinear Dynamics*, (2020):1-14.
- [7] Fanelli, D. and Piazza, F. Analysis and forecast of COVID-19 spreading in China, Italy and France. *Chaos, Solitons and Fractals*, **134**(2020):1-5.
- [8] A. Atangana. Modelling the spread of COVID-19 with new fractal-fractional operators: Can the lockdown save mankind before vaccination ?, *Chaos, Solitons and Fractals* **136**(2020):1-38
- [9] Shaikh, A.S., Jadhav, K.S., Timol, V.S., Nisar, M.G. and Khan I. Analysis of the COVID-19 Pandemic Spreading in India by an Epidemiological Model and Fractional Differential Operator. *Preprints*, 2020:1-16.
- [10] Naveed, M., Rafiq, M., Raza, A., Ahmed, N., Khan, I., Nisar, K.S., Soori, A. H. Mathematical Analysis of Novel Coronavirus (2019-nCov) Delay Pandemic Model, *CMC-Computers, Materials Continua*, 64(3), 1401–1414, doi:10.32604/cmc.2020.011314

- [11] Shaikh, A.S., Shaikh, I.N. Nisar, K.S. A mathematical model of COVID-19 using fractional derivative: outbreak in India with dynamics of transmission and control. *Advances in Difference Equation* 2020, 373 (2020). <https://doi.org/10.1186/s13662-020-02834-3>
- [12] I. Ahmed, Isa A. Baba, A. Yusuf, P. Kumam, W. Kumam. Analysis of Caputo fractional-order model for COVID-19 with lockdown. *Advances in Difference Equations* (2020) 2020:394 <https://doi.org/10.1186/s13662-020-02853-0>
- [13] Wikipedia Website. <https://en.wikipedia.org/wiki/Coronavirus/>
- [14] Qun Li, Xuhua Guan, Peng Wu, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus Infected Pneumonia. *New England Journal of Medicine*, 2020.doi: 10.1056/NEJMoa2001316.
- [15] Weijie Guan, Zhengyi Ni, Yu Hu, et al. Clinical characteristics of 2019 novel coronavirus infection in China. *medRxiv*,2020.doi: 10.1056/NEJMoa2002032.
- [16] Yang Yang, Qingbin Lu, Mingjin Liu, et al. Epidemiological and clinical features of the 2019 novel coronavirus outbreak in China. *medRxiv*, 2020.doi: <https://doi.org/10.1101/2020.02.10.20021675>.
- [17] Diagnosis and treatment of novel coronavirus pneumonia. (trial version Sixth, in Chinese.)
- [18] Wrapp D, Wang N, Corbett K, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science*, 2020.doi: 10.1126/science.abb2507.
- [19] Rothe C, Schunk M, Sothmann P, et al. Transmission of 2019-nCoV infection from an asymptomatic contact in Germany. *New England Journal of Medicine*, 2020.doi: 10.1056/NEJMc2001468.
- [20] J. Singh, D. Kumar, D. Baleanu, On the analysis of fractional diabetes model with exponential law, *Adv. Differential Equations* 2018 (2018) 231.

- [21] S. Qureshi, A. Yusuf. Modeling chickenpox disease with fractional derivatives: From caputo to atangana-baleanu. *Chaos, Solitons and Fractals* 122 (2019) 111–118.
- [22] S. Qureshi, A. Yusuf. Mathematical modeling for the impacts of deforestation on wildlife species using Caputo differential operator. *Chaos, Solitons and Fractals* 126 (2019) 32–40.
- [23] A. Jajarmi, A. Yusuf, D. Baleanu, M. Inc. A new fractional HRSV model and its optimal control: A non-singular operator approach. *Physica A* 547 (2020) 123860.
- [24] M. Abdulhameed, M.M. Muhammad, A.Y. Gital, D.G. Yakubu, I. Khan, Effect of fractional derivatives on transient MHD flow and radiative heat transfer in a micro-parallel channel at high zeta potentials, *Physica A* 519 (2019) 42–71.
- [25] V.P. Dubey, R. Kumar, D. Kumar, Analytical study of fractional Bratu-type equation arising in electro-spun organic nanofibers elaboration, *Physica A* 521 (2019) 762–772.
- [26] A. Chang, H. Sun, Y. Zhang, C. Zheng, F. Min, Spatial fractional Darcy’s law to quantify fluid flow in natural reservoirs, *Physica A* 519 (2019) 119–126.
- [27] A.G. Goulart, M.J. Lazo, J.M.S. Suarez, A new parameterization for the concentration flux using the fractional calculus to model the dispersion of contaminants in the Planetary Boundary Layer, *Physica A* 518 (2019) 38–49.
- [28] Jajarmi, A., Ghanbari, B., & Baleanu, D. (2019). A new and efficient numerical method for the fractional modeling and optimal control of diabetes and tuberculosis co-existence. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 29(9), 093111.
- [29] Baleanu, D., Aydogan, S. M., Mohammadi, H., & Rezapour, S. (2020). On modelling of epidemic childhood diseases with the Caputo-Fabrizio derivative by using the Laplace Adomian decomposition method. *Alexandria Engineering Journal*.

- [30] Iqbal, Z., Ahmed, N., Baleanu, D., Adel, W., Rafiq, M., Rehman, M. A. U., & Alshomrani, A. S. (2020). Positivity and boundedness preserving numerical algorithm for the solution of fractional nonlinear epidemic model of HIV/AIDS transmission. *Chaos, Solitons & Fractals*, 134, 109706.
- [31] Ameen, I., Baleanu, D., & Ali, H. M. (2020). An efficient algorithm for solving the fractional optimal control of SIRV epidemic model with a combination of vaccination and treatment. *Chaos, Solitons & Fractals*, 137, 109892.
- [32] Kumar, S., Ahmadian, A., Kumar, R., Kumar, D., Singh, J., Baleanu, D., & Salimi, M. (2020). An efficient numerical method for fractional SIR epidemic model of infectious disease by using Bernstein wavelets. *Mathematics*, 8(4), 558.
- [33] Raza, A., Rafiq, M., Baleanu, D., & Arif, M. S. (2020). Numerical simulations for stochastic meme epidemic model. *Advances in Difference Equations*, 2020(1), 1-16.
- [34] Arshad, S., Defterli, O., & Baleanu, D. (2020). A second order accurate approximation for fractional derivatives with singular and non-singular kernel applied to a HIV model. *Applied Mathematics and Computation*, 374, 125061.
- [35] Ullah, S., Khan, M. A., Farooq, M., Hammouch, Z., & Baleanu, D. (2020). A Fractional Model for the Dynamics of Tuberculosis Infection Using Caputo-Fabrizio Derivative.
- [36] Khan, H., Li, Y., Khan, A., and Khan, A. Existence of solution for a fractional-order lotka-volterra reaction-diusion model with mittag-leer kernel. *Mathematical Methods in the Applied Sciences* 42, 9 (2019), 3377-3387.
- [37] Ulam, S. M. A collection of mathematical problems, vol. 8. Interscience Publishers, 1960.
- [38] Ulam, S. M. Problems in modern mathematics. Courier Corporation, 2004.
- [39] J. K. K. Asamoah, Z. Jin, G. Q. Sun, and M. Y. Li. A Deterministic Model for QFever Transmission Dynamics within Dairy Cattle Herds: Using Sensitivity Analysis and Optimal Controls.

Figures

Figure 1

(a) Best data fitting for the COVID-19 daily cases (black circles) in Pakistan from April 01 to May 15, 2020 with the infectious class $I(t)$ (solid blue line) of the model under the classical case ($\nu = 1$) and the (b) respective residuals.

Figure 2

Best data fitting for the COVID-19 daily cases (black circles) in Pakistan from April 01 to May 15, 2020 with the infectious class $I(t)$ (solid blue line) of the model under the Caputo operator ($\nu = 8.561591 \times 10^{-1}$) and the (b) respective residuals.

A

B

Figure 3

Normalized Local Sensitivity Indices of R_0 for each parameter

Figure 4

Behavior of susceptible population under varying values of β and γ .

Figure 5

Behavior of infectious population under varying values of β and γ

Figure 6

Behavior of recovered population under varying values of β and γ .

Figure 7

Behavior of susceptible population for different values of the fractional order parameter ν .

Figure 8

Behavior of infectious population for different values of the fractional order parameter ν .

Figure 9

Behavior of recovered population for different values of the fractional order parameter ν .

Figure 10

Behavior of basic reproductive number.