Integrated miRNA-mRNA Expression Profile Analysis Reveals a Role of miR-146a-3p/TRAF6 in Plasma from Gestational Diabetes Mellitus Patients

Min Chen

Fujian Medical University

Jianying Yan (✉ jyy363@163.com)

Fujian Medical University

Research

Keywords: Gestational diabetes mellitus, molecular markers, signaling pathway

Posted Date: January 2nd, 2020

DOI: <https://doi.org/10.21203/rs.2.19652/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

BACKGROUND: By utilizing an integrative strategy, we screened noninvasive molecular markers for the early detection of GDM and constructed miRNA-mRNA regulatory networks associated with GDM.

METHODS: A total of 3 microarray datasets (GSE98043, GSE19649 and GSE92772) of plasma samples comparing GDM pregnant women and healthy control pregnant women were downloaded from the GEO database. The GEO2R online platform was used to identify the differentially expressed genes (DEmRNAs) and the differentially expressed miRNAs (DEmiRNAs). The target genes of DEmiRNAs were identified using two independent and complementary types of information: computational target predictions and expression relationships. KEGG pathway annotation was performed for target genes and DEmiRNAs. An interaction network was constructed to identify hub genes of GDM. Another dataset (GSE92772) was used to externally verify the predictive ability of the hub genes. Gene set enrichment analysis (GSEA) was performed to explore the biological function of hub genes.

RESULTS: A total of 264 DEmiRNAs and 1217 DEmRNAs were identified by comparing the microarray data of serum samples of GDM patients and healthy pregnant women. Hsa-miR-146a-3p ranked first because of its lowest P value and was selected for further analysis. A total of 47 target genes, including TRAF6, were shared between the computational target predictions and DEmRNAs and were identified as target genes of hsa-miR-146a-3p. Enrichment analysis indicated that GDM-related miRNAs were mainly enriched in the glypican pathway, proteoglycan syndecan-mediated signaling events, and syndecan-1-mediated signaling events. In addition, the glypican pathway was also one of the pathways regulated by hsa-miR-146a-3p. The interaction network analysis of DEmRNAs identified TRAF6, CASP8, PTPN6, and CHD3 as hub genes involved in the pathophysiological process of GDM. Next, these 4 hub genes were selected for independent external validation using the GSE19649 dataset. The expression of TRAF6, CASP8 and CHD3 in 8 pairs of GDM blood samples was confirmed to be higher than that in healthy pregnant women blood samples. However, the expression of PTPN6 in the blood samples of GDM patients was similar to that of healthy pregnant women blood samples. The AUC for predicting GDM was 0.813, 0.828, 0.813, and 0.703 for CHD3, PTPN6, and CASP8, respectively. GSEA showed that 9 hallmark gene sets of metabolism processes were enriched in TRAF6 function.

CONCLUSIONS: Three hub genes, TRAF6, CASP8, and CHD3, were identified and independently externally validated as potential GDM noninvasive serum markers. Enrichment analysis indicated that GDM-related miRNAs were mainly enriched in the glypican pathway, proteoglycan syndecan-mediated signaling events, and syndecan-1-mediated signaling events. In addition, integrated miRNA-mRNA expression profile analysis showed that miR-146a-3p/TRAF6 might play a central role in the pathogenesis of gestational diabetes mellitus by involving the above pathways.

1 | Background

In China, the incidence of gestational diabetes mellitus (GDM) has increased from approximately 5% to more than 16% since the implementation of diagnostic criteria for GDM in December 2011[1]. Compared with healthy pregnant women, pregnant women with GDM have a higher incidence of pre-eclampsia and a cumulative incidence of type 2 diabetes within 5–16 years after delivery[2]. In addition, the risk of macrosomia, abortion, hypoglycemia, cardiac dysplasia, obesity, and metabolic syndrome in infants of mothers with GDM is also increased[3]. Due to the high prevalence of GDM and its adverse outcomes, it is necessary to explore noninvasive markers for the early detection of GDM.

In addition to common risk factors, some new environmental factors, such as passive smoking[4] and excessive fruit intake in mid-pregnancy[5], significantly influence the predisposition to GDM. Genomic studies have also revealed that GDM is a multigenic disease. At present, many genes, including protein-coding RNA and noncoding RNA, are considered to be involved in the pathogenesis of GDM. MicroRNAs (miRNAs) are a class of small noncoding RNAs with 18–22 nucleotides in length[6]. They modulate gene expression at the posttranscriptional level by interacting with mRNA targets[7]. Serum levels of fibroblast growth factor 23 (FGF23) are upregulated in pregnant women with GDM compared with healthy pregnant women and could be a useful marker of cardiovascular disease in GDM[8]. Serum CysC is significantly and independently associated with insulin resistance, which is helpful in identifying the risk of GDM in Chinese pregnant women[9]. In addition, the combined detection of plasma microRNA-16-5p, -17-5p and -20a-5p is expected to provide early diagnosis for pregnant women with GDM[10]. However, these studies are only based on a single gene. To date, studies providing systematic insight into the composition and expression of the serum mRNA and miRNA transcriptome in GDM are still lacking. Therefore, in the present study, we utilized an integrative strategy to construct functional miRNA-mRNA regulatory networks by combining the reverse expression relationships between miRNAs and targets and computational predictions. Our study may provide clues to a better understanding of the mechanism of GDM and help in screening noninvasive molecular markers for the early detection of GDM.

2 | Methods

2.1 | Study samples and datasets

A total of 3 microarray datasets of plasma samples comparing GDM pregnant women and healthy control pregnant women were downloaded from the GEO database (<http://www.ncbi.nlm.nih.gov/geo/>). The GSE98043 dataset included 2 GDM pregnant women and 2 healthy control pregnant women. The platform used was GPL21575 (Agilent-070156 Human MicroRNA v21.0 Microarray 046064). The GSE19649 dataset includes plasma and placental mRNA microarray data. Since this study was designed to explore noninvasive plasma markers, we used only plasma mRNA data, which included the pooled blood tissues from 5 GDM women and one healthy control pregnant woman. The platform used was GPL7350 (Aalborg University Illumina Human-6 v2.0 Expression BeadChip). Another independent dataset, GSE92772, was used for validation of the expression of hub genes identified from the GSE19649 dataset[11]. The GSE92772 dataset included blood RNA-seq data from 8 pregnant GDM women and 8

healthy controls matched for age and BMI. Venous blood was collected during pregnancy at week 24–32. The platform used was GPL16791 (Illumina HiSeq 2500).

2.2 | Screening of molecular markers

The GEO2R online platform was used to identify the differentially expressed genes (DEmRNAs) in the GSE19649 dataset and the differentially expressed miRNAs (DEmiRNAs) in the GSE98043 dataset between serum samples of GDM pregnant women and healthy control pregnant women. GEO2R is an interactive web tool that allows gene expression analysis of microarray datasets using the limma R package[12]. $P < 0.05$ and fold change > 1.5 or < 0.5 were set as the cutoff criteria.

2.3 | Functional enrichment analysis

The target genes of microRNAs were predicted by the mirDIP online platform[13], and the target genes with high scores (top 5%) were selected for further analysis. The computationally predicted target genes and DEmRNAs were represented in a Venn diagram to identify real target genes. KEGG pathway annotation was performed for the real target genes and DEmiRNAs, respectively. All pathway analyses were performed using Funrich 3.1.3 software[14].

2.4 | Interaction network analysis

Funrich 3.1.3 software was used to systematically analyze and construct the interaction network of DEmRNAs. Then, the hub genes, defined by the genes with the largest number of connecting nodes, were identified.

2.5 | External hub gene verification

Another dataset (GSE92772) was used as a testing set to verify the differential expression of hub genes between GDM women and healthy controls. Continuous variables were compared using paired t tests. A receiver operating characteristic (ROC) curve analysis was generated, and the area under the curve (AUC) was calculated to evaluate the ability of the hub gene to predict GDM.

2.6 | Gene set enrichment analysis (GSEA)

To explore the biological function of hub genes, GSEA was performed in patients with high and low hub gene expression levels in the GSE92772 dataset. The high and low expression groups were separated by the median gene expression. Annotated gene sets c2.cp.kegg.v5.2.symbols. The GMT pathway database was used as the reference. $P < 0.05$ and |enrichment score (ES)| > 0.25 were set as the cutoff values.

A schematic overview of the proposed approach in this study is shown in Fig. 1.

3 | Results

By comparing the microarray data of serum samples of GDM patients and those of healthy pregnant women, 264 DEmiRNAs were identified in the GSE98043 dataset, of which 137 genes were upregulated and 127 genes were downregulated. In addition, a total of 1217 DEmRNAs were identified in the GSE19649 dataset. The top 10 DEmiRNAs and DEmRNAs are listed in Tables 1–2. Hsa-miR-146a-3p ranked first because of its lowest P value and was selected for further analysis.

Table 1
Top 10 DEmiRNAs

Probe ID	MicroRNA	Log(fold change)	P Value
9861	hsa-miR-146a-3p	-12.76	0.0002
55274	hsa-miR-146a-5p	10.82	0.0004
30074	hsa-miR-6831-3p	9.97	0.0008
12711	hsa-miR-1285-5p	9.99	0.0008
32345	hsa-miR-146b-3p	-5.54	0.0008
31670	hsa-miR-1911-5p	6.04	0.0009
17343	hsa-miR-4726-5p	8.89	0.0012
52744	hsa-miR-4690-5p	8.23	0.0012
29799	hsa-miR-548w	8.24	0.0012
56049	hsa-miR-4756-5p	-14.99	0.0012

Table 2
Top 10 DEmRNAs

Probe ID	MicroRNA	Log(fold change)	P Value
ILMN_4582	KRT2B	-2.41	0.000306
ILMN_31624	TRPC2	-2.45	0.000751
ILMN_25067	NR1H4	-2.46	0.000964
ILMN_28015	LGALS13	-2.47	0.000993
ILMN_24642	DENND1B	-2.47	0.001123
ILMN_17357	ZNF606	-2.48	0.001284
ILMN_26171	WDR64	-2.48	0.001289
ILMN_11089	AGRP	-2.49	0.001331
ILMN_1266	DSC1	-2.51	0.001709
ILMN_7518	DPYSL4	-2.51	0.001783

The target genes of hsa-miR-146a-3p were identified using two independent and complementary types of information: computational target predictions and expression relationships. For computational target predictions, a total of 1366 mRNAs were predicted to be the target genes of hsa-miR-146a-3p in the mirDIP online platform. These 1366 target genes were further mapped to the 1217 DEmRNAs screened from the GSE19649 dataset. Finally, a total of 47 target genes were shared between the computational target predictions and DEmRNAs and were identified as real target genes of hsa-miR-146a-3p (Fig. 2).

Enrichment analysis of the signaling pathways of DEmiRNAs indicated that GDM-related miRNAs were mainly enriched in the glypican pathway ($P < 0.001$), proteoglycan syndecan-mediated signaling events ($P < 0.001$), and syndecan-1-mediated signaling events ($P < 0.001$) (Fig. 3, Table 3). In addition, analysis of the signaling pathways of the real target genes of hsa-miR-146a-3p found that the glypican pathway was also one of the pathways regulated by hsa-miR-146a-3p ($P < 0.001$). In addition, the DEmRNAs TRAF6, MEF2A, TNFAIP3, SKIL, FLT1 and IRS2 were also targeted by hsa-miR-146a-3p in the present study and were enriched in the glypican pathway.

Table 3
Enrichment analysis of signal pathways involved in DEmiRNAs

Biological pathway	Fold enrichment	P-value	Q-value
Glypican pathway	1.769653	6.84E-13	1.99E-09
Proteoglycan syndecan-mediated signaling events	1.760422	1.05E-12	1.99E-09
Syndecan-1-mediated signaling events	1.763494	2.21E-12	1.99E-09
VEGF and VEGFR signaling network	1.758073	2.81E-12	1.99E-09
IFN-gamma pathway	1.757313	3.46E-12	1.99E-09

To explore the hub DErnAs involved in GDM, we further performed an interaction network analysis of DErnAs screened from the GSE19649 dataset (Fig. 4). The results indicated that TRAF6, with a total of 16 connecting nodes, was the top hub gene involved in the pathophysiological process of GDM. Other hub genes included CASP8 (number of connecting nodes: 6), PTPN6 (number of connecting nodes: 6), and CHD3 (number of connecting nodes: 6). Next, these 4 hub genes were selected for independent external validation using the GSE19649 dataset. The expression of TRAF6, CASP8 and CHD3 in 8 pairs of GDM blood samples was confirmed to be higher than that in healthy pregnant women blood samples. However, the expression of PTPN6 in the blood samples of GDM patients is similar to that of healthy pregnant women blood samples (Fig. 5). The AUC for predicting GDM was 0.813, 0.828, 0.813, and 0.703 for CHD3, PTPN6, TRAF6, and CASP8, respectively (Fig. 6, Table 4). Furthermore, TRAF6 was involved in the following three core pathways of GDM: glypican pathway, proteoglycan syndecan-mediated signaling events, and syndecan-1-mediated signaling events.

Table 4
ROC analysis of predicting abilities of
hub genes

Gene	AUC	95%CI	
CHD3	0.813	0.592	1.000
PTPN6	0.828	0.621	1.000
TRAF6	0.813	0.600	1.000
CASP8	0.703	0.427	0.979

To determine the potential mechanism of TRAF6 in GDM, GSEA was conducted (Table 5). A total of 9 hallmark gene sets of metabolism processes were enriched, including cysteine and methionine metabolism, pyruvate metabolism, sphingolipid metabolism, beta alanine metabolism, propanoate metabolism, galactose metabolism, starch and sucrose metabolism, butanoate metabolism, and arginine and proline metabolism processes.

Table 5

The gene sets that were significantly associated with TRAF6 by Gene set enrichment analysis (GSEA)

NAME	SIZE	ES	FWER p-value
KEGG_CYSTEINE_AND_METHIONINE_METABOLISM	23	0.465224	0
KEGG_PYRUVATE_METABOLISM	26	0.348111	0
KEGG_SPHINGOLIPID_METABOLISM	24	0.339434	0
KEGG_BETA_ALANINE_METABOLISM	15	0.332718	0
KEGG_PROPANOATE_METABOLISM	26	0.331809	0
KEGG_GALACTOSE_METABOLISM	18	0.324297	0
KEGG_STARCH_AND_SUCROSE_METABOLISM	25	0.299221	0
KEGG_BUTANOATE_METABOLISM	18	0.29276	0
KEGG_ARGININE_AND_PROLINE_METABOLISM	25	0.276111	0

4 | Discussion

Because the symptoms of GDM mostly occur in late pregnancy, women with first-time GDM often do not have enough time for clinical intervention[15]. Since accumulated evidence has demonstrated that GDM can be prevented by diet, medicine and exercise intervention started in early pregnancy, early detection of women with a high risk of GDM is essential[16]. To identify noninvasive plasma molecular markers of GDM, we first screened DE miRNAs and DE mRNAs by comparing serum samples of GDM patients and those of healthy pregnant women. A total of 264 DE miRNAs and 1217 DE mRNAs were identified, and these differentially expressed markers might serve as noninvasive plasma molecular markers for predicting GDM in early and mid-pregnancy. Further research is needed to evaluate their predictive value. To date, there have been two studies on identifying noninvasive markers of GDM in early and mid-term pregnancy based on miRNAomics. The expression of several plasma miRNAs including miRNA-29a and miRNA-16 was found to be associated with GDM, and both were also differentially expressed in our present study (miRNA-29a, $P = 0.003$; miRNA-16, $P = 0.008$)[15, 17].

The results of the present study also provide insights into the molecular mechanisms that underlie GDM. miRNAs are involved in many biological processes through posttranscriptional regulation[18]. They regulate gene expression and transcription by complementary binding to the 3' end of target genes[18]. However, to date, little is known about the regulatory process between miRNAs and mRNAs in GDM. Thus, we performed an integrative analysis to construct functional miRNA-gene regulatory networks by combining the expression relationships between miRNAs and targets and computational predictions. First, enrichment analysis of the involved signaling pathways was carried out based on DE miRNAs. The

metabolic pathways glypican pathway, proteoglycan syndecan-mediated signaling events, and syndecan-1-mediated signaling events, were the core pathways regulated by miRNAs in GDM. Glypicans belong to the family of membrane-bound heparan sulfate proteoglycans that are linked to the cell surface and are involved in the regulation of growth factor activity[19]. Proteoglycan families are involved in a wide range of diseases, including obesity and diabetes[20]. Impaired glucose metabolism in GDM may alter the expression of proteoglycans, which may impair the biological functions of the placenta[21]. A previous study demonstrated that the expression of a member of the proteoglycan family, endocan, was increased in the human placenta from obese women with GDM and in response to proinflammatory stimuli[20]. In addition, the metabolic effect of high glucose and high osmotic pressure of GDM patients may contribute to the increased proteoglycan perlecan expression in diabetic placentas[22]. Leelalertlauw et al.[23] found that serum glypican 4 levels were increased with increasing degrees of obesity. Furthermore, Ussar et al. [24] demonstrated that circulating glypican-4 levels correlate with BMI and insulin sensitivity in humans. Moreover, glypican-4 interacts with the insulin receptor, enhances insulin receptor signaling, and increases adipocyte differentiation. These results suggest that abnormal expression of proteoglycan family genes in GDM is helpful for understanding the pathogenesis of GDM.

In the present study, hsa-miR-146a-3p, ranked first in the altered DEmiRNAs of GDM because of its lowest P value, was downregulated with a log (fold change) of -12.76 in the blood samples of GDM patients. Changes in miR-146a in diabetic patients and animals were reported in a previous study. Liu et al.[25] demonstrated that treatment of diabetic mice with miR-146a mimics robustly reduced diabetic peripheral neuropathy. Then, the target genes of hsa-miR-146a-3p were identified based on the parallel DEmRNAs in the present study. The results indicated that TRAF6 was not only identified as a target of hsa-miR-146a-3p but also a hub gene involved in the pathophysiological process of GDM. TRAF6 has been demonstrated to be a target gene of miR-146a at the experimental level in previous immune studies[26]. Regarding diabetes, Kamali et al found that miR-146a antagomir significantly increased TRAF6 mRNA levels in human umbilical vein endothelial cells under hyperglycemic conditions. In addition, previous research revealed that culture of aortic endothelial cells in high glucose medium significantly increased TRAF6 expression in a time-dependent manner. In addition, TRAF6 mediated high glucose-induced endothelial dysfunction via NF-KB and AP-1-dependent signaling[27]. To date, the role of miR-146a and its target gene TRAF6 in the pathogenesis of GDM has never been explored. In the present study, GSA revealed that several metabolic processes were enriched in the mechanism of TRAF6 in GDM. Moreover, TRAF6 was involved in all three core pathways regarding glypican metabolism in GDM based on pathway analysis of DEmiRNAs. Furthermore, we confirmed the altered expression of TRAF6 in 8 pairs of GDM blood samples to be higher than that in healthy pregnant blood samples in another independent patient group. Thus, serum TRAF6 may serve as a novel biomarker for GDM.

Other core genes, such as CASP8 and CHD3, were confirmed to be higher in 8 pairs of GDM blood samples than in healthy pregnant women blood samples. At present, there is no research on the relationship between CASP8 or CHD3 and GDM. Wu et al.[27] found that maternal type 2 diabetes mellitus causes heart defects in the developing embryo manifested by excessive apoptosis in heart cells,

with increasing apoptosis markers of cleaved caspase 3 and 8. Whether CASP8 can be used as a marker of fetal heart defects in pregnant women with GDM needs to be further studied.

There are limitations to this study. The serum GDM markers screened from the GSE19649 and GSE98043 datasets were independently externally validated using the GSE19649 dataset. The sample size was relatively small. However, the clinical information, such as age and BMI, of GDM and healthy pregnant women were matched in the GSE19649 dataset, which improved the test effectiveness. In addition, other GDM noninvasive markers explored in this study still need to be further verified by expanding the sample size in the future.

In summary, a series of potential GDM noninvasive serum markers were identified by integrated miRNA-mRNA expression profile analysis. Enrichment analysis indicated that GDM-related miRNAs were mainly enriched in the glypican pathway, proteoglycan syndecan-mediated signaling events, and syndecan-1-mediated signaling events. In addition, miR-146a-3p/TRAF6 might play a central role in the pathogenesis of GDM through the above pathways.

Declarations

Acknowledgements

Not applicable.

Authors' contributions

M. and Y. J. designed experiments; C. M. carried out experiments, analyzed data and experimental results. C. M. and Y. J. wrote the manuscript.

Funding

This work was supported by the Fujian Science and Technology Leading Project (2016Y0060), the Key Clinical Specialty Discipline Construction of Fujian (2015 no. 593), Fujian Science and Technology Leading Project (2018Y0005), National Health and Family Planning Commission Science Foundation(2019-WJ-04), Fujian Medical Innovation Subject (2017-CX-11) and Fujian Provincial Maternity and Children's Health Hospital Foundation (YCXM 18-26).

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declared no competing interests.

Availability of data and materials

All data generated or analyzed during this study are included in this published article. The data of this study are available upon request.

References

1. WQ H, Y L, M X, J H, YX S, CX Z: **Excessive fruit consumption during the second trimester is associated with increased likelihood of gestational diabetes mellitus: a prospective study.** *Scientific reports* 2017, **7**(undefined):43620.
2. Zhang Q, He M, Wang J, Liu S, Cheng H, Cheng Y: **Predicting of disease genes for gestational diabetes mellitus based on network and functional consistency.** *Eur J Obstet Gynecol Reprod Biol*, **186**:91-96.
3. Robitaille J, Grant AM: **The genetics of gestational diabetes mellitus: evidence for relationship with type 2 diabetes mellitus.** *Genetics in Medicine* 2008, **10**(4):240.
4. Loeken MR: **Passive smoking as an independent risk factor for gestational diabetes that synergizes with prepregnancy obesity in urban Chinese women.** *Diabetes/metabolism research and reviews* 2017, **33**(6):e2910.
5. Huang W-Q, Lu Y, Xu M, Huang J, Su Y-X, Zhang C-X: **Excessive fruit consumption during the second trimester is associated with increased likelihood of gestational diabetes mellitus: a prospective study.** *Scientific reports* 2017, **7**:43620.
6. Sayed D, Abdellatif M: **MicroRNAs in development and disease.** *Physiological reviews* 2011, **91**(3):827-887.
7. Poirier C, Desgagné V, Guérin R, Bouchard L: **MicroRNAs in pregnancy and gestational diabetes mellitus: emerging role in maternal metabolic regulation.** *Current diabetes reports* 2017, **17**(5):35.
8. Kizilgul M, Kan S, Beysel S, Apaydin M, Ozcelik O, Caliskan M, Ozbek M, Ozdemir S, Cakal E: **Is fibroblast growth factor 23 a new cardiovascular risk marker in gestational diabetes?** *Archives of endocrinology and metabolism* 2017, **61**(6):562-566.
9. Zhao W, Pan J, Li H, Huang Y, Liu F, Tao M, Jia W: **Relationship between high serum cystatin C levels and the risk of gestational diabetes mellitus.** *PloS one* 2016, **11**(2):e0147277.
10. Cao YL, Jia YJ, Xing BH, Shi DD, Dong XJ: **Plasma microRNA-16-5p,-17-5p and-20a-5p: Novel diagnostic biomarkers for gestational diabetes mellitus.** *Journal of Obstetrics and Gynaecology Research* 2017, **43**(6):974-981.
11. Stirn L, Huypens P, Sass S, Batra R, Fritsche L, Brucker S, Abele H, Hennige AM, Theis F, Beckers J: **Maternal whole blood cell miRNA-340 is elevated in gestational diabetes and inversely regulated by glucose and insulin.** *Scientific reports* 2018, **8**(1):1366.

12. Davis S, Meltzer PS: **GEOquery: a bridge between the Gene Expression Omnibus (GEO) and BioConductor.** *Bioinformatics* 2007, **23**(14):1846-1847.
13. Tokar T, Pastrello C, Rossos AE, Abovsky M, Hauschild A-C, Tsay M, Lu R, Jurisica I: **mirDIP 4.1—integrative database of human microRNA target predictions.** *Nucleic acids research* 2017, **46**(D1):D360-D370.
14. Pathan M, Keerthikumar S, Ang CS, Gangoda L, Quek CY, Williamson NA, Mouradov D, Sieber OM, Simpson RJ, Salim A: **FunRich: An open access standalone functional enrichment and interaction network analysis tool.** *Proteomics* 2015, **15**(15):2597-2601.
15. Zhu Y, Tian F, Li H, Zhou Y, Lu J, Ge Q: **Profiling maternal plasma microRNA expression in early pregnancy to predict gestational diabetes mellitus.** *International Journal of Gynecology & Obstetrics* 2015, **130**(1):49-53.
16. Zhang C, Tobias DK, Chavarro JE, Bao W, Wang D, Ley SH, Hu FB: **Adherence to healthy lifestyle and risk of gestational diabetes mellitus: prospective cohort study.** *Bmj* 2014, **349**:g5450.
17. Zhao C, Dong J, Jiang T, Shi Z, Yu B, Zhu Y, Chen D, Xu J, Huo R, Dai J: **Early second-trimester serum miRNA profiling predicts gestational diabetes mellitus.** *PloS one* 2011, **6**(8):e23925.
18. Bartel DP: **MicroRNAs: genomics, biogenesis, mechanism, and function.** *cell* 2004, **116**(2):281-297.
19. Filmus J, Selleck SB: **Glypicans: proteoglycans with a surprise.** *The Journal of clinical investigation* 2001, **108**(4):497-501.
20. Murthi P, Sarkis R, Lim R, Nguyen-Ngo C, Pratt A, Liong S, Lappas M: **Endocan expression is increased in the placenta from obese women with gestational diabetes mellitus.** *Placenta* 2016, **48**:38-48.
21. Chen C-P, Chang S-C, Yang W-CV: **High glucose alters proteoglycan expression and the glycosaminoglycan composition in placentas of women with gestational diabetes mellitus and in cultured trophoblasts.** *Placenta* 2007, **28**(2-3):97-106.
22. Yang W-C, Su T-H, Yang Y-C, Chang S-C, Chen C-Y, Chen C-P: **Altered perlecan expression in placental development and gestational diabetes mellitus.** *Placenta* 2005, **26**(10):780-788.
23. Leelalertlauw C, Korwutthikulrangsri M, Mahachoklertwattana P, Chanprasertyothin S, Khlairit P, Pongratanakul S, Poomthavorn P: **Serum glypican 4 level in obese children and its relation to degree of obesity.** *Clinical endocrinology* 2017, **87**(6):689-695.
24. Ussar S, Bezy O, Blüher M, Kahn CR: **Glypican-4 enhances insulin signaling via interaction with the insulin receptor and serves as a novel adipokine.** *Diabetes* 2012, **61**(9):2289-2298.
25. Liu XS, Fan B, Szalad A, Jia L, Wang L, Wang X, Pan W, Zhang L, Zhang R, Hu J: **MicroRNA-146a mimics reduce the peripheral neuropathy in type 2 diabetic mice.** *Diabetes* 2017, **66**(12):3111-3121.
26. Park H, Huang X, Lu C, Cairo MS, Zhou X: **MicroRNA-146a and microRNA-146b regulate human dendritic cell apoptosis and cytokine production by targeting TRAF6 and IRAK1 proteins.** *Journal of Biological Chemistry* 2015, **290**(5):2831-2841.
27. Liu R, Shen H, Wang T, Ma J, Yuan M, Huang J, Wei M, Liu F: **TRAF6 mediates high glucose-induced endothelial dysfunction.** *Experimental cell research* 2018, **370**(2):490-497.

Figures

Figure 1

Schematic overview of the proposed approach in this study.

Figure 4

Analysis of the real target genes of hsa-miR-146a-3p. The computationally predicted target genes and DEmRNAs were integrated in a Venn diagram to identify the real target genes. DEmRNAs, differentially expressed genes

Figure 6

Enrichment analysis of signaling pathways involved in DEmiRNAs.

Figure 8

Interaction network analysis of DEmRNAs.

Figure 10

Independent validation of 4 hub genes using the GSE19649 dataset. The expression levels of (A) TRAF6, (B) CASP8 and (C) CHD3 in 8 pairs of GDM blood samples were confirmed to be higher than those in healthy pregnant women blood samples. (D) The expression of PTPN6 in the blood samples of GDM patients is similar to that of healthy pregnant women blood samples.

Figure 12

The receiver operating characteristic (ROC) curve analysis validated that all four hub genes were able to correctly predict GDM in the external dataset GSE19649.