

Transcriptome of Visceral Adipose Tissue Identifies an Inflammation-Related ceRNA Network That Regulates Obesity

GanBin Li

Beijing Chaoyang Hospital <https://orcid.org/0000-0002-2576-1100>

Haoyu Zhang

Beijing Chaoyang Hospital

Ke Cao

Beijing Chaoyang Hospital

Zhiwei Zhai

Beijing Chao-Yang Hospital: Beijing Chaoyang Hospital

Zhenjun Wang

Beijing Chaoyang Hospital

Lei Yang

Beijing Chao-Yang Hospital: Beijing Chaoyang Hospital

JiaGang Han (✉ ligb9508@163.com)

Beijing Chaoyang Hospital

Research Article

Keywords: Noncoding RNA, Visceral adipose tissue, RNA sequencing, competing endogenous RNA

Posted Date: November 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1029192/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Molecular and Cellular Biochemistry on January 22nd, 2022. See the published version at <https://doi.org/10.1007/s11010-022-04362-y>.

Abstract

Obesity is becoming an epidemic of widespread concern, but the underlying causes remain elusive. In this study, whole transcriptome RNA sequencing (RNA-seq) revealed differential profiles of noncoding (nc) RNAs and mRNAs in visceral adipose tissue (VAT) from obese (BMI >32.5 kg/m²) and lean (BMI < 20 kg/m²) individuals, with 1920 differentially expressed (DE) genes, 1466 long noncoding (lnc)RNAs, 122 micro (mi)RNAs, and 52 circular (circ)RNAs identified. Gene Set Enrichment Analysis, Gene Ontology analysis, Kyoto Encyclopedia of Genes and Genomes analysis revealed that these ncRNAs were involved in inflammation-related pathways that included cytokine-cytokine receptor interaction, the tumor necrosis factor and nuclear factor kappa B signaling pathways. The results indicated a critical role of inflammation in the pathogenesis of obesity. The network interaction of lncRNA, circRNA, and miRNA revealed a competing endogenous (ce)RNA network that was associated with inflammation. The ceRNA network included circORC5/miR-197-5p/TNFRSF10D and circNTRK2/miR-760/LAT, which were dysregulated in obese patients. Notably, the overexpression of *TNFRSF10D* and *LAT* were significantly positively associated with body weight in obese patients. In conclusion, this whole transcriptome study provided a pool of data that will be useful for identifying biomarkers of obesity and identified an obesity-associated ceRNA network that is regulated by circORC5 and circNTRK2.

Introduction

Obesity is increasing worldwide and is associated with metabolic diseases, such as diabetes, hypertension, cardiovascular disease, as well as an increased risk of malignancy^[1, 2]. Obesity is characterized by an imbalance between caloric intake and energy expenditure resulting in excessive accumulation of fat in visceral adipose tissue (VAT)^[3]. In addition to energy storage, VAT is an endocrine organ, secreting adipose-derived cytokines, leptin, visfatin, and adiponectin^[4]. Obesity is accompanied by hyperplasia and enlargement of adipocytes and increased secretion of multiple adipokines and proinflammatory cytokines, such as interleukin (IL)-1, IL-6, and tumor necrosis (TNF)- α that lead to chronic, low-grade local inflammation and insulin resistance^[5, 6]. Chronic inflammation and insulin resistance are the main characteristics of obesity, but details of mechanisms involved in either the inflammation or obesity remain to be uncovered^[2, 3, 5]

Noncoding (nc)RNAs, including micro (mi)RNA, long noncoding (lnc)RNA, and circular (circ)RNA, are active in metabolic disease^[7]. miRNAs are small endogenous ncRNAs with 20–22 bps. The major function of miRNAs is dependent on complementary interaction with the target mRNAs via complete or incomplete matching^[7]. Several recent studies reported that dysregulated miRNAs enhanced chronic inflammation, insulin resistance, adipogenesis, and angiogenesis in VAT and thus promoted the process of obesity^[8, 9]. On the contrary, lncRNAs are transcripts that are more than 200 nt in length and lack an apparent protein-coding capacity^[10, 11]. For example, lncRNA-GAS5 regulates adipogenesis of mesenchymal stem cells by preventing miR-18a from decreasing the expression of connective tissue growth factor (CTGF)^[11]. CircRNAs are ncRNAs with a closed loop structure and reverse splicing of its 5'

and 3' ends by covalent bonding, which is more stable than its linear RNA transcript^[12, 13]. SAMD4A, is a circRNA that is upregulated in VAT to activate the miR-138-5p/ZEH2 axis to promote the pathogenesis of obesity^[13].

To identify regulatory mechanisms influencing obesity, we used whole transcriptome RNA sequencing (RNA-seq) to uncover the profiles of differentially expressed (DE) mRNAs and ncRNAs. The dysregulated profiles were then evaluated by Gene Set Enrichment Analysis (GSEA), Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment to reveal the functions of the ncRNAs. The aim was to construct a competing endogenous (ce)RNA network that includes DE circRNA-miRNA-mRNA to explore the regulatory mechanisms of the ncRNAs and the potential relationship with obesity.

Materials And Methods

Sample collection

Clinical samples were collected from patients diagnosed with obesity or chronic appendicitis and treated by laparoscopic appendectomy at our hospital between January and May 2021. Patients with a body mass index (BMI) ≥ 32.5 kg/m² were included in an obese group and those with a BMI ≤ 20 kg/m² were included in a normal-weight group. About 100 milligrams of VAT (**Figure S1**) was obtained during surgery, histologically confirmed, immediately frozen in liquid nitrogen, and stored at -80°C until RNA extraction. Thirty patients were enrolled in an obesity group and 30 in a normal-weight group. Differences in BMI, abdomen circumference, systolic pressure, diastolic pressure, fasting blood glucose, cholesterol, and triglycerides between the two groups were significant (all $P < 0.001$, **Table S1**). The study was approved by the ethics committee of our hospital (2021-scientific-367) and all patients gave their informed consent.

Whole transcriptome RNA-seq

Six pairs of VAT samples from obese patients and four pairs from normal-weight patients were selected for RNA-seq, which was performed by Biomarker Technologies (Beijing, China). Total RNA was extracted with TRIzol (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions. The RNA quality and purity were determined with a Nanodrop 2000 Spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA). The cDNA libraries for circRNA, lncRNA, and small RNA were constructed separately. Briefly, 1.5 μg RNA samples were used as input material for rRNA removal using Ribo-Zero rRNA Removal Kits (Epicentre, Madison, WI, USA). Sequencing libraries were generated with Next Ultra Directional RNA Library Prep kits (Illumina, New England Biolabs, USA) following the manufacturer's instructions. First-strand cDNA was synthesized using random hexamer primer and reverse transcriptase. After adenylation of the 3' ends of DNA fragments, Nebnext hairpin loop structure adapters were ligated to prepare for hybridization. The library fragments were purified with AMPure XP Beads (Beckman Coulter, Beverly, MA, USA) to select insert fragments of 150–200 bp in length. The cDNA was treated with 3 μL uracil DNA glycosylase and DNA glycosylase-lyase endonuclease VIII (USER) enzyme (New England Biolabs, USA) at 37°C for 15 min before PCR. PCR was performed with Phusion High-Fidelity DNA

polymerase, universal PCR primers and Index(X) Primer. After cluster generation, the library preparations were sequenced on an Illumina HiSeq platform at Biomarker Technologies (Beijing, China), and paired-end reads were generated.

The transcriptome was assembled using Cufflinks v2.1.1^[14] and Scripture^[15], based on the reads mapped to the reference genome (GRCh38). Differential expression analysis of the two groups was performed using the DESeq R package v.1.10.1 (<https://bioconductor.org/packages/release/bioc/html/DESeq2.html>). RNAs found by RNA-seq with an adjusted *P*-value < 0.05 and an absolute value of log₂ fold change of >1 were considered as DE.

GSEA, GO and KEGG enrichment analysis

Gene sets were downloaded from the MsigDB database^[16], and broad hallmarks and KEGG symbols were included for GSEA, and pathways with corrected *P*-values of < 0.05 were considered significantly enriched. GO and KEGG enrichment analysis were conducted by DAVID (<https://david.ncifcrf.gov>) and KOBAS (<http://kobas.cbi.pku.edu.cn/kobas3>). GO includes three items, biologic processes (BPs), molecular function (MF), and cellular components (CCs). The DE mRNAs that were targets of DE miRNAs, DE lncRNAs, or DE circRNAs were enriched by GSEA, GO or KEGG analysis. The top ten enriched GO items and KEGG pathways were selected and visualized by SangerBox (<http://sangerbox.com>).

Protein-protein interaction (PPI) analysis

STRING (<https://www.string-db.org>) was used to construct PPI networks that showed the gene interactions involved in selected KEGG pathways. An interaction score > 0.4 was used as the extraction cutoff for the PPI pairs. CytoHubba (<https://apps.cytoscape.org/apps/cytohubba>) was used to select the top ten hub genes in the PPI network. The relative results were visualized by Cytoscape v3.7.0^[17].

ceRNA network

lncRNA-miRNA-mRNA and circRNA-miRNA-mRNA networks were constructed using current hypothetical models of ceRNA regulation and the RNA-seq results. DE miRNA targets of DE lncRNAs were predicted using TargetScan (<http://www.targetscan.org>). DE miRNA targets of DE circRNAs were predicted by miRanda (<http://www.mirbase.org>). The target DE genes of DE miRNAs were predicted by the miRBase database (<http://www.mirbase.org>). Finally, one downregulated lncRNA and four dysregulated circRNAs were selected for inclusion in the ceRNA network, and the results were visualized by Cytoscape v3.7.0.

Quantitative real-time (qRT)-PCR validation

Total RNA was extracted from VAT with TRIzol (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions and then reverse transcribed to cDNA with FastKing RT Kits and gDNase (Tiangen Biotech, Beijing, China). qRT-PCR was performed with SuperReal PreMix Plus (SYBR Green; Tiangen Biotech, Beijing, China) on an ABI 7500 RT-PCR system (Applied Biosystems, Foster city, CA, USA) following the protocol provided by the manufacturer. Relative RNA expression was calculated by the

$2^{-\Delta\Delta Ct}$ method, and 18s rRNA (for circRNA), GAPDH (for mRNA), and U6 (for miRNA) were used as internal controls. The primers that were used are listed in **Table S2**.

Statistical analysis

SPSS 25.0 (IBM Corp., Armonk, NY, USA) was used for the statistical analysis and GraphPadPrism7.0 (GraphPad Software, La Jolla, CA, USA) was used to draw the figures. Normally distributed variables were reported as means \pm standard deviation and categorical data were reported as numbers and percentages (%). Between-group differences were compared by paired *t*-tests or chi-square tests. *P*-values < 0.05 were considered statistically significant

Results

Obesity-related transcriptomic profiles in VAT from obese and normal-weight individuals

The RNA-seq results included 32,639 genes, 69,045 lncRNAs, 2020 miRNAs, and 6029 circRNAs (**Figure 1A**). Among those, 1920 mRNAs, 1466 lncRNAs, 122 miRNAs, and 52 circRNAs were DE (fold change ≥ 1 or ≤ -1 , *P*-value < 0.01 , **Figure 1B**). The chromosomal locations of the DE genes are shown in **Figure 1C**. Chromosomes 5, 6 and X carried the circRNAs with the greatest difference in expression (**Figure. 1D**), and the five most upregulated and downregulated DE ncRNAs are listed in **Table 1**. The primary RNA-Seq datasets are available in <https://submit.ncbi.nlm.nih.gov>, with a grant number PRJNA763819.

The analysis identified 827 upregulated and 639 downregulated lncRNAs. The top three upregulated lncRNAs were MSTRG.28247.73, MSTRG.30278.66, MSTRG.9074.6, and the top three downregulated lncRNAs were MSTRG.68830.10, MSTRG.22811.9, MSTRG.17871.15. Of the DE circRNAs, 25 were upregulated and 27 were downregulated. The top three upregulated circRNAs were has-circ-0086376, has-circ-0001654, and has-circ-0003239; the top three downregulated circRNAs were 6:31271073|31355592 (a novel circRNA not previously identified), hsa-circ-0000277, and hsa-circ-0006114. There were 69 upregulated and 53 downregulated miRNAs (**Figures 2A–F, Table 1**).

Bioinformatic analysis of enriched obesity-associated inflammation pathways

GSEA enrichment (**Figure 3A–C**) identified 130 upregulated gene sets associated with obesity, 37 of which were significantly upregulated (*P* < 0.05). Of 48 downregulated obesity-associated gene sets, the difference in expression was significant in six (*P* < 0.05). The top three enriched gene sets were chemokine signaling pathways (NSE = 1.636, *P* = 0.001), apoptosis (NSE = 1.546, *P* = 0.002), and adipogenesis (NSE = 1.137, *P* = 0.03).

GO enrichment confirmed that inflammatory responses were the most affected BPs (**Figure 4A**). The enriched MFs and CCs identified in the GO analysis are listed in **Figure 4A**. KEGG pathway enrichment analysis found that cytokine-cytokine receptor interaction; the PI3K-Akt, chemokine, TNF, and NF-kappa B signaling pathways were the most likely to be associated with obesity (**Figure 4B**). GO and KEGG

integrative analysis to screen the key genes associated with obesity defined 4658 genes that were involved in pathways related cancer cell proliferation, such as the PI3K-Akt, NF-kappa B, and TNF signaling pathways, and cytokine-cytokine receptor interaction (**Figure S2A–B**).

Genes involved in three key obesity-associated pathways

We found overlapping involvement of 52 genes in obesity-associated pathways and interactions, 29 involved in cytokine-cytokine receptor interactions, 11 in the TNF signaling pathway, and 12 in the NF-kappa B signaling pathway (**Table 2**). The analysis suggested that obesity was a disease associated with chronic inflammation. The PPI network for the genes involved in the pathways was constructed with STRING online tools (**Figure 5A–C**).

CeRNA mechanisms associated with obesity

In order to further explore the potential regulatory involvement of DE circRNAs in the pathogenesis of obesity, we constructed three ceRNA networks that were most likely associated with inflammation. The dysregulated circRNAs, included in the networks were hsa_circ_0003000, hsa_circ_0054228, hsa_circ_0000545, and hsa_circ_0139142. In addition, hsa_circ_0003000 was targeted by eight miRNAs, miR-1226-3p, miR-197-5p, miR-6780-5p, miR-3188, miR-1228-5p, miR-760, miR-6837-3p, and miR-6843-3p (**Figure 6A**). The miRNAs regulated 229 DE genes. The overlapping genes in the three selected KEGG pathways and PPI network included *CSF3R*, *LAT*, and *LTB*. They were in the ceRNA network and regulated by has-circ-0000545 and has-circ-0139142. has-circ-003000 sponged miR-1226-3p and miR-6780a-5p as well as miR-197-5p. The targeted genes included *XCR1*, *CARD11*, and *TNFRSF10D* (**Figure. 6B**).

Validation of obesity-related ncRNAs in the ceRNA network

We performed qRT-PCR in human samples to validate the DE findings. circ-0003000 and circ-0139142 were significantly downregulated in VAT from obese compared with normal-weight individuals (Figure 7A). Consistent with the ceRNA network, miR-760 and miR-197-5p were targeted by circ-0003000 and circ-0139142, which were significantly upregulated in obese VAT (Figure 7B). The expression of *LAT* and *TNFRSF10D* mRNA was significantly downregulated in obese VAT. Those genes were key genes for circ-0003000 and circ-0139142 regulation in the ceRNA networks (Figure 7C). Our analysis confirmed that the inflammation-related ceRNA network had important roles in regulating obesity. Because the host genes of circ-0003000 and circ-0139142 were *ORC5* and *NTRK2* we name them circORC5 and circNTRK2 respectively. We further validated the association between circNTRK2 and circNTRK2 expression and BMI, negative correlations of both circRNAs and BMI, indicating that downregulated obesity-associated ceRNA networks promoted obesity

Discussion

Accumulating evidence confirms that excessive energy storage in VAT leads to dysregulated secretion of adipokines and cytokines and state of chronic, low-grade inflammation and insulin resistance associated

with obesity^[18, 19]. In this study, the VAT transcriptome revealed that DE genes and DE ncRNAs have an important role in the pathogenesis of obesity by modulating the process of inflammation and adipogenesis.

Numerous studies have illustrated the key role of miRNAs in obesity^[20]. The expression of miR-28-3p^[21], miR-125-5p^[22], and miR-103-5p^[23] was shown to be downregulated, and the expression of miR-27^[24], miR-142-3p^[24], miR-486, and miR-806^[25] upregulated in the VAT of obese patients. The DE miRNAs were shown to be involved in obesity-associated biological processes, including adipogenesis, angiogenesis, inflammation, insulin resistance, and macrophage polarization^[21, 23]. Our RNA-seq analysis screened novel DE miRNAs that were associated with obesity and might eventually serve as biomarkers or targets for the treatment and diagnosis of obesity.

We identified both DE lncRNAs and circRNAs that regulated the pathogenesis of obesity^[26]. A ceRNA model of circRNA or lncRNA has been associated the inhibition function of miRNA downregulating mRNA^[26, 27]. Previous studies revealed that most lncRNAs and circRNAs function as ceRNAs to regulate gene expression and biological processes^[28, 29]. lncRNA/MEG3 was found to be downregulated during adipogenesis, which significantly promoted adipogenesis by modulating an miR-140-5P/PPAR- γ axis^[28]. CircRNA is emerging as an interesting area because of its specific structure and function in regulating obesity^[30]. At present, a total of 52 DE circRNAs have been identified, and the involvement of most of them in regulating obesity has not been investigated.

We used bioinformatic analysis to investigate the roles DE circRNAs active in a ceRNA network to regulate genes associated with chronic inflammation by regulating IL-1, IL-6, and TNF- α ^[31, 32]. Inflammatory factors were increased in patients with obesity^[33, 34]. Conversely, IL-13 and IL-33, which are anti-inflammatory cytokines, were decreased in obesity^[35]. A high-fat diet with IL-33 injection improved insulin sensitivity in an animal study, which indicated that anti-inflammatory treatment could decrease insulin resistance in obesity^[35, 36]. In this study, KEGG enrichment showed that pathways including cytokine-cytokine receptor interactions, and TNF and NF- κ B pathway signaling, regulated obesity-related inflammation and insulin resistance^[37-39]. CXCL6, IL-6, IL-1B, TNFAIP3, and CARD11 have been identified as inflammatory factors involved in glucose intolerance and insulin resistance^[40]. Our study demonstrated that those genes regulated the PPI and ceRNA network associated with obesity. CXCL6 was reported to be upregulated in high-fat diet-induced obese mice and promoted cell proliferation and immune responses in addition to inflammation^{[38] [39]}. In our study, *CARD11* was identified as the target of circORC5 (hsa_circ_000300), which is a member of caspase recruitment domain family and promotes inflammation and insulin resistance by activated NF- κ B signaling^[40].

ceRNA networks including lncRNA GAS5/miR-18a/CTGF, and the circRNA SAMD4A/miR-138-5p/ZEH2 axis were shown to regulate adipogenesis of mesenchymal stem cells^[24]. Here, we identified ceRNA networks that included a total of either six (*CSF3R*, *LAT*, *LTB*, *XCR1*, *CARD11*, *TNFRSF10D*) or five (*PRLR*, *LTB*, *CARD11*, *LAT* and *CD27*) overlapped genes. *TNFRSF10D* and *LAT* were increased in obese VAT and

were affected by the ceRNA network regulated by circORC5 and circNTRK2. CircORC5 is an ncRNA with a spliced length of 612 bp that was active in the ceRNA of miR-197-5p, miR-6780a-5p, and miR-1226-3p. In addition, *CARD11*, *XCR1*, and were DE in obese VAT and served the targets of iR-197-5p, miR-6780a-5p and miR-1226-3p.

In summary, we identified and constructed profiles of DE mRNAs, lncRNAs, circRNAs, and miRNAs in VAT from obese individuals. Bioinformatic analysis revealed that the dysregulated ncRNAs were associated with inflammatory responses, insulin resistance, adipogenesis, and apoptosis. The KEGG pathways included cytokine-cytokine receptor interaction, the TNF signaling pathway and the NF-kappa B signaling pathway, which were significantly associated with chronic inflammation. We developed ceRNA networks that involved the inflammatory response. The molecular details are the subject of future study.

Declarations

Institutional Review Board Statement

The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Ethics Committee of BeiJing ChaoYang Hospital, Capital Medical University (2021-scientific-367)

Informed Consent Statement

Informed consent was obtained from all subjects involved in the study, written informed consent has been obtained from the patient(s) to publish this paper.

Data Availability Statement

The primary RNA-Seq datasets are available in <https://submit.ncbi.nlm.nih.gov>, with a grant number PRJNA763819.

Conflicts of Interest

The authors declare no conflict of interest.

Author Contributions

Conceptualization, Gan-Bin Li and Zhi Wei Zhai; Data curation, Gan-Bin Li, Hao Yu Zhang , Ke Cao, Zhi Wei Zhai and Zhen Jun Wang; Formal analysis, Gan-Bin Li and Zhi Wei Zhai; Funding acquisition, Gan-Bin Li, Lei Yang and Jia-Gang Han; Investigation, Hao Yu Zhang and Lei Yang; Methodology, Gan-Bin Li, Ke Cao, Zhi Wei Zhai, Zhen Jun Wang, Lei Yang and Jia-Gang Han; Project administration, Zhen Jun Wang; Resources, Hao Yu Zhang , Ke Cao and Lei Yang; Software, Gan-Bin Li, Ke Cao and Jia-Gang Han; Supervision, Hao Yu Zhang and Zhi Wei Zhai; Validation, Gan-Bin Li, Hao Yu Zhang and Zhen Jun Wang; Writing – original draft, Gan-Bin Li and Lei Yang; Writing – review & editing, Zhen Jun Wang and Jia-Gang Han.

References

1. Dixon JB. Obesity in 2015: Advances in managing obesity. *Nature reviews Endocrinology* 2016; **12**(2): 65-66. <https://doi.org/10.1038/nrendo.2015.221>
2. El-Sayed Moustafa JS, Froguel P. From obesity genetics to the future of personalized obesity therapy. *Nature reviews Endocrinology* 2013; **9**(7): 402-413. <https://doi.org/10.1038/nrendo.2013.57>
3. Mouton AJ, Li X, Hall ME, Hall JE. Obesity, Hypertension, and Cardiac Dysfunction: Novel Roles of Immunometabolism in Macrophage Activation and Inflammation. *Circulation research* 2020; **126**(6): 789-806. <https://doi.org/10.1161/circresaha.119.312321>
4. Petrus P, Lecoutre S, Dollet L, Wiel C, Sulen A, Gao H, Tavira B, Laurencikiene J, Rooyackers O, Checa A, Douagi I, Wheelock CE, Arner P, McCarthy M, Bergo MO, Edgar L, Choudhury RP, Aouadi M, Krook A, Rydén M. Glutamine Links Obesity to Inflammation in Human White Adipose Tissue. *Cell metabolism* 2020; **31**(2): 375-390.e311. <https://doi.org/10.1016/j.cmet.2019.11.019>
5. Kawashima K, Maeda K, Saigo C, Kito Y, Yoshida K, Takeuchi T. Adiponectin and Intelectin-1: Important Adipokine Players in Obesity-Related Colorectal Carcinogenesis. *International journal of molecular sciences* 2017; **18**(4) . <https://doi.org/10.3390/ijms18040866>
6. Deng T, Lyon CJ, Bergin S, Caligiuri MA, Hsueh WA. Obesity, Inflammation, and Cancer. *Annual review of pathology* 2016; **11**: 421-449. <https://doi.org/10.1146/annurev-pathol-012615-044359>
7. Carpi S, Scoditti E, Massaro M, Polini B, Manera C, Digiacoimo M, Esposito Salsano J, Poli G, Tuccinardi T, Doccini S, Santorelli FM, Carluccio MA, Macchia M, Wabitsch M, De Caterina R, Nieri P. The Extra-Virgin Olive Oil Polyphenols Oleocanthal and Oleacein Counteract Inflammation-Related Gene and miRNA Expression in Adipocytes by Attenuating NF- κ B Activation. *Nutrients* 2019; **11**(12). <https://doi.org/10.3390/nu11122855>
8. Rohm B, Holik AK, Kretschy N, Somoza MM, Ley JP, Widder S, Krammer GE, Marko D, Somoza V. Nonivamide enhances miRNA let-7d expression and decreases adipogenesis PPAR γ expression in 3T3-L1 cells. *Journal of cellular biochemistry* 2015; **116**(6): 1153-1163. <https://doi.org/10.1002/jcb.25052>
9. Zhao XY, Xiong X, Liu T, Mi L, Peng X, Rui C, Guo L, Li S, Li X, Lin JD. Long noncoding RNA licensing of obesity-linked hepatic lipogenesis and NAFLD pathogenesis. *Nature communications* 2018; **9**(1): 2986. <https://doi.org/10.1038/s41467-018-05383-2>
10. Li M, Xie Z, Wang P, Li J, Liu W, Tang S, Liu Z, Wu X, Wu Y, Shen H. The long noncoding RNA GAS5 negatively regulates the adipogenic differentiation of MSCs by modulating the miR-18a/CTGF axis as a ceRNA. *Cell death & disease* 2018; **9**(5): 554. <https://doi.org/10.1038/s41419-018-0627-5>
11. Wang Y, Liu J, Ma J, Sun T, Zhou Q, Wang W, Wang G, Wu P, Wang H, Jiang L, Yuan W, Sun Z, Ming L. Exosomal circRNAs: biogenesis, effect and application in human diseases. *Molecular cancer* 2019; **18**(1): 116. <https://doi.org/10.1186/s12943-019-1041-z>
12. Zhou ZB, Huang GX, Fu Q, Han B, Lu JJ, Chen AM, Zhu L. circRNA.33186 Contributes to the Pathogenesis of Osteoarthritis by Sponging miR-127-5p. *Molecular therapy : the journal of the*

- American Society of Gene Therapy* 2019; **27**(3): 531-541.
<https://doi.org/10.1016/j.ymthe.2019.01.006>
13. Liu Y, Liu H, Li Y, Mao R, Yang H, Zhang Y, Zhang Y, Guo P, Zhan D, Zhang T. Circular RNA SAMD4A controls adipogenesis in obesity through the miR-138-5p/EZH2 axis. *Theranostics* 2020; **10**(10): 4705-4719. <https://doi.org/10.7150/thno.42417>
 14. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nature protocols* 2012; **7**(3): 562-578. <https://doi.org/10.1038/nprot.2012.016>
 15. Guttman M, Garber M, Levin JZ, Donaghey J, Robinson J, Adiconis X, Fan L, Koziol MJ, Gnirke A, Nusbaum C, Rinn JL, Lander ES, Regev A. Ab initio reconstruction of cell type-specific transcriptomes in mouse reveals the conserved multi-exonic structure of lincRNAs. *Nature biotechnology* 2010; **28**(5): 503-510. <https://doi.org/10.1038/nbt.1633>
 16. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences of the United States of America* 2005; **102**(43): 15545-15550. <https://doi.org/10.1073/pnas.0506580102>
 17. Doncheva NT, Morris JH, Gorodkin J, Jensen LJ. Cytoscape StringApp: Network Analysis and Visualization of Proteomics Data. *Journal of proteome research* 2019; **18**(2): 623-632. <https://doi.org/10.1021/acs.jproteome.8b00702>
 18. Rohm TV, Fuchs R, Müller RL, Keller L, Baumann Z, Bosch AJT, Schneider R, Labes D, Langer I, Pilz JB, Niess JH, Delko T, Hruz P, Cavelti-Weder C. Obesity in Humans Is Characterized by Gut Inflammation as Shown by Pro-Inflammatory Intestinal Macrophage Accumulation. *Frontiers in immunology* 2021; **12**: 668654. <https://doi.org/10.3389/fimmu.2021.668654>
 19. Kiran S, Kumar V, Kumar S, Price RL, Singh UP. Adipocyte, Immune Cells, and miRNA Crosstalk: A Novel Regulator of Metabolic Dysfunction and Obesity. *Cells* 2021; **10**(5) . <https://doi.org/10.3390/cells10051004>
 20. Prats-Puig A, Ortega FJ, Mercader JM, Moreno-Navarrete JM, Moreno M, Bonet N, Ricart W, López-Bermejo A, Fernández-Real JM. Changes in circulating microRNAs are associated with childhood obesity. *The Journal of clinical endocrinology and metabolism* 2013; **98**(10) <https://doi.org/10.1210/jc.2013-1496>
 21. Villard A, Marchand L, Thivolet C, Rome S. Diagnostic Value of Cell-free Circulating MicroRNAs for Obesity and Type 2 Diabetes: A Meta-analysis. *Journal of molecular biomarkers & diagnosis* 2015; **6**(6) <https://doi.org/10.4172/2155-9929.1000251>
 22. Benito-Vicente A, Uribe KB, Rotllan N, Ramírez CM, Jebari-Benslaiman S, Goedeke L, Canfrán-Duque A, Galicia-García U, Saenz De Urturi D, Aspichueta P, Suárez Y, Fernández-Hernando C, Martín C. miR-27b Modulates Insulin Signaling in Hepatocytes by Regulating Insulin Receptor

- Expression. *International journal of molecular sciences* 2020; **21**(22)
<https://doi.org/10.3390/ijms21228675>
23. Al-Rawaf HA. Circulating microRNAs and adipokines as markers of metabolic syndrome in adolescents with obesity. *Clinical nutrition (Edinburgh, Scotland)* 2019; **38**(5): 2231-2238
<https://doi.org/10.1016/j.clnu.2018.09.024>
24. Tay Y, Rinn J, Pandolfi PP. The multilayered complexity of ceRNA crosstalk and competition. *Nature* 2014; **505**(7483): 344-352 <https://doi.org/10.1038/nature12986>
25. Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP. A ceRNA hypothesis: the Rosetta Stone of a hidden RNA language? *Cell* 2011; **146**(3): 353-358 <https://doi.org/10.1016/j.cell.2011.07.014>
26. Li Z, Jin C, Chen S, Zheng Y, Huang Y, Jia L, Ge W, Zhou Y. Long non-coding RNA MEG3 inhibits adipogenesis and promotes osteogenesis of human adipose-derived mesenchymal stem cells via miR-140-5p. *Molecular and cellular biochemistry* 2017; **433**(1-2): 51-60.
<https://doi.org/10.1007/s11010-017-3015-z>
27. Zhang F, Yang Y, Chen X, Liu Y, Hu Q, Huang B, Liu Y, Pan Y, Zhang Y, Liu D, Liang R, Li G, Wei Q, Li L, Jin L. The long non-coding RNA β Faar regulates islet β -cell function and survival during obesity in mice. *Nature communications* 2021; **12**(1): 3997 <https://doi.org/10.1038/s41467-021-24302-6>
28. Xu X, Zhang J, Tian Y, Gao Y, Dong X, Chen W, Yuan X, Yin W, Xu J, Chen K, He C, Wei L. CircRNA inhibits DNA damage repair by interacting with host gene. *Molecular cancer* 2020; **19**(1): 128
<https://doi.org/10.1186/s12943-020-01246-x>
29. Alexaki VI. The Impact of Obesity on Microglial Function: Immune, Metabolic and Endocrine Perspectives. *Cells* 2021; **10**(7) <https://doi.org/10.3390/cells10071584>.
30. Kim JN, Han SN, Kim HK. Anti-Inflammatory and Anti-Diabetic Effect of Black Soybean Anthocyanins: Data from a Dual Cooperative Cellular System. *Molecules (Basel, Switzerland)* 2021; **26**(11) <https://doi.org/10.3390/molecules26113363>
31. Ghanbari M, Momen Maragheh S, Aghazadeh A, Mehrjuyan SR, Hussen BM, Abdoli Shadbad M, Dastmalchi N, Safaralizadeh R. Interleukin-1 in obesity-related low-grade inflammation: From molecular mechanisms to therapeutic strategies. *International immunopharmacology* 2021; **96**: 107765 <https://doi.org/10.1016/j.intimp.2021.107765>
32. Wang X, Ba T, Cheng Y, Zhang P, Chang X. Probiotics alleviate adipose inflammation in high-fat diet-induced obesity by restoring adipose invariant natural killer T cells. *Nutrition (Burbank, Los Angeles County, Calif)* 2021; **89**: 111285. <https://doi.org/10.1016/j.nut.2021.111285>
33. Miller AM, Asquith DL, Hueber AJ, Anderson LA, Holmes WM, McKenzie AN, Xu D, Sattar N, McInnes IB, Liew FY. Interleukin-33 induces protective effects in adipose tissue inflammation during obesity in mice. *Circulation research* 2010; **107**(5): 650-658 <https://doi.org/10.1161/circresaha.110.218867>
34. Zeyda M, Wernly B, Demyanets S, Kaun C, Hämmerle M, Hantusch B, Schranz M, Neuhofer A, Itariu BK, Keck M, Prager G, Wojta J, Stulnig TM. Severe obesity increases adipose tissue expression of interleukin-33 and its receptor ST2, both predominantly detectable in endothelial cells of human

- adipose tissue. *International journal of obesity (2005)* 2013; **37**(5): 658-665.
<https://doi.org/10.1038/ijo.2012.118>
35. Law YY, Lee WF, Hsu CJ, Lin YY, Tsai CH, Huang CC, Wu MH, Tang CH, Liu JF. miR-let-7c-5p and miR-149-5p inhibit proinflammatory cytokine production in osteoarthritis and rheumatoid arthritis synovial fibroblasts. *Aging* 2021; **13**(13): 17227-17236. <https://doi.org/10.18632/aging.203201>
36. Wang J, Luo X, Cai S, Sun J, Wang S, Wei X. Blocking HOTAIR protects human chondrocytes against IL-1 β -induced cell apoptosis, ECM degradation, inflammatory response and oxidative stress via regulating miR-222-3p/ADAM10 axis. *International immunopharmacology* 2021; **98**: 107903
<https://doi.org/10.1016/j.intimp.2021.107903>
37. Li W, Pan X, Cheng W, Cheng Y, Yin Y, Chen J, Xu G, Xie L. Serum biochemistry, histology and transcriptomic profile analysis reflect liver inflammation and damage following dietary histamine supplementation in yellow catfish (*Pelteobagrus fulvidraco*). *Fish & shellfish immunology* 2018; **77**: 83-90. <https://doi.org/10.1016/j.fsi.2018.03.036>
38. Wang X, Dai Y, Zhang X, Pan K, Deng Y, Wang J, Xu T. CXCL6 regulates cell permeability, proliferation, and apoptosis after ischemia-reperfusion injury by modulating Sirt3 expression via AKT/FOXO3a activation. *Cancer biology & therapy* 2021; **22**(1): 30-39. <https://doi.org/10.1080/15384047.2020.1842705>
39. Mauer J, Chaurasia B, Goldau J, Vogt MC, Ruud J, Nguyen KD, Theurich S, Hausen AC, Schmitz J, Brönneke HS, Estevez E, Allen TL, Mesaros A, Partridge L, Febbraio MA, Chawla A, Wunderlich FT, Brüning JC. Signaling by IL-6 promotes alternative activation of macrophages to limit endotoxemia and obesity-associated resistance to insulin. *Nature immunology* 2014; **15**(5): 423-430 .
<https://doi.org/10.1038/ni.2865>
40. Yamamoto-Furusho JK, Fonseca-Camarillo G, Furuzawa-Carballeda J, Sarmiento-Aguilar A, Barreto-Zuñiga R, Martínez-Benitez B, Lara-Velazquez MA. Caspase recruitment domain (CARD) family (CARD9, CARD10, CARD11, CARD14 and CARD15) are increased during active inflammation in patients with inflammatory bowel disease. *Journal of inflammation (London, England)* 2018; **15**: 13.
<https://doi.org/10.1186/s12950-018-0189-4>

Tables

Table 1. The top 5 up- and down-regulated non-coding RNAs between Obesity and Noraml group

Non-coding RNAs	RNA name	<i>log2FC</i>	<i>P</i> value	Regualtion		
miRNAs	miR-342	9.05	0.0012	up		
	miR-395	7.99	0.0002			
	miR-122-5p	7.54	0.0001			
	miR-3	6.19	0.0107			
	miR-31	5.15	0.0381			
	miR_258	-4.50	0.0112	down		
	miR-449c-5p	-4.33	0.0099			
	miR_370	-4.17	0.0229			
	miR-196b-5p	-3.95	0.0001			
	miR-4670-3p	-3.68	0.0413			
	circRNAs	hsa_circ_0086376	7.33		0.0476	up
hsa_circ_0001654		7.06	0.0006			
hsa_circ_0003239		7.01	0.0007			
hsa_circ_0006382		7.00	0.0219			
hsa_circ_0001707		6.98	0.0290			
6:31271073 31355592		-11.18	0.0008	down		
hsa_circ_0000277		-7.44	0.0022			
hsa_circ_0006114		-6.74	0.0049			
hsa_circ_0093991		-6.69	0.0007			
hsa_circ_0023936		-6.68	0.0225			
lncRNAs		MSTRG.28247.73	12.23		0.0001	up
		MSTRG.30278.66	10.48		0.0001	
	MSTRG.9074.6	8.17	0.0001			
	MSTRG.46982.20	9.79	0.0001			
	MSTRG.66485.11	8.29	0.0001			
	MSTRG.68830.10	-13.99	0.0001	down		
	MSTRG.22811.9	-11.78	0.0001			
	MSTRG.17871.15	-11.38	0.0001			

MSTRG.83382.44	-9.44	0.0001
MSTRG.53248.38	-8.53	0.0001

Table 2. Common shared genes in three KEGG pathways

KEGG pathway	Common genes
Cytokine-cytokine receptor interaction	CXCL6, CSF3R, CXCL8, IL20RA, CX3CL1, CCL8, CXCR1, CXCR2, IL21R, CCL2, CCR7, IL12RB2, XCR1, CCL21, LIF, IL18, IFNLR1, PPBP, PRLR, TNFRSF10D, BMP2, IL6, CD40LG, IL1B, LEP, IL2RB, CD27, LTB, IL7R
TNF signaling pathway	EDN1, LIF, TNFAIP3, SELE, CXCL2, CX3CL1, ICAM1, NFKBIA, IL6, IL1B, CCL2
NF-kappa B signaling pathway	CXCL8, CCL21, TNFAIP3, ICAM1, NFKBIA, ZAP70, CD40LG, LCK, IL1B, LTB, LAT, CARD11

Figures



Figure 1

The high-throughput sequencing results between Obesity and Normal group. A The general expression profiles of mRNAs, lncRNAs, miRNAs and circRNAs; B Differentially expressed mRNAs, lncRNAs, miRNAs and circRNAs between Obesity and Normal group; C The relative location information of non-coding RNAs in chromosome, the outermost circle represents chromosome location and then mRNAs, lncRNAs, circRNAs and miRNAs from outside to inside, respectively. The expression profiles of up- and down-

regulated RNAs are marked red and blue respectively, and the height of each line represents differential expression profiles ($\log_{10} P$ or FDR value); D Different RNA expression ratio in related chromosomes, each bar with different color represents different RNA, and the X-axis represents 23 chromosomes, and the Y-axis represents RNA ratio (the number of RNA in related chromosome/ the total number of RNA), the higher the bar, the higher the RNA ratio.




Figure 2

Differential expression profiles of non-coding RNAs in visceral fatty tissue between Obesity and Normal body weight individuals. A~C The hierarchical clustering of DElncRNAs, DEMiRNAs and DEcircRNAs, A1~A6 represents Obesity while B1~B4 represents lean group; D~F Volcano plot of DElncRNAs, DEMiRNAs and DEcircRNAs, respectively, the up- and down-regulated DEncRNAs are marked red and green respectively, the top 5 up- and down-regulated DEncRNAs are marked with RNA symbols.




Figure 3

GSEA analysis of expressed genes between Obesity and Normal body weight individuals. A Significantly enriched GSEA plot of Chemokine signaling pathway; B and C Significantly enriched GSEA plots of Apoptosis and Adipogenesis.




Figure 4

GO and KEGG enrichment analysis of DE genes. A The top 10 significantly enriched GO items of BP, MF and CC. B The top 10 significantly enriched KEGG pathways of DE genes.




Figure 5

PPI networks. A PPI network of top 10 most enriched proteins in the KEGG pathway of “Cytokine-cytokine receptor interaction”; B and C PPI networks of top 10 most enriched proteins in the KEGG pathway of “TNF signaling pathway” and “NF-kappa B signaling pathway”, respectively.




Figure 6

ceRNA networks of DE circRNAs. A The ceRNA network of the mostly down-regulated circRNA of has-circ-000300, has-circ-0054228, has-circ-0139142, has-circ-0000545 and its associated DE target miRNAs and mRNAs; B Five pairs of ceRNA networks of “circRNA-miRNA-mRNA” by selecting common shared genes between genes enriched in three KEGG pathways and DE target genes of related circRNA;




Figure 7

qRT-PCR validation of ncRNAs. A~C The relative expression of circRNAs, miRNAs and mRNA in VAT between obesity and normal body weight individuals, respectively, “***” means P<0.01, and “****” means P<0.001, “NS” refers to no significance; D and E revealed that the expression profile of circORC5 and circNTRK2 were negatively associated with BMI.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [FigureS1.tif](#)
- [FigureS2.tif](#)
- [TableS1.docx](#)
- [TableS2.docx](#)