Association of plasma apolipoprotein B with cerebrospinal fluid Alzheimer's disease biomarkers in patients with subjective cognitive decline

Hao Hu

Qingdao University Medical College

Lan Tan

Qingdao University Medical College

Yan-Lin Bi

Qingdao Municipal Hospital Group

Wei Xu

Qingdao Municipal Hospital Group

Lin Tan

Qingdao Municipal Hospital Group

Xue-Ning Shen

Fudan University

Xiao-He Hou

Qingdao University Medical College

Ya-Hui Ma

Qingdao University Medical College

Qiang Dong

Fudan University

Jin-Tai Yu (✉ jintai_yu@fudan.edu.cn)

Huashan Hospital, Fudan University <https://orcid.org/0000-0002-7686-0547>

Research

Keywords: Apolipoprotein, Alzheimer's disease, SCD, plasma biomarker

Posted Date: December 30th, 2019

DOI: <https://doi.org/10.21203/rs.2.19696/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

The preclinical stage of Alzheimer's disease (AD) has become a key target stage for future AD prevention trials. Plasma apolipoprotein, an important lipid-related factor, has been found to be involved in the pathogenesis of AD. This study was to examine whether apolipoprotein B (ApoB), apolipoprotein A-1 (ApoA1) or the ratio of ApoB and ApoA1 (ApoB/A1) were associated with early changes of cerebrospinal fluid (CSF) AD biomarkers in elderly adults with subjective cognitive decline.

Methods

This study finally included 201 cognitive normal (CN) elderly adults and 101 participants with subjective cognitive decline (SCD) from the Chinese Alzheimer's Biomarker and Lifestyle (CABLE) database. The One-way analysis of variance, t-test, Kruskal-Wallis test and Chi-square test were applied in the intergroup comparisons. Multiple linear regression models were used to examine the cross-sectional associations of plasma ApoB, ApoA1 and ApoB/A1 levels with CSF AD-related biomarkers.

Results

Compared with the control group, SCD participants with significant AD biological characteristics had lower ApoB levels. In the total participants, higher level of plasma ApoB was associated with increases in CSF A β 42 ($p = 0.0009$) and A β 42/40 ($p = 0.0038$) as well as decreases in CSF t-tau/A β 42 ($p < 0.0001$) and p-tau/A β 42 ($p < 0.0001$), independent of *APOE* ϵ 4 status. In the further subgroup analysis, these associations were much more significant in the SCD participants. In addition, higher levels of plasma ApoB were also found associated with decreases in CSF t-tau ($p = 0.0224$), p-tau ($p = 0.0086$) and A β 40 ($p = 0.0297$) in the SCD subgroup. Furthermore, we found that these protective associations between plasma ApoB and CSF AD core biomarkers were much more significant in the overweight participants. Results showed no association between ApoA1 and all CSF biomarkers in either total participants or subgroups.

Conclusions

This study is the first to find protective associations of plasma ApoB, but not ApoA1, with CSF AD core biomarkers in elderly adults, and these associations were more significant in SCD individuals. This finding indicated that ApoB may play a protective role in the preclinical stage of AD. Identifying the underlying mechanisms may contribute to the discovery of new pathogenic mechanisms and therapeutic targets.

Background

It has been widely accepted that neuropathological processes associated with Alzheimer's disease (AD) begin a decade or more before the emergence of obvious objective cognitive impairment [1-3]. Biomarker research proved that levels of cerebrospinal fluid (CSF) β -amyloid 42 (A β 42), phosphorylated tau (p-tau) and total tau (t-tau) have been significantly abnormal in the preclinical stage of AD [3-5]. The 2018 National Institute on Aging and Alzheimer's Association (NIAA) research framework further included the three CSF biomarkers in the ATN framework, reflecting brain amyloidosis (A: A β 42), tau pathology (T: p-tau) and neurodegeneration (N: t-tau) [6]. All these theories shifted the definition of AD from a syndromic to a biological construct and led us to focus on disease's earlier stage such as preclinical stage.

Subjective cognitive decline (SCD) in individuals without objective cognitive impairment was recognized as a transitional stage between a fully asymptomatic stage and a mild cognitively impaired (MCI) stage of the disease [7, 8]. It has been suggested to be a possible first symptomatic expression of preclinical AD and a target stage for future AD prevention trials. Therefore, finding some available plasma biomarkers associated with early pathological changes of AD biomarkers in SCD individuals may contribute to creating more practical early diagnostic methods or finding some new factors involved in AD early pathogenesis.

Apolipoproteins are key proteins in the lipid metabolism of peripheral blood and central nervous system and also play an important role in the transport of some key proteins [9, 10]. Previous studies have indicated that Apolipoprotein B (ApoB) and Apolipoprotein A-1 (ApoA1) may be involved in the pathogenesis of AD. Some studies have shown that AD patients had higher levels of circulating ApoB and lower levels of circulating ApoA1 compared with healthy controls [11, 12]. ApoB may promote the formation of amyloid plaques and vascular pathological changes [13, 14]. ApoA1 may have a neuroprotective effect by forming a complex with the A β peptide [12]. However, some studies showed different results [15]. Overall, the number of studies on these factors is still limited. Moreover, to our

knowledge, no studies have been reported on the associations of plasma ApoB, ApoA1 and ApoB/A1 with CSF AD biomarkers, especially in the preclinical stage.

The purpose of our study was to examine whether ApoB, ApoA1, or the ratio of ApoB and ApoA1 (ApoB/A1) can reflect the early pathological changes of CSF AD biomarkers in elderly adults with subjective cognitive decline and to find some potential candidate early biomarkers in blood for AD.

Methods

Participants

The participants without objective cognitive impairment in our study came from the Chinese Alzheimer's Biomarker and Lifestyle (CABLE) database. The CABLE database is an independent replication cohort which has been approved by the Institutional Review Board of Qingdao Municipal Hospital. Since 2017, CABLE has been an ongoing large-scale study majorly focused on AD's risk factors and biomarkers in Chinese Northern Han population. CABLE aimed to determine the genetic and environmental modifiers of AD biomarkers and their utility in early diagnosis. The samples in CABLE study were recruited at Qingdao Municipal Hospital, consisting of cognitively normal individuals without subjective and objective cognitive decline (CN) and individuals with SCD, MCI or AD.

The diagnosis of AD dementia and MCI was made by experienced clinicians according to the criteria of National Institute on Aging-Alzheimer's Association (NIA-AA) workgroups criteria [16, 17]. All clinical evaluations were conducted using standardized questionnaires. Specifically, the general cognitive function was assessed by a China-Modified Mini-Mental State Examination (CM-MMSE) and Montreal Cognitive Assessment (MoCA). The cut-off values after correcting the years of education were used in the MMSE (≤ 24 for 6+ y of education, ≤ 20 for 6- y of education, ≤ 17 for 0 y of education) and MOCA (< 26 for 12+ y of education, < 25 for 12- y of education). The basic living ability was assessed by basic Activities of Daily Living score (ADL). Behavioral or psychological symptoms were assessed by Geriatric Depression Scale (GDS), Hamilton Rating Scale for Depression (HAMD) and Hamilton Rating Scale for Anxiety (HAMA). The GDS and HAMD scores of participants in CABLE database were less than seven. Vascular factors were assessed by Hachinski Inchemic Score (HIS). The HIS scores of participants in our study were less than four. Subjective cognitive decline was assessed by a subjective cognitive decline scale (SCDS) which was designed based on SCD-I recommendations [18, 19]. We distinguished participants with SCD accompanied by particular concerns (worries) from those without objective cognitive decline by the first section of SCDS which included a dichotomous question. Then we chose SCD participants who met the SCD-plus criteria as much as possible [19].

All enrolled participants in the CABLE were Han Chinese aged between 40 to 90 years old. The exclusion criteria include: (1) central nervous system infection, head trauma, epilepsy, multiple sclerosis or other major neurological disorders; (2) major psychological disorders; (3) severe systemic diseases; (4) family history of genetic diseases. All participants underwent biochemical testing as well as blood and CSF sample collection. Demographic information, AD risk factor profile and medical history were collected via a structured questionnaire and an electronic medical record system.

Finally, we included 302 participants without objective cognitive impairment from CABLE database, including 201 CN and 101 SCD participants. All the participants, who had no history of taking lipid-lowering drugs provided both blood and CSF samples.

CSF sample collection and measurements

Collection of fasting lumbar CSF samples and blood samples was performed at Qingdao Municipal Hospital. CSF was drawn and processed within two hours after collection. Each tube was centrifuged at 2000×g for 10 min, and CSF was separated and stored in enzyme free EP tube at -80 centigrade until further use in this study. The thaw/freezing cycle was limited and did not surpass two times. Baseline CSF A β 42, A β 40, t-tau, and p-tau181 were determined with the ELISA kit (Innotest β -AMYLOID (1-42), β -AMYLOID (1-40), hTAU-Ag, and PHOSPHO-TAU (181p); Fujirebio, Ghent, Belgium) on the microplate reader (Thermo Scientific Multiskan MK3). The within-batch precision values were $< 5\%$ (4.5% for A β 42, 3.5% for A β 40, 4.4% for t-tau, and 2.5% for p-tau181). The inter-batch CVs were $< 15\%$ (9.6% for A β 42, 8.1% for A β 40, 12.2% for t-tau, and 11.0% for p-tau181).

Blood sample collection and measurements

Blood samples of all the participants were drawn after an overnight fast. Serum levels of ApoB and ApoA1 were assayed by immunonephelometry using a BN II analyzer (Siemens Healthcare, Marburg, Germany).

QIAamp® DNA Blood Mini Kit (250) was used to extract DNA from Blood samples. And the extracted DNA was separated and stored in enzyme free EP tube at -80 centigrade until the apolipoprotein E (*APOE*) 4 genotyping was completed in this study. Specific loci were selected for genotyping with restriction fragment length polymorphism (RFLP) technology, including the two loci related to *APOE*4 status (rs7412 and rs429358).

Standard protocol approvals, registrations, and patient consents

The CABLE database was conducted in accordance with the Helsinki declaration, and the research program was approved by the Institutional Ethics Committee of Qingdao Municipal Hospital. All study participants or their caregivers provided written informed consent directly.

Statistical analysis

One-way analysis of variance (ANOVA), t-test, Kruskal-Wallis test and Chisquare test were applied in the comparison among groups. Amyloid imaging and neuropathological studies showed that approximately one-third of older adults without objective cognitive impairment had AD pathology in their brains [20-23]. Therefore, in our study, CSF biomarker positive participants were defined as having CSF A β 42 levels in the lower one-third of the distribution of participants (A+: ≤ 116.11 pg/mL) or having p-tau (T+: ≥ 39.17 pg/mL) or t-tau (N+: ≥ 180.59 pg/mL) levels in the upper one-third of the distribution. All plasma and CSF variables were log-transformed to normalize the distributions. Multiple linear regression models were used to examine the cross-sectional associations of baseline plasma ApoB, ApoA1 and ApoB/A1 ratio with CSF biomarkers, adjusting for age, sex, education and *APOE*4 status. Statistical analyses were conducted using R, version 3.5.1. A two-tailed $p < 0.05$ was considered significant.

Results

Characteristics of participants

For the current study, we analyzed data from 302 participants of the CABLE cohort, including 201 CN and 101 SCD (see Table 1 for participants' characteristics). The average age of participants was 63 years; 111 (63%) participants were male; and 40 (13%) participants were *APOE*4 positive. We found no significant difference in age, sex distribution or level of education between two subgroups. Frequency of positive *APOE*4 status showed an increasing trend in SCD subgroup (CN: 11.94%, SCD: 15.84%), although these results were not statistically significant.

As for cognitive levels, we did not find a significant cognitive difference between CN subgroup and SCD subgroup (CM-MMSE: $p = 0.3812$; MOCA: $p = 0.0925$). As for CSF biomarkers, compared with CN subgroup, SCD subgroup had lower levels of A β 42 ($p = 0.0080$) and A β 42/40 ($p < 0.0001$) and higher levels of p-tau ($p = 0.0389$), t-tau/A β 42 ($p < 0.0001$) and p-tau/A β 42 ($p < 0.0001$).

Levels of plasma ApoB and ApoA1 in different diagnostic groups

As for levels of plasma ApoB, in a clinical diagnostic construct, we did not find difference between CN and SCD subgroups (Figure 1a). In an ATN biological construct, all CSF biomarker positive subgroups had lower levels of plasma ApoB compared with CSF biomarker negative subgroups (Figure 1d, e, f). Then, according to the 2018 NIAA research framework, we chose A+T-N-, A+T+N- and A+T+N+ as AD continuum and chose A-T-N- as control group. Results showed that A+T+N+ subgroup showed significantly lower levels of plasma ApoB than A-T-N- ($p < 0.0001$) and A+T-N- ($p = 0.0038$) subgroups. Furthermore, in a diagnostic structure combining clinical diagnosis and biomarkers, SCD participants with A+ had significantly lower levels of plasma ApoB than CN participants without pathologic changes in CSF biomarkers ($p = 0.0018$) (Figure 1c). As for ApoA1, we didn't find any difference in both clinical construct and biological construct (results were not shown).

Associations of ApoB, ApoA1, and ApoB/A1 with CSF AD core biomarkers in total participants without objective cognitive impairment

The results on associations of ApoB, ApoA1, and ApoB/A1 with CSF biomarkers in total participants without objective cognitive impairment were shown in Table 2. Results showed that higher levels of plasma ApoB were significantly associated with increased levels of CSF A β 42 ($\beta = 0.34$, $p = 0.0009$) and A β 42/40 ($\beta = 0.35$, $p = 0.0038$) but not with A β 40 (Table 2, Figure 2). Although no association of ApoB was found with t-tau and p-tau, higher levels of plasma ApoB were significantly associated with decreased levels of CSF t-tau/A β 42 ($\beta = -0.47$, $p < 0.0001$) and p-tau/A β 42 ($\beta = -0.41$, $p < 0.0001$) (Table 2, Figure 2). Similar associations were found of ApoB/A1 with these A β -related biomarkers (A β 42, A β 42/40, t-tau/A β 42 and p-tau/A β 42) (Table 2), but these associations seemed less significant than those

of ApoB (Table 2). We did not find any association between ApoA1 and all CSF biomarkers (A β 42, t-tau, p-tau, A β 40, A β 42/40, t-tau/A β 42 and p-tau/A β 42).

Furthermore, we added the interaction between *APOE* ϵ 4 status and ApoB into multiple linear regression models. Results showed that these associations between ApoB and A β -related biomarkers were independent of *APOE* ϵ 4 status. In these models, both ApoB and *APOE* ϵ 4 status were associated with A β -related biomarkers, but their interactions were not (see Additional file 1). Moreover, subgroup analyses of *APOE* ϵ 4 status also showed that associations between ApoB and A β -related biomarkers were similar for different *APOE* ϵ 4 statuses (see Additional file 2).

Associations of ApoB, ApoA1, and ApoB/A1 with CSF AD core biomarkers in different diagnostic subgroups

Results from subgroup analyses of different diagnostic groups were shown in Table 3. As for ApoB, these associations between ApoB and A β -related biomarkers (A β 42, A β 42/40, t-tau/A β 42 and p-tau/A β 42) still remained significant in both CN and SCD subgroups (Table 3, Figure 3). It is worth noting that these associations were more significant in the SCD subgroup than in the CN subgroup (Table 3). In addition, we found that higher levels of ApoB were also associated with decreased CSF t-tau (β = -0.37, p = 0.0224), p-tau (β = -0.24, p = 0.0086) and A β 40 (β = -0.43, p = 0.0297) in SCD subgroup, while these associations were not found in the CN subgroup (Table 3, Figure 3). As for ApoB/A1, significant associations between ApoB/A1 and all CSF biomarkers (A β 42, t-tau, p-tau, A β 40, A β 42/40, t-tau/A β 42 and p-tau/A β 42) were found only in the SCD subgroup but not in the CN subgroup (Table 3). We did not find any association between ApoA1 and all CSF biomarkers (A β 42, t-tau, p-tau, A β 40, A β 42/40, t-tau/A β 42 and p-tau/A β 42).

Given that there may be different metabolic states of apolipoproteins between overweight and normal populations, we further tested our results found above in different weight participants that were classified according to Body Mass Index (BMI) (normal weight participants: BMI < 25, overweight participants: BMI $\geq$ 25). To our surprise, in both CN subgroup and SCD subgroup, almost all of associations found above existed only in the overweight participants but not in normal weight participants (Table 4, Table 5).

Discussion

This study is the first to systematically explore cross-sectional associations of CSF AD core biomarkers with ApoB, ApoA1, and ApoB/A1 in elderly adults without objective cognitive impairment. The main finding of this study was that plasma ApoB may have some protective effects on CSF AD core biomarkers in elderly adults without objective cognitive impairment independently of *APOE* ϵ 4 genotype, especially in the SCD overweight population. This finding was novel and potentially important, because it suggested that levels of plasma ApoB added information beyond *APOE* ϵ 4 status on the level of CSF AD core biomarkers in the preclinical stage, which may contribute to the discovery of new pathogenic mechanisms and therapeutic targets.

First of all, we analyzed the characteristics of SCD participants included in our study. Results showed that, compared with the CN subgroup, although there was no significant cognitive decline, there were obvious changes in CSF AD core biomarkers (especially A β -related biomarkers) in the SCD subgroup, which were consistent with previous studies [8, 24]. In fact, accumulating cross-sectional or longitudinal evidence has supported that SCD occurred at the preclinical stage of AD and might serve as a symptomatic indicator of preclinical AD [7, 8, 24-30]. Our analysis of population characteristics showed that, in our study, SCD participants had preclinical AD characteristics.

Then we explored the distribution of ApoB and ApoA in the diagnostic and biological constructs. Results showed that SCD participants with significant AD biological characteristics had a lower ApoB level (Figure 1) suggesting that the level of ApoB may decrease in the preclinical stage of AD.

Finally we detailed the associations of ApoB, ApoA1, and ApoB/A1 with CSF AD core biomarkers in total participants or different diagnostic subgroups and analyzed the factors that might affect these associations. We firstly found that plasma ApoB and ApoB/A1 were associated with CSF AD core biomarkers, especially with the A β -related biomarkers (A β 42, A β 42/40, t-tau/A β 42 and p-tau/A β 42). It is worth noting that these associations were much more significant in SCD participants. Surprisingly, inconsistent with previous studies, above associations showed that plasma ApoB and ApoB/A1 appeared to have some protective effects that may alleviate the early pathological changes of AD. To be precise, higher ApoB and ApoB/A1 were associated with increased CSF A β 42 and A β 42/40, as well as with decreased CSF t-tau, p-tau, A β 40, t-tau/A β 42 and p-tau/A β 42. No previous studies have focused on the associations of plasma ApoB and ApoB/A1 with CSF AD core biomarkers in cognitively normal or SCD participant. However, two autopsy studies in AD patients indicated that ApoB may be a risk factor for the late stage of AD. One showed that serum ApoB were positive associated with the amount

of A β 42 in AD brains [14]. The other showed that ApoB immunoreactivity was positively associated with cerebral amyloids [13]. The exact causes of these seemingly contradictory results were still unclear, but we thought that there were several possible reasons as follows.

Firstly, the different stages of the disease might contribute to this contradiction. Previous studies have found that changes in the blood-brain barrier contributed differently to the dynamic changes of CSF AD core biomarkers in different stages of disease. CSF AD core biomarker (especially A β 42) may be temporarily elevated and its significant deposition in brain may occur when the blood-brain barrier was significantly impaired [31]. As a possible transporter of CSF biomarkers, ApoB may facilitate the processes of early clearance and late deposition of biomarkers in brain. In addition, in the stage of the disease, atherosclerosis and other vascular damages caused by ApoB-LDL may also aggravate the pathological changes, which may not be apparent in the early stage of the disease.

Secondly, genetic factors may also contribute to this contradiction. In this study, we tested the influences of *APOE* ϵ 4 status on above associations. We found that these protective associations were independent of the *APOE* ϵ 4 status, based on the following evidences. a) Both plasma ApoB and *APOE* ϵ 4 status were associated with CSF AD core biomarkers, but their interaction was not associated with these biomarkers. b) The results of the subgroups analysis were similar among population with different *APOE* ϵ 4 status. This independence on *APOE* ϵ 4 was also seen in a previous study on the association between serum ApoB and amount of A β 42 in AD brains [14]. Collectively, these data provided evidences that ApoB and *APOE* ϵ 4 status may contribute to the changes of CSF AD core biomarkers, independently of each other. However, a recent study found that rare genetic coding variants of *APOB* gene were strongly associated with familial AD, which indicated that *APOB* gene might also harbor protective and deleterious variants in sporadic AD similar to the *APOE* ϵ 2 and *APOE* ϵ 4 alleles [32]. Identifying such a protective coding variant of *APOB* gene would greatly strengthen the link between ApoB and AD pathogenesis.

In addition, some other factors may also contribute to this contradiction, such as changes involved low density lipoprotein cholesterol metabolism (the low density lipoprotein receptor and proprotein convertase subtilisin/kexin type 9). So given the concept that normal weight populations and overweight populations often have different lipid metabolism status, we further tested our results found above in different weight participants. It is worth noting that these protective effects of plasma ApoB on CSF AD core biomarkers were particularly stronger in overweight participants compared with the normal weight participants. These results indicated that different lipid metabolism status may also influence above associations, which may contribute to the contradiction. Moreover, growing body of studies suggested that overweight in older age may be a protective factor for AD. Overweight older adults had a lower incidence or a later onset of AD than those of normal weight or low weight [33-35]. Our results indicated that the protective effect of plasma ApoB on CSF AD core biomarkers may be the potential mechanism. In other words, the plasma ApoB's ability to alleviate early pathological changes may contribute to slow the progression of the disease in overweight older adults, but not in normal weight older adults.

Overall, we found protective associations of plasma ApoB with CSF AD core biomarkers, these associations may be affected by disease's stages and lipid metabolism status or some other indefinite factors such as genetic factors. In this study we did not find any associations between ApoA1 and CSF AD core biomarkers, which indicated that ApoA1 may not have significant effects on the early pathological changes in population without objective cognitive impairment.

There were some potential limitations in our study. Firstly, this was a cross-sectional study and the results still needed to be tested in larger longitudinal cohort. Secondly, the sample sizes of some subgroups were limited, which may affect the results to some extent.

Conclusions

This study is the first to find some protective associations of plasma ApoB, but not ApoA1, with CSF AD core biomarkers in the preclinical stage of AD. This finding indicated that ApoB may play different roles in different stages of AD. Further studies to identify the underlying mechanisms would greatly strengthen the link between ApoB and AD pathogenesis, which may contribute to the discovery of new pathogenic mechanisms and therapeutic targets.

Abbreviations

AD: Alzheimer disease; ADL: Activities of Daily Living score; A β 42: β -amyloid 42; ANOVA: One-way analysis of variance; ApoA1: Apolipoprotein A-1; ApoB: Apolipoprotein B; APOE: apolipoprotein E; BMI: Body Mass Index; CABLE: Chinese Alzheimer's Biomarker and Lifestyle; CM-MMSE: China-Modified Mini-Mental State Examination; CSF: cerebrospinal fluid; GDS: Geriatric Depression Scale; HAMA: Hamilton Rating Scale for Anxiety; HAMD: Hamilton Rating Scale for Depression; HIS: Hachinski Inchemic Score; MCI: mild cognitively

impaired; MoCA: Montreal Cognitive Assessment; NIAA: National Institute on Aging and Alzheimer's Association; p-tau: phosphorylated tau; SCD: Subjective cognitive decline; t-tau: total tau

Declarations

Acknowledgements

Colleagues who have made contributions to build the CABLE cohort.

Funding

This study was supported by grants from the National Natural Science Foundation of China (91849126), the National Key R&D Program of China (2018YFC1314700), Shanghai Municipal Science and Technology Major Project (No.2018SHZDZX01) and ZHANGJIANG LAB, Tianqiao and Chrissy Chen Institute, and the State Key Laboratory of Neurobiology and Frontiers Center for Brain Science of Ministry of Education, Fudan University.

Authors' contributions

HH, Lan Tan, YLB, and JTY did the manuscript preparation and drafting. HH, LT, YLB, XW, XNS, XHH, YHM, and JTY did the clinical assessments and data acquisition. LT and JTY did the clinical diagnosis. HH, LT, YLB, XW, XNS, XHH, YHM, QD, and JTY did the data analysis and interpretation. JTY is responsible for the study conception and design. All authors have contributed to the manuscript revising and editing critically for important intellectual content and given final approval of the version and agreed to be accountable for all aspects of the work presented here.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

The study was approved by the local ethics committee, and all individuals gave written informed consent to participate.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests

References

1. Selkoe, D.J. and J. Hardy, *The amyloid hypothesis of Alzheimer's disease at 25 years*. EMBO Mol Med, 2016. **8**(6): p. 595-608.
2. Lane, C.A., J. Hardy, and J.M. Schott, *Alzheimer's disease*. Eur J Neurol, 2018. **25**(1): p. 59-70.
3. Weiner, M.W., et al., *The Alzheimer's disease neuroimaging initiative: progress report and future plans*. Alzheimers Dement, 2010. **6**(3): p. 202-11 e7.
4. Olsson, B., et al., *CSF and blood biomarkers for the diagnosis of Alzheimer's disease: a systematic review and meta-analysis*. Lancet Neurol, 2016. **15**(7): p. 673-684.
5. Molinuevo, J.L., et al., *Current state of Alzheimer's fluid biomarkers*. Acta Neuropathol, 2018. **136**(6): p. 821-853.
6. Jack, C.R., Jr., et al., *NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease*. Alzheimers Dement, 2018. **14**(4): p. 535-562.
7. Glodzik-Sobanska, L., et al., *Subjective memory complaints: presence, severity and future outcome in normal older subjects*. Dement Geriatr Cogn Disord, 2007. **24**(3): p. 177-84.
8. Wolfsgriber, S., et al., *Prevalence of abnormal Alzheimer's disease biomarkers in patients with subjective cognitive decline: cross-sectional comparison of three European memory clinic samples*. Alzheimers Res Ther, 2019. **11**(1): p. 8.
9. Brown, M.S. and J.L. Goldstein, *How LDL receptors influence cholesterol and atherosclerosis*. Sci Am, 1984. **251**(5): p. 58-66.

10. Emerging Risk Factors, C., et al., *Lipid-related markers and cardiovascular disease prediction*. JAMA, 2012. **307**(23): p. 2499-506.
11. Raygani, A.V., et al., *Association between apolipoprotein E polymorphism and serum lipid and apolipoprotein levels with Alzheimer's disease*. Neurosci Lett, 2006. **408**(1): p. 68-72.
12. Merched, A., et al., *Decreased high-density lipoprotein cholesterol and serum apolipoprotein AI concentrations are highly correlated with the severity of Alzheimer's disease*. Neurobiol Aging, 2000. **21**(1): p. 27-30.
13. Namba, Y., H. Tsuchiya, and K. Ikeda, *Apolipoprotein B immunoreactivity in senile plaque and vascular amyloids and neurofibrillary tangles in the brains of patients with Alzheimer's disease*. Neurosci Lett, 1992. **134**(2): p. 264-6.
14. Kuo, Y.M., et al., *Elevated low-density lipoprotein in Alzheimer's disease correlates with brain abeta 1-42 levels*. Biochem Biophys Res Commun, 1998. **252**(3): p. 711-5.
15. Tynkkynen, J., et al., *Apolipoproteins and HDL cholesterol do not associate with the risk of future dementia and Alzheimer's disease: the National Finnish population study (FINRISK)*. Age (Dordr), 2016. **38**(5-6): p. 465-473.
16. Albert, M.S., et al., *The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease*. Alzheimers Dement, 2011. **7**(3): p. 270-9.
17. McKhann, G.M., et al., *The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease*. Alzheimers Dement, 2011. **7**(3): p. 263-9.
18. Jessen, F., et al., *A conceptual framework for research on subjective cognitive decline in preclinical Alzheimer's disease*. Alzheimers Dement, 2014. **10**(6): p. 844-52.
19. Molinuevo, J.L., et al., *Implementation of subjective cognitive decline criteria in research studies*. Alzheimers Dement, 2017. **13**(3): p. 296-311.
20. Reiman, E.M., et al., *Fibrillar amyloid-beta burden in cognitively normal people at 3 levels of genetic risk for Alzheimer's disease*. Proc Natl Acad Sci U S A, 2009. **106**(16): p. 6820-5.
21. Bennett, D.A., et al., *Neuropathology of older persons without cognitive impairment from two community-based studies*. Neurology, 2006. **66**(12): p. 1837-44.
22. Hulette, C.M., et al., *Neuropathological and neuropsychological changes in "normal" aging: evidence for preclinical Alzheimer disease in cognitively normal individuals*. J Neuropathol Exp Neurol, 1998. **57**(12): p. 1168-74.
23. Knopman, D.S., et al., *Neuropathology of cognitively normal elderly*. J Neuropathol Exp Neurol, 2003. **62**(11): p. 1087-95.
24. Miebach, L., et al., *Which features of subjective cognitive decline are related to amyloid pathology? Findings from the DELCODE study*. Alzheimers Res Ther, 2019. **11**(1): p. 66.
25. Jessen, F., et al., *Prediction of dementia by subjective memory impairment: effects of severity and temporal association with cognitive impairment*. Arch Gen Psychiatry, 2010. **67**(4): p. 414-22.
26. Reisberg, B., et al., *Outcome over seven years of healthy adults with and without subjective cognitive impairment*. Alzheimers Dement, 2010. **6**(1): p. 11-24.
27. Wang, Y., et al., *Altered default mode network connectivity in older adults with cognitive complaints and amnesic mild cognitive impairment*. J Alzheimers Dis, 2013. **35**(4): p. 751-60.
28. Perrotin, A., et al., *Subjective cognition and amyloid deposition imaging: a Pittsburgh Compound B positron emission tomography study in normal elderly individuals*. Arch Neurol, 2012. **69**(2): p. 223-9.
29. Pritchep, L.S., et al., *Prediction of longitudinal cognitive decline in normal elderly with subjective complaints using electrophysiological imaging*. Neurobiol Aging, 2006. **27**(3): p. 471-81.
30. van Harten, A.C., et al., *Cerebrospinal fluid Abeta42 is the best predictor of clinical progression in patients with subjective complaints*. Alzheimers Dement, 2013. **9**(5): p. 481-7.
31. Potter, R., et al., *Increased in vivo amyloid-beta42 production, exchange, and loss in presenilin mutation carriers*. Sci Transl Med, 2013. **5**(189): p. 189ra77.
32. Wingo, T.S., et al., *Association of Early-Onset Alzheimer Disease With Elevated Low-Density Lipoprotein Cholesterol Levels and Rare Genetic Coding Variants of APOB*. JAMA Neurol, 2019. **76**(7): p. 809-817.
33. Hughes, T.F., et al., *Association between late-life body mass index and dementia: The Kame Project*. Neurology, 2009. **72**(20): p. 1741-6.

34. Lee, J.Y., et al., *Risk of Incident Dementia According to Metabolic Health and Obesity Status in Late Life: A Population-Based Cohort Study*. J Clin Endocrinol Metab, 2019. **104**(7): p. 2942-2952.
35. Tolppanen, A.M., et al., *Midlife and late-life body mass index and late-life dementia: results from a prospective population-based cohort*. J Alzheimers Dis, 2014. **38**(1): p. 201-9.

Tables

Table 1. Characteristics of study participants from CABLE database

Variable	CN	SCD	Total	P
N	201	101	302	-
AGE (year) mean (SD)	62.08 (11.42)	64.37 (10.00)	62.85 (11.00)	0.0783
SEX (Female/Male)	71/130	40/61	111/191	0.5475
Education (year) mean (SD)	9.90 (4.49)	10.49 (4.34)	10.09 (4.44)	0.2427
<i>APOE</i> ε4 N (%)	24 (11.94)	16 (15.84)	40 (13.25)	0.4450
CSF Aβ42 (pg/mL) mean (SD)	181.96 (130.45)	149.75 (81.13)	171.19 (117.16)	0.0080
CSF t-tau (pg/mL) mean (SD)	175.83 (82.36)	186.86 (85.62)	179.52 (83.49)	0.0509
CSF p-tau (pg/mL) mean (SD)	37.55 (9.90)	39.02 (8.55)	38.04 (9.48)	0.0389
CSF Aβ40 (pg/mL) mean (SD)	6525.83 (3070.02)	6998.23 (2339.81)	6683.82 (2851.52)	0.1386
CSF Aβ42/40 mean (SD)	0.03 (0.03)	0.02 (0.01)	0.03 (0.03)	<0.0001
CSF t-tau/Aβ42 mean (SD)	1.16 (0.69)	1.57 (1.07)	1.30 (0.86)	<0.0001
CSF p-tau/Aβ42 mean (SD)	0.25 (0.10)	0.30 (0.13)	0.27 (0.11)	<0.0001
CM-MMSE mean (SD)	27.82 (2.13)	27.50 (2.50)	27.71 (2.26)	0.3812
MOCA mean (SD)	26.92 (0.99)	26.74 (0.98)	26.86 (0.99)	0.0925
ApoB (mg/mL) mean (SD)	1.04 (0.25)	1.02 (0.27)	1.03 (0.26)	0.6097
ApoA1 (mg/mL) mean (SD)	1.45 (0.30)	1.43 (0.28)	1.45 (0.29)	0.9588
ApoB/A1 mean (SD)	0.74 (0.20)	0.74 (0.25)	0.74 (0.22)	0.7560
BMI mean (SD)	25.51 (3.28)	25.24 (3.25)	25.34 (3.40)	0.8450

Abbreviations: CN, cognitively normal participants; SCD, participants with subjective cognitive decline; *APOE*, apolipoprotein E gene; CSF, cerebrospinal fluid; Aβ, amyloid-β; t-tau, total tau protein; p-tau, phosphorylated tau protein; CM-MMSE, China-Modified Mini-Mental State Examination; MOCA, Montreal Cognitive Assessment; ApoB, Apolipoprotein B; ApoA1, Apolipoprotein A-1; BMI, Body Mass Index; SD, standard deviation
 Bold indicated that the results were statistically significant.

Table 2. Results of cross-sectional associations in total participants

Variable	ApoB		ApoA1		ApoB/A1	
	β	P	β	P	β	P
CSF Aβ42	0.34	0.0009	0.07	0.5930	0.22	0.0135
CSF t-tau	-0.13	0.1556	-0.06	0.6207	-0.07	0.3656
CSF p-tau	-0.07	0.1823	-0.01	0.9055	-0.05	0.2823
CSF Aβ40	-0.06	0.5620	-0.03	0.8170	-0.03	0.7260
CSF Aβ42/40	0.35	0.0038	0.17	0.2920	0.19	0.0179
CSF t-tau/Aβ42	-0.47	<0.0001	-0.13	0.3918	-0.29	0.0034
CSF p-tau/Aβ42	-0.41	<0.0001	-0.08	0.5331	-0.27	0.0015

Abbreviations: CSF, cerebrospinal fluid; A β , amyloid- β ; t-tau, total tau protein; p-tau, phosphorylated tau protein; ApoB, Apolipoprotein B; ApoA1, Apolipoprotein A-1

Bold indicated that the results were statistically significant.

Multiple linear regression models were used to examine the crosssectional associations between baseline plasma ApoB, ApoA1 and ApoB/A1 levels with CSF biomarkers, adjusting for age, sex, education and *APOE*4 status.

Table 3. Results of cross-sectional associations in different diagnostic subgroups

Variable	CN						SCD					
	ApoB		ApoA1		ApoB/A1		ApoB		ApoA1		ApoB/A1	
	β	P	β	P	β	P	β	P	β	P	β	P
CSF A β 42	0.33	0.0133	0.06	0.7210	0.23	0.0531	0.50	0.0084	0.09	0.8010	0.39	0.0258
CSF t-tau	-0.07	0.5586	-0.08	0.5941	-0.01	0.8840	-0.37	0.0224	-0.19	0.5183	-0.25	0.0110
CSF p-tau	-0.01	0.9937	-0.04	0.6545	0.02	0.7604	-0.24	0.0086	-0.10	0.5762	-0.17	0.0394
CSF A β 40	-0.02	0.8860	-0.14	0.4460	0.05	0.6860	-0.43	0.0297	0.01	0.9750	-0.35	0.0489
CSF A β 42/40	0.35	0.0150	0.20	0.2869	0.18	0.1622	0.70	0.0009	0.46	0.2510	0.46	0.0208
CSF t-tau/A β 42	-0.40	0.0051	-0.14	0.4501	-0.25	0.0524	-0.87	0.0001	-0.28	0.5200	-0.64	0.0024
CSF p-tau/A β 42	-0.33	0.0073	-0.10	0.5296	-0.21	0.0548	-0.74	<0.0001	-0.19	0.6080	-0.56	0.0011

Abbreviations: CN, cognitively normal participants; SCD, participants with subjective cognitive decline; CSF, cerebrospinal fluid; A β , amyloid- β ; t-tau, total tau protein; p-tau, phosphorylated tau protein; ApoB, Apolipoprotein B; ApoA1, Apolipoprotein A-1

Bold indicated that the results were statistically significant.

Multiple linear regression models were used to examine the crosssectional associations between baseline plasma ApoB, ApoA1 and ApoB/A1 levels with CSF biomarkers, adjusting for age, sex, education and *APOE*4 status.

Table 4. Results of cross-sectional associations among different weight participants in CN subgroup

Variable	Normal-weight (BMI < 25) N = 89						Overweight (BMI $\geq$ 25) N = 114					
	ApoB		ApoA1		ApoB/A1		ApoB		ApoA1		ApoB/A1	
	β	P	β	P	β	P	β	P	β	P	β	P
CSF A β 42	0.16	0.3920	0.28	0.2090	-0.03	0.8430	0.39	0.0395	-0.09	0.7390	0.36	0.0371
CSF t-tau	0.03	0.8749	0.10	0.6211	-0.03	0.8163	-0.13	0.4190	-0.28	0.1989	0.02	0.9070
CSF p-tau	-0.01	0.9330	0.07	0.5612	-0.04	0.6141	0.02	0.8650	-0.15	0.2670	0.08	0.3750
CSF A β 40	-0.19	0.3650	0.02	0.4180	-0.24	0.1670	0.09	0.6580	-0.42	0.1180	0.26	0.1510
CSF A β 42/40	0.35	0.0847	0.09	0.7140	0.21	0.2320	0.30	0.1490	0.34	0.2360	0.10	0.6050
CSF t-tau/A β 42	-0.14	0.5080	-0.18	0.4480	-0.01	0.9970	-0.52	0.0086	-0.19	0.4885	-0.34	0.0578
CSF p-tau/A β 42	-0.17	0.3580	-0.21	0.3290	-0.01	0.9510	-0.37	0.0263	-0.06	0.7971	-0.28	0.0673

Abbreviations: CN, cognitively normal participants; BMI, Body Mass Index; N, number of samples; CSF, cerebrospinal fluid; Aβ, amyloid-β; t-tau, total tau protein; p-tau, phosphorylated tau protein; ApoB, Apolipoprotein B; ApoA1, Apolipoprotein A-1

Bold indicated that the results were statistically significant.

Multiple linear regression models were used to examine the crosssectional associations between baseline plasma ApoB, ApoA1 and ApoB/A1 levels with CSF biomarkers, adjusting for age, sex, education and *APOE*ε4 status.

Table 5. Results of cross-sectional associations among different weight participants in SCD subgroup

Variable	Normal-weight (BMI < 25) N = 49						Overweight (BMI ≥ 25) N = 52					
	ApoB		ApoA1		ApoB/A1		ApoB		ApoA1		ApoB/A1	
	β	P	β	P	β	P	β	P	β	P	β	P
CSF	0.08	0.8350	0.38	0.4610	-0.08	0.7850	0.64	0.0052	-0.59	0.2790	0.65	0.0017
Aβ42												
CSF t-tau	-0.41	0.2590	-0.26	0.5900	-0.15	0.5910	-0.51	0.0012	-0.46	0.2301	-0.36	0.0189
CSF p-tau	-0.18	0.3280	-0.05	0.8190	-0.08	0.5420	-0.32	0.0051	-0.30	0.2740	-0.23	0.0417
CSF Aβ40	-0.62	0.1270	-0.23	0.6755	-0.29	0.3674	-0.54	0.0099	-0.29	0.5637	-0.42	0.0366
CSF Aβ42/40	0.70	0.1410	0.61	0.3330	0.20	0.5910	0.77	0.0017	0.51	0.3998	0.58	0.0147
CSF t-tau/Aβ42	-0.50	0.3440	-0.64	0.3470	-0.07	0.8670	-1.14	<0.0001	0.13	0.8440	-1.01	<0.0001
CSF p-tau/Aβ42	-0.26	0.4976	-0.44	0.3760	0.01	0.9980	-0.96	<0.0001	0.29	0.6380	-0.88	<0.0001

Abbreviations: SCD, participants with subjective cognitive decline; BMI, Body Mass Index; N, number of samples; CSF, cerebrospinal fluid; Aβ, amyloid-β; t-tau, total tau protein; p-tau, phosphorylated tau protein; ApoB, Apolipoprotein B; ApoA1, Apolipoprotein A-1

Bold indicated that the results were statistically significant.

Multiple linear regression models were used to examine the crosssectional associations between baseline plasma ApoB, ApoA1 and ApoB/A1 levels with CSF biomarkers, adjusting for age, sex, education and *APOE*ε4 status.

Figures

Figure 1

Levels of plasma ApoB in different groups (a) Levels of plasma ApoB in a clinical diagnostic construct. CN (N = 201): cognitively normal individuals without subjective and objective cognitive decline; SCD (N = 101): individuals with subjective cognitive decline; (b) Levels of plasma ApoB in an ATN biological construct. A-T-N- (N = 120), A+T-N- (N = 63), A+T+N- (N = 5), A+T+N+ (N = 28); (c) Levels of plasma ApoB in a diagnostic structure combining clinical diagnosis and biomarkers. CN- (N = 90): cognitively normal individuals without subjective and objective cognitive decline and with three negative CSF biomarkers [A-T-N-]; CN+ (N = 57): cognitively normal individuals without subjective and objective cognitive decline and with positive CSF A β 42 [A+]; SCD+ (N = 44): individuals with subjective cognitive decline and positive CSF A β 42 [A+]. (d) (e) (f) Levels of plasma ApoB in A or T or N construct respectively. All intergroup comparisons were tested by ANOVA or t-test.

Figure 2

Associations between ApoB and CSF AD biomarkers in total participants Abbreviations: CSF, cerebrospinal fluid; A β , amyloid- β ; t-tau, total tau protein; p-tau, phosphorylated tau protein; ApoB, Apolipoprotein B Multiple linear regression models were used to examine the cross-sectional associations between baseline plasma ApoB levels with CSF biomarkers, adjusting for age, sex, education and APOE ϵ 4 status.

Figure 3

Associations between ApoB and CSF AD biomarkers in SCD participants Abbreviations: CSF, cerebrospinal fluid; A β , amyloid- β ; t-tau, total tau protein; p-tau, phosphorylated tau protein; ApoB, Apolipoprotein B Multiple linear regression models were used to examine the crosssectional associations between baseline plasma ApoB levels with CSF biomarkers, adjusting for age, sex, education and APOE ϵ 4 status.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile.docx](#)