Evaluation of Early Innate and Adaptive Immune Responses to the TB Vaccine Mycobacterium Bovis BCG and Vaccine Candidate BCGdBCG1419c

Manuja Gunasena

The Ohio State University

Rajni Kant Shukla

The Ohio State University

Naiquan Yao

The Ohio State University

Oscar Rosas Mejia

The Ohio State University

Michael D Powell

The Ohio State University

Kenneth J Oestreich

The Ohio State University

Michel de Jesús Aceves-Sánchez

Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco

Mario Alberto Florez Valdez

Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco

Namal P M Liyanage

The Ohio State University

Richard T Robinson (✉ richard.robinson@osumc.edu)

The Ohio State University

Research Article

Keywords: Mycobacterium bovis, lungs, higher frequencies, inflammatory, TB vaccine candidate, BCG efficacy

Posted Date: November 11th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1032864/v1>

License: (cc) (i) This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The vaccine *Mycobacterium bovis* Bacillus Calmette–Guérin (BCG) elicits an immune response that is protective against certain forms of tuberculosis (TB); however, because BCG efficacy is limited it is important to identify alternative TB vaccine candidates. Recently, the BCG deletion mutant and vaccine candidate BCGDBCG1419c was demonstrated to survive longer in intravenously infected BALB/c mice due to enhanced biofilm formation, and better protected both BALB/c and C57BL/6 mice against TB-induced lung pathology during chronic stages of infection, relative to BCG controls. BCGDBCG1419c-elicited protection also associated with lower levels of proinflammatory cytokines (i.e. IL6, TNF α) at the site of infection in C57BL/6 mice. Given the distinct immune profiles of BCG- and BCGDBCG1419c-immunized mice during chronic TB, we set out to determine if there are early immunological events which distinguish these two groups, using multi-dimensional flow cytometric analysis of the lungs and other tissues soon after immunization. Our results demonstrate a number of innate and adaptive response differences between BCG- and BCGDBCG1419c-immunized mice which are consistent with the latter being longer lasting and potentially less inflammatory, including lower frequencies of exhausted CD4⁺ T helper (T_H) cells and higher frequencies of IL10-producing T cells, respectively. These studies suggest the use of BCGDBCG1419c may be advantageous as an alternative TB vaccine candidate.

Introduction

Tuberculosis (TB) is the leading cause of death from an infectious disease ¹. TB is caused by aerogenic transmission of *Mycobacterium tuberculosis* (Mtb), a bacterial species which infects alveolar macrophages and can disseminate to extrapulmonary tissues via lymphatic or hematogenous spread ². TB has a spectrum of clinical manifestations that vary depending on the host response, including severe, active, chronic, subclinical and latent forms ³. Improved socioeconomic conditions, public health practices and the use of effective drug treatment have reduced global TB rates; however, these rates fell short of the World Health Organization (WHO) goal of reversing TB incidence by 2015 ⁴, and are not on target to achieve WHO benchmarks for ending the TB epidemic by 2030 ⁵. For these reasons, it is important for there to be a safe vaccine which effectively promotes TB resistance.

The TB vaccine Bacillus Calmette-Guerin (BCG) is a live-attenuated strain of *Mycobacterium bovis* whose human virulence has been attenuated by multiple passages in artificial culture. BCG is administered throughout the world as a neonatal vaccine against severe forms of childhood TB ⁶; however, the use of BCG in children is controversial due to its variable efficacy ⁷, as well as the number of adverse events which stem from its early immunogenicity ⁸. Within the first two weeks post-immunization, nearly all BCG recipients experience mild adverse events, ranging from papule formation at the site of vaccination, which may become ulcerated or leave a scar, to swelling of the draining lymph nodes. Severe adverse events include more pronounced forms of skin lesions (e.g. BCG lupus vulgaris), lymphadenitis with suppuration, and osteitis. Due to the number of adverse inflammatory events which accompany BCG-immunization, and increased vaccine hesitancy among parents in TB-endemic countries ⁹⁻¹² who are

reluctant to expose their children to inflammatory vaccines (regardless of their efficacy), it is important to identify TB vaccine candidates which—relative to BCG—have reduced early reactogenicity but comparable or better protective efficacy against TB.

Recently, the vaccine candidate BCGDBCG1419c was demonstrated to confer equal and/or better protection against experimental TB compared to its parental strain, BCG Pasteur, in a variety of mouse models (BALB/c^{13,14}, C57BL/6^{15,16}, and B6D2F1 [C57BL/6J × DBA/2J]¹³), including a type 2 diabetic BALB/c model¹⁷, while in guinea pigs it was more effective than BCG in reducing hematogenous spread at 2 months post-infection¹⁸. The *M. bovis* BCG1419c gene encodes a phosphodiesterase activity that hydrolyzes bis-(3'-5')-cyclic dimeric GMP (c-di-GMP), a molecule which reduces bacterial motility and increases biofilm formation (biofilms are sessile bacteria encased within a matrix of an extracellular polymeric substance)¹⁹. As a consequence of BCG1419c deletion, BCGDBCG1419c strain has an increased capacity for biofilm production and persists longer in immunocompetent mice relative to BCG Pasteur after intravenous infection²⁰. Importantly, BCGDBCG1419c-immunized mice also exhibit less lung pathology post Mtb-infection, relative to the BCG-immunized mice that were later infected with Mtb¹⁵. Interestingly, at 60 days post-vaccination, BCGDBCG1419c-immunized mice showed increased numbers of T CD3⁺ CD4⁺ and of T CD3⁺CD4⁺CD44⁺ cells in BALF obtained from C57BL/6 as compared with parental BCG¹⁶. Also, in spleens of BALB/c mice, BCGDBCG1419c increased T CD3⁺CD4⁺IFNγ⁺ and T CD3⁺CD8⁺IFNγ⁺ cells with respect to parental BCG¹³. Moreover, we have observed that BCGDBCG1419c promote a differential presence in lungs of IFNγ-producing T cells, and activated macrophages in a BALB/c model of active and chronic TB¹³ while it changed the relative presence in lungs of B, CD8⁺, and dendritic cells, in a BALB/c model of TB-type 2 diabetes¹⁷. All these findings show that BCG and BCGDBCG1419c affect the relative composition of adaptive immune cells pre- and post-infection in diverse organs and tissues of vaccinated hosts.

Given that BCGDBCG1419c is an effective vaccine in multiple models of experimental TB, outperforming BCG in preventing TB disease measured as lung pathology, and spread to spleen, we carried out a series of experiments to determine if early immunogenicity of BCGDBCG1419c is comparable or distinct from that elicited by BCG, the adverse inflammatory effects of which are typically observed within 2-3 weeks post-immunization. Specifically, we immunized wild type mice and a transgenic IL10 reporter mouse strain with BCG or BCGDBCG1419c, and subsequently performed deep immunophenotyping to comprehensively identify the innate and adaptive immune events that occur within the first 2-3 weeks post-immunization, as well the cellular sources of IL10²²⁻²⁵. Our data demonstrate heretofore unknown, early immunological events induced by BCG, as well as phenotypic and functional differences between the immune responses elicited by BCG and BCGDBCG1419c. The relevance of our data to BCG mechanisms of early immunogenicity, as well as understanding of BCGDBCG1419c efficacy, are discussed.

Methods

Institutional approvals and guidelines. This study and its associated experiments were approved by The Ohio State University (OSU) Institutional Biosafety Committee (IBC), as well as the OSU Institutional Animal Care and Use Committee (IACUC). The study was performed in accordance with relevant institutional guidelines and is reported in accordance with ARRIVE guidelines (<https://arriveguidelines.org>).

BCG culture. Frozen vials of *M. bovis* BCG Pasteur strain (BCG) and its isogenic derivative, BCGDBCG1419c, which lacks a cyclic di-GMP phosphodiesterase encoded by the gene *BCG1419c*²⁰, were received from Colorado State University (Ft. Collins, CO) from a previous study¹⁵, and stored at -80°C. Upon thawing, the bacterial suspension (1 mL) was transferred into 9 mL 7H9 media (supplemented with OADC, 0.1% Tween 80), in a sterile 150x25 mm screw-capped glass culture tube; a very small, plastic coated magnetic stir bar was added for culture aeration and agitation. This bacteria/7H9 suspension was placed on a magnetic stirrer and cultured for 15 days @ 37°C 5% CO₂ (the minimum magnetic force needed to agitate the stir bar was applied during this period). After this initial 15 day period, 5 mL of the bacteria/7H9 suspension was transferred into 50 mL of synthetic Proskauer Beck media (modified from Youmans and Karlson²⁶, PB: 0.5% KH₂PO₄, 0.5% L-asparagine monohydrate, 0.06% MgSO₄ · 7H₂O, 0.25% magnesium citrate, 2% glycerol, 0.05% Tween 80) in a sterile glass culture bottle and cultured for an additional 17 days @ 37°C 5% CO₂ (cultures were gently rocked at 30 RPM during this period). At the end of this 17-day period, BCG and BCGDBCG1419c were aliquoted and rapidly frozen in screw-capped microcentrifuge tubes at -80°C. After 5 days of being frozen at -80°C, individual aliquots of BCG and BCGDBCG1419c were thawed and serial dilutions plated on 7H10 (supplemented with OADC, 0.1% L-asparagine) and TCA (to test for contamination). Colony counts demonstrated both BCG and BCGDBCG1419c aliquots to be at 5 x 10⁴/mL; TCA plates showed no growth (i.e. neither BCG nor BCGDBCG1419c contained contaminants).

Mouse strains. C57BL/6 mice were acquired from The Jackson Laboratory (Bar Harbor, ME); IL10 reporter mice (C57BL/6 background) were generously provided by Dr. Weiguo Cui (Blood Research Institute, Blood Center of Wisconsin, Milwaukee WI)²⁷. All mice were maintained within The Ohio State University (OSU) University Laboratory Animal Resources (ULAR), and supplied with sterile and water ad libitum. All methods and procedures were performed in accordance with OSU Institution Animal Care and Use Committee (IACUC) approved protocols and procedures.

Immunization and tissue collection. On the day of immunization (day 0), aliquots of BCG and BCGDBCG1419c were thawed and directly loaded into tuberculin syringes. Mice were immunized subcutaneously (s.c.) at the back of their neck with 1 x 10⁴ CFU of either BCG or BCGDBCG1419c (i.e. 200 µL of a 5 x 10⁴ CFU/mL aliquot). On post-immunization day 14, the cervical lymph nodes and spleen were collected from euthanized IL10 reporter mice and used for flow cytometric analysis. On post-immunization day 18, blood, spleen and lungs were collected from euthanized C57BL/6 mice and used for flow cytometric analysis. As described in our Results, our analysis of IL10 reporter mice was performed before that of C57BL/6 mice (day 14 vs day 18) to determine whether IL10 expression

differences precede the phenotypic differences between BCG- and BCGDBCG1419c-immunized C57BL/6 mice.

Cell isolation. Spleen and lymph node tissues were gently pressed through a mesh screen using the plunger portion of a 1 ml syringe. Lungs were similarly processed following collagenase/DNase treatment. The resulting cell suspensions were transferred to a 15 ml tube by filtering through 100-micron filter using ice-cold R10 media (RPMI 1640 by Gibco® Thermo Fisher, 10% FBS, 2 mM L-glutamine, 100 U/ml penicillin G, 100 µg/ml streptomycin). Tubes were spun down at 250x g for 7 min at 4 °C. The cell pellet was resuspended with R10 media. After discarding supernatant, the cell pellet was washed again and incubated with 1ml of RBC lysis buffer at room temperature. The RBC lysis reaction was stopped after 10 minutes using 1ml of R10 media and spun down at 250x g for 7 min at 4 °C.

Flow cytometry cell staining. Freshly isolated cells from spleen, lung and blood tissues were transferred to pre-warmed R10 media and washed. Cells were then resuspended at 1–2 million cells per ml in R10 media and stimulated in FACs tubes with PMA/Ionomycin (final concentration of 2 µg/ml) in the presence of GolgiPlug, at a final concentration of 10 µg/ml; (BD Biosciences, San Jose, California) for 12 h in a 37 °C and 5% CO₂ incubator. At the end of the incubation, intracellular staining procedure was performed. The following monoclonal antibodies were used: FITC anti-CD3 (clone 17A2), Alexa Fluor 532 anti-CD4 (clone RM4-5), PE anti-CD127 (clone SB/199), PE-eFluor 610 anti-PD1 (clone J43), PE-Cy5 anti-CD11b (clone M1/70), PerCP-Cy5.5 anti-CD44 (clone IM7), PerCP-eFluor 710 anti-CXCR5 (clone SPRCL5), PE-Cy7 anti-CD117 (clone 2B8), Alexa Fluor 647 anti-CRTH2 (clone No3m1scz), Alexa Fluor 700 anti-TNFα (clone MP6-XT22), Brilliant Violet 421 anti-CD45 (clone 30-F11), Super Bright 436 anti-CD62L (clone MEL-14), eFluor 450 anti-Granzyme b (clone NGZB), BD Horizon BV480 anti-IFNγ (clone XMG1.2),

Brilliant Violet 570 anti-CD8 (clone 53-6.7), Brilliant Violet 605 anti-IL17 (clone TC11-18H10.1), Brilliant Violet 650 anti-Gr1 (clone RB6-8C5), Brilliant Violet 711 anti-CD19 (clone 6D5), Brilliant Violet 750 anti-CD27 (clone LG.3A10), Brilliant Violet 785 anti-NK1.1 (clone PK136). Samples were stained with Fixability Viability Dye (Aqua Live/Dead stain) and incubated for 15 minutes. Then cells were washed and followed by adding of surface antibody cocktail. After incubating for 30 minutes cells were washed and fixed with fixation buffer (Cat No:420801 Biolegend) for 20 minutes and permeabilized with 1x Intracellular Staining Permeabilization Wash Buffer (Cat No: 421002 Biolegend). Recommended amounts of directly conjugated primary antibodies for detection of intracellular antigens were added and incubated for 25 minutes at 4 °C in dark. All washing steps were carried out at 700g for 6 minutes at 4 °C. All antibodies were previously titrated to determine the optimal concentration. Finally, after washing and filtering cells through strainer capped FACs tubes, samples were acquired on a Cytex Aurora or FACS Canto flow cytometer and analyzed using FlowJo version 10.6.1 (Becton Dickinson, Ashland, OR).

Statistics. Figures were prepared using GraphPad Prism, version 7. Statistical analyses used the bundled software. Bars in the figures show means plus standard deviations (SD). Numbers shown between data points represent *p* values for the comparisons indicated on the figures. Prior to performing statistical comparisons, the Shapiro Wilk test was used to determine if data distribution was normal. Statistical

comparisons involving more than two experimental groups used analysis of variance (ANOVA). All other statistical comparisons used Student's t test.

Results

BCG and BCGDBCG1419c elicit distinct immune responses early post-immunization.

To identify the early innate and adaptive immune populations which respond to BCG, as well as identify what effect the deletion of *BCG1419c* has on these responses, we immunized C57BL/6 mice with 10^4 CFU of live BCG Pasteur and its isogenic derivative, BCGDBCG1419c (**FIG 1A**). Mice immunized with PBS alone served as negative controls. At post-immunization day 18, multiple tissues (blood, spleen and lung) were collected, processed into single cell suspensions and analyzed by multi-dimensional flow cytometry to identify phenotypic and functional differences between innate and adaptive immune subsets (**FIG 1B**).

To comprehensively assess whether deletion of *BCG1419c* alters the BCG-induced immune landscape in mice, a 27-color flow cytometry panel was designed to characterize the pulmonary and systemic immune populations. t-distributed stochastic embedding (tSNE) dimensionality reduction plots of CD45+ cells from the spleen, lung and blood allowed the identification of 19 immune cell populations (**FIG 2**). The clustering algorithm allowed for the automatic definition of 11 adaptive immune metaclusters based on expression of cell surface markers CD3 and CD19 (**FIG 2A**), and 8 innate immune metaclusters wherein CD3 and CD19 expression was absent (**FIG 2B**; the same tSNE plots shown in **FIG 2A** minus the T- and B-cell overlay). Distinctions between lung, spleen and blood cluster distributions can be observed by comparing the top, middle and bottom rows of **FIG2A-B**, as well as between PBS control, BCG and BCGDBCG1419c immunized mice (compare left, middle and right columns of **FIG2-B**). Collectively, tSNE analysis demonstrated that BCG and BCGDBCG1419c elicit distinct immune responses early post-immunization.

BCG and BCGDBCG1419c elicited innate immune responses.

Whereas tSNE analysis permits high-level data visualization, we used traditional flow cytometry analysis to more closely and quantitatively examine the innate and adaptive populations that were differentially affected by BCG and BCGDBCG1419c. Since tSNE analysis demonstrated the pulmonary and systemic responses to vaccination were distinct, differences among spleen (left panels of **FIG 3-6**), lung (center panels of **FIG 3-6**) and blood (right panels of **FIG 3-6**) were considered separately.

Among the innate lineages (ILC3, ILC1, NK cells, monocytes, and Gr1 cells), nearly all changed in response to BCG and/or BCGDBCG1419c immunization, the degree to which varied between spleen, lung and blood (**FIG 3**). In all tissues, NK cell frequencies declined following BCG or BCGD1419c immunization (**FIG 3C**), while ILC1 frequencies were unchanged (**FIG 3B**). Other changes were tissue-specific: whereas the frequencies of spleen ILC3s, monocytes, and Gr1 cells were unaffected by immunization, their frequencies were decreased in the lung of BCG or BCGDBCG1419c immunized animals (**FIG 3A, D, E**). The only innate frequencies which were positively affected by immunization were blood monocytes in

BCGDBCG1419c immunized animals (**FIG 3D**), and blood Gr1 cells in BCG immunized animals (**FIG 3E**). Collectively, these data demonstrate that BCG and BCGDBCG1419c immunization each have generally negative effects on innate cell frequency in the spleen and lung, but unique effects on monocyte and Gr1 cell frequencies in the blood.

BCG and BCGDBCG1419c elicited adaptive immune responses.

Conventional ab T cells are critical for the long term efficacy of BCG vaccination^{28,29}, and although B cells have historically been considered dispensable for BCG efficacy their presence may shape the T cell response in an antibody-independent manner^{30,31}. Lung CD4+ and CD8+ T cell frequencies increased in BCG and BCGDBCG1419c immunized animals, with corresponding declines in the spleen and blood (likely a consequence of T cell redistribution from the circulation) (**FIG 4A-B**); the ability of subcutaneously administered BCG to increase lung T cell numbers have been previously reported³². B cells' response to immunization was likewise tissue-specific: while no differences were observed in the lung or blood across groups, increased B cell frequencies being observed in the spleens of BCG and BCGDBCG1419c-immunized compared to PBS controls (**FIG 4C**). Collectively, these data demonstrate that BCG and BCGDBCG1419c immunization causes similar increases in lung T cell and spleen B cell frequencies.

BCG and BCGDBCG1419c elicited T cell memory.

An important component of BCG-mediated immunity is T cells' capacity for memory development. Unlike other pulmonary infection models, BCG-mediated immunity lay within T cells that maintain a naïve, antigen-inexperienced surface phenotype³². Shown in **FIG 5** and **FIG 6**, respectively, are the frequencies of CD4 and CD8 T cells which exhibit a naïve (CD62L^{HI}CD44^{LO}), effector (CD62L^{LO}CD44^{HI}) and central memory (CD62L^{HI}CD44^{HI}) surface phenotype. The frequency of follicular T_H cells (i.e. Tfh cells) was also examined.

Relative to PBS controls, higher proportions of lung CD4 T cells in BCG and BCGDBCG1419c immunized mice expressed either a naïve surface phenotype (**FIG 5A**) or memory surface phenotype (**FIG 5C**), with corresponding declines in spleen and blood. BCG and BCGDBCG1419c both elicited significant declines in blood effector T_H cell (**FIG 5B**) and spleen Tfh frequencies (**FIG 5E**); however and interestingly, BCG and BCGDBCG1419c had different effects on the proportion of lung effector and exhausted T_H cells: whereas only BCGDBCG1419c elicited a significant decline in lung effector T_H cell frequencies (**FIG 5B**), only BCG elicited a significant increase in the lung exhausted T_H cell frequencies (**FIG 5D**). Similar patterns were observed in the CD8 T cell (T_C) compartment: in the lung compartment, BCG and BCGDBCG1419c both elicited an increased frequency of CD8 T cells with a naïve surface phenotype (and corresponding decrease in the spleen), as well as decreases in effector CD8 frequencies in the lung and blood (**FIG 6**). Collectively, these data demonstrate that while both BCG and BCGDBCG1419c elicit increases in the proportion of T cells known to retain memory, BCG-elicited T cells were more likely to have an exhausted phenotype, and BCGDBCG1419c-elicited T cells were less likely to have an effector phenotype.

BCG and BCGDBCG1419c elicited T cell cytokine responses.

Although there is no universally-accepted cytokine correlate of BCG-elicited protection against *M. tuberculosis* infection³³, lymphocytes' capacity to secrete IFN γ , IL17 and TNF α likely contribute to BCG vaccine efficacy³⁴. To compare the early cytokine profile of BCG- and BCGDBCG1419c-elicited lymphocytes, the same cell preparations used for surface phenotype analysis (**FIG 2-6**) were stimulated with a polyclonal immunogen (PMA/Ionomycin) and subsequently used for intracellular cytokine staining (ICS) of the NK cell (**FIG 7**), CD4 T cell (**FIG 8**) and CD8 T cell (**FIG 9**) lymphocyte compartments. The results for each compartment are described below.

NK cells: NK cell frequencies decline following BCG- or BCGDBCG1419c-immunization (**FIG 3C**); nevertheless, the cytokine profile of those which remain are distinct from those of PBS controls, insomuch there is an expansion of IFN γ ⁺TNF α ⁺ NK cells in spleen, lung and blood of BCG- and BCGDBCG1419c-immunized mice (**FIG 7**). BCGDBCG1419c-immunized mice had more single IFN γ ⁺ cells and BCG-immunized mice had more double IFN γ ⁺TNF α ⁺ cells in lungs, while BCG induced a higher proportion of single IFN γ ⁺ cells and BCGDBCG1419c a more balanced presence of single IFN γ ⁺, single TNF α ⁺, and double IFN γ ⁺TNF α ⁺ cells in blood.

CD4 T cells: Depicted in **FIG8A-C** are our gating profiles to identify and enumerate T_H cells expressing IFN γ (**FIG 8A**), IL17 (**FIG 8B**) and TNF α (**FIG 8C**). Based upon these flow cytometry data, we were able to distinguish 7 cytokine-producing T_H cells: those that were single positive for an individual cytokine (i.e. IFN γ ⁺, IL17⁺, or TNF α ⁺), double positive for two cytokines (IFN γ ⁺IL17⁺, IFN γ ⁺TNF α ⁺, IL17⁺TNF α ⁺), or triple positive for all three cytokines (IFN γ ⁺IL17⁺TNF α ⁺). The relative proportion of each subset among cytokine producing CD4⁺ T_H cells is depicted in **FIG 8D** for the spleen, lung and blood of each experimental group. At baseline (i.e. in PBS control mice), cytokine-producing T_H cells in all tissues were primarily TNF α ⁺, with tissue-specific differences in which subsets were of secondary abundance (spleen: IFN γ ⁺TNF α ⁺; lung and blood: IFN γ ⁺). In the spleen, BCG immunization expanded the representation of IFN γ ⁺IL17⁺TNF α ⁺ cells; interestingly, the expansion of IFN γ ⁺IL17⁺TNF α ⁺ was not as large in BCGDBCG1419c immunized animals and was accompanied by a contraction in IFN γ ⁺TNF α ⁺ (**FIG 8D top row**). A similar contraction of IFN γ ⁺TNF α ⁺ T_H cells was observed in the blood relative to BCG-immunized animals (**FIG 8D bottom row**). In the lung, BCG and BCGDBCG1419c similarly resulted in an expansion of IFN γ ⁺IL17⁺TNF α ⁺ and IFN γ ⁺TNF α ⁺, and contraction of IFN γ ⁺. IL17⁺TNF α ⁺ cells also appeared in BCG and BCGDBCG1419c lung relative to PBS. Collectively, our analysis of T_H cell cytokine profiles demonstrates the induction of systemic IFN γ ⁺IL17⁺TNF α ⁺ and IFN γ ⁺TNF α ⁺ populations was less robust in BCGDBCG1419c-immunized animals, relative to BCG-immunized animals, yet in lungs, BCGDBCG1419c-immunized animals increased the presence of IFN γ ⁺IL17⁺ and IFN γ ⁺TNF α ⁺ populations than BCG.

CD8 T cells: Depicted in **FIG9A-B** are our gating profiles to identify and enumerate T_C cells expressing IFN γ (**FIG 9A**) and TNF α (**FIG 9B**); unlike CD4 T cells, no IL17 expressing CD8 T cells were detected. In

PBS controls, cytokine producing CD8 T cells in the spleen were primarily TNF α +, and secondarily IFN γ +, in response to BCG and BCGDBCG1419c immunization, the proportion of CD8 T cells that were TNF α + expanded further alongside IFN γ +TNF α +, at the expense of IFN γ + CD8 T cells. Similar expansions of TNF α + and IFN γ +TNF α + cells (and accompanying contractions of IFN γ + CD8 T cells) were observed in the lung and blood of BCG and BCGDBCG1419c immunized animals. Similar to what we observed for NK cells, BCGDBCG1419c-immunized mice had more single IFN γ + cells and BCG-immunized mice had more double IFN γ +TNF α + cells in lungs.

The altered early immune response of BCGDBCG1419c-immunized animals is associated with elevated IL10 production.

IL10 suppresses T_H1/T_H17 differentiation following BCG immunization²¹. At post-immunization day 18, BCGDBCG1419c-immunization elicits fewer systemic IFN γ - and IL17-producing T_H cells (**FIG 8D**), as well as a greater shift away from the more pro-inflammatory “effector” T_H phenotype (**FIG 5B**); therefore, we predicted that BCGDBCG1419c-immunized mice would—relative to BCG-immunized mice—have higher frequencies of systemic IL10-producing cells prior to day 18. To test this prediction, we immunized transgenic VertX mice that express surface Thy1.1 concomitant to IL10 expression²⁷, permitting the identification of IL10 producing cells by flow cytometry staining with anti-Thy1.1 (**FIG 10A**). VertX mice were immunized with BCG or BCGDBCG1419c in a manner identical to that used for C57BL/6 mice; the spleen was removed at 14 days post-immunization. The results of this study are shown in **FIG 10** and demonstrate that BCGDBCG1419c elicits higher frequencies of IL10+ cells in the spleen (**FIG 10B**), and that a larger proportion of these IL10+ cells were CD4+ T cells and CD8+ T cells (**FIG 10C**). Overall, these data demonstrate that BCGDBCG1419c elicits higher frequencies of IL10 producing T cells, which may account for less inflammatory response of BCGDBCG1419c-immunized animals relative to wild type BCG-immunized animals.

Discussion

BCG was developed as a TB vaccine in the early 1900s by Calmette and Guérin, following serial passage of *M. bovis* for eleven years (~230 passages) on bile-imbibed potato slices³⁵. Relative to *M. bovis*, the resulting strain (i.e. BCG) was avirulent in multiple animal species and unable to cause tuberculosis. We now know this attenuation stems in part from the loss of three genomic regions of difference (RD1, RD2 and RD3) which encode a number of virulence factors (e.g. ESX)³⁶. Despite several starts and stops due to safety concerns, human BCG trials moved forward in a number of countries (UK, US, India) with differing conclusions regarding its protective efficacy, which ranges from 0 to 80% depending on the study³⁷. Still a matter of debate, the reasons for BCG’s variable efficacy in different countries and populations include different preparation methods, genetic drift over time, and environmental mycobacteria which themselves confer some protection against TB and thus mask any effect of BCG. Regardless of the reason(s), there is near-universal agreement that despite BCG’s success a TB vaccine with better efficacy is needed to meet WHO benchmarks for reducing TB prevalence⁵. Partly due to lack

of full efficacy of current BCG, an estimated 2 billion people are thought to harbor *M. tuberculosis* latent infection (LTBI). It has been estimated that by 2050, up to 75% percent of new TB cases will be due to reactivation from LTBI, rather than new infections, in China, one of the countries with the highest number of TB cases (including MDR-TB)³⁸. In fact, a model that includes data from China, South Africa, and India, suggested that vaccines preventing disease in *M. tuberculosis*-infected populations would have the greatest impact by 2050 (10-year, 70% efficacy against disease, incidence rate reduction 51%, 52%, and 54% in China, South Africa, and India, respectively)³⁹.

Like many mycobacterial species, BCG is capable of forming biofilms which impact upon *in vitro* behavior and *in vivo* persistence. Biofilms comprise a community of bacteria surrounded by an extracellular matrix produced by the constituent species, are generally more resistant to environmental stress, and enable persistence in environmental reservoirs and biomaterials³⁸. *In vitro*, mycobacteria biofilm formation manifests as pellicles at the media-air interface^{39,40}, which in BCG follow a defined genetic program⁴¹. Pellicles have an altered transcriptional profile relative to planktonic bacteria and are enriched in antibiotic drug resistant cells or persisters⁴²⁻⁴⁴. The *in vivo* contribution of mycobacteria biofilm formation to lung disease is often a topic of debate since their presence is not obvious upon histo-pathological examination of infected lungs (e.g. by acid fast staining), and mycobacterial pathogenesis involves intracellular survival within alveolar phagocytes (at least during early phases of disease) rather than extracellular survival on airway surfaces, which is more typical of *Pseudomonas aeruginosa* and other canonical biofilm-forming pathogens. That said, mycobacteria which are deficient in biofilm formation—due to either genetic manipulation or natural variation—generally exhibit attenuated virulence and elicit different inflammatory responses during *in vivo* infection models^{39,45}. Our understanding of the relationship between biofilm formation and mycobacterial pathogenesis is therefore far from complete.

The BCGDBCG1419c strain was generated in part to test the relationship between BCG's ability to form biofilms and its vaccine efficacy²⁰. The *BCG1419c* gene encodes a cyclic di-GMP phosphodiesterase (PDE) that normally functions to hydrolyze Bis-(3'-5')-cyclic dimeric GMP (c-di-GMP), a second messenger which promotes the biofilm phase of bacterial growth⁴⁶. In the absence of *BCG1419c*, the deletion mutant BCGDBCG1419c exhibits an altered colony morphology, higher biofilm production and extended survival in the lungs and spleen of immunocompetent mice, relative to parental BCG or complemented mutants²⁰ when used for intravenous infection of BALB/c mice. Relative to BCG immunized controls and at 6 months post-Mtb challenge, those immunized with BCGDBCG1419c also exhibit significantly reduced lung histopathological scores, as well as reduced lung IL6 and TNF α protein levels¹⁵. Since IL-6 and TNF- α may contribute to lung pathology in chronic disease settings, these data implicate BCG1419c-dependent hydrolysis of c-di-GMP in conditioning the long-term TB response of vaccinated animals.

Given that BCGDBCG1419c immunized mice have less lung pathology following TB challenge, suggesting an improvement to prevent disease, we undertook the above study to determine if there are early immunological events which coincide with its long term effects. Our results demonstrate that while

there are many similarities in the early innate and adaptive response to BCG and BCGDBC1419c, there are also notable differences in the innate and adaptive compartments. Among innate lineages, whereas both BCG and BCGDBC1419c elicited strong reductions in NK cell frequency across all tissues examined (**Fig 3C**), only BCGDBC1419c elicited significant increases in circulating monocytes (**Fig 3D**) and reductions in lung Gr1 cells (**Fig 3E**). Among adaptive lineages, both BCG and BCGDBC1419c elicited significant increases in lung CD4⁺ T cell frequencies (**Fig 4A**), those induced by BCG were more likely to exhibit an exhausted phenotype relative to those induced by BCGDBC1419c (**Fig 5D**). BCGDBC1419c elicited CD4 T cells were also less likely to exhibit a pro-inflammatory effector phenotype (**Fig 5B**). Importantly, the CD4 and CD8 T cell response elicited by BCGDBC1419c were also more likely to contain IL10-producing cells (**Fig 10C**), which may underlie the less inflammatory nature of BCGDBC1419c during chronic TB stages. We have shown that BCGDBC1419c produced less DnaK, HbhA, PstS2, 35KDa antigen, and AcpM (a protein involved in synthesis of mycolic acids) in biofilm cultures compared with parental BCG¹⁴, possibly contributing to its less inflammatory nature. Still to be determined is whether the immune response elicited by BCGDBC1419c differs from that elicited by BCG due to either prolonged antigen presentation or altered lipid profiles, which contribute to mycobacteria immunogenicity⁴⁷.

As with any bacteria-driven inflammatory response, the BCG-immunized host must balance the generation of robust immunity with the potential for detrimental, non-specific effects of excess cytokine production. Here, we used the mouse model to demonstrate that the early immunological events associated with BCGDBC1419c immunization are less inflammatory relative to BCG. This is important because BCGDBC1419c's efficacy in controlling *M. tuberculosis* replication in lungs is comparable or superior to that of BCG, and more protective against TB-associated immunopathology in mice. Given the reactogenicity of BCG in immunized individuals, which commonly manifests as fever, headache and swollen lymph nodes, BCGDBC1419c's unique immunological profile may prove important in TB endemic regions where parents of young children are increasingly hesitant of inflammatory vaccine administration, regardless of the vaccine efficacy⁹⁻¹².

Declarations

ACKNOWLEDGEMENTS

This work was supported by The Ohio State University (OSU), the National Institutes of Health (R01 AI134972 to K.J.O.; K22 AI127072 to N.P.M.L.; R01 AI121212 to R.T.R.), and CONACYT Grant 86396 (to MAFV and MJAS) allowed construction of the BCGDBC1419c strain.

References

1. Guinn KM, Rubin EJ. Tuberculosis: Just the FAQs. *mBio* 2017;**8** doi: 10.1128/mBio.01910-17
2. Russell DG. Mycobacterium tuberculosis and the intimate discourse of a chronic infection. *Immunol Rev* 2011;**240**:252-268. doi: 10.1111/j.1600-065X.2010.00984.x

3. Lin PL, Flynn JL. The End of the Binary Era: Revisiting the Spectrum of Tuberculosis. *J Immunol* 2018;**201**:2541-2548. doi: 10.4049/jimmunol.1800993
4. Raviglione M, Marais B, Floyd K, Lonnroth K, Getahun H, Migliori GB, Harries AD, Nunn P, Lienhardt C, Graham S, Chakaya J, Weyer K, Cole S, Kaufmann SH, Zumla A. Scaling up interventions to achieve global tuberculosis control: progress and new developments. *Lancet* 2012;**379**:1902-1913. doi: 10.1016/S0140-6736(12)60727-2
5. Marais B, Zumla A. Advancing global tuberculosis control after the UNGA-HLM. *Lancet* 2018;**392**:1096-1097. doi: 10.1016/S0140-6736(18)32361-4
6. Trunz BB, Fine P, Dye C. Effect of BCG vaccination on childhood tuberculous meningitis and miliary tuberculosis worldwide: a meta-analysis and assessment of cost-effectiveness. *Lancet* 2006;**367**:1173-1180. doi: 10.1016/S0140-6736(06)68507-3
7. Izzo AA. Tuberculosis vaccines - perspectives from the NIH/NIAID Mycobacteria vaccine testing program. *Curr Opin Immunol* 2017;**47**:78-84. doi: 10.1016/j.coi.2017.07.008
8. Lahey T, von Reyn CF. Mycobacterium bovis BCG and New Vaccines for the Prevention of Tuberculosis. *Microbiol Spectr* 2016;**4** doi: 10.1128/microbiolspec.TNMI7-0003-2016
9. Silveira MF, Buffarini R, Bertoldi AD, Santos IS, Barros AJD, Matijasevich A, Menezes AMB, Goncalves H, Horta BL, Barros FC, Barata RB, Victora CG. The emergence of vaccine hesitancy among upper-class Brazilians: Results from four birth cohorts, 1982-2015. *Vaccine* 2020;**38**:482-488. doi: 10.1016/j.vaccine.2019.10.070
10. Sankaranarayanan S, Jayaraman A, Gopichandran V. Assessment of Vaccine Hesitancy among Parents of Children between 1 and 5 Years of Age at a Tertiary Care Hospital in Chennai. *Indian J Community Med* 2019;**44**:394-396. doi: 10.4103/ijcm.IJCM_351_18
11. Dasgupta P, Bhattacharjee S, Mukherjee A, Dasgupta S. Vaccine hesitancy for childhood vaccinations in slum areas of Siliguri, India. *Indian J Public Health* 2018;**62**:253-258. doi: 10.4103/ijph.IJPH_397_17
12. Brown AL, Sperandio M, Turssi CP, Leite RMA, Berton VF, Succi RM, Larson H, Napimoga MH. Vaccine confidence and hesitancy in Brazil. *Cad Saude Publica* 2018;**34**:e00011618. doi: 10.1590/0102-311X00011618
13. Pedroza-Roldan C, Guapillo C, Barrios-Payan J, Mata-Espinosa D, Aceves-Sanchez Mde J, Marquina-Castillo B, Hernandez-Pando R, Flores-Valdez MA. The BCGDeltaBCG1419c strain, which produces more pellicle in vitro, improves control of chronic tuberculosis in vivo. *Vaccine* 2016;**34**:4763-4770. doi: 10.1016/j.vaccine.2016.08.035
14. Segura-Cerda CA, Aceves-Sanchez MJ, Marquina-Castillo B, Mata-Espinoza D, Barrios-Payan J, Vega-Dominguez PJ, Pedroza-Roldan C, Bravo-Madriral J, Vallejo-Cardona AA, Hernandez-Pando R, Flores-Valdez MA. Immune response elicited by two rBCG strains devoid of genes involved in c-di-GMP metabolism affect protection versus challenge with M. tuberculosis strains of different virulence. *Vaccine* 2018;**36**:2069-2078. doi: 10.1016/j.vaccine.2018.03.014

15. Flores-Valdez MA, Pedroza-Roldan C, Aceves-Sanchez MJ, Peterson EJR, Baliga NS, Hernandez-Pando R, Troudt J, Creissen E, Izzo L, Bielefeldt-Ohmann H, Bickett T, Izzo AA. The BCGDeltaBCG1419c Vaccine Candidate Reduces Lung Pathology, IL-6, TNF-alpha, and IL-10 During Chronic TB Infection. *Front Microbiol* 2018;**9**:1281. doi: 10.3389/fmicb.2018.01281
16. Sathkumara HD, Pai S, Aceves-Sanchez MJ, Ketheesan N, Flores-Valdez MA, Kupz A. BCG Vaccination Prevents Reactivation of Latent Lymphatic Murine Tuberculosis Independently of CD4(+) T Cells. *Front Immunol* 2019;**10**:532. doi: 10.3389/fimmu.2019.00532
17. Segura-Cerda CA, Marquina-Castillo B, Lozano-Ordaz V, Mata-Espinosa D, Barrios-Payan JA, Lopez-Torres MO, Aceves-Sanchez MJ, Bielefeldt-Ohmann H, Hernandez-Pando R, Flores-Valdez MA. BCG and BCGDeltaBCG1419c protect type 2 diabetic mice against tuberculosis via different participation of T and B lymphocytes, dendritic cells and pro-inflammatory cytokines. *NPJ Vaccines* 2020;**5**:21. doi: 10.1038/s41541-020-0169-6
18. Aceves-Sanchez MJ, Flores-Valdez MA, Shanley C, Orme I, Bielefeldt-Ohmann H. Vaccination of guinea pigs with BCGDeltaBCG1419c transiently reduces hematogenous spread of M. tuberculosis to the spleen. *Pathog Dis* 2018;**76** doi: 10.1093/femspd/fty088
19. Tamayo R, Pratt JT, Camilli A. Roles of cyclic diguanylate in the regulation of bacterial pathogenesis. *Annu Rev Microbiol* 2007;**61**:131-148. doi: 10.1146/annurev.micro.61.080706.093426
20. Flores-Valdez MA, Aceves-Sanchez Mde J, Pedroza-Roldan C, Vega-Dominguez PJ, Prado-Montes de Oca E, Bravo-Madrigal J, Laval F, Daffe M, Koestler B, Waters CM. The Cyclic Di-GMP Phosphodiesterase Gene Rv1357c/BCG1419c Affects BCG Pellicle Production and In Vivo Maintenance. *IUBMB Life* 2015;**67**:129-138. doi: 10.1002/iub.1353
21. Pitt JM, Stavropoulos E, Redford PS, Beebe AM, Bancroft GJ, Young DB, O'Garra A. Blockade of IL-10 signaling during bacillus Calmette-Guerin vaccination enhances and sustains Th1, Th17, and innate lymphoid IFN-gamma and IL-17 responses and increases protection to Mycobacterium tuberculosis infection. *J Immunol* 2012;**189**:4079-4087. doi: 10.4049/jimmunol.1201061
22. Noguchi N, Nakamura R, Hatano S, Yamada H, Sun X, Ohara N, Yoshikai Y. Interleukin-21 Induces Short-Lived Effector CD8(+) T Cells but Does Not Inhibit Their Exhaustion after Mycobacterium bovis BCG Infection in Mice. *Infect Immun* 2018;**86** doi: 10.1128/IAI.00147-18
23. Paidipally P, Tripathi D, Van A, Radhakrishnan RK, Dhiman R, Venkatasubramanian S, Devalraju KP, Tvinnereim AR, Valluri VL, Vankayalapati R. Interleukin-21 Regulates Natural Killer Cell Responses During Mycobacterium tuberculosis Infection. *J Infect Dis* 2018;**217**:1323-1333. doi: 10.1093/infdis/jiy034
24. Cheekatla SS, Tripathi D, Venkatasubramanian S, Paidipally P, Welch E, Tvinnereim AR, Nurieva R, Vankayalapati R. IL-21 Receptor Signaling Is Essential for Optimal CD4(+) T Cell Function and Control of Mycobacterium tuberculosis Infection in Mice. *J Immunol* 2017;**199**:2815-2822. doi: 10.4049/jimmunol.1601231
25. Bickett TE, McLean J, Creissen E, Izzo L, Hagan C, Izzo AJ, Silva Angulo F, Izzo AA. Characterizing the BCG Induced Macrophage and Neutrophil Mechanisms for Defense Against Mycobacterium

- tuberculosis. *Front Immunol* 2020;**11**:1202. doi: 10.3389/fimmu.2020.01202
26. Youmans GP, Karlson AG. Streptomycin sensitivity of tubercle bacilli; studies on recently isolated tubercle bacilli and the development of resistance to streptomycin in vivo. *Am Rev Tuberc* 1947;**55**:529-535.
 27. Xin G, Zander R, Schauder DM, Chen Y, Weinstein JS, Drobyski WR, Tarakanova V, Craft J, Cui W. Single-cell RNA sequencing unveils an IL-10-producing helper subset that sustains humoral immunity during persistent infection. *Nat Commun* 2018;**9**:5037. doi: 10.1038/s41467-018-07492-4
 28. Ladel CH, Daugelat S, Kaufmann SH. Immune response to Mycobacterium bovis bacille Calmette Guerin infection in major histocompatibility complex class I- and II-deficient knock-out mice: contribution of CD4 and CD8 T cells to acquired resistance. *Eur J Immunol* 1995;**25**:377-384. doi: 10.1002/eji.1830250211
 29. Ladel CH, Hess J, Daugelat S, Mombaerts P, Tonegawa S, Kaufmann SH. Contribution of alpha/beta and gamma/delta T lymphocytes to immunity against Mycobacterium bovis bacillus Calmette Guerin: studies with T cell receptor-deficient mutant mice. *Eur J Immunol* 1995;**25**:838-846. doi: 10.1002/eji.1830250331
 30. Maglione PJ, Chan J. How B cells shape the immune response against Mycobacterium tuberculosis. *Eur J Immunol* 2009;**39**:676-686. doi: 10.1002/eji.200839148
 31. Moliva JI, Turner J, Torrelles JB. Immune Responses to Bacillus Calmette-Guerin Vaccination: Why Do They Fail to Protect against Mycobacterium tuberculosis? *Front Immunol* 2017;**8**:407. doi: 10.3389/fimmu.2017.00407
 32. Kipnis A, Irwin S, Izzo AA, Basaraba RJ, Orme IM. Memory T lymphocytes generated by Mycobacterium bovis BCG vaccination reside within a CD4 CD44^{lo} CD62 ligand(hi) population. *Infect Immun* 2005;**73**:7759-7764. doi: 10.1128/IAI.73.11.7759-7764.2005
 33. Flores-Valdez MA. New potential biomarkers to assess efficacy of protection afforded by vaccine candidates against tuberculosis. *Clin Microbiol Infect* 2019;**25**:656-658. doi: 10.1016/j.cmi.2019.02.010
 34. Bhatt K, Verma S, Ellner JJ, Salgame P. Quest for correlates of protection against tuberculosis. *Clin Vaccine Immunol* 2015;**22**:258-266. doi: 10.1128/CI.00721-14
 35. Luca S, Mihaescu T. History of BCG Vaccine. *Maedica (Buchar)* 2013;**8**:53-58.
 36. Mahairas GG, Sabo PJ, Hickey MJ, Singh DC, Stover CK. Molecular analysis of genetic differences between Mycobacterium bovis BCG and virulent M. bovis. *J Bacteriol* 1996;**178**:1274-1282. doi: 10.1128/jb.178.5.1274-1282.1996
 37. Mangtani P, Abubakar I, Ariti C, Beynon R, Pimpin L, Fine PE, Rodrigues LC, Smith PG, Lipman M, Whiting PF, Sterne JA. Protection by BCG vaccine against tuberculosis: a systematic review of randomized controlled trials. *Clin Infect Dis* 2014;**58**:470-480. doi: 10.1093/cid/cit790
 38. Esteban J, Garcia-Coca M. Mycobacterium Biofilms. *Front Microbiol* 2017;**8**:2651. doi: 10.3389/fmicb.2017.02651

39. Chakraborty P, Kumar A. The extracellular matrix of mycobacterial biofilms: could we shorten the treatment of mycobacterial infections? *Microb Cell* 2019;**6**:105-122. doi: 10.15698/mic2019.02.667
40. Ojha A, Anand M, Bhatt A, Kremer L, Jacobs WR, Jr., Hatfull GF. GroEL1: a dedicated chaperone involved in mycolic acid biosynthesis during biofilm formation in mycobacteria. *Cell* 2005;**123**:861-873. doi: 10.1016/j.cell.2005.09.012
41. Flores-Valdez MA, Aceves-Sanchez MJ, Peterson EJR, Baliga N, Bravo-Madrigal J, De la Cruz-Villegas MA, Ares MA, Born S, Voskuil M, Perez-Padilla NA, Burciaga-Flores M, Camacho-Villegas TA, Espinoza-Jorge MG. Transcriptional portrait of *M. bovis* BCG during biofilm production shows genes differentially expressed during intercellular aggregation and substrate attachment. *Sci Rep* 2020;**10**:12578. doi: 10.1038/s41598-020-69152-2
42. Trivedi A, Mavi PS, Bhatt D, Kumar A. Thiol reductive stress induces cellulose-anchored biofilm formation in *Mycobacterium tuberculosis*. *Nat Commun* 2016;**7**:11392. doi: 10.1038/ncomms11392
43. Ackart DF, Hascall-Dove L, Caceres SM, Kirk NM, Podell BK, Melander C, Orme IM, Leid JG, Nick JA, Basaraba RJ. Expression of antimicrobial drug tolerance by attached communities of *Mycobacterium tuberculosis*. *Pathog Dis* 2014;**70**:359-369. doi: 10.1111/2049-632X.12144
44. Ojha AK, Baughn AD, Sambandan D, Hsu T, Trivelli X, Guerardel Y, Alahari A, Kremer L, Jacobs WR, Jr., Hatfull GF. Growth of *Mycobacterium tuberculosis* biofilms containing free mycolic acids and harbouring drug-tolerant bacteria. *Mol Microbiol* 2008;**69**:164-174. doi: 10.1111/j.1365-2958.2008.06274.x
45. Flores-Valdez MA. Vaccines Directed Against Microorganisms or Their Products Present During Biofilm Lifestyle: Can We Make a Translation as a Broad Biological Model to Tuberculosis? *Front Microbiol* 2016;**7**:14. doi: 10.3389/fmicb.2016.00014
46. Hengge R. Principles of c-di-GMP signalling in bacteria. *Nat Rev Microbiol* 2009;**7**:263-273. doi: 10.1038/nrmicro2109
47. Robinson RT, Huppler AR. The Goldilocks model of immune symbiosis with *Mycobacteria* and *Candida* colonizers. *Cytokine* 2017;**97**:49-65. doi: 10.1016/j.cyto.2017.05.015

Figures

Figure 1

Overview of experiment. (A) Vaccination schedule and (B) gating strategy to identify innate and adaptive subsets.

Figure 3

Innate immune subset frequencies in BCG- and BCGBCG1419c-immunized mice. Innate immune subset (i.e. ILC, NK cell, monocyte and Gr1 cell) frequencies in BCG- and BCGBCG1419c-immunized mice, as well as PBS controls, were measured in the spleen, lung and blood on post-immunization day 18. Shown are the frequencies of (A) ILC3s, (B) ILC1s, (C) NK cells, (D) monocytes and (E) Gr1 cells in the spleen (left column), lung (middle column) and blood (right column), as gated off of live single CD45⁺ cells. ILCs were gated off CD3-CD19-CD127⁺, and different subsets were determined according to the expression of CRTH2 vs cKit: ILC1, CRTH2-cKit⁻; ILC2, CRTH2+cKit^{+/+}; ILC3, CRTH-cKit⁺. Boxes represent data from 3 mice per group; asterisks represent a statistically significant difference between indicated rows (*, p0.05; **, p0.005; ***, p0.0005).

Figure 4

Adaptive immune subset frequencies in BCG- and BCGBCG1419c-immunized mice. Adaptive immune subset (i.e. T and B cell) frequencies in BCG- and BCGBCG1419c-immunized mice, as well as PBS controls, were measured in the spleen, lung and blood on post-immunization day 18. Shown are the frequencies of (A) CD4+ T cells, (B) CD8+ T cells and (C) B cells in the spleen (left column), lung (middle column) and blood (right column), as gated off of live single CD45⁺ cells. Boxes represent data from 3 mice per group; asterisks represent a statistically significant difference between indicated rows (*, p0.05; **, p0.005; ***, p0.0005).

Figure 5

CD4⁺ T cell subset frequencies in BCG- and BCGBCG1419c-immunized mice. CD4⁺ T cell (T_H) subset frequencies in BCG- and BCGBCG1419c-immunized mice, as well as PBS controls, were measured in the spleen, lung and blood, based on their expression of CD62L, CD44, PD1 and CXCR5 (naïve, CD62L^{hi}CD44^{lo}; effector, CD62L^{lo}CD44^{hi}; memory, CD62L^{hi}CD44^{hi}; exhausted, PD1⁺; follicular, CXCR5⁺PD1⁺). Shown are the frequencies of (A) naïve T_H cells, (B) effector T_H cells, (C) memory T_H cells, (D) exhausted T_H cells, and (E) follicular T_H cells in the spleen (left column), lung (middle column) and blood (right column), as gated off of live single CD45⁺ cells. Boxes represent data from 3 mice per group; asterisks represent a statistically significant difference between indicated rows (*, p0.05; **, p0.005; ***, p0.0005).

Figure 6

CD8⁺ T cell subset frequencies in BCG- and BCGBCG1419c-immunized mice. CD8 T cell (T_C) subset frequencies in BCG- and BCGBCG1419c-immunized mice, as well as PBS controls, were measured in the spleen, lung and blood, based on their expression of CD62L and CD44 (naïve, CD62L^{HI}CD44^{LO}; effector, CD62L^{LO}CD44^{HI}; memory, CD62L^{HI}CD44^{HI}; exhausted, PD1⁺). Shown are the frequencies of (A) naïve T_C cells, (B) effector T_C cells, (C) memory T_C cells and (D) exhausted T_C cells in the spleen (left column), lung (middle column) and blood (right column), as gated off of live single CD45⁺ cells. Boxes represent data from 3 mice per group; asterisks represent a statistically significant difference between indicated rows (*, p0.05; **, p0.005; ***, p0.0005).

Figure 8

CD4 T cell cytokine production in BCG- and BCGBCG1419c-immunized mice. C57BL/6 mice were s.c. immunized with 104 CFU of either BCG or BCGBCG1419c, or PBS as a control; 18 days later, mononuclear cells from the spleen, lung and blood were prepared and used for intracellular cytokine staining, following stimulation with PMA/Ionomycin. Shown are (A) representative IFN staining, (B) representative TNF staining and (C) representative IL17 staining of unstimulated and stimulated mononuclear cells, gated off CD4 T cells; (D) Spice analysis of single positive (IFN+, TNF+, IL17+), double positive (IFN+IL17+, IFN+TNF+, IL17+TNF+), and triple positive (IFN+, TNF+, IL17+) CD4 T cells in the spleen, lung and blood.

Figure 10

BCGBCG1419c-immunization elicits a higher frequency of IL10-producing T cells. IL10 reporter mice were s.c. immunized with 104 CFU of BCG or BCGBCG1419c; 14 days later, splenocytes from each group were stained for Thy1.1 (the transgene reporter of IL10 production) and T cell markers CD4 and CD8. Shown are (A) representative side scatter (SSC) and Thy1.1 staining from BCG-immunized mice, (B) the frequency of IL10+ splenocytes in BCG- and BCGBCG1419c-immunized mice (mean+SD; 3 mice per group), and (C) the relative proportion of IL10+ cells that are CD4 T cells, CD8 T cells or non-T cells. Asterisks represent a statistically significant difference between indicated comparison (*, p0.05).