

Cone-beam Computed Tomography Investigation of Middle Mesial Canals and Isthmuses in Mandibular First Molars in a Chinese Population

Shuaimei Xu

Stomatological Hospital, Southern Medical University

Junfeng Dao

Stomatological Hospital, Southern Medical University

Zhongjun Liu

Stomatological Hospital, Southern Medical University

Zhiyong Zhang

Stomatological Hospital, Southern Medical University

Yu Lu

Stomatological Hospital, Southern Medical University

Xiongqun Zeng (✉ xiongqun_zeng@163.com)

Stomatological Hospital, Southern Medical University <https://orcid.org/0000-0001-6226-7809>

Research article

Keywords: Middle mesial canal; isthmus; mandibular molar; cone-beam computed tomography

Posted Date: May 28th, 2019

DOI: <https://doi.org/10.21203/rs.2.9841/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Oral Health on May 8th, 2020. See the published version at <https://doi.org/10.1186/s12903-020-01126-2>.

Abstract

Background While there is ample research into the anatomy of mandibular molars, little is known regarding isthmuses and middle mesial (MM) canals in Chinese populations. The goal of this study was to determine the prevalence of MM canals and isthmuses in the mesial root of mandibular first molars using Cone-beam Computed Tomography. **Methods** Cone-beam Computed Tomography images of 357 mature mandibular first molars were retrospectively analyzed. Data regarding demographics, side of mandible, presence of separated distal-lingual roots, presence of isthmuses and MM canals, and the length of isthmuses in the mesial root were recorded. **Results** Of these 357 teeth, 209 showed evidence of either complete or partial communication in the mesial root. Of these, 11(3.08%) exhibited true MM canals while 198(55.46%) exhibited isthmuses. Sex or side of mandible was not correlated with the incidence of communication (isthmuses or MM canals) ($P > 0.05$). However, there was a significant association between the presence of a distal-lingual root and the incidence of such communication ($P < 0.05$). The average length of isthmuses was 4.26 ± 3.14 mm. **Conclusions** We detected high rates of mandibular molars with isthmuses, which is important as such areas should be identified and cleaned during root canal treatment.

Background

Disinfecting the root canal system is essential during endodontic treatment to ensure optimal patient outcomes, with particular efforts made to remove any infected pulp or debris from areas of operation. Owing to the presence of canals and communications between canals, however, fully sterilizing this environment is often impossible [1]. As the positioning and number of such canals vary on a patient-by-patient basis, this further increases the complexity for clinicians aiming to achieve a safe and sterile operation.

An isthmus is identified as a narrow, ribbon-shaped communication between two root canals that contains pulp tissue. Isthmuses in the root canal system, specifically of the maxillary and mandibular molars, may contain necrotic debris, tissue remnants, or organic substrates that facilitate the growth of microorganisms. Rates of treatment success have been found to be lower for teeth with a prepared isthmus than for teeth without any isthmus, and as such, there is a key need to explore isthmus anatomy for each individual tooth to better understand and control surgical outcomes and procedures in patients [2].

Many researchers to date have investigated the complexities of mesial root anatomy in mandibular molar teeth [3-5]. However, the incidence of isthmuses or the middle mesial (MM) canals in the mesial root of these teeth still remain controversial.

Indeed, many approaches have been employed for assessing the presence of MM canals, with estimated frequencies from 0% to 46.2% [6-8]. Isthmus frequencies have been found to range from 20-70% [2, 9-11]. These very wide ranges may result from individual study variations in investigative methods, patient age,

or patient ethnicity. However, little is specifically known regarding isthmuses and middle mesial (MM) canals in Chinese populations. As such this study had 3 objectives:

1. Using cone-beam computed tomographic (CBCT) images to determine the prevalence of MM canals and isthmuses in the mesial root of mandibular first molar.
2. To correlate these prevalence rates with key demographic variables, and with mandible side as well as with separated distal-lingual root presence.
3. To assess the isthmus length and the type of MM canals in the mesial root of mandibular first molar.

Methods

2.1 Sample

The Institutional Review Board and Ethical Committee of Stomatological Hospital, Southern Medical University approved this study protocol (Date of approval: March 12, 2018. Approval number: 201808). A total of 357 CBCT images of mandibular first molars from 334 native Chinese patients (Ages 9-81) were randomly selected from the database of the oral and maxillofacial radiology department, which were taken between January 1, 2016, and June 1, 2017. All chosen teeth exhibited complete root formation and closed apices without any crown restorations and root canal therapy.

Patients were divided into 4 groups based on age: group A (<20 years), group B was (20-39 years), group C (40-59 years), and group D (≥60 years).

2.2 Radiographic image acquisition and analysis

A New Tom VGI (QR Srl, Verona, Italy) CBCT unit with an isotropic voxel size of 30µm was used for this study, with 110kVp and 2.79 mA as exposure parameters and a 3.6s exposure time. In a dimly lit room, the Carestream Dental Imaging Software 3D module v2.4 (Carestream Health, Inc, Rochester, NY) was used to inspect CBCT images on a Dell Professional P2213 workstation (Dell, Round Rock, TX) using a 22-inch monitor at a resolution of 1680 x1050. Two independent endodontic specialists who had been calibrated based on study criteria before each evaluation session reviewed all images, and came to a consensus concerning radiographic findings.

The report of Mehrnaz Tahmasbi [6] was used as a guide for establishing MM canal configurations. An isthmus was identified based on the presence of a thin ribbon-shaped communication between the ML and MB canals when inspected in the axial view. A true MM canal was determined to be present if there was a clear, round cross-section visual in the radiographic image located between the MB and ML canals, which was used to identify MM canals without regard for the presence or absence of an isthmus.

The descriptions of Pomeranz [10] were used to identify MM type. Type I: Independent, three independent root canals were detectable extending to the root apex from the chamber; Type II: Confluent, an MM canal joined to either the mesiolingual or mesiobuccal root canal on its trajectory towards the apex; Type III: Fin,

the MMC orifice was connected to the orifices of the mesiobuccal or mesiolingual canals, but ended in a separate foramen.

2.3 Statistical analysis

All collected data were entered into Microsoft Excel (Microsoft, Redmond, WA), then subsequently analyzed using SPSS v25.0 (SPSS Inc, Chicago, IL). Chi-squared tests were used to compare differences in MM canal and isthmus presence, with $P < 0.05$ as the threshold of significance.

Results

We analyzed a total of 357 mandibular first molars from 334 patients (mean age = 41.24 years). Of these teeth, 198 (55.46%) had isthmuses (Fig 1), while only 11 (3.08%) had true MM canals (Fig 2, Table 1). This indicated a total incidence rate of isthmus and MM of 58.54%.

No significant sex-dependent differences were detected in terms of MM canal/isthmus prevalence, nor did incidence rates differ significantly based on the side of the mandible examined ($P > 0.05$). There was a significantly different prevalence of isthmuses and MM canals based on age, with patients in group A (<20 years) and group B (20-39 years) exhibiting higher rates of these conformations than those patients in group C (40-60 years) and group D (>60 years) ($P < 0.05$). With regard to group D, the incidence of isthmuses and MM canals in group C was also significantly higher ($P < 0.05$) (Table 2). When a separated distal-lingual root was present, the frequency of isthmuses and MM canals was 45.2%, which was significantly lower than that when such a root was not present ($P < 0.05$) (Table 2).

Next, the 198 detected isthmus canals were analyzed based on the level at which the isthmus was no longer evident, allowing us to classify these canals into seven different types, the prevalence and length of which was then measured (Table 3). In 47.89% of cases, isthmuses were present in the cervical third, while in 18.48% of cases they were present in the middle third, and in 17.36% of cases they were present in the apical third (Table 3). In 28 of these cases, complete communication between mesial-buccal canal and mesial-lingual canal was detectable. The average isthmus length in the mesial root was 4.26 ± 3.14 mm.

True MM canal prevalence was only 3.08 % in first mandibular molars. Of these 11 true MM canal cases, 2 exhibited a separate orifice distinct from the MB and ML canals, while 6 shared the same orifice with either the MB or ML canal, and the remaining cases branched off from either the middle or the apical third of the MB or ML canals (Table 4).

Discussion

We found the prevalence of isthmuses to be roughly consistent with the findings of previous studies, although we detected a lower than expected incidence of MM canals. There may be a few different potential explanations for this finding. For one, MM canals are defined differently in various studies. For

example, Mehrnaz Tahmasbi [6] defined MM canals as we did in the present study, identifying them based on the presence of a distinct round cross-sectional area in radiographic image located in-between the MB and ML canals without regard for whether or not an isthmus was present. In other studies, however, MM canals were considered to be present if 3 canals were evident in the mesial roots. While most studies define isthmuses based on the presence of thin ribbon-shaped communications visible in between the MB and ML canals, others such as Hsu and Kim [12] define types of isthmuses, with a type I isthmus being defined as having either 2 or 3 canals with no noticeable communication, types II and III isthmuses containing 2 and 3 canals, respectively, each with clear communication between the main canals, type IV isthmuses having canals extending into the isthmus area, and type V isthmuses having extensive connections throughout the region. By this definition, MM canals would thus be classified as a form of isthmus, thereby potentially throwing off prevalence rates for these two types of features in certain studies.

We chose to define isthmuses and MM canals based on the criteria outlined in the study of Mehrnaz Tahmasbi, making effort to distinguish between isthmuses and true MM canals. We classified isthmuses and true MM canals as specific kinds of communication between the MB and ML canals, and once we established prevalence rates of these features, we analyzed them separately.

An additional factor potentially complicating comparisons between studies is that a range of different techniques have been employed to explore the morphology of mandibular molar root canals, and different approaches have their unique advantages and disadvantages. Previously used methods include plastic casts [11], staining and clearing [13], an operating microscope [14], and micro-CT imaging [3, 4, 15]. These different methods have all yielded different prevalence rates with respect to isthmuses and MM canals. Past findings and relative advantages of some of these approaches are detailed in the following paragraphs.

Microscope-based examinations have determined MM canal frequencies to be from 1 – 37.5% [7, 8, 10, 16, 17]. In a study by Sunil Kim [2], a total of 72/106 teeth exhibited isthmuses, with no evidence of such communications in the remaining 34 teeth. In that same study, cumulative 4-year survival rates were 61.5% and 87.4% when isthmuses were present and absent, respectively. In certain clinical studies, including the above research, MM canals were determined to be present based on the detection of 3 canals upon treatment. However, as certain canal conformations such as a Type IV canal can be present without any true MM canal in evidence, this approach is insufficient as a means of accurately determining the rates of MM canal prevalence.

With respect to micro-CT, this approach has more success than microscopic and clearing-based techniques for accurately observing fine anatomic structures such as MM canals in the resultant 2D and 3D images. Indeed, in some studies highly complex configurations of mandibular first molar root canals were detected, defying standardized classification efforts [18]. When patients were separated by age and assessed for isthmus presence, frequencies for individuals aged 20 - 39 years, 40 - 59 years, and ≥ 60 years were 50%, 41%, and 24%, respectively [19].

Mehrnaz Tahmasbi used CBCT-based imaging and reported that an MM canal prevalence of 26% in the mesial roots of mandibular first molars, while the frequency of isthmuses was 64.7%. Another study employing CBCT imaging detected an MM canal prevalence rate of just 0.35% (5/1435 teeth), while 26/1435 teeth showed evidence of only a single mesial canal (1.81%) [9]. These and other results suggest that CBCT imaging results may be less accurate than micro-CT based imaging strategies, however as micro-CT images are often only available from older patients or those with serious periodontal diseases, the results from such studies cannot be construed as being representative of the general population. CBCT is primarily utilized in pre-treatment clinical contexts, making it an extremely valuable technique for establishing the presence of isthmuses and MM canals in areas of operative interest. Indeed, given its superiority over traditional periapical radiography, it has been proposed as an ideal means of identifying root canals [9]. For these reasons, we selected CBCT imaging for use in the present retrospective analysis.

We found frequencies of communications containing isthmuses and MM canals to be 59.54% in males and 57.61% in females. Incidence rates of these anatomical features on the left and right sides were 61.21% and 56.25%, respectively. Neither of these differences was significant, and these prevalence rates were consistent with previous studies [7]. We did, however, identify a significant relationship between the presence of additional mesial-lingual roots and the incidence of MB-ML communication. Mandibular first molars most typically have a total of 2 roots and 3 canals in the mesial and distal roots [20]. Different populations seem to exhibit different rates of additional mandibular molar roots, with rates of 22.7 % in Japanese patients [21], 15 % in patients from Hong Kong [22], and 29 % in those from China [23]. A lower rate of just 3.64% was detected in a Turkish Cypriot population. We found in our assessment of a Chinese population that the incidence of additional distal roots was 20.4%. Of the 73 individual analyzed cases in which a separated distal-lingual root was evident, 41.1% of these showed evidence of either an isthmus or MM canal. A significantly higher rate was detected in those cases without this separated distal-lingual root. The reason for this significant variation between these two groups remains uncertain and warrants further investigation.

Our findings indicate that younger individuals are significantly more likely to have MM canals and isthmuses than their older counterparts, with an incidence of MB and ML communication in mandibular first molars of 71.4% in those age 20 or less. In their micro-CT reconstruction-based study of isthmus anatomy, Gu et al. [19] found that isthmuses were present significantly more often in younger patients (20-39 years old; 50%) relative to older patients (60+ years old; 24%), with an increasing ratio of partial to complete isthmuses increasing with age [19]. This was also true in a study conducted by Ali Nosrat [8], and are consistent with our results in the present study. Together these results indicate that clinicians should take additional time to explore and assess the pulp chamber floor area between MB and ML canals to identify any isthmuses which may be present, especially in younger patients.

When assessing types and lengths of isthmuses, we diverged from the Vertucci standard and instead divided roots into three portions: the cervical, middle, and apical sections. As different sections of the isthmus can have different effects and clinical significance, and this stratification can have important

clinical implications. In clinical settings, particularly using microscopic enhancement, cervical sections can be readily dealt with using ultrasonic tips. Such isthmuses were evident in 47.89% of cases, suggesting that isthmuses in the cervical third should be routinely treated under the microscope via standard RCT approaches. The apical portion of the RCS has been considered critical for therapeutic and pathogenic reasons [24], with cleaning isthmuses in this region being essential to effective long-term treatment outcomes. Even when appropriate irrigation and preparation techniques are used, preventing and controlling infections in this site can be challenging. We determined that 17.36% of cases in our study exhibited isthmuses in the apical third region, although limitations in the CBCT technology may mean that the true prevalence rate is even higher as the apical structure is relatively subtle. Fortunately, treatment of this region via apical surgery typically yields favorable clinical outcomes. The treatment of the apical area by apical surgery often achieves ideal clinical results, although surgical success rates are highest in teeth that do not have an isthmus relative to those with a prepared isthmus. Given these success rates and the fact that there is a risk of weakened roots following preparation, efforts must be made to improve isthmus preparation techniques for isthmuses located in the apical region. Enhanced efforts to understand how such isthmuses affect surgical outcomes will make it possible to ensure better patient results [2]. Isthmuses located in the middle third region are relatively more difficult to effectively clean in the process of apical surgery or RCT. Substantial amounts of tissue may be destroyed to access isthmuses in this location, potentially significantly increasing fracture rates. We found that the prevalence rate of such isthmuses was 18.48%, which is fairly low. While it is important to clean isthmuses located in the middle third section, they are not as prevalent as those in cervical or apical regions. Given the risks of operating in this site, conservative measures should be taken during treatment.

Given the relatively strict definition which we used, we detected a very low prevalence of true MM canals by CBCT. Our and others' results suggest that true MM canals should only be identified when three distinct independent canals are present. Detecting and managing isthmuses presents as an important but complex challenge. Navigating both isthmuses and MM canals can be accomplished using a combination of imaging approaches, tactile search techniques, and an operative microscope to carefully and accurately navigate these delicate and complex structures. Future studies assessing long-term patient outcomes are needed to better explore the effects of MM canal preparation on non-surgical endodontic treatments in mandibular first and second molars.

Conclusions

In this retrospective study of CBCT images, we identified a complex range of mandibular first molar canal configurations among a Chinese population. As a leading cause of the failure of endodontic treatments for mandibular molars is the presence of a canal isthmus [12], these results will provide important anatomic insights valuable to endodontists seeking to locate and identify isthmuses and MM canals, thereby potentially avoiding the unnecessary removal of tooth structures.

Abbreviations

CBCT: Cone Beam Computed Tomography; MM: Mesial Middle; M: Mesial; D: Distal; L: Lingual; B: Buccal; mm: millimeter; RCT: Root Canal Therapy.

Declarations

Ethics approval and consent to participate

This study was approved by the Institutional Review Board and Ethical Committee of Stomatological Hospital, Southern Medical University. Informed consent was waived by the study subjects because all the patient data were anonymized and de-identified before the analysis.

Consent for publication

Not applicable.

Availability of data and material

The data analysed during this present study are available from corresponding author on request.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was supported by Scientific research start-up project of Southern Medical University in 2017 (grant no. PY2017N037) and Youth Innovation Talents Project of "Innovation to improve School " in Guangdong Province (2017KQNCX029) in relation to English proofreading.

Authors' contributions

XZ and ZZ conceived the original idea and designed the experiment. SX and ZL performed the experiment and contributed to the interpretation of the results. JD wrote the manuscript with support from YL. All authors read and approved the final manuscript.

Acknowledgements

Not applicable.

References

1. Walton RE: Histologic evaluation of different methods of enlarging the pulp canal space. *J Endod* 1976, 2(10):304-311.
2. Kim S, Jung H, Kim S, Shin SJ, Kim E: The Influence of an Isthmus on the Outcomes of Surgically Treated Molars: A Retrospective Study. *J Endod* 2016, 42(7):1029-1034.
3. Villas-Boas MH, Bernardineli N, Cavenago BC, Marciano M, Del Carpio-Perochena A, de Moraes IG, Duarte MH, Bramante CM, Ordinola-Zapata R: Micro-computed tomography study of the internal anatomy of mesial root canals of mandibular molars. *J Endod* 2011, 37(12):1682-1686.
4. Wolf TG, Paque F, Zeller M, Willershausen B, Briseno-Marroquin B: Root Canal Morphology and Configuration of 118 Mandibular First Molars by Means of Micro-Computed Tomography: An Ex Vivo Study. *J Endod* 2016, 42(4):610-614.
5. Versiani MA, Ordinola-Zapata R, Keles A, Alcin H, Bramante CM, Pecora JD, Sousa-Neto MD: Middle mesial canals in mandibular first molars: A micro-CT study in different populations. *Arch Oral Biol* 2016, 61:130-137.
6. Tahmasbi M, Jalali P, Nair MK, Barghan S, Nair UP: Prevalence of Middle Mesial Canals and Isthmi in the Mesial Root of Mandibular Molars: An In Vivo Cone-beam Computed Tomographic Study. *J Endod* 2017, 43(7):1080-1083.
7. Azim AA, Deutsch AS, Solomon CS: Prevalence of middle mesial canals in mandibular molars after guided troughing under high magnification: an in vivo investigation. *J Endod* 2015, 41(2):164-168.
8. Nosrat A, Deschenes RJ, Tordik PA, Hicks ML, Fouad AF: Middle mesial canals in mandibular molars: incidence and related factors. *J Endod* 2015, 41(1):28-32.
9. Kim SY, Kim BS, Woo J, Kim Y: Morphology of mandibular first molars analyzed by cone-beam computed tomography in a Korean population: variations in the number of roots and canals. *J Endod* 2013, 39(12):1516-1521.
10. Pomeranz HH, Eidelman DL, Goldberg MG: Treatment considerations of the middle mesial canal of mandibular first and second molars. *J Endod* 1981, 7(12):565-568.
11. Skidmore AE, Bjorndal AM: Root canal morphology of the human mandibular first molar. *Oral Surg Oral Med Oral Pathol* 1971, 32(5):778-784.
12. Hsu YY, Kim S: The resected root surface. The issue of canal isthmuses. *Dent Clin North Am* 1997, 41(3):529-540.
13. Gulabivala K, Aung TH, Alavi A, Ng YL: Root and canal morphology of Burmese mandibular molars. *Int Endod J* 2001, 34(5):359-370.

14. Karapinar-Kazandag M, Basrani BR, Friedman S: The operating microscope enhances detection and negotiation of accessory mesial canals in mandibular molars. *J Endod* 2010, 36(8):1289-1294.
15. Harris SP, Bowles WR, Fok A, McClanahan SB: An anatomic investigation of the mandibular first molar using micro-computed tomography. *J Endod* 2013, 39(11):1374-1378.
16. Fabra-Campos H: Three canals in the mesial root of mandibular first permanent molars: a clinical study. *Int Endod J* 1989, 22(1):39-43.
17. Goel NK, Gill KS, Taneja JR: Study of root canals configuration in mandibular first permanent molar. *J Indian Soc Pedod Prev Dent* 1991, 8(1):12-14.
18. Kim Y, Perinpanayagam H, Lee JK, Yoo YJ, Oh S, Gu Y, Lee SP, Chang SW, Lee W, Baek SH et al: Comparison of mandibular first molar mesial root canal morphology using micro-computed tomography and clearing technique. *Acta Odontol Scand* 2015, 73(6):427-432.
19. Gu L, Wei X, Ling J, Huang X: A microcomputed tomographic study of canal isthmuses in the mesial root of mandibular first molars in a Chinese population. *J Endod* 2009, 35(3):353-356.
20. Nur BG, Ok E, Altunsoy M, Aglarci OS, Colak M, Gungor E: Evaluation of the root and canal morphology of mandibular permanent molars in a south-eastern Turkish population using cone-beam computed tomography. *Eur J Dent* 2014, 8(2):154-159.
21. de Souza-Freitas JA, Lopes ES, Casati-Alvares L: Anatomic variations of lower first permanent molar roots in two ethnic groups. *Oral Surg Oral Med Oral Pathol* 1971, 31(2):274-278.
22. Fabra-Campos H: Unusual root anatomy of mandibular first molars. *J Endod* 1985, 11(12):568-572.
23. Zhang R, Yang H, Yu X, Wang H, Hu T, Dummer PM: Use of CBCT to identify the morphology of maxillary permanent molar teeth in a Chinese subpopulation. *Int Endod J* 2011, 44(2):162-169.
24. Ricucci D, Siqueira JF, Jr.: Biofilms and apical periodontitis: study of prevalence and association with clinical and histopathologic findings. *J Endod* 2010, 36(8):1277-1288.

Tables

Table 1. Incidence rate of MM and isthmus

	Isthmus	MM	Without communication	Total
Number	198	11	138	357
Incidence	55.46%	3.08%	41.46%	100%

Table 2. Variables associated with communications (isthmus and MM)

Independent variables	Group	Number	Incidence%	P value
Age	<20 years old	28	(20) 71.4%	< 0.05
	20-39 years	137	(96) 70.1%	
	40-59 years	143	(82) 57.3%	
	≥60years	49	(14) 28.6%	
Sex	Male	173	(103) 59.5%	> 0.05
	Female	184	(109) 59.2%	
Position	36	167	(104) 62.3%	> 0.05
	46	190	(108) 56.8%	
Seperated distal-lingual root canal	present	73	(33) 45.2%	<0.05
	not present	284	(179) 63.0%	

Table 3. Length and distribution of isthmuses based on the isthmus beginning and end

	Type1	Type2	Type3	Type4	Type5	Type6	Type7	Total
Cervical third	+	+	+	+	-	-	-	
Middle third	+	+	-	-	+	-	+	
Apical third	+	-	+	-	+	+	-	
Number	28	24	11	108	10	13	4	198
Prevalence of isthmus %	7.84%	6.72%	3.08%	30.25%	2.80%	3.64%	1.12%	55.46%
Length(mm)	9.92±	5.86±1.16	5.69±	2.42±	6.16±	2.47±	2.11±	4.26±
	3.25		1.70	0.93	1.93	1.46	1.48	3.14

+, present; -, absent

Table 4. Type distribution of MM canals based on descriptions of Pomeranz

MM	Number	Incidence (%)
Type I	3	0.84%
Type II	6	1.68%
Type III	2	0.56%
Total	11	3.08%

Figures



Figure 1

An axial CBCT image showing the isthmus between MB canal and ML canal of tooth #30. The arrow points to the isthmus.




Figure 2

(A) An axial CBCT image showing the MM canal of tooth #19. The arrow points to the MM canal. (B) An axial CBCT image showing the MM canal of tooth #30. The arrow points to the MM canal.