

# The regulatory role of the RANKL/RANK/OPG signaling pathway in the mechanisms of tooth eruption in patients with impacted teeth

**Petro Flis**

Natsional'nii meditsnii universitet imeni O O Bogomoltsa

**Ludmila Brodetska**

Natsional'nii meditsnii universitet imeni O O Bogomoltsa

**Olexandr Kaniura**

Natsional'nii meditsnii universitet imeni O O Bogomoltsa

**Olha Lisakovska** (✉ [o.lisakovskaya@gmail.com](mailto:o.lisakovskaya@gmail.com))

Institut biohimii imeni O V Palladina Nacional'na akademija nauk Ukraini <https://orcid.org/0000-0002-5844-1453>

**Larysa Natrus**

Natsional'nii meditsnii universitet imeni O O Bogomoltsa

**Irina Skrypnyk**

Natsional'nii meditsnii universitet imeni O O Bogomoltsa

---

## Research article

**Keywords:** impacted teeth, cytokine system RANKL/RANK/OPG, nuclear factor kappa-B, bone remodeling, tooth eruption, osteoclastogenesis, apoptosis

**Posted Date:** December 30th, 2019

**DOI:** <https://doi.org/10.21203/rs.2.19720/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on September 18th, 2020. See the published version at <https://doi.org/10.1186/s12903-020-01251-y>.

# Abstract

**Background** Tooth impaction is a common problem in orthodontic practice and usually accompanied by pain and pathological changes of surrounding teeth. Understanding the cellular and molecular mechanisms underlying tooth impaction allows finding the most effective way of orthodontic treatment for patients with impacted teeth (IT). The RANK (receptor activator of NF- $\kappa$ B) / RANKL (RANK ligand) / OPG (osteoprotegerin) signaling pathway controls bone resorption and may be involved in the regulation of tooth eruption. The aim of the study was to evaluate the bone remodeling based on the assessment of the RANKL/RANK/OPG status in patients with IT.

**Methods** Bone samples from 20 patients (mean age 15.27, SD = 3.83) were divided into 3 groups: 1 – bone tissue of healthy persons (control group); 2 – bone tissue taken near the healthy tooth in patients with tooth impaction; 3 – bone tissue taken near the IT. Levels of RANKL, RANK, OPG, osteocalcin (OC), NF- $\kappa$ B p65 subunit, NFATc1 and caspase-3 were determined by Western blotting. The difference between groups was assessed by one-way ANOVA. P-values less than 0.05 were considered statistically significant.

**Results** We established a 1.73-fold elevation of RANK level in the IT area vs. control, indicating the recruitment of preosteoclasts. An increase in RANKL, OPG and OC content was demonstrated (by 1.46, 1.48, and 1.42 fold respectively), reflecting high activity of osteoblasts near the IT. Despite the activation of the RANKL/RANK/OPG system in the impaction area, NF- $\kappa$ B and NFATc1 levels did not change compared vs. control, indicating a blocked/delayed process of osteoclastogenesis. We found a decrease in the content of procaspase-3 (1.28-fold), while the level of its active form p17 increased by 2.26 fold near the healthy tooth in patients with IT compared with control. In the area of IT we observed an increase in procaspase-3 and p17 levels (1.32 and 1.78 fold). This reflects impairments of caspase-3 activation and accumulation of its inactive form in the IT area that may contribute to the tooth eruption failure.

**Conclusions** Tooth impaction may be associated with the disturbances in the caspase-3 cascade activation and the imbalance in the RANKL/RANK/OPG system, and as a result, blocked bone resorption.

## Background

Impacted teeth (IT) are those that remain partially or completely in the bone tissue or under the mucous membrane for 2 years after the term of their physiological eruption. Tooth impaction is usually accompanied by pain, pathological resorption of the surrounding teeth, and even an increased risk of malignancies [1]. Delayed or failed tooth eruption is a common problem in daily orthodontic practice, therefore it is important to understand the cellular and molecular mechanisms underlying blocking tooth from breaking through the gum, in order to find the most effective way of orthodontic treatment for patients with IT. The presence of unerupted teeth requires a thorough diagnostics and a balanced approach to the choice of treatment method, which depends on the overall clinical situation, the patient's age and general somatic status.

The process of tooth eruption is a complex highly regulated process in which the cells of both the tooth organ and the surrounding alveolus are involved. Eruption of the teeth and their exit into the oral cavity is considered to be a valuable model for studying the process of bone remodeling, since it involves tightly coordinated processes of bone formation and bone resorption with the participation of osteoblasts and osteoclasts respectively [2]. For an adequate eruption process, mononuclear cells (osteoclast precursors) have to be recruited into the dental follicle during the preruptive phase [3]. The process of their fusion occurs to form active multinuclear osteoclasts that are responsible for the formation of an eruptive pathway in the alveolar bone for tooth movement.

More recently, cytokine systems that are involved in the regulation of tooth eruption, has been intensively explored, in particular the main efforts has been focused on the studying the signaling pathway RANK (receptor activator of nuclear factor kappa-B) / RANKL (RANK ligand) / OPG (osteoprotegerin).

RANKL is one of the tumor necrosis factor (TNF) superfamily members and acts through RANK as a potent inducer of osteoclast formation and bone resorption. RANKL is produced by bone marrow stromal cells, osteoblasts, activated dendritic cells and T-lymphocytes [4]. RANK is expressed as a transmembrane heterotrimer on the surface of haematopoietic osteoclasts progenitors, mature osteoclasts, chondrocytes and other cells. Osteoprotegerin (OPG) is a decoy receptor for RANKL and prevents its osteoclastogenic action. The adequate balance between RANKL and OPG determines the rate of bone resorption [5]. An impaired OPG to RANKL ratio can be considered as an essential pathogenetic factor for the development of bone tissue diseases.

RANKL/RANK/OPG axis regulates the maturation of osteoclasts and bone resorption, acting through the activation of the transcription factor NF- $\kappa$ B [6]. However, cells that are responsible for bone formation—osteoblasts—can influence the course of the tooth eruption by modulating the state of the RANKL/RANK/OPG axis as well, since they synthesize both osteoprotegerin, which is an inhibitor of bone resorption, and RANKL, which acts as its inducer. The role of the RANKL/RANK/OPG signaling pathway in the process of tooth eruption has been largely demonstrated on knockout rodents [7–9]. Nevertheless there is a lack of experimental studies related to the role of this system in tooth eruption process in humans due to the difficulties with obtaining the written permissions from patients and an approval from the local ethics committee.

Understanding the cellular and molecular mechanisms underlying tooth eruption and its complications allows developing and implementing the most effective way of orthodontic treatment for patients with impacted teeth in clinical practice. In view of this, the aim of the study was to evaluate the status of bone remodeling based on the assessment of changes in the RANKL/RANK/OPG axis in patients with tooth impaction.

## Methods

***Ethical approval.*** The study was registered with the research center of Bogomolets National Medical University. Ethical approval for the study was obtained from Ethics and Research Committee of

Bogomolets National Medical University (Protocol No 95 from 05/05/2016). Written informed consent was obtained from each patient. Any uncooperative patients and patients who had not provided consent to participate in the study, were excluded from the study.

***Patients and sample collection.*** Among 20 participants, consisting of 9 males and 11 females, the average age was 15.27 years.

Bone samples were divided into three groups: 1 – bone tissue of healthy persons from the area of third molar that has been removed according to orthodontic prescriptions; in patients of this group all teeth were normally erupted based on the data of X-ray diagnostics (control group); 2 – bone tissue of patients with tooth impaction taken from the area of normally erupted third molar that has been removed according to orthodontic prescriptions; 3 – bone tissue of patients with tooth impaction taken near the impacted tooth. After collection bone samples were frozen in liquid nitrogen in order to prepare total protein lysates.

***Western blot analysis.*** Levels of RANKL, RANK, OPG, osteocalcin (OC), NF- $\kappa$ B p65 subunit, NFATc1 and caspase-3 in bone tissue were determined by Western blot analysis. Bone tissue samples (100 mg) were ground in a porcelain mortar with liquid nitrogen. After that samples were lysed by an incubation on ice for 20 min in 0.9 ml of RIPA buffer (20 mM Tris-HCl, pH 7.5; 150 mM NaCl; 1% Triton X-100; 1 mM EGTA; 0.1% SDS, 1% sodium deoxycholate, 10 mM sodium pyrophosphate) containing protease inhibitor cocktail (PIC, Sigma, USA), sonicated and then centrifuged at 14000 g for 20 min at +4°C. Protein concentrations in the lysates were measured by Lowry's method. 30  $\mu$ g of protein were loaded onto 10% SDS polyacrylamide gels. After SDS-PAGE, the proteins were transferred onto nitrocellulose membrane (350 mA, 1 h). Immunoblotting was conducted with primary antibodies against RANKL and OPG (1:250; Santa Cruz, USA), RANK, OC and caspase-3 (1:500; Santa Cruz, USA), NF- $\kappa$ B p65 subunit (1:500; Thermo Fisher Scientific, USA), NFATc1 (1:1000; Thermo Fisher Scientific, USA) and  $\beta$ -actin (1:10000; Sigma-Aldrich, USA). Primary-antibody-bound membranes were incubated with peroxidase-conjugated secondary antibodies: anti-mouse IgG (Fab Specific)-Peroxidase (1:2500; Sigma, USA), anti-rabbit IgG (H+L)-HRP conjugate (1:4000; Bio-Rad Laboratories, Inc., USA) or anti-goat IgG (H+L) (1:2500; Invitrogen, USA). Thereafter the membranes were developed with chemiluminescent agents: p-coumaric acid (Sigma-Aldrich, USA) and luminol (AppliChem GmbH). Tissue levels of all proteins were normalized to  $\beta$ -actin. The immunoreactive bands were quantified using [Gel-Pro Analyzer32](#) software.

***Statistical analysis.*** The results are expressed as the mean  $\pm$  SEM. Data distribution was analyzed using the Kolmogorov-Smirnov normality test. Statistical differences between groups were analyzed by the one-way ANOVA and Tukey's post hoc test. Differences were considered to be statistically significant when p-value was less than 0.05. All statistical analyses were carried out using Origin Pro 8.5 (OriginLab Corporation, Northampton, MA, USA).

## Results

## ***Assessment of the RANKL/RANK/OPG axis and ratio between its components in bone tissue of patients***

As the RANKL/RANK/OPG signaling axis regulates the interplay between osteoclasts and osteoblasts that can influence bone remodeling [6], we first determined the content of the RANKL/RANK/OPG cytokine system components in the bone tissue of patients with tooth impaction. The results of the study presented in Fig. 1 show that there is no difference in the relative protein content of RANK (Fig. 1 A, B) and osteocalcin (Fig. 1 A, D) between the samples of healthy persons (control group) and samples taken near the healthy tooth in patients with tooth impaction. At the same time, it was shown a slight tendency to decreased content of osteoprotegerin (by 1.2 times) in the bone tissue near the healthy tooth in patients with tooth impaction compared with the control (Fig. 1C) and we observed a significant decrease in the RANKL content (membrane-bound form - by 1.6 fold, and soluble form - by 10 fold) (Fig. 2C, 2D).

**Fig. 1** Immunoblotting analysis of RANK, OPG and osteocalcin in bone tissue samples: representative immunoblots are shown (A) and quantified using  $\beta$ -actin as a loading control. The bar graphs of RANK (B), OPG (C) and osteocalcin (D) are presented as arithmetic means  $\pm$  SEM (n = 6/group): 1 - bone tissue of healthy persons (control group); 2 - bone tissue taken near the healthy tooth in patients with tooth impaction; 3 - bone tissue taken near the impacted tooth. \* p<0.05 vs. control, # p<0.05 vs. bone tissue taken near the healthy tooth in patients with tooth impaction.

Completely different results were obtained when studying the area of the impacted tooth of the same patients. Western blot analysis data revealed a 1.73-fold elevation of the RANK level in the IT area vs. control (Fig. 1B), indicating an accumulation of osteoclast precursors. Moreover, a significant increase in the relative content of OPG (Fig. 1C), osteocalcin (Fig. 1D) and total RANKL (Fig. 2B) proteins (by 1.48, 1.42, and 1.46 fold, respectively) was demonstrated as well, that could correlate, at least partially, with the high activity of osteoblasts and/or its precursors in the area of IT.

**Fig. 2** Immunoblotting analysis of RANKL in bone tissue samples: representative immunoblots are shown (A) and quantified using  $\beta$ -actin as a loading control. The bar graphs of total content of RANKL (B), its membrane-bound (C) and soluble (D) forms are shown as arithmetic means  $\pm$  SEM (n = 6/group): 1 - bone tissue of healthy persons (control group); 2 - bone tissue taken near the healthy tooth in patients with tooth impaction; 3 - bone tissue taken near the impacted tooth. \* $p < 0.05$  vs. control, # $p < 0.05$  vs. bone tissue taken near the healthy tooth in patients with tooth impaction.

Physiological effects of OPG and RANKL on the bone tissue metabolism are known to be largely determined by the ratio of their synthesis [10]. Based on the measurement of protein levels of RANKL/RANK/OPG signaling pathway components by the Western blot analysis, we obtained the ratios applied for assessing the status of local bone remodeling in the areas near the healthy and impacted teeth in patients with tooth impaction. It was established a significant decrease in the the RANK/RANKL (2.6-fold) and OPG/RANKL (2.5-fold) ratios in the area near the impacted tooth compared with the healthy tooth (Table 1). These reduced ratios may reflect the violation of local interrelation between bone resorption and bone formation in the IT area.

Table 1 Ratios of the RANKL/RANK/OPG signaling pathway components in the bone tissue of patients, in a.u. (arbitrary units)

Ratio, a.u.	Bone tissue taken near the healthy tooth in patients with tooth impaction	Bone tissue taken near the impacted tooth
RANK	0.97 $\pm$ 0.06	1.73 $\pm$ 0.11*
RANKL	0.32 $\pm$ 0.02	1.46 $\pm$ 0.12*
OPG	0.8 $\pm$ 0.07	1.48 $\pm$ 0.13*
RANK/RANKL	3.03	1.18*
OPG/RANKL	2.5	1.01*

\* $p < 0.05$  vs. bone tissue taken near the healthy tooth in patients with tooth impaction

### ***NF- $\kappa$ B-mediated signaling and apoptosis in the bone tissue***

Binding RANKL to RANK leads to the further activation of NF- $\kappa$ B and translocation of this transcription factor to the nucleus in order to stimulate osteoclastogenesis [11]. To

address whether excessive RANKL and RANK levels in bone tissue near the area of tooth impaction are involved in the downstream transcriptional activation, we explored the involvement of NF- $\kappa$ B in the mechanisms of tooth eruption.

We demonstrated a slight decrease in the NF- $\kappa$ B (Fig. 3 A, B) and NFATc1 (Fig. 3 A,C) levels by 17 and 20% respectively in the bone tissue selected near the healthy tooth compared vs. control patients. Interestingly, it was first demonstrated that despite the activation of the RANKL/RANK/OPG system in the impaction area, the content of both NF- $\kappa$ B and NFATc1 did not significantly change compared vs. control group, indicating a blocked or delayed process of osteoclastogenesis near the IT although an accumulation of osteoclast precursors in this area was observed.

**Fig. 3** Immunoblotting analysis of NF- $\kappa$ B, NFATc1 and caspase-3 in bone tissue samples: representative immunoblots are shown (A, D) and normalized to  $\beta$ -actin. The bar graphs of NF- $\kappa$ B (B), NFATc1 (C), procaspase-3 (E) and its active subunit p17 (F) are demonstrated as arithmetic means  $\pm$  SEM (n = 6/group): 1 - bone tissue of healthy persons (control group); 2 - bone tissue taken near the healthy tooth in patients with tooth impaction; 3 - bone tissue taken near the impacted tooth. \*p<0.05 vs. control, #p<0.05 vs. bone tissue taken near the healthy tooth in patients with tooth impaction.

Since *caspase cascades* play a key role in an *activation* of cell apoptosis [12] and, moreover, may be triggered by NF- $\kappa$ B, we determined the levels of procaspase-3 (32 kDa) and its active subunit p17 (17 kDa). It was shown a decrease in the content of procaspase-3 (32 kDa) in the area of healthy tooth of patients with tooth impaction by 28%, while in the area of impacted tooth its content was increased by 1.32 fold compared with control (Fig. 3E). Considering the active caspase-3 enzyme is a heterodimer consisting of two p17 and two p11 caspase-3 subunits [13], it was also important to evaluate the content of the activated p17 form. We detected an increase of its protein level in the area of healthy tooth by 2.26 fold compared vs. control, and an elevation in the area of impacted tooth by 1.78 fold (Fig. 3F) on the background of an accumulation of inactive procaspase-3 32 kDa. These observations indicate the cleavage of procaspase-3 and its activation in the area of

healthy tooth of patients with IT, that promote the tooth eruption process. We also suggest the impairment of caspase-3 activation and the accumulation of an inactive form in the area of tooth impaction may lead to the tooth eruption failure.

Our data demonstrate that tooth impaction may be associated with blocked NF- $\kappa$ B signaling and NFATc1 expression and with the disturbances in the cascade activation of caspase-3.

## Discussion

The cellular and molecular mechanisms underlying tooth impaction still remain poorly understood. It is assumed that an imbalance between the osteosynthesis and bone resorption is one of the factors playing an important role in the development of tooth eruption abnormalities [14]. A key role in controlling the process of bone remodeling belongs to cytokine systems. The regulatory protein osteocalcin, which is produced by osteoblasts [15], is involved in the process of osteosynthesis and considered a valid marker of bone formation. In turn, bone resorption is regulated by the osteotropic cytokine system RANKL/RANK/OPG [5].

The role of the RANKL/RANK/OPG signalling pathway in tooth eruption has been demonstrated in knockout rats and mice in greater extent [16]. Nevertheless, there is a lack of experimental studies dedicated to the role of RANKL/RANK/OPG axis in tooth eruption process in humans under the normal and pathological conditions due to the complexity of sampling and obligatoriness of confirmation from patients and ethics committee approval. The small number of studies dealing exclusively with the changes in the state of the cytokine system RANKL/RANK/OPG in bone tissue of patients with impacted teeth is available [17, 18]. Therefore, the purpose of our research was to determine the levels of its key components as indicators of the processes of bone resorption and bone formation.

The current findings showed no significant difference in RANK and OC levels between the bone tissue samples of healthy persons and samples taken near the healthy tooth in patients with tooth impaction. A slight tendency to decreased OPG level in the bone tissue near the healthy tooth in patients with tooth impaction compared with the control was observed on a background of significant decrease in the content of membrane-bound and soluble forms of RANKL. These findings clearly indicate a slight deviation in the production of cytokines by osteoblasts. Nevertheless since we demonstrated a simultaneous decrease in the content of both cytokines RANKL and OPG on the background of unchanged level of the preosteoclast marker RANK and marker of osteosynthesis osteocalcin, in general, this ratio of locally synthesized cytokines ensured a normal process of tooth eruption.

Interestingly, we obtained completely different results concerning the state of the RANKL/RANK/OPG system in the area of the impacted tooth of the same patients. Findings of the present study demonstrated a significant increase in the relative content of RANK, OPG, both forms of RANKL and

osteocalcin in the IT area vs. control. Based on these observations we can assume an accumulation of osteoclast precursors as well as active osteoblastic cells, which excessively produce cytokines in the area of impacted tooth. In addition, since monocytes/macrophages are RANKL producers [19], high RANKL level may be explained by the recruitment of immunocompetent cells and the development of the local proinflammatory reaction near the impaction area—the formation of mononuclear infiltrations, accumulation of preosteoclasts, preosteoblasts and fibroblasts, that actively produce cytokines. Nevertheless, despite the high levels of cytokines tooth eruption does not occur in this area.

Therefore, it may be important to consider the ratios of the components of the RANKL/RANK/OPG system that are determined its physiological effects on the bone tissue metabolism. We used two indices for assessing the state of local bone remodeling in the areas near the healthy and impacted teeth in patients with tooth impaction: RANK/RANKL as an indicator of bone resorption and OPG/RANKL that mainly reflects the process of osteosynthesis. In the area of impacted tooth we found a decreased RANK/RANKL and OPG/RANKL ratios compared with the healthy tooth that may reflect inhibited bone remodeling process and, as a result, disturbances in tooth eruption process.

This method, based on comparison of the RANK/RANKL and OPG/RANKL ratios during the normal and pathological eruption process, allows to estimate the intensity of the processes of bone resorption and bone formation in patients and may be recommended for implementation in orthodontic practice for the detection of local cytokine dysregulation in the impaction area and for choosing an effective method for correction of this condition.

In last decades, considerable progress has been made in understanding the molecular basis of osteoclastogenesis. It has been discovered that the process of osteoclast formation requires the involvement of NF- $\kappa$ B signaling, which is activated in response to several major “osteoclastogenic” cytokines, in particular RANKL and macrophage colony-stimulating factor 1 (M-CSF1) [20]. RANKL and other proinflammatory cytokines such as tumor necrosis factor alpha (TNF $\alpha$ ), activate NF- $\kappa$ B-associated signaling pathways, inducing c-Fos and nuclear activated T-cell factor (NFATc1), and also inhibit NFATc1 repressors, thereby positively regulating formation and activation of osteoclasts [21].

In addition to the importance of the balance between osteosynthesis and bone resorption, it is known that reorganization into the periodontal membrane is also required for an adequate course of the tooth eruption [22]. This process involves the activation of apoptosis of the inner cell layer of the periodontium and the formation of moving layer of apoptotic cells. Recently, it has been suggested that inappropriate process of apoptosis of osteoblasts/osteocytes accounted for, at least in part, the imbalance of bone remodeling. Accumulated evidence suggests that caspase-3 is a critical enzyme for apoptosis and cell survival; however the exact role of caspase-3 in bone remodeling and development of skeletal diseases is largely unknown. To clarify why despite the high osteoclastogenic potential of cells in the impaction area, tooth eruption does not occur, the next step was to estimate the state of NF- $\kappa$ B signaling as a trigger factor for apoptosis in bone tissue of patients with impacted teeth.

Since the RANKL level was declined in the bone tissue selected near the healthy tooth of patients with tooth impaction compared vs. control patients, it was logically to observe a slight decrease in the NF- $\kappa$ B level and, as a result, in the NFATc1 and procaspase-3 (32 kDa) levels. Nevertheless, the content of caspase-3 subunit p17 was significantly higher thereby ensuring the reorganization into the periodontal membrane and normal course of the tooth eruption.

In the present study, we found that there was no difference in the NF- $\kappa$ B and NFATc1 levels compared vs. control group despite the established activation of the RANKL/RANK/OPG signaling axis in the impaction area. Although an accumulation of osteoclast precursors in the IT area was observed, our results indicate a blocked or delayed process of osteoclastogenesis in the bone tissue near the impacted tooth. These data indicate that there is no adequate maturation and activation of osteoclasts, therefore resorption in the area of the impacted tooth does not normally occur, that may explain the tooth impaction and/or delayed tooth eruption. Understanding the molecular basis of impaired bone resorption in the IT area is required further investigation.

While assessing the process of apoptosis we detected an increase in caspase-3 p17 subunit level on the background of declined procaspase-3 content in the area of healthy tooth. At the same time, we obtained different results in the area of impacted tooth and found a simultaneous elevation of both procaspase-3 and caspase-3 p17 subunit. Accordingly to these findings, a ratio between subunit p17 and procaspase-3 was lower in the area of impacted tooth compared with the bone tissue taken near the healthy tooth in patients with tooth impaction. These observations indicate the enhanced cleavage of procaspase-3 and its activation in the area of healthy tooth of patients with IT, that promote the tooth eruption process. We have to emphasize that the impairment of caspase-3 activation and the accumulation of inactive form in the area of tooth impaction may lead to the tooth eruption failure. In addition, it may be assumed that apoptosis activation in the impaction area may possibly indicate the death of functional cells—preosteoclasts and osteoblasts, and explain the absence of NF- $\kappa$ B activation after the observed activation of the RANKL/RANK/OPG system near the IT.

The obtained experimental results may be beneficial for practical medicine, especially for the new field of molecular orthodontics, as they extend the current understanding of the cellular and molecular mechanisms of tooth impaction in humans and emphasize the role of the NF- $\kappa$ B-associated RANKL/RANK/OPG signaling pathway in disturbances of tooth eruption. Based on our data, therapeutic approaches for patients with failed tooth eruption should be directed to restoring the normal course of the resorption process in order to facilitate tooth eruption. Thus, to improve the functioning of local cytokine pathways responsible for osteoregulation in patients with impacted teeth, mechanical stimulation for local therapy may be beneficial. A removable orthodontic apparatus, developed earlier by Flis et al. [23] is allow improving blood circulation within the area of the impacted tooth and may be recommended for the impacted teeth management. The therapeutic use of this apparatus may allow mechanically stimulating the formation of eruptive pathway, setting the direction for mature impacted tooth with high cellular potential and exiting into the oral cavity.

## Conclusions

Our findings suggest that tooth impaction may be associated with blocked NF- $\kappa$ B signaling, disturbances in the cascade activation of caspase-3 and the imbalance in the ratio of components of the RANKL/RANK/OPG signaling axis, and as a consequence, the blocked or delayed process of bone resorption. Based on the measurement in the bone tissue of the RANKL/RANK/OPG signaling pathway components the method for assessing the status of local bone remodeling in the area of impacted tooth has been developed.

## Abbreviations

IT—impacted tooth

M-CSF 1—macrophage colony-stimulating factor 1

NFATc1—nuclear factor of activated T-cells 1

NF- $\kappa$ B—nuclear factor kappa B

OC—osteocalcin

OPG—osteoprotegerin

RANK—receptor activator of NF $\kappa$ B

RANKL—receptor activator of NF- $\kappa$ B ligand

TNF $\alpha$ —tumor necrosis factor alpha

## Declarations

*Availability of data and materials.* The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

*Ethics approval and consent to participate*

The study was approved by the Ethics and Research Committee of Bogomolets National Medical University (Protocol No 95 from 05/05/2016). Written informed consent was obtained after the details of the study were explained to the participants.

*Consent for publication.* Not applicable.

*Competing interests.* The authors declare that they have no competing interests.

*Funding.* No funding was obtained for this study.

*Authors' contributions.* FP, BL and NL designed the study. KO, SI, BL recruited study subjects. SI and BL collected the bone tissue samples. LO performed the Western blot analysis. LO and BL statistically analyzed the data. LO, NL and BL prepared the manuscript. All authors reviewed the original draft as well as read and approved the final manuscript.

*Acknowledgements.* Not applicable.

*Authors' information*


<sup>1</sup> Department of [Orthodontics and propaedeutics of Orthopedic Dentistry](#), Bogomolets National Medical University, Kyiv, Ukraine. <sup>2</sup> Department of Biochemistry of Vitamins and Coenzymes, Palladin Institute of Biochemistry, Kyiv, Ukraine. <sup>3</sup> Research Institute of Experimental and Clinical Medicine, Bogomolets National Medical University, Kyiv, Ukraine.

## References

1. [Santosh P](#). Impacted mandibular third molars: review of literature and a proposal of a combined clinical and radiological classification. [Ann Med Health Sci Res](#). 2015;5(4):229-34. doi: 10.4103/2141-9248.160177.
2. [Wise GE](#), [Frazier-Bowers S](#), [D'Souza RN](#). Cellular, molecular, and genetic determinants of tooth eruption. [Crit Rev Oral Biol Med](#). 2002;13(4):323-34.
3. [Wise GE](#), [King GJ](#). Mechanisms of tooth eruption and orthodontic tooth movement. [J Dent Res](#). 2008;87(5):414-34.
4. [Ponzetti M](#), [Rucci N](#). Updates on osteoimmunology: what's new on the cross-talk between bone and immune system. [Front Endocrinol \(Lausanne\)](#). 2019;10:236. doi: 10.3389/fendo.2019.00236.
5. [Nagy V](#), [Penninger JM](#). The RANKL-RANK Story. [Gerontology](#). 2015;61(6):534-42. doi: 10.1159/000371845.
6. [Wada T](#), [Nakashima T](#), [Hiroshi N](#), [Penninger JM](#). RANKL-RANK signalling in osteoclastogenesis and bonedisease. [Trends Mol Med](#). 2006;12(1):17-25.
7. [Marks SC Jr](#). Osteopetrosis in the toothless (t1) rat: presence of osteoclasts but failure to respond to parathyroid extract or to be cured by infusion of spleen or bone marrow cells from normal littermates. [Am J Anat](#). 1977;149(2):289-97.
8. [Seifert MF](#), [Popoff SN](#), [Marks SC Jr](#). Skeletal biology in the toothless (osteopetrotic) rat. [Am J Anat](#) 1988;183:158-165.
9. [Marks SC Jr](#), [Lane PW](#). Osteopetrosis, a new recessive skeletal mutation on chromosome 12 of the mouse. [J Hered](#). 1976;67:11-18.
10. [Florencio-Silva R](#), [Sasso GR](#), [Sasso-Cerri E](#), [Simões MJ](#), [Cerri PS](#). Biology of bone tissue: structure, function, and factors that influence bone cells. [Biomed Res Int](#). 2015;2015:421746. doi: 10.1155/2015/421746.

11. Novack DV. Role of NF- $\kappa$ B in the skeleton. *Cell Res.* 2011;21(1):169-82. doi: 10.1038/cr.2010.159.
12. Porter AG, Jänicke RU. Emerging roles of caspase-3 in apoptosis. *Cell Death Differ.* 1999;6(2):99-104.
13. Tawa P, Hell K, Giroux A, Grimm E, Han Y, Nicholson DW, Xanthoudakis S. Catalytic activity of caspase-3 is required for its degradation: stabilization of the active complex by synthetic inhibitors. *Cell Death Differ.* 2004;11(4):439-47.
14. Frazier-Bowers SA, Puranik CP, Mahaney MC. The etiology of eruption disorders - further evidence of a 'genetic paradigm'. *Semin Orthod.* 2010;16(3):180-185.
15. Zoch ML, Clemens TL, Riddle RC. New insights into the biology of osteocalcin. *Bone.* 2016;82:42-9. doi: 10.1016/j.bone.2015.05.046.
16. Kong YY, Yoshida H, Sarosi I, Tan HL, Timms E, Capparelli C, et al. OPG is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. *Nature.* 1999;397(6717):315-23. doi:10.1038/16852.
17. Heinrich J, Bsoul S, Barnes J, Woodruff K, Abboud S. CSF-1, RANKL and OPG regulate osteoclastogenesis during murine tooth eruption. *Arch Oral Biol.* 2005;50(10):897-908.
18. Tyrovola JB, Spyropoulos MN, Makou M, Perrea D. Root resorption and the OPG/RANKL/RANK system: a mini review. *J Oral Sci.* 2008;50(4):367-76.
19. Lampiasi N, Russo R, Zito F. The alternative faces of macrophage generate osteoclasts. *Biomed Res Int.* 2016;2016:9089610. doi: 10.1155/2016/9089610.
20. Boyce BF, Xiu Y, Li J, Xing L, Yao Z. NF- $\kappa$ B-mediated regulation of osteoclastogenesis. *Endocrinol Metab (Seoul).* 2015;30(1):35-44. doi: 10.3803/EnM.2015.30.1.35.
21. Matsuo K, Galson DL, Zhao C, Peng L, Laplace C, Wang KZ, Bachler MA, Amano H, Aburatani H, Ishikawa H, Wagner EF. Nuclear factor of activated T-cells (NFAT) rescues osteoclastogenesis in precursors lacking c-Fos. *J Biol Chem.* 2004;279(25):26475-80.
22. Kjær I. Mechanism of human tooth eruption: review article including a new theory for future studies on the eruption process. *Scientifica (Cairo).* 2014;2014:341905. doi: 10.1155/2014/341905.
23. Flis PS, Brodetska LO, Tormakhov MM, Grygorenko OYa. Orthodontic efforts in the treatment of impacted teeth complicated by lack of space in the dentition. *Novyny Stomatolohiyi.* 2019;4(101):69-73.


## Figures



**Figure 1**

Immunoblotting analysis of RANK, OPG and osteocalcin in bone tissue samples: representative immunoblots are shown (A) and quantified using  $\beta$ -actin as a loading control. The bar graphs of RANK (B), OPG (C) and osteocalcin (D) are presented as arithmetic means  $\pm$  SEM ( $n = 6$ /group): 1 – bone tissue of healthy persons (control group); 2 – bone tissue taken near the healthy tooth in patients with tooth


impaction; 3 – bone tissue taken near the impacted tooth. \*  $p < 0.05$  vs. control, #  $p < 0.05$  vs. bone tissue taken near the healthy tooth in patients with tooth impaction.



## Figure 2

Immunoblotting analysis of RANKL in bone tissue samples: representative immunoblots are shown (A) and quantified using  $\beta$ -actin as a loading control. The bar graphs of total content of RANKL (B), its membrane-bound (C) and soluble (D) forms are shown as arithmetic means  $\pm$  SEM (n = 6/group): 1 –

bone tissue of healthy persons (control group); 2 – bone tissue taken near the healthy tooth in patients with tooth impaction; 3 – bone tissue taken near the impacted tooth. \* $p < 0.05$  vs. control, # $p < 0.05$  vs. bone tissue taken near the healthy tooth in patients with tooth impaction.



**Figure 3**

Immunoblotting analysis of NF-κB, NFATc1 and caspase-3 in bone tissue samples: representative immunoblots are shown (A, D) and normalized to β-actin. The bar graphs of NF-κB (B), NFATc1 (C),

procaspase-3 (E) and its active subunit p17 (F) are demonstrated as arithmetic means  $\pm$  SEM (n = 6/group): 1 – bone tissue of healthy persons (control group); 2 – bone tissue taken near the healthy tooth in patients with tooth impaction; 3 – bone tissue taken near the impacted tooth. \*p<0.05 vs. control, #p<0.05 vs. bone tissue taken near the healthy tooth in patients with tooth impaction.