Assessment of the Effects of Stretch-Injury on Primary Rat Microglia

Mike Shaughness (✉ michael.shaughness.ctr@usuhs.edu)

Uniformed Services University: Uniformed Services University of the Health Sciences

<https://orcid.org/0000-0003-4138-3311>

Kimberly Byrnes

Uniformed Service University

Research

Keywords: Microglia, mechanical stretch-injury, in vitro, inflammation, TBI, migration, morphology

Posted Date: November 13th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-104466/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Molecular Neurobiology on March 24th, 2021. See the published version at <https://doi.org/10.1007/s12035-021-02362-5>.

Abstract

Background: Mechanical stretch-injury is a prominent force involved in the etiology of traumatic brain injury (TBI). It is known to directly cause damage and dysfunction in neurons, astrocytes, and endothelial cells. However, the deleterious effects of stretch-injury on microglia, the brain's primary immunocompetent cell, are currently unknown.

Methods: The Cell Injury Controller II (CICII), a validated model of cellular neurotrauma, was used to induce a mechanical stretch-injury in primary rat microglia. Statistical analysis utilized student t-test and one and two way-ANOVAs with Tukey's and Sidak's multiple comparisons, respectively.

Results: Cells exposed to stretch-injury showed no signs of membrane permeability, necrosis, or apoptosis, as measured by media derived lactate dehydrogenase (LDH) and cleaved-caspase 3 immunocytochemistry, respectively. Interestingly, injured cells displayed a functional deficit in production nitric oxide (NO), identified by media assay and immunocytochemistry, at 6, 12, 18, and 48 hours post-injury. Furthermore, gene expression analysis revealed the expression of inflammatory cytokines IL-6 and IL-10 and enzyme arginase-1 was significantly down-regulated at 12 hours post-injury. Time course evaluation of migration, using a cell exclusion zone assay, showed stretch-injured cells display decreased migration into the exclusion zone at 48 and 72 hours post-stretch. Lastly, coinciding with the functional immune deficits, was a significant change in morphology, with process length decreasing and cell diameter increasing following an injury at 12 hours.

Conclusions: Taken together, the data demonstrate that stretch-injury produces significant alterations in microglial function, which may have marked impact on their response to injury or their interaction with other cells.

Background

Traumatic brain injury (TBI) is a growing severe public and military health concern, with 1.7 million cases accruing an economic burden near 76.8 billion annually (1). TBI is divided into two injury phases, primary and secondary (2). The primary injury results from the impact or acceleration events generating mechanical forces that act upon the sensitive brain tissue, resulting in immediate, irreparable damage and dysfunction. Secondary injury processes resulting from the initial injury evolve over a period of time and contribute to the destruction of brain tissue. A prominent secondary injury process is the inflammatory response, primarily mediated by activated microglia, the brain's immunocompetent cell (3). The initial immune response is thought to be a protective and restorative process; however, there are multiple experimental models (4–7) and human (8, 9) cases of TBI that implicate prolonged microglia activation in deleterious outcomes. Currently, the mechanism causing prolonged microglial activation is unknown.

Mechanical strain produced from the primary injury generates shearing forces that stretch and deform the cytoarchitecture and cells, contributing to death and dysfunction. Stretch-injury *in vitro* models allows

for the study of the effects of primary injury forces on individual cells. To date, *in vitro* models of neurotrauma show neuronal (10, 11), astrocyte (12), and endothelial (13) pathologies that mimic those injuries observed in *in vivo* TBI models. For example, rat primary cortical neurons exposed to a uniaxial stretch-injury with a strain of 10% displayed focal axonal swellings, similar to swellings observed in diffuse axonal injury (DAI) in experimental models and human cases of TBI (10). Human astrocytes exposed to a biaxial stretch-injury with a strain of 20% displayed retracted end-feet processes and activation c-Jun NH₂ –terminal kinase (JNK), replicating astrocyte associated dysfunction in blood-brain barrier (BBB) integrity. Furthermore, murine endothelial cells exposed to a stretch-injury with a 20% strain displayed markedly retracted, swollen and deformed morphology with notable intercellular spaces between cells (13). To date, the effects of stretch-injury on microglia viability, phenotype, and function are currently unknown.

Following the initial trauma, microglia interact with trauma associated-factors and become activated; this transformative process is necessary to mediate an inflammatory response. This is characterized by a morphological and phenotypic transformation known as activation (14). The morphological changes involve a transition from a ramified to amoeboid shape, and functional changes such as up-regulation of nitric oxide (NO) (3) and reactive oxygen species (ROS) production (14), the release of chemokine/cytokines (15), alteration in cell surface immune receptors and functional enzymes, and an increase in phagocytic and migratory behavior (16). Persistent, or chronic, activation of microglia, following a TBI, can occur if the initial response is left unresolved, and this can lead to functional deficits and neurodegeneration (17). To date, the mechanism(s) underlying chronic post-traumatic microglia activation remain elusive, and given that the primary insult can induce cellular pathologies in other cell types, an investigation into the effects of primary forces on microglia function is warranted.

In this study, we aimed to evaluate the impact of stretch-injury on microglial viability, function, and morphology. Additionally, the effect of stretch-injury on microglial priming was examined by examining microglia viability and function after microglia were treated with a known inflammogen after a stretch-injury. First, time-course studies were conducted using media assays to assess viability (LDH) and function (NO). From this, we focused on a single post-injury time point to conduct an evaluation of inflammatory associated cytokines and enzymes, reactive oxygen species and morphology analysis. Lastly, a time-course study assessed the priming effect of stretch-injury on microglial viability and function. This study is the first to comprehensively study the effect of stretch-injury on microglial viability, morphology, and function.

Methods

Cell culture

Primary microglia cultures were isolated from 2-day-old Sprague-Dawley rat pups by shaking mixed glia cultures at 100 RPM for 1 hour at 37 °C yielding 90% purity as previously described (18). Cells were maintained at 37 °C and a humidified atmosphere augmented with 5% CO₂ in media containing

Dulbecco's modified Eagle medium (Gibco), 1% l-glutamine (Gibco), 1% sodium pyruvate (Gibco), 10% FBS (Hyclone), and 1% penicillin and streptomycin (Thermo Fisher Scientific). Media replacement occurred every 3–5 days.

Stretch-injury

Cells were seeded at a density of 3×10^5 cells/mL on Pronectin coated silicone-based deformable membrane BIOFLEX® culture plates (BF-3001P Flexcell International, Hillsborough, NC). Cells were incubated for 24 hours in growth media to allow for adherence (37°C, 5% CO₂), after which a stretch-injury was administered, as previously described (19). Briefly, a stretch-injury was produced using the Cell injury controller II (CIC II), which utilizes nitrogen gas to pressurize a BIOFLEX® well individually. The CIC II regulates the duration and pressure of gas delivered to the well by using a rubber tube that feeds through a plastic circular well adaptor plug that attaches to the top of the well, creating a hermetically sealed system. Upon discharging the gas, the well becomes pressurized, and a bi-axial deformation of the flexible membrane, resulting in the cultured cells experiencing a stretch-injury. All stretch-injury experiments used identical parameters with 60psi regulator pressure, for a duration of 50 mSec, resulting in a 5psi pressurization of the well that created a 20% stretch event in the center of the well.

Lipopolysaccharide (LPS) and Etoposide treatment

To evaluate the potential priming effect of stretch-injury, primary microglia were treated with 100ng/mL concentration of LPS (Sigma, L8274) 15 minutes after stretch-injury(3). Additionally, an LPS treated-only group served as an activation control group. A dose of 15uM of Etoposide (catalog no. E1383; Sigma) was used to induce apoptosis and elicit a robust cleaved caspase-3 response to serve as a positive control (20).

Reactive oxygen species (ROS) assays

To detect ROS intermediates, cells were treated with 10 uM concentration of 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA, Thermo Fisher); a separate experiment was conducted to evaluate ROS intermediates at 12 hours post-injury. Before treatment H₂DCFDA, cells were washed with warm PBS, then cells were incubated for 45 min with the assay components, and fluorescence emission was measured on a FLUOstar (BMG LABTECH Inc., Cary, NC) plate reader.

Cell death assays

Cell culture media was assessed for lactate dehydrogenase (LDH) at 1, 6, 12, 18, 24, 48, and 72 hours post-injury using the Cytotox 96 assay (Promega) according to the manufacturer's instructions. Absorbance at 496nm was measured using a ChroMate microplate reader (Awareness Technology Inc., Palm City, FL).

Nitric oxide (NO) measurement

NO was measured in primary microglia exposed to stretch-injury at 1, 6, 12, 18, 24, 48, and 72 hours post-injury. NO content in the media was quantified using a Griess reagent assay kit (Invitrogen). The assay was performed per manufacturer's recommendations and as previously described (21). Colorimetric changes in 96-well plates were quantified with a Chroma Plate Reader (Midwest Scientific, St. Louis, MO, USA) at 545 nm.

Immunocytochemistry

Cells were fixed with 4% paraformaldehyde and rinsed twice with 1× PBS. Primary antibodies included rabbit anti-arginase-1 (20 µl/ml; Abcam), rabbit anti-NOS (10 µl/ml; Abcam, Cambridge, MA) and rabbit anti-cleaved caspase-3 (10 µl/ml; Abcam). Also, cell nuclei were stained with DAPI (0.5 µl/ml; Life Technologies, Carlsbad, CA). Fluorescently tagged secondary antibodies (Invitrogen) were visualized with an Olympus BX43 fluorescent microscope with a CellSens Standard imaging program. Five randomly selected images were taken per sample, and each image was qualitatively assessed for pixel density and normalized to the total number of DAPI stained cells. Images were taken from the center of the membrane.

RNA isolation and gene analysis

TRIzol® (Invitrogen, Carlsbad, CA) was used to extract mRNA from the cells, and mRNA concentrations were measured by a Nanodrop system (Thermo Scientific). A Veriti thermal cycler (Applied Biosciences) and a complementary DNA (cDNA) conversion kit (Applied Biosciences, Waltham, MA) was used to convert 1 µg of mRNA into cDNA, according to the manufacturer's suggested protocol. All quantitative real-time PCR (qRT-PCR) data was procured through the StepOnePlus Real-Time PCR System (Applied Biosciences), according to the manufacturer's recommended protocol. All primers were designed by Sigma-Aldrich (Darmstadt, Germany) and listed in Table 1.

Migration assay and morphological evaluation

A cellular exclusion zone assay was used to quantify migration. Cells were seeded at a density of 3×10^5 /mL on Pronectin coated BioFlex well plates with a 1.4 cm x .1 cm strip of silicone membrane placed in the center of the well and allowed to attach for 24 hours. A graded grid was drawn onto the underside of the membrane as a reference. Microphotographs were taken using a phase-contrast microscope (Zeiss, PrimoVert) with a USB adapted ocular (Swiftcam 5.0-megapixel digital camera microscopes, Amazon.com) for capturing images. Images were captured at pre-injury baseline and 12, 24, 48, and 72 hours post-injury. ImageJ (NIH, open-access software) was used to quantify cell exclusion zone diameter. Twenty-four hours after plating, the barrier strip of silicone was removed, and a baseline image was taken to define the cell exclusion zone. Then a stretch-injury was administered, and cells were returned to the incubator. At pre-stretch, 12, 24, 48, and 72 hours post-injury time points, cells were

counted in the exclusion zone. Cells were counted if they matched criteria, 1) dark coloring, 2) circular or branched phenotype. In some experiments, cells were able to attach underneath the silicone barrier in the exclusion zone. These cells were counted and subtracted from the total number of migrating cells in subsequent post-injury assessment time points, 12, 24, 48, and 72 hours.

Morphology was evaluated using the same devices, and images were taken at pre-injury and twelve hours post-injury. ImageJ (NIH, open-access software) was used to quantify cellular characteristics, i.e., number and length of processes and cell diameter.

Phagocytosis assay

At 12 hours after stretch-injury, cells were assessed for phagocytic activity using the Phagocytosis Assay Kit (Cayman Chemical, Ann Arbor, MI). Briefly, at 8 hours after injury, latex beads with rabbit IgG-FITC conjugates (1:500) were incubated with control and injured cells for 4 hours, followed by a 1-minute incubation with trypan blue to quench non-phagocytosed bead fluorescence and cell fixation with 4% paraformaldehyde. Cells were counterstained with DAPI, and five images at 20X magnification were collected from each coverslip. The number of DAPI cells containing FITC labeled beads were counted and normalized to the total number of DAPI. This assay was repeated in triplicate.

Statistical analysis

Statistical analysis was conducted using Graphpad Prism Software version 6.01. (GraphPad Software, San Diego, CA). NO, and LDH assays were performed in triplicate for each trial, and the trials were replicated four times to generate a sample size of $n = 4$ per group. A two way ANOVA with Sidak's multiple comparisons test was conducted on the groups which controls were set to 1.0. The stretch-injured values were normalized to control values for each trial for NO and LDH assay measures. ROS assays were conducted singly for each trial, and the trials were replicated three times to generate a sample size of $n=3$. A one way ANOVA with Tukey's multiple comparisons test was conducted. Western blot analysis was performed in pooled samples from triplicates for each trial, and the trials were replicated three times to generate a sample size of $n=3$. Immunocytochemistry experiments were conducted singly and repeated three times to create a sample size of $n=3$. Western blot and immunocytochemistry results were analyzed using an unpaired student's t-test. The migration assay was conducted singly for each trial and replicated three times for an $n=3$. The total number of migrating injured cells was normalized to control cells at each time point. A two way ANOVA with Sidak's multiple comparisons test was conducted for assay measures. The morphology analysis was assessed from five images per trial, and each trial was replicated three times to generate a sample size of $n=3$. A total of 15 images were analyzed for the control and stretch-injured condition. Percent averages for each group were assessed using a two-way ANOVA with Sidak's multiple comparisons. For all statistical tests described, a p -value < 0.05 was considered statistically significant. Data are presented as mean $\pm$ standard deviation (SD).

Results

Stretch-injury does not increase LDH at multiple time points or cleaved caspase-3 at 12 hours post-injury

A time-course study of media-derived LDH was conducted to determine the effect of stretch-injury on microglial viability and membrane permeability at acute and long-term time points. LDH is typically confined in the cytosol and is detectable in the event of cell lysis, necrosis, or compromised membrane permeability. Following a 20% stretch-injury, media was assayed for LDH at 1, 6, 12, 18, 24, 48, and 72 hours post-injury and showed no significant differences in media LDH content between stretch-injured and control cells at any time point (Fig. 1A). To further assess microglia viability, an immunocytochemistry evaluation of cleaved caspase-3, a marker of apoptosis, was conducted on fixed cells at the 12 hours post-injury time-point and revealed no significant difference between injured and non-injured cells (Fig. 1B). The observation that LDH media levels were unchanged between controls and injured cells suggests that a 20% stretch-injury does not alter microglia viability or membrane permeability. Furthermore, there was no significant difference between control and injured cells of cleaved caspase-3 expression, a marker of apoptosis, demonstrating that 20% stretch-injury does not induce apoptosis. Taken together, these results suggest that a 20% stretch-injury does not significantly affect microglia viability or membrane permeability at 12 hours post-injury.

NO release and iNOS expression is significantly down-regulated following stretch-injury

To examine the acute and long-term effect of stretch-injury on microglia function, a time course study, using assessment of NO in the media, was conducted at 1, 6, 12, 18, 24, 48, and 72 hours post-injury. Stretch-injury significantly down-regulated release of NO at 6, 12, 18, and 48 hours post-injury (Fig. 2A; 6, 12, and 18 hours $p < 0.05$, and 48 hours $p < 0.01$, two way ANOVA, Sidak's multiple comparisons post-test). Immunocytochemistry evaluated the primary enzyme responsible for catalyzing NO production, iNOS, at 12 hours post-injury and showed that stretch-injury significantly decreased iNOS expression compared to non-stretched control cells (Fig. 2B; $p < 0.01$ unpaired student's t-test). Compared to controls, stretch-injury did not alter gene expression of iNOS at 12 hours post-injury. The observed decreases in NO release and iNOS protein expression following a 20% stretch-injury suggest that a mechanical stimulus can reduce microglia function and that this reduction is not related to a decrease in cellular viability.

Oxidative Stress Is Unaffected By Stretch-injury

ROS production was assessed at 12 and 24 hours post-stretch to determine if this microglial activation product and subsequent oxidative stress aligned with NO release. Evaluation of ROS at 12 and 24 hours post-injury showed that there is no significant effect of stretch-injury on ROS production at either time point compared to control cells (Fig. 3A). There was no effect of stretch-injury on microglia-mediated oxidative stress, evaluated using western blot analysis for 3-NT, at 12 hours post-injury (Fig. 3B). To further investigate ROS production, the expression of the NOX2 membrane-bound enzyme component, gp91, was evaluated at 12 hours post-injury using western blot, and its expression, compared to controls

cells, was not significantly altered (Fig. 3C). The data shows that ROS production and oxidative stress do not track with decreases in NO release.

Genes associated with microglia activation are significantly down-regulated or unaffected following exposure to stretch-injury

The expression of genes associated with microglial activation in TBI was evaluated using RT-PCR at 12 hours post-injury. The 12 hour time point was chosen based on preliminary data identifying it as the first time point after post-injury to display a significant decrease in microglia function, NO release. Pro-inflammatory associated genes, IL-6 (4), CD86 (22), and IL-1 β (4), were selected for their observed up-regulation in activated microglia in the secondary injury cascade in experimental models of TBI. Anti-inflammatory associated genes were chosen for their microglia-associated roles in TBI: immunomodulation (IL-10) (23), tissue repair (arginase-1) (24), homeostatic maintenance (TGF β) (25), and pathogen recognition and capture (CD206) (2). Pro-inflammatory associated cytokine gene IL-6 was significantly down-regulated compared to controls, 12 hours after stretch-injury (Fig. 4E; $p < 0.05$ unpaired student's t-test). Interestingly, the anti-inflammatory associated gene coding for the enzyme arginase-1 was also significantly down-regulated at the 12 hour post-injury time point (Fig. 4A $p < 0.05$ unpaired student's t-test). Immunocytochemistry analysis of arginase-1 confirmed the genetic findings (Fig. 4I; $p < 0.05$ unpaired student's t-test). There was no significant difference in gene expression of pro-inflammatory associated genes, CD86 and IL-1 β (Fig. 4F-G) or anti-inflammatory associated genes CD206, IL-10, and TGF β (Fig. 4B-D) between injured and control cells at 12 hours post-injury. Interestingly, both pro and anti-inflammatory genes were significantly down-regulated at 12 hours post-injury, suggesting suppression of basal level gene expression

Morphology Is Significantly Altered In Response To Stretch-injury

Following a TBI, microglia undergo a morphological transformation from a small, ramified, cell body to a broader, rounded amoeboid phenotype. To determine the effect of mechanical strain on morphology, phase-contrast microscopy images were taken from the center of the well twelve hours post-injury. The cell count, cell body area, number, and length of processes were evaluated using Image J analysis software. Following stretch-injury, microglia display a significant increase in their cell body area, which is defined by the total area of the cell except for their processes, compared to controls at twelve hours after injury (Fig. 5D $p < .01$ unpaired student's t-test). Furthermore, stretch-injured cell processes length is significantly decreased compared to control cells twelve hours following injury. (Fig. 5E $p < .05$ unpaired student's t-test)

Migration Is Impaired Substantially Following Stretch-injury

In response to injury, microglia localize to the site to help contain the damage and mediate the immune response. To assess this behavior following injury, we developed and conducted a longitudinal examination of migratory behavior using a cell exclusion zone assay. The results showed significantly fewer cells migrated into the exclusion zone in the injured condition at forty-eight and seventy-two hours after injury than control cells (Fig. 6 $p < .001$ and $p < .0001$ two way ANOVA with Sidak's multiple comparisons).

Phagocytosis Is Not Affected By Injury

To assess the phagocytic activity of microglial cells in response to injury, primary cells were exposed to a stretch-injury, and eight hours after the measurement of phagocytosis of FITC-IgG conjugated latex beads was assessed. Interestingly, there was no difference between bead uptake between the control and injured cells, demonstrating that stretch does not affect phagocytosis. An LPS treated (100 ng/mL) was included as a positive control. Briefly, cells were incubated with LPS for 44 hours and treated with identical conditions as the control and injured cells. The LPS treated group displayed a robust trend towards increased bead uptake (Fig. 7. $p = .0501$ One-way ANOVA with multiple comparisons) compared to controls.

Stretch-injury does not prime or depress microglial NO release or alter LDH release when exposed to a subsequent LPS challenge

To evaluate the potential priming effect of stretch-injury, primary microglia were exposed to a stretch-injury then treated with LPS (100 ng/mL) fifteen minutes after the injury. Following treatment, cells were assayed for viability and function using LDH and NO media assays, respectively. Compared to LPS only treated cells, stretch-injury did not significantly alter viability when subsequently treated with LPS at 24, 48, or 72 hours post-injury (Fig. 8A). There was a significant increase in media LDH in both the LPS treated only and LPS plus stretch-injury at 24, 48 and 72 (Fig. 8A $p < .001$, $p < .001$ and $p < .05$) hours compared to control cells. Stretch-injury does not prime or depress NO release when treated with LPS fifteen minutes after injury compared to LPS treated only cells at 24, 48, or 72 hours post-injury (Fig. 8B). Cells that were exposed to a stretch-injury plus LPS and LPS treated only displayed a significant increase in NO release at 48 (Fig. 8B $p < .0001$ One-way ANOVA with Dunnett's post-test) and 72 ($p < .01$ One-way ANOVA with Dunnett's post-test) hours after LPS treatment. Surprisingly, stretch-injury antecedent to LPS treatment did not affect NO release at any post-injury time point, despite the finding that stretch-injury decreases NO release at 48 hours. However, several factors may have contributed to this observation, such as the potency of LPS, schedule of subsequent treatment, and outcome measure of microglia function.

Discussion

The findings presented here address a significant gap in the microglia literature in the context of TBI, specifically the consequences of the primary-injury on microglia viability, morphology, and function. We

demonstrate that rat primary microglia remain viable while displaying significant functional and morphological perturbations in response to a stretch-injury with a 20% strain magnitude.

The longitudinal assessment of media LDH from stretch-injured cells showed no significant changes at any post-injury time point examined compared to controls, demonstrating that stretch-injury does not result in altered membrane integrity or necrosis. Media levels of the intracellular enzyme LDH are an indicator of compromised membrane integrity and necrosis (26, 27). For example, stretch-injured rat cortical astrocytes exposed to injury parameters similar to those used in this study exhibit increased media LDH up to twenty-four hours after injury, suggestive of injury-induced membrane permeability (27). Additionally, stretch-injury does not induce apoptosis, as indicated by no changes in cleaved caspase-3 expression between injured and control cells. These findings are counter to what is found in stretch-injured astrocytes (12) and neurons (28). They can be attributed to differences in cellular size, morphology, and functional roles in the brain. Astrocytes are significantly larger than microglia and, as a consequence, have more surface area exposed to the strain forces, leading to increased membrane alterations (29). Similar to astrocytes, neurons vary significantly in cell diameter and morphology compared to microglia, with a larger cell body and long, delicate axons that are susceptible to damaging strain forces (10). Furthermore, microglia may be less sensitive to injurious strain forces than astrocytes and neurons because they are constantly surveying the brain through migratory and motile behavior. In contrast, astrocytes and neurons are adherents, stationary and non-migratory in non-pathological conditions, rendering them more susceptible to mechanical deformations. Taken together, these results suggest that stretch-injury does not alter viability or membrane permeability.

Stretch-injury significantly downregulated iNOS expression and decreased media levels of NO. These findings contrast to what is observed in patients and experimental models of TBI. They can be attributed to fundamental modeling differences and may indicate an innate response of microglia to the injury itself. To assess the effects of primary injurious forces on microglia, mono cultured are exposed to damage in a sterile environment, free from other damaged cells and damage-associated molecular patterns (DAMPS). Because of this unique environment, microglia activation and subsequent up-regulation of iNOS and NO production are not observed. Microglia upregulate the immune-response enzyme iNOS increasing the production of NO, an inflammatory mediator. Multiple experimental models of TBI (30–33) show a significant upregulation of iNOS following injury. Continuous production of NO, through increased iNOS expression and activity, can lead to cytotoxic oxidants, such as peroxynitrite, that contribute to oxidative stress (34–36). Multiple aspects of the process involved in NO production were evaluated in the current study. Media NO levels were significantly down-regulated at six, twelve, eighteen, and forty-eight hours after injury compared to uninjured cells.

Consistent with the decreased NO in the media, the expression of iNOS protein was significantly down-regulated twelve hours following injury. However, mRNA transcripts were unaffected by injury at the same time points compared to non-stretched cells. Interestingly, a rat model using a controlled cortical impact (CCI) showed that tissue NO is significantly decreased at two, six, and twelve hours following injury near the injury site. However, opposite to our iNOS findings, there was a significant increase in NADPH-

diaphorase-positive cells, indicative of NOS activity, in the same regions where the NO decrease was detected. The authors attributed the reduction in NO to it being converted into peroxynitrite {Ziaja, 2007 #76}. Additionally, consistent with the findings of the *in vivo* study, we observed a significant decrease in media NO levels following injury; however, in contrast to the animal model, we found that the protein expression of iNOS is significantly decreased. This can be explained by the context of the *in vitro* environment, which lacks damaged associated factors that lead to microglial activation and increase iNOS expression. Future studies will evaluate the mechanism underlying the marked reduction in iNOS expression and determine if stretch-injury alters NO production in activated microglia. Our findings show that a stretch-injury suppresses media NO levels and down-regulates iNOS protein expression, suggesting that basal level NO production is affected by stretch-injury

The finding that stretch alters media NO and iNOS expression, but not iNOS mRNA transcription, is puzzling. A possible explanation could be that stretch-injury disrupts protein synthesis by inducing endoplasmic reticulum stress, which has been observed in blast (37) and fluid percussion injury (38) animal models of TBI. Additionally, rats exposed to a CCI model of TBI showed differentially methylated mRNA transcripts (39), a prominent post-transcriptional modifier of RNA, and has been shown to inhibit translation (40). Collectively, the data indicate that stretch-injury significantly down-regulated NO media production at multiple time points, and that this is mediated by a significant decrease in iNOS expression, which demonstrates that stretch-injury produces functional deficits in immune function.

Experimental models of TBI show that chronically activated microglia contribute to neuronal damage and dysfunction by producing free radical oxygen species that lead to cytotoxic oxidative stress (36, 41, 42). Following activation, microglia upregulate NOX2 and increase the production of ROS, that when combined with RNS, can form cytotoxic factors, such as peroxynitrite, that lead to protein nitrosylation resulting in cellular dysfunction or death. To evaluate the effect of stretch-injury on microglial ROS generation and oxidative stress, assessments of ROS, oxidative stress and the NOX2 enzyme were performed. Production of ROS was conducted by evaluating media and cellular levels twelve hours after injury. Interestingly, there was no effect of stretch-injury on ROS production compared to control cells. Next, western blot analysis showed that the expression of gp91, the membrane-bound component of the NOX2 complex, was unaffected by stretch-injury. Lastly, oxidative stress was evaluated by western blot for 3-NT, a marker of protein nitrosylation, specifically tyrosine residues. Consistent with the other ROS-related outcome measures, stretch-injury did not affect the oxidative stress response; specifically, protein nitrosylation, indicated by no changes between control and injured cells using western blot analysis of 3-NT. These results are surprising, given that multiple lines of research show both NO and ROS production are increased following microglial activation (43–45). Contrary to the RNS data, ROS does not appear to be affected by stretch-injury.

Following a TBI, activated microglia release both pro and anti-inflammatory associated cytokines (46). Multiple converging lines of research show that the post-insult inflammatory response in the brain presents with a pro and anti-inflammatory phenotype, and this duality varies temporally. For example, a rodent model of diffuse brain injury, where strain-induced injuries are prevalent, showed an increase in

both pro (IL-1 β , CD14 and iNOS) and anti (arginase-1) inflammatory mRNA transcripts at twenty-four hours post-injury (47). The effect(s) of the primary forces on the expression of microglial inflammatory-related genes is undefined. To address this gap in the literature, mRNA transcripts of inflammatory-associated genes were evaluated using RT-PCR at twelve hours post-stretch-injury. Interestingly, we found that both pro-inflammatory (IL-6) and anti-inflammatory (IL-10 and arginase1) associated genes were down-regulated following injury. This is counter to what is observed in experimental models and patients with TBI, where these genes are commonly upregulated. For example, plasma levels of IL-6 are significantly elevated acutely in patients who sustain a mTBI (48). In a rat model of experimental TBI, there was an immediate increase in brain IL-10 mRNA transcripts, peaking at four hours post-injury and continuing to rise until 20 hours post injury (49). In a mouse model of TBI, arginase-1 mRNA transcripts were significantly upregulated twenty-four hours following injury in both young (three-month old) and old (24-months old) animals (50). However, an important distinction between our findings and these in vivo observations is that, in our study, 1) only microglia are exposed to the stretch-injury, and 2) they do not interact with DAMPs or damaged cells and do not appear to be activated, either morphologically or functionally. Therefore, the upregulation of inflammatory genes would not be expected. Stretch-injury suppresses multiple genes associated with the inflammatory response following injury; the consequence and long-term ramification of this effect on microglial inflammatory response is an important question since dysfunctional inflammatory responses are observed in the chronically injured brain.

Stretch-injury decreases basal level microglial migration. Following the initial injury, microglia sense factors from the trauma and migrate to the injury site to remove debris and damaged cells (51). To date, the effect of stretch-injury on migration in the biomechanically injured brain is unexplored in the literature. To address this gap in the literature, we assessed migration following stretch-injury by using a cell exclusion assay and found that stretch-injury significantly reduced the number of cells that migrated into the exclusion zone at forty-eight and seventy-two hours post-injury. These findings show that a mechanical stimulus can reduce an essential functional behavior integral to the response to injury and necessary for homeostatic function. Future studies will elucidate the extent these migratory deficits play in a physiologically relevant setting.

Microglial morphology is significantly altered following exposure to stretch-injury. Following the primary injury, microglia undergo a morphological transformation characterized by a significant increase in soma diameter and reduction of processes to take on an amoeboid phenotype (52). Phase-contrast imaging showed that stretch-injury significantly reduced process length twelve hours after injury, demonstrating that a deformation event alone can rearrange the cytoskeleton. This is consistent with morphological alteration in neurons (53, 54), astrocytes (55), and endothelial (56) cells exposed to stretch-injury. The stretch-induced reduction in process length may contribute to the decreased number of cells migrated into the exclusion zone. Since migration involves remodeling of the cytoskeleton, one explanation could be that stretch-injury alters the cytoskeleton and impairs its remodeling during migration.

The microglial inflammatory response is not altered following treatment with LPS post-stretch-injury. To extend our model into a more physiologically relevant setting, where to mechanical forces are followed by

exposure to inflammatory challenges, stretch-injured microglia were treated with LPS, a potent microglial activator, fifteen minutes following injury. Then media levels of NO and LDH were assessed at twenty-four, forty-eight, and seventy-two hours post-treatment to determine viability and activation. Surprisingly, the suppressive effect of stretch-injury on NO media levels was not observed in the post-stretch, LPS treated cells. Consistent with the LDH results from the time course assessment, LDH media levels were unchanged between any groups. The unexpected NO media result could be attributable to multiple issues. The first could be the potency of LPS as an inflammatory challenge.

Given that it is an endotoxin found on the outer membrane of gram-negative bacteria, it is expected that elicit a more robust inflammatory response than factors that are released during a TBI. For example, primary rat microglia treated with interferon- γ (IFN γ) and tumor necrosis factor- α (TNF α), factors associated with brain trauma, display a varied inflammatory response compared to LPS. Compared to the IFN γ and TNF α treatment, LPS treated cells expressed significantly more NOS and arginase-1 mRNA transcripts and protein expression. Additionally, media NO levels were also considerably higher. The second issue could be the timing of the challenge post-injury. The effects of stretch-injury may not have enough time to alter the response to LPS. Future studies will challenge stretch-injured cells at immediate and delayed time points. These results suggest that stretch-injury does not change microglial inflammatory response when subsequently challenged; however, as noted above, multiple factors could account for these findings.

Phagocytosis is unaffected following stretch-injury. Multiple studies show that activated microglia significantly increase their phagocytic function compared to their resting cells {Fan, 2018 #78}{Hornik, 2016 #80}{Yi, 2020 #81}. A critical role of activated microglia in the post-injured brain is phagocytosing debris and dead and damaged cells {Loane, 2016 #5}. The assessment of phagocytosis on stretch-injured cells showed no difference in fluorescently label conjugated beads uptake between the control and stretch-injured group, twelve hours post-injury. Given that basal level phagocytosis in resting microglia is already low, it was not expected to see an effect of a stretch injury. An interesting observation was the relatively low number of phagocytosed beads in the non-activated cells cultured on Pronectin. This finding is consistent with a study showing that when primary rat microglia are incubated with fibronectin, there is a reduction in phagocytosis, as measure by the engulfment of zymogen particles. Furthermore, the same study showed that morphological phenotypes were consistent with our control cells, with cells displaying a reduced cell body diameter and an increased number of processes {Chamak, 1991 #79}.

Conclusions

The results presented here are the first to examine the consequences of the primary injury on microglia by characterizing the effects of stretch-injury on viability, morphology and function. The data shows that microglia remain viable after the injury and display morphological alterations, with decreased process length and functional deficits, with reduced migration and nitric oxide production, at twelve hours after injury. Additionally, stretch-injury significantly down-regulates inflammatory associated gene transcripts

at 12 hours. These findings will contribute to the understanding of microglia in the context of TBI, and future studies will elucidate the molecular mechanism(s) that are responsible for these deficits. Furthermore, studies will examine the contributions of these effects of dysfunctional microglia response observed in experimental models and individuals with TBI.

Abbreviations

TBI

Traumatic brain injury

CIC II

Cell injury controller II

NO

Nitric oxide

LDH

Lactate dehydrogenase

DAI

Diffuse axonal injury

JNK

c-Jun NH² terminal kinase

BBB

Blood brain barrier

ROS

Reactive oxygen species

LPS

Lipopolysaccharide

Declarations

Ethics approval and consent to participate:

Not applicable

Consent for publication

Not applicable

Availability of data and material:

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interest:

The authors declare that they have no competing interest

Funding:

Funding provided by the Center for the study of traumatic stress (CSTS) at USUHS.

Author's contribution:

MC designed and performed experiments, analyzed data and drafted the manuscript. KB co-conceived the study, participated in its design and coordination and helped draft the manuscript. All authors have read and approved the final version of the manuscript.

Acknowledgements:

Not applicable

References

1. Byrnes KR, Wilson CM, Brabazon F, von Leden R, Jurgens JS, Oakes TR, et al. FDG-PET imaging in mild traumatic brain injury: a critical review. *Front Neuroenergetics*. 2014;5:13.
2. Kumar A, Alvarez-Croda D-M, Stoica BA, Faden AI, Loane DJ. Microglial/Macrophage Polarization Dynamics following Traumatic Brain Injury. *J Neurotrauma*. 2016;33(19):1732–50.
3. Brabazon F, Bermudez S, Shaughness M, Khayrullina G, Byrnes KR. The effects of insulin on the inflammatory activity of BV2 microglia. *PLoS One*. 2018;13(8):e0201878.
4. Kumar A, Stoica BA, Loane DJ, Yang M, Abulwerdi G, Khan N, et al. Microglial-derived microparticles mediate neuroinflammation after traumatic brain injury. *J Neuroinflammation*. 2017;14(1):47-.
5. Henry RJ, Ritzel RM, Barrett JP, Doran SJ, Jiao Y, Leach JB, et al. Microglial Depletion with CSF1R Inhibitor During Chronic Phase of Experimental Traumatic Brain Injury Reduces Neurodegeneration and Neurological Deficits. *The Journal of Neuroscience*. 2020;40(14):2960–74.
6. Gao T, Chen Z, Chen H, Yuan H, Wang Y, Peng X, et al. Inhibition of HMGB1 mediates neuroprotection of traumatic brain injury by modulating the microglia/macrophage polarization. *Biochem Biophys Res Commun*. 2018;497(1):430–6.
7. Villapol S, Loane DJ, Burns MP. Sexual dimorphism in the inflammatory response to traumatic brain injury. *Glia*. 2017;65(9):1423–38.
8. Coughlin JM, Wang Y, Munro CA, Ma S, Yue C, Chen S, et al. Neuroinflammation and brain atrophy in former NFL players: An in vivo multimodal imaging pilot study. *Neurobiol Dis*. 2015;74:58–65.
9. Ramlackhansingh AF, Brooks DJ, Greenwood RJ, Bose SK, Turkheimer FE, Kinnunen KM, et al. Inflammation after trauma: microglial activation and traumatic brain injury. *Ann Neurol*. 2011;70(3):374–83.
10. Hemphill MA, Dabiri BE, Gabriele S, Kerscher L, Franck C, Goss JA, et al. A possible role for integrin signaling in diffuse axonal injury. *PLoS One*. 2011;6(7):e22899.

11. López-García I, Gerő D, Szczesny B, Szoleczky P, Olah G, Módis K, et al. Development of a stretch-induced neurotrauma model for medium-throughput screening in vitro: identification of rifampicin as a neuroprotectant. *Br J Pharmacol*. 2018;175(2):284–300.
12. Augustine C, Cepinskas G, Fraser DD. Traumatic injury elicits JNK-mediated human astrocyte retraction in vitro. *Neuroscience*. 2014;274:1–10.
13. Au -SE, Au - Neuhaus W. Au - Foerster C. Stretch in Brain Microvascular Endothelial Cells (cEND) as an In Vitro Traumatic Brain Injury Model of the Blood Brain Barrier. *JoVE*. 2013(80):e50928.
14. Yauger YJ, Bermudez S, Moritz KE, Glaser E, Stoica B, Byrnes KR. Iron accentuated reactive oxygen species release by NADPH oxidase in activated microglia contributes to oxidative stress in vitro. *J Neuroinflammation*. 2019;16(1):41.
15. Kumar A, Alvarez-Croda DM, Stoica BA, Faden AI, Loane DJ. Microglial/Macrophage Polarization Dynamics following Traumatic Brain Injury. *J Neurotrauma*. 2016;33(19):1732–50.
16. Qu WS, Liu JL, Li CY, Li X, Xie MJ, Wang W, et al. Rapidly activated epidermal growth factor receptor mediates lipopolysaccharide-triggered migration of microglia. *Neurochem Int*. 2015;90:85–92.
17. Faden AI, Wu J, Stoica BA, Loane DJ. Progressive inflammation-mediated neurodegeneration after traumatic brain or spinal cord injury. *Br J Pharmacol*. 2016;173(4):681–91.
18. Tamashiro TT, Dalgard CL, Byrnes KR. Primary microglia isolation from mixed glial cell cultures of neonatal rat brain tissue. *J Vis Exp*. 2012(66):e3814-e.
19. López-García I, Gerő D, Szczesny B, Szoleczky P, Olah G, Módis K, et al. Development of a stretch-induced neurotrauma model for medium-throughput screening in vitro: identification of rifampicin as a neuroprotectant. *Br J Pharmacol*. 2018;175(2):284–300.
20. Ciacci-Zanella J, Stone M, Henderson G, Jones C. The latency-related gene of bovine herpesvirus 1 inhibits programmed cell death. *J Virol*. 1999;73(12):9734–40.
21. Byrnes KR, Stoica B, Loane DJ, Riccio A, Davis MI, Faden AI. Metabotropic glutamate receptor 5 activation inhibits microglial associated inflammation and neurotoxicity. *Glia*. 2009;57(5):550–60.
22. Izzy S, Liu Q, Fang Z, Lule S, Wu L, Chung JY, et al. Time-Dependent Changes in Microglia Transcriptional Networks Following Traumatic Brain Injury. *Front Cell Neurosci*. 2019;13:307-.
23. Lobo-Silva D, Carriche GM, Castro AG, Roque S, Saraiva M. Balancing the immune response in the brain: IL-10 and its regulation. *J Neuroinflammation*. 2016;13(1):297-.
24. Loane DJ, Kumar A. Microglia in the TBI brain: The good, the bad, and the dysregulated. *Exp Neurol*. 2016;275:316–27.
25. Butovsky O, Jedrychowski MP, Moore CS, Cialic R, Lanser AJ, Gabriely G, et al. Identification of a unique TGF- β -dependent molecular and functional signature in microglia. *Nat Neurosci*. 2014;17(1):131–43.
26. Benson BA, Vercellotti GM, Dalmaso AP. IL-4 and IL-13 induce protection from complement and melittin in endothelial cells despite initial loss of cytoplasmic proteins: membrane resealing impairs

- quantifying cytotoxicity with the lactate dehydrogenase permeability assay. *Xenotransplantation*. 2015;22(4):295–301.
27. Ellis EF, McKinney JS, Willoughby KA, Liang S, Povlishock JT. A new model for rapid stretch-induced injury of cells in culture: characterization of the model using astrocytes. *J Neurotrauma*. 1995;12(3):325–39.
 28. Xu Z, Liu Y, Yang D, Yuan F, Ding J, Chen H, et al. Sesamin protects SH-SY5Y cells against mechanical stretch injury and promoting cell survival. *BMC Neurosci*. 2017;18(1):57-.
 29. Vainchtein ID, Molofsky AV. Astrocytes and Microglia: In Sickness and in Health. *Trends Neurosci*. 2020;43(3):144–54.
 30. Fenn AM, Skendelas JP, Moussa DN, Muccigrosso MM, Popovich PG, Lifshitz J, et al. Methylene blue attenuates traumatic brain injury-associated neuroinflammation and acute depressive-like behavior in mice. *J Neurotrauma*. 2015;32(2):127–38.
 31. Kumar A, Stoica BA, Sabirzhanov B, Burns MP, Faden AI, Loane DJ. Traumatic brain injury in aged animals increases lesion size and chronically alters microglial/macrophage classical and alternative activation states. *Neurobiol Aging*. 2013;34(5):1397–411.
 32. Kuwar R, Rolfe A, Di L, Xu H, He L, Jiang Y, et al. A novel small molecular NLRP3 inflammasome inhibitor alleviates neuroinflammatory response following traumatic brain injury. *J Neuroinflammation*. 2019;16(1):81.
 33. Kumar A, Henry RJ, Stoica BA, Loane DJ, Abulwerdi G, Bhat SA, et al. Neutral Sphingomyelinase Inhibition Alleviates LPS-Induced Microglia Activation and Neuroinflammation after Experimental Traumatic Brain Injury. *J Pharmacol Exp Ther*. 2019;368(3):338–52.
 34. Shin N, Kim HG, Shin HJ, Kim S, Kwon HH, Baek H, et al. Uncoupled Endothelial Nitric Oxide Synthase Enhances p-Tau in Chronic Traumatic Encephalopathy Mouse Model. *Antioxid Redox Signal*. 2019;30(13):1601–20.
 35. Günther M, Al Nimer F, Piehl F, Risling M, Mathiesen T. Susceptibility to Oxidative Stress Is Determined by Genetic Background in Neuronal Cell Cultures. *eNeuro*. 2018;5(2).
 36. Cooney SJ, Bermudez-Sabogal SL, Byrnes KR. Cellular and temporal expression of NADPH oxidase (NOX) isoforms after brain injury. *J Neuroinflammation*. 2013;10:155.
 37. Lucke-Wold BP, Logsdon AF, Turner RC, Huber JD, Rosen CL. Endoplasmic Reticulum Stress Modulation as a Target for Ameliorating Effects of Blast Induced Traumatic Brain Injury. *J Neurotrauma*. 2017;34(S1):62-s70.
 38. Tan HP, Guo Q, Hua G, Chen JX, Liang JC. Inhibition of endoplasmic reticulum stress alleviates secondary injury after traumatic brain injury. *Neural Regen Res*. 2018;13(5):827–36.
 39. Yu J, Zhang Y, Ma H, Zeng R, Liu R, Wang P, et al. Epitranscriptomic profiling of N6-methyladenosine-related RNA methylation in rat cerebral cortex following traumatic brain injury. *Mol Brain*. 2020;13(1):11.
 40. Castro A, Bartos D, Kutas M, Weiss G. A new microcolumn for steroid separation in radioimmunoassay. *Clin Biochem*. 1974;7(1):64–7.

41. Henry RJ, Ritzel RM, Barrett JP, Doran SJ, Jiao Y, Leach JB, et al. Microglial Depletion with CSF1R Inhibitor During Chronic Phase of Experimental Traumatic Brain Injury Reduces Neurodegeneration and Neurological Deficits. *J Neurosci*. 2020;40(14):2960–74.
42. Henry RJ, Doran SJ, Barrett JP, Meadows VE, Sabirzhanov B, Stoica BA, et al. Inhibition of miR-155 Limits Neuroinflammation and Improves Functional Recovery After Experimental Traumatic Brain Injury in Mice. *Neurotherapeutics*. 2019;16(1):216–30.
43. Brown GC. Mechanisms of inflammatory neurodegeneration: iNOS and NADPH oxidase. *Biochem Soc Trans*. 2007;35(Pt 5):1119–21.
44. Chhor V, Le Charpentier T, Lebon S, Oré MV, Celador IL, Josserand J, et al. Characterization of phenotype markers and neuronotoxic potential of polarised primary microglia in vitro. *Brain Behav Immun*. 2013;32:70–85.
45. Chuang DY, Simonyi A, Kotzbauer PT, Gu Z, Sun GY. Cytosolic phospholipase A2 plays a crucial role in ROS/NO signaling during microglial activation through the lipoxygenase pathway. *J Neuroinflammation*. 2015;12:199.
46. Dalgard CL, Cole JT, Kean WS, Lucky JJ, Sukumar G, McMullen DC, et al. The cytokine temporal profile in rat cortex after controlled cortical impact. *Front Mol Neurosci*. 2012;5:6.
47. Fenn AM, Skendelas JP, Moussa DN, Muccigrosso MM, Popovich PG, Lifshitz J, et al. Methylene blue attenuates traumatic brain injury-associated neuroinflammation and acute depressive-like behavior in mice. *J Neurotrauma*. 2015;32(2):127–38.
48. Edwards KA, Pattinson CL, Guedes VA, Peyer J, Moore C, Davis T, et al. Inflammatory Cytokines Associate With Neuroimaging After Acute Mild Traumatic Brain Injury. *Front Neurol*. 2020;11:348.
49. Kamm K, VanderKolk W, Lawrence C, Jonker M, Davis AT. The Effect of Traumatic Brain Injury Upon the Concentration and Expression of Interleukin-1 β and Interleukin-10 in the Rat. *Journal of Trauma and Acute Care Surgery*. 2006;60(1).
50. Kumar A, Stoica BA, Sabirzhanov B, Burns MP, Faden AI, Loane DJ. Traumatic brain injury in aged animals increases lesion size and chronically alters microglial/macrophage classical and alternative activation states. *Neurobiol Aging*. 2013;34(5):1397–411.
51. Herzog C, Pons Garcia L, Keatinge M, Greenald D, Moritz C, Peri F, et al. Rapid clearance of cellular debris by microglia limits secondary neuronal cell death after brain injury in vivo. *Development*. 2019;146(9).
52. Caplan HW, Cardenas F, Gudenkauf F, Zelnick P, Xue H, Cox CS, et al. Spatiotemporal Distribution of Microglia After Traumatic Brain Injury in Male Mice. *ASN Neuro*. 2020;12:1759091420911770.
53. Ji J, Tyurina YY, Tang M, Feng W, Stolz DB, Clark RSB, et al. Mitochondrial injury after mechanical stretch of cortical neurons in vitro: biomarkers of apoptosis and selective peroxidation of anionic phospholipids. *J Neurotrauma*. 2012;29(5):776–88.
54. Saykally JN, Hatic H, Keeley KL, Jain SC, Ravindranath V, Citron BA. Withania somnifera Extract Protects Model Neurons from In Vitro Traumatic Injury. *Cell Transplant*. 2017;26(7):1193–201.

55. Wanner IB. An in vitro trauma model to study rodent and human astrocyte reactivity. *Methods Mol Biol.* 2012;814:189–219.
56. Salvador E, Neuhaus W, Foerster C. Stretch in brain microvascular endothelial cells (cEND) as an in vitro traumatic brain injury model of the blood brain barrier. *J Vis Exp.* 2013(80):e50928-e.

Tables

Table 1. Selected gene sequences for RT-PCR

Gene	Symbol	Forward (5'-3')	Reverse (5'-3')
Transforming growth factor beta 1	TGF-β-1	GGAAATCAATGGGATCAGTC	CTGAAGCAGTAGTTGGTATC
Interleukin-1, beta	IL-1β	TAAGCCAACAAGTGGTATTC	AGGTATAGATTCTTCCCCTTG
Interleukin-4	IL-4	GAACCAGGTCACAGAAAAAG	GGGAAGTAAAATTTGCGAAG
Interleukin-6	IL-6	CAGAGTCATT CAGAGCAATAC	CTTTCAAGATGAGTTGGATGG
Interleukin-12	IL-12	CTTCTTTGATGATGACCCTG	AATAGCCATCAGCATGTTTC
Interleukin-10	IL-10	TCTCCCCTGTGAGAAATAAAAG	GAAAGGAAAGTTCCCAGATG
Cluster of differentiation 86	CD86	CACAAGCTTTGATAGGGATAAC	TTCTTGTTTCATTCTGAGCC
Mannose receptor	CD206	GGAGTTGTGGAGCAGATGGAAG	CTTGAATGGAAATGCACAGAC
Arginase-1	Arg1	GAAGGAAAGTTCCCAGATG	CTTGTCCACTTCAGTCATTG
Inducible nitric oxide synthase	iNOS	TTCCTCAGGCTTGGGTCTTGT	GAGGAACTGGGGGAAACCATT
Phosphoglycerate kinase 1	PGK1	GGAGATGTCTATGTCAATGATG	TTTAGCTCCTCCCAAGATAG

Figures

Figure 1

Stretch-injury does not affect primary microglia viability. A) Primary microglia exposed to a stretch-injury of 5psi show no significant difference in media LDH when compared to control cells. Injured condition normalized to controls (set as 1.0). B) Representative sample of one trial presented. Scale bar is 50µm. C) Immunocytochemistry evaluation of cleaved caspase-3 indicates no significant difference between injured and control cells. Points represent individual data points. N=3/group. Error bar represent $\pm$ SD.

Figure 1

Stretch-injury does not affect primary microglia viability. A) Primary microglia exposed to a stretch-injury of 5psi show no significant difference in media LDH when compared to control cells. Injured condition normalized to controls (set as 1.0). B) Representative sample of one trial presented. Scale bar is 50µm. C) Immunocytochemistry evaluation of cleaved caspase-3 indicates no significant difference between injured and control cells. Points represent individual data points. N=3/group. Error bar represent $\pm$ SD.

Figure 2

Stretch-injured primary microglia display decreased NO production and iNOS expression. A) Media assays show injured cells produce significantly less NO when compared to uninjured cells at 6, 12, 18, and 48 hours after injury. Injured condition normalized to controls set at 1.0. All groups were compared using two-way ANOVA with Sidak's post-test. * $p < 0.05$, ** $p < 0.01$. $N=3/\text{group}$. Error bars represent $\pm\text{SD}$. B) Stretch-injury does not significantly alter iNOS gene expression twelve hours post-injury. Points represent individual data points. C) Representative sample of one trial presented. Scale bar is $50\mu\text{m}$. D) Quantification of pixel density shows stretch-injury significantly down-regulates iNOS protein expression at 12 hours post-injury. $N = 3/\text{group}$. Points represent individual data points. Error bars represent $\pm\text{SD}$. Groups were compared using an unpaired t-test, ** $p < 0.01$.

Figure 2

Stretch-injured primary microglia display decreased NO production and iNOS expression. A) Media assays show injured cells produce significantly less NO when compared to uninjured cells at 6, 12, 18, and 48 hours after injury. Injured condition normalized to controls set at 1.0. All groups were compared using two-way ANOVA with Sidak's post-test. * $p < 0.05$, ** $p < 0.01$. $N=3/\text{group}$. Error bars represent $\pm\text{SD}$. B) Stretch-injury does not significantly alter iNOS gene expression twelve hours post-injury. Points represent individual data points. C) Representative sample of one trial presented. Scale bar is $50\mu\text{m}$. D) Quantification of pixel density shows stretch-injury significantly down-regulates iNOS protein expression at 12 hours post-injury. $N = 3/\text{group}$. Points represent individual data points. Error bars represent $\pm\text{SD}$. Groups were compared using an unpaired t-test, ** $p < 0.01$.

Figure 3

ROS production is not altered by stretch-injury. A) Cellular and media ROS production is not altered at 12 or 24 hours after stretch-injury. All groups were compared using One-way ANOVA. N=3/group. Experimental groups were normalized to control values. B) Oxidative stress, as measured by 3-NT, is not significantly altered at twelve hours following stretch-injury. C) Representative blots for 3NT. D) Stretch-injury does not alter expression of NOX2/gp91 twelve hours after injury. E) Representative blots for gp91. N=3/group. All groups were compared using unpaired t-test. Points represent individual data points. Error bar represent $\pm$ SD.

Figure 3

ROS production is not altered by stretch-injury. A) Cellular and media ROS production is not altered at 12 or 24 hours after stretch-injury. All groups were compared using One-way ANOVA. N=3/group. Experimental groups were normalized to control values. B) Oxidative stress, as measured by 3-NT, is not significantly altered at twelve hours following stretch-injury. C) Representative blots for 3NT. D) Stretch-injury does not alter expression of NOX2/gp91 twelve hours after injury. E) Representative blots for gp91. N=3/group. All groups were compared using unpaired t-test. Points represent individual data points. Error bar represent $\pm$ SD.

Figure 4

Injured cells display decreases in both pro and anti-inflammatory genes. Stretch-injury decreased expression of pro-inflammatory associated gene IL-6 (A) and reduced expression of anti-inflammatory associated gene ARG1 (B) and IL-10 (C) at 12 hours after injury. Gene expression of experimental groups normalized to controls set as 1.0. Stretch-injury did not alter anti-inflammatory genes TGFβ (D) and CD206 (E), or pro-inflammatory genes CD86 (F) and IL-1β (G). H) Representative sample of one trial presented. Scale bar is 50μm. I) Quantification of pixel density shows stretch-injury, compared to control cells, significantly down-regulates arginase-1 protein expression at 12 hours post-injury. N = 3/group. Points represent individual data points. Error bars represent ±SD. Groups were compared using an unpaired t-test, * p<.05, **p < 0.01.

Figure 4

Injured cells display decreases in both pro and anti-inflammatory genes. Stretch-injury decreased expression of pro-inflammatory associated gene IL-6 (A) and reduced expression of anti-inflammatory associated gene ARG1 (B) and IL-10 (C) at 12 hours after injury. Gene expression of experimental groups normalized to controls set as 1.0. Stretch-injury did not alter anti-inflammatory genes TGFβ (D) and CD206 (E), or pro-inflammatory genes CD86 (F) and IL-1β (G). H) Representative sample of one trial presented. Scale bar is 50μm. I) Quantification of pixel density shows stretch-injury, compared to control cells, significantly down-regulates arginase-1 protein expression at 12 hours post-injury. N = 3/group. Points represent individual data points. Error bars represent ±SD. Groups were compared using an unpaired t-test, * p<.05, **p < 0.01.

Figure 5

Stretch-injury significantly alters microglia morphology at 12 hours post-injury. A) Representative images of phase contrast micrographs twelve hours post-injury. Scale bar is set to 50µm. The total cell count (B) and average process number per cell (C) are unaffected following stretch-injury. D) Injured cells display a significantly larger cell body area compared to controls. E) The average process length is significantly decreased compared to controls. N=3/group. All groups were compared using unpaired t-test, * p<.05, **p<.01. Bars represent mean +/- SD.

Figure 5

Stretch-injury significantly alters microglia morphology at 12 hours post-injury. A) Representative images of phase contrast micrographs twelve hours post-injury. Scale bar is set to 50µm. The total cell count (B) and average process number per cell (C) are unaffected following stretch-injury. D) Injured cells display a significantly larger cell body area compared to controls. E) The average process length is significantly decreased compared to controls. N=3/group. All groups were compared using unpaired t-test, * p<.05, **p<.01. Bars represent mean +/- SD.

Figure 6

Migration is significantly decreased following stretch injury at 48 and 72 hours after injury. A) Representative phase contrast micrographs of the cell exclusion zone assay. B) Stretch-injury significantly decreases the number of cells that migrate into the exclusion zone at 48 and 72 hours post injury when compared to the control group. N=3/group. Two-way ANOVA with Sidak's multiple comparison test, *** $p < .001$, **** $p < .0001$. Bars represent mean $\pm$ SD

Figure 6

Migration is significantly decreased following stretch injury at 48 and 72 hours after injury. A) Representative phase contrast micrographs of the cell exclusion zone assay. B) Stretch-injury significantly decreases the number of cells that migrate into the exclusion zone at 48 and 72 hours post injury when compared to the control group. N=3/group. Two-way ANOVA with Sidak's multiple comparison test, *** $p < .001$, **** $p < .0001$. Bars represent mean $\pm$ SD

Figure 7

Stretch-injury does not alter phagocytic activity. A) Representative images of latex beads conjugated with FITC-IgG (green) were incubated for 4 hours with control, stretch-injured and LPS treated primary microglia cells. Scale bar set to 50µm. B) Quantitation of phagocytosed beads demonstrated that stretch-injury did not alter phagocytic activity when compared to control, uninjured cells, at 12 hours post-injury (8 hour following stretch-injury cells were treated with FITC-IgG for 4 hours). The LPS positive control (100ng/mL) displayed a strong trend towards phagocytosing significantly more beads when assessed at 48 hours post treatment (44 hours after LPS treatment cells were incubated with FITC-IgG for 4 hours).

Figure 7

Stretch-injury does not alter phagocytic activity. A) Representative images of latex beads conjugated with FITC-IgG (green) were incubated for 4 hours with control, stretch-injured and LPS treated primary microglia cells. Scale bar set to 50µm. B) Quantitation of phagocytosed beads demonstrated that stretch-injury did not alter phagocytic activity when compared to control, uninjured cells, at 12 hours post-injury (8 hour following stretch-injury cells were treated with FITC-IgG for 4 hours). The LPS positive control (100ng/mL) displayed a strong trend towards phagocytosing significantly more beads when assessed at 48 hours post treatment (44 hours after LPS treatment cells were incubated with FITC-IgG for 4 hours).

Figure 8

Stretch-injury does not alter microglia NO or LDH responses to subsequent LPS stimulation. A) Stretch-injury prior to LPS treatment does not induce significant difference in LDH release between LPS treated and stretch-injury plus LPS treatment. Both LPS and LPS plus stretch-injury both significantly increase LDH release at 24, 48, and 72 hour post treatment. B) Pre-stretch-injury does not alter NO responses when subsequently exposed to LPS. LPS and stretch-injury plus LPS both significantly upregulate NO at 48 and 72 hours post treatment. N=3/group. One-way ANOVA with Dunnett's post-test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Experimental groups are normalized to control group set at 1.0. Error bars represent mean $\pm$ SD.

Figure 8

Stretch-injury does not alter microglia NO or LDH responses to subsequent LPS stimulation. A) Stretch-injury prior to LPS treatment does not induce significant difference in LDH release between LPS treated and stretch-injury plus LPS treatment. Both LPS and LPS plus stretch-injury both significantly increase LDH release at 24, 48, and 72 hour post treatment. B) Pre-stretch-injury does not alter NO responses when subsequently exposed to LPS. LPS and stretch-injury plus LPS both significantly upregulate NO at 48 and 72 hours post treatment. N=3/group. One-way ANOVA with Dunnett's post-test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Experimental groups are normalized to control group set at 1.0. Error bars represent mean $\pm$ SD.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Westernblot3NTandbetaactin.pdf](#)
- [Westernblot3NTandbetaactin.pdf](#)
- [Westernblotgp91andbetaactin.pdf](#)
- [Westernblotgp91andbetaactin.pdf](#)