

# Long Acting Reversible Contraception use and associated factors among postpartum women in Hossana Town, Southern Ethiopia: cross sectional study

Biruktawit Fekade Woldu (✉ [biruktifekade@gmail.com](mailto:biruktifekade@gmail.com))

Wachemo University

Tadesse Lelago Ermolo

Wachemo University

Lidiya Gutema Lemu

Mizan-Tepi University

Negeso Gebeyehu Gejo

Wachemo University

---

## Research

**Keywords:** Long acting, contraception, Postpartum, Prevalence, Factors, Hossana

**Posted Date:** March 27th, 2020

**DOI:** <https://doi.org/10.21203/rs.2.19848/v3>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

**Version of Record:** A version of this preprint was published on August 5th, 2020. See the published version at <https://doi.org/10.1186/s40834-020-00117-6>.

## Abstract

Background: In low and middle-income countries, 95% of postpartum women want to avoid a pregnancy for 2 years, but 70% are not using contraception. Delay in use of contraception by couples during postpartum period can result in many unwanted pregnancies. Long-acting reversible contraception (LARC) is ideal for postpartum women. Therefore this study aimed at assessing the prevalence and factors associated with LARC use among postpartum women. Methods: Facility based cross sectional study was conducted from July 23-Aug 25, 2018. Systematic random sampling technique was employed to recruit a total of 381 women in extended postpartum period visiting Child Immunization service in hosanna health institutions. Pretested structured questionnaire was used for data collection. Data was analyzed by SPSS version 20. Binary and multivariate logistic regression analysis was done. The presence and strength of association was determined using AOR with its 95% CI. Variables with P value less than 0.05 were considered as statistical significant. Results: The prevalence of LARC use was 36.5% (95%CI (33.05-39.95)). The main reason for not using LARC was fear of side effect and false information. Previous use of LARC (AOR=3.3, 95%CI (1.7-6.5)) and have ever discussed with health providers on LARC (AOR=2.5, 95%CI (1.1-5.74)) were found to be significantly associated with LARC use. Conclusions: The utilization of LARC among postpartum women was found to be higher than other studies in Ethiopia. Increase women's access to realistic information on LARC, mainly on IUD need to be emphasized to enhance its uptake.

## Background

After one-time placement of a device, Long-acting reversible contraception (LARC) methods provide reliable long-term highly effective prevention of pregnancy. LARC methods include subdermal hormonal implant and Intra-Uterine Devices (hormonal IUDs and non-hormonal copper-containing IUDs) [1]. Regardless of considerable advances in contraceptive technologies, unintended pregnancies remain a significant global public health issue. In developing countries 214 million reproductive age women who desire to avoid pregnancy are not using a modern contraceptive method [2-3].

Modern contraceptive use has increased around the world. However, less than 15 percent of married women of reproductive age use a modern method in many of the poorest countries. The use of LARC among modern contraception users is still low [4]. In Ethiopia, Contraceptive prevalence rate (CPR) among currently married women has increased steadily from 14 percent in 2005 to 41 percent in 2019. The largest increase is observed in the use of injectable. The use of LARC remains low at 11% (9%-Implants & 2%-IUD) [5]. In southern region, where the study area is located, 44.6 % of currently married women using a modern family planning. Among these only 9.2% use LARC (7.7%-Implants & 1.5%-IUD). Utilization of LARC was found to have significant association with women's socio-demographic and reproductive characteristics. Knowledge, attitude and previous LARCs use was also found to have significant association with its use [6-14].

In the first year after childbirth, 95% of women in low and middle-income countries want to avoid pregnancy within the next 2 years; however 70% are not using contraception. Delay in practice of contraception by couples during postpartum period can result in many unwanted or mistimed pregnancies. A woman conceives within/less than 24 months of delivery has higher chances of complications like abortions, pre-term labor, low birth weight babies, postpartum hemorrhage, neonatal morbidity and mortality[15].

Postpartum family planning (PPFP) is defined as the prevention of unintended pregnancy and closely spaced pregnancies through the first 12 months following childbirth [16]. Postpartum period is one of the important and crucial times when women and couples are highly motivated and receptive to family planning methods. LARC methods are safe for almost all women, including those in a postpartum period. As they do not contain estrogen and have few contraindications, they are ideal for use in postpartum women[17]. According to a study in five low-income countries, almost all women wish to delay or prevent a future pregnancy at six weeks postpartum. Despite this, usage of modern contraceptive methods at 42 days post-partum varied widely. The uptake of the most effective LARC (intrauterine devices) is very low even amongst users of modern contraceptives [18]. Different studies conducted in different parts of Ethiopia also indicated lower prevalence of postpartum LARC use [19-24]. Study on LARC use among postpartum women is lacking in the study area. Therefore the objective of this study was to assess the prevalence and factors associated with LARC use among women in extended postpartum period visiting child immunization in Hosanna health institutions, Southern Ethiopia.

## Methods

### Study setting and population

Facility based cross sectional study was conducted from July 23-Aug 25, 2018 in hosanna health institutions, Hadiya zone Southern Ethiopia. Hossana town comprises of three sub cities and eight sub kebeles(the lowest governmental administrative units). According to CSA population count of May 2007, the total number of Hossana town population was 69,959 for the year 2007/8. Among these 34454 (49.25%) were females. There is 1 teaching referral hospital and 3 health centers in the town which provide postpartum family planning service. Women in extended postpartum period visiting child immunization service in the four health institutions during the study period were eligible for enrollment. Those women who are pregnant and unable to speak or hear were excluded from the study.

### Sample size determination and Sampling technique

Sample size was calculated using single population proportion formula by taking account of Proportion of LARC use among postpartum women in Durame town 36.7%[13], CI=95% Margin of error =5% and 10% non-response rate. A total of 392 women on extended postpartum period (6 month to 1 year after delivery) were invited to participate. Systematic random sampling technique was employed to select participants. The total number of women who visit child Immunization (N) was estimated by considering the immediate previous 1 month record of EPI chart. Then number of sampling intervals (k) was determined by dividing N by sample size (n=392). Then lottery method was used to select the first participant. Then after every 3<sup>rd</sup> unit was included until the required sample reached. In case of refusal or not meeting inclusion criteria, the next person was selected. The sample size for each institution was distributed proportionally.

## Data collection

Data were collected using pretested, interviewer administered structured questionnaire. These questionnaires were adapted from different literatures by considering the local situation of the study subjects. Information on participants' socio-demographic, reproductive characteristics, knowledge, attitudes and practice towards LARC was collected. The data collection tool prepared in English was translated in to Amharic and then returned back to English to ensure accuracy. Before the actual data collection, Pre-test was done on 5% women in a health center located outside the town and necessary corrections made accordingly. Four trained diploma midwifery graduates collected the data supervised by one BSc midwife working in university. Completeness of the questionnaires was checked on the spot by supervisors.

## Measurement

LARC use: was defined as utilization of any of LARCs (IUD or subdermal hormonal implant) by women on extended postpartum period.

The mean score was used to categorize women's knowledge towards LARC; women who scored above mean score were considered to have good knowledge and women who scored below the mean were considered to have poor knowledge. Women's attitude towards LARC was categorized in the same approach. Those women who scored above the mean were considered to have positive attitude and scored below the mean considered to have negative attitude towards LARC.

## Data analysis

Data was analyzed by using SPSS version 20. Descriptive statistics were computed. Binary logistic regression analysis was done and all explanatory variables with p value less than 0.2 were entered to Multivariate logistic regression model. The second analytical step used a stepwise backward model. Then association was assessed using AOR and its 95%CI. And p value of <0.05 was considered as statistically significant. Goodness of model fitness was tested by using Hosmer and Lemeshow test.

## Results

Data were analyzed from 381 women, making response rate of 97.1%. Mean age of participants were 27(SD±4.6). Majority of participants were urban resident (89%). Less than one fifth of respondents attend diploma and above (17.1%). Half of the women were housewife (50.4%) (Table 1).

Table 1: Sociodemographic characteristics of postpartum women in Hossana health institutions, 2018(n=381)

	Variable	Frequency	Percentage
Age (years)	16-20	31	8.1
	21-25	124	32.5
	26-30	154	40.4
	31-35	50	13.1
	36-40	22	5.8
	Residence	Urban	339
Marital status	Rural	42	11.0
	Married	369	96.9
	Single	7	1.8
Educational status	Separated	5	1.3
	Can't read and write	15	3.9
	Can Read & Write	19	5.0
	Primary	141	37.0
	Secondary	141	37.0
	Diploma & Above	65	17.1
Working status	Housewife	192	50.4
	Merchant	51	13.4
	Government Employee	67	17.6
	Private Employee	58	15.2
	Other	13	3.4

## Reproductive characteristics of respondents

More than two-third (71%) of participants had two or more births. Median number of alive children per women and birth interval were 2 IQR [1,4] and 36 months IQR [24,48] respectively. Around one-third (28.3%) participants do not want any more children for the future.

Less than one fifth (18.1%) intended to give birth within 2 years. More than one-fourth (28.6%) had never discussed with health professionals and 52.4% were advised on LARC during the immediate postpartum period (Table 2).

Table 2: Reproductive characteristics of postpartum women in Hossana health institutions, 2018(n=381)

	Variable	LARC use	
		Yes n(%)	No n(%)
Parity	1	72(65.5)	38(34.5)
	2-4	137(61.7)	85(38.3)
	≥5	33(67.3)	16(32.7)
Number of alive children	1	75(65.8)	39(34.2)
	2-4	135(60.8)	87(39.2)
	≥5	32(71.1)	13(28.9)
Place of last birth	Home	15(71.4)	6(28.6)
	Health institution	227(63.1)	133(36.9)
Birth interval(in month) (n=272)	0-23	28(73.7)	10(26.3)
	24-35	56(64.4)	31(35.6)
	36 & above	87(59.2)	60(40.8)
Number of children want for the future	0	74(68.5)	34(31.5)
	1-3	131(59.8)	88(40.2)
	4-6	37(68.5)	17(31.5)
Intended to give birth within 2years (n=273)	Yes	49(71.0)	20(29.0)
	No	119(58.3)	85(41.7)
Ever discussed with health professionals	Yes	123(45.2)	149(54.8)
	No	16(14.7)	93(85.3)
Advised during immediate postpartum	Yes	95(47.5)	105(52.5)
	No	44(24.3)	137(75.7)

### Knowledge, Attitude and previous LARC use

More than two-third (71.4%) of the women had high level of knowledge towards LARC. Around half of them (53.2%) had positive attitude towards LARC. Two-third (68.5%) of participant's partner had supportive attitude towards LARC. Two-third (66.4%) of the respondents had ever used at least one type of modern contraception and one-fourth (26%) had ever used LARC before delivery of last child (Table 3).

Table 3: Knowledge, attitude and previous LARC use among postpartum women in Hossana health institutions, 2018(n=381)

	Variable	LARC use	
		Yes	No
Knowledge towards LARC	High level of knowledge	121(44.5%)	151(55.5%)
	Low level of Knowledge	18(16.5%)	91(83.5%)
Attitude towards LARC	Positive	77(41.8%)	107(58.2%)
	Negative	61(38.4%)	98(61.6%)
Previous LARC use	Used before	58(59.2%)	40(40.8%)
	Never used	81(28.6%)	202(71.4%)

### Prevalence of postpartum LARC use

Two-third (66.6%) of postpartum women had started to use modern family planning method. LARC method use among participants was 36.5% (implant 34.1%, IUCD-2.4%). The main reason for not using LARC was fear of side effect (39.3%) followed by false rumors towards LARC (it cause harm to uterus, infertility, interfere with work, fly/lost in the body) (21.9%) (Table 4).

Table 4: method-mix adopted among postpartum women in Hossana health institutions, 2018(n=381)

Contraception method	Frequency	Percentage
Injectable	98	25.7
Implants	130	34.1
IUCD	9	2.4
Pills	15	3.9
Condom	2	0.5
Non users	127	33.4

### Factors associated with LARC use

Previous LARC use and ever discussed with Health Professionals on LARC were found to be significantly associated with LARC use. Those women who have ever used LARC before were 3.3 times more likely to use LARC on postpartum period (AOR=3.3, 95%CI(1.7-6.5). Postpartum women who have ever discussed with health professionals on LARC were 2.5 times more likely to use LARC (AOR=2.5, 95% CI (1.1-5.7)) (Table 5).

Table 5: Bivariate and Multivariate analysis of factors associated with LARC use among postpartum women in hossana health institutions, 2018 (n=381)

Variable		LARC use		COR (95%CI)	AOR (95%CI)
		Yes (n%)	No (n%)		
Residence	Urban	130(38.3)	209(61.7)	1	
	Rural	9(21.4)	33(78.6)	0.43 (.20-.94)	0.53 (0.19-1.42)
working status	Working	75(39.7)	114(60.3)	1.31 (.86-1.99)	
	Not working	64(33.3)	128(66.7)	1	
Intention to give birth within 2 years	Yes	20(29.0)	49(71.0)	1	
	No	85(41.7)	119(58.3)	1.75(.97-3.15)	1.82 (0.94-3.54)
Knowledge towards LARC	Good	121(44.5)	151(55.5)	4.01(2.31-7.08)	0.95 (0.43-2.10)
	Poor	18(16.5)	91(83.5)	1	
Partner attitude	Supportive	118(45.2)	143(54.8)	2.59 (1.48-4.54)	0.50 (0.24-1.03)
	Not supportive	20(24.1)	63(75.9)	1	
Previous LARC use	Yes	58(59.2)	40(40.8)	3.61 (2.24-5.83)	3.34(1.70-6.55)*
	No	81(28.6)	202(71.4)	1	
Advised during immediate postpartum	Yes	95(47.5)	105(52.5)	2.81 (1.81-4.36)	1.26 (0.66-2.42)
	No	44(24.3)	137(75.7)	1	
Ever discussed on LARC with professionals	Yes	123(45.2)	149(54.8)	4.79 (2.68-8.58)	2.56 (1.14-5.74)**
	No	16(14.7)	93(85.3)	1	

\*Significant at <0.05 \*\* significant at <0.001

## Discussion

In this study the prevalence of LARC use among postpartum women was found to be 36.5% (95%CI (33.05-39.95)), which is similar with a study done in Durame [13]. However it is higher than a study in Northern Ethiopia[24]. The difference can be due to variation in participants' characteristics and institution based nature of our study.

Majority of LARC users opted for Implant(93.5%) and very few used IUCD(6.5%). Which is consistent with different studies [6, 14] where strong preference for implants over IUD were observed. These can be due to prevailing misinformation towards IUCD.

The main reason for not using LARC was fear of side effect (39.3%) followed by false information towards LARC (it cause harm to uterus and infertility, interfere work, fly/lost in the body)(21.9%) which is consistent with other study[8]. This indicates the need of community based health education and awareness creation programs.

In this study previous LARC use was found to be significantly associated with postpartum LARC use (AOR=3.34 (1.70-6.55)). Similar finding were indicated by other studies [10,13]. The possible explanation can be women possibly will have adequate information on the contraception method they used before, which can motivate them to use the same method repetitively.

Those women who ever discussed with health professionals on LARC were found to be more likely to use LARC (AOR=2.569(1.148-5.7470)). It is in line with another study [6]. This can possibly explained by the fact that access to clear factual information on the benefits and limitations of contraceptive methods is imperative to promote informed choice and to overcome misperceptions and myths more broadly perpetuated in the community.

## Limitation

Quantitative nature of the study hinders indepth exploration of women's perception and barrieres for LARC use. As this study was conducted in urban facilities, the finding might not be representative for the rural communities.

## Conclusion

The utilization of LARC among postpartum women was found to be higher than other studies done in Ethiopia. Majority participants opted for implants over IUCD. The main reason for not using LARC was fear of side effect and rumors. Increase women's access to realistic information on LARC mainly on IUCD need to be emphasized. Further studies need to be conducted to assess women's knowledge, perceptions and attitudes towards PPIUCD.

## Declarations

### Ethics approval and consent to participate

Ethical approval was obtained from wachemo university ethical review committee. Cooperation letter submitted to health institutions. The objective, benefit and risks of the study were explained and informed written consent obtained from all respondents. The attainment of confidentiality was assured for participants.

### Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

### Competing interests

The authors proclaim that they have no competing interests.

## Funding

This study was conducted by the fund obtained from Wachemo University.

## Authors' contributions

BFW, TLE, and LGL contributed to the initiation of the study, design, participating in organizing the data collection process and analysis of the result. BFW, TLE, LGL and NGG involved in writing of the manuscript. All authors read and approved the final manuscript.

## Acknowledgements

We are indebted to data collectors and all women who agreed to participate in this study. We want to thank health institution administrators for their great collaboration.

## Consent for publication

Not applicable(2)

## References

1. Winner B, Peipert JF, Zhao Q, Buckel C, Madden T, Allsworth J et al. Effectiveness of long-acting reversible contraception. *N Engl J Med.* 2012;366:1998-2007.
2. Family planning/Contraception Fact sheet. In: WHO, editor. 2017.
3. Curtis KM, Peipert Long-Acting Reversible Contraception. *N Engl J Med.* 2017;376:461-8. DOI: 10.1056/NEJMcp1608736
4. Family Planning Worldwide 2019 Data Sheet [www.prb.org/pdf19/familyplanning-2019-datasheet\\_eng.pdf](http://www.prb.org/pdf19/familyplanning-2019-datasheet_eng.pdf) Accessed December 11, 2019.
5. Central Statistical Agency (CSA) [Ethiopia] and ICF. 2019. Ethiopia Mini Demographic and Health Survey 2019: Key Indicators Report. Addis Ababa, Ethiopia, and Rockville, Maryland, USA. CSA and ICF. <https://dhsprogram.com/pubs/pdf/PR120/PR120.pdf>
6. Taye A, Woldie M, Sinaga M. Predictors of Long Acting Reversible Contraceptive use among Married Women Visiting Health Facilities in Jimma Town. *J Women's Health Care.* 2014; 4: 217. doi:10.4172/2167-0420.1000217. ([https://www.researchgate.net/profile/Mirkuzie\\_Woldie2/publication/273128776\\_Predictors\\_of\\_Long\\_Acting\\_Reversible\\_Contraceptive\\_use\\_among\\_Married-of-Long-Acting-Reversible-Contraceptive-use-among-Married-Women-Visiting-Health-Facilities-in-Jimma-Town.pdf](https://www.researchgate.net/profile/Mirkuzie_Woldie2/publication/273128776_Predictors_of_Long_Acting_Reversible_Contraceptive_use_among_Married-of-Long-Acting-Reversible-Contraceptive-use-among-Married-Women-Visiting-Health-Facilities-in-Jimma-Town.pdf))
7. Shegaw G, Mohammed AA, Nadew K, Tamrat K, Zeru G, Desta H, Yinager W. Long Acting Contraceptive Method Utilization and Associated Factors among Reproductive Age Women in Arba Minch Town, Ethiopia. *Greener Journal of Epidemiology and Public Health.* 2014;2 (1): 023-031.
8. Gebremichael H, Haile F, Dessie A, Birhane A, Alemayehu M, Yebyo H. Acceptance of Long ActingContraceptive Methods and Associated Factors among Women in Mekelle City, Northern Ethiopia. *Science Journal of Public Health.* 2014;2(4):239-45.
9. Kabalo Utilization of reversible long acting family planning methods among married 15-49 years women in Areka town, Southern Ethiopia. *International Journal of Scientific Reports.* 2016;2(1):1-6 <http://www.sci-rep.com>
10. MelkaAS, Tekelab T, Wirtu D. Determinants of long acting and permanent contraceptive methods utilization among married women of reproductive age groups in western Ethiopia: a cross-sectional study. *Pan African Medical.* 2015;21 10.11604(246).
11. Austad K, Shah P, Rohloff P. Correlates of long-acting reversible contraception uptake among rural women in Guatemala. *PLoS ONE.* 2018;13(6): e0199536. <https://doi.org/10.1371/pone.0199536>
12. Teferra AS, Wondifraw AA. Determinants of Long Acting Contraceptive Use among Reproductive Age Women in Ethiopia: Evidence from EDHS 2011. *Science Journal of Public Health.* 2014; 3(1), 143-149.doi: 10.11648/j.sjph.20150301.33
13. Tamrie YE, Hanna EG, Argaw M.D. Determinants of Long Acting Reversible Contraception Method Use among Mothers in Extended Postpartum Period, Durame Town, Southern Ethiopia: A Cross Sectional Community Based Survey. *Health,* 2015; 7, 1315-1326. <http://dx.doi.org/10.4236/health.2015.710146>
14. Gudaynhe SW, Zegeye DT, Tarekegn A, Kibret GD. Factors Affecting the use of Long-Acting Reversible Contraceptive Methods among Married Women in DebreMarkos Town, Northwest Ethiopia. *Global Journal of Medical Research.* 2014; 14(5), 13-14.
15. Shoupe LARC methods: entering a new age of contraception and reproductive health *Contraception and Reproductive Medicine.* 2016;1(4). DOI 10.1186/s40834-016-0011-8(<http://creativecommons.org/publicdomain/zero/1.0/>).
16. World Health Organization. Programming strategies for postpartum family planning. 2013. ([https://www.who.int/reproductivehealth/publications/family\\_planning/ppfp\\_strategies/en/](https://www.who.int/reproductivehealth/publications/family_planning/ppfp_strategies/en/))
17. Parks C, Peipert F. Eliminating Health Disparities in Unintended Pregnancy with Long-Acting Reversible Contraception (LARC) *Am J Obstet Gynecol.* 2016; 214(6):681688.doi:10.1016/j.ajog.2016.02.017. (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4884485/>)
18. Pasha O, Goudar SS, Patel A, Garces A, Esamai F, Chomba E, et al. Postpartum contraceptive use and unmet need for family planning in five low-income countries. *Reproductive health.* 2015;12(2):S11. (<https://www.ncbi.nlm.nih.gov/pubmed/26063346>)
19. Abera Y, Mengesha ZB, Tessema GA. Postpartum contraceptive use in Gondar town, Northwest Ethiopia: a community based cross-sectional study. *BMC women's health.* 2015;15(1):19.
20. Gebremedhin AY, Kebede Y, Gelagay AA, Habitu YA. Family planning use and its associated factors among women in the extended postpartum period in Addis Ababa, Ethiopia. *Contraception and reproductive medicine.* 2018;3(1):1.

21. Abraha TH, Gebrezgiabher BB, Aregawi BG, Belay DS, Tikue LT, Welay GM. Predictors of postpartum contraceptive use in rural Tigray region, northern Ethiopia: a multilevel analysis. *BMC public health*. 2018;18(1):1017.
22. Gebremariam A, Gebremariam H. Contraceptive use among lactating women in Ganta-Afeshum District, Eastern Tigray, Northern Ethiopia, 2015: a cross sectional study. *BMC pregnancy and childbirth*. 2017;17(1):421.
23. Demie T, Demissew T, Huluka T, Workineh F, Libanos H. Postpartum family planning utilization among postpartum women in public health institutions of Debre Berhan town, Ethiopia. *J Women's Health Care*. 2018;7(426):2167-0420.
24. Mengesha ZB, Worku AG, Feleke SA. Contraceptive adoption in the extended postpartum period is low in Northwest Ethiopia. *BMC pregnancy and childbirth*. 2015;15(1):160. (<https://bmcpregnancychildbirth.biomedcentral.com/articles/10.1186/s12884-015-0598-9>)