

VATS right posterior segmentectomy with anomalous bronchi and pulmonary vessels: a case report and literature review

Jianbin Zhang (✉ fishing11@163.com)

HuZhou Central Hospital, Affiliated Central Hospital of Huzhou University <https://orcid.org/0000-0001-6007-1247>

Yilv Zhu

Affiliated Hospital of Huzhou Normal University: Huzhou Central Hospital

Hongwei Li

Affiliated Hospital of Huzhou Normal University: Huzhou Central Hospital

Caihua Yu

Affiliated Hospital of Huzhou Normal University: Huzhou Central Hospital

Weiwei Min

Affiliated Hospital of Huzhou Normal University: Huzhou Central Hospital

Case report

Keywords: Video-assisted thoracoscopic surgery (VATS), three-dimensional computed tomography bronchography and angiography (3D-CTBA), segmentectomy

Posted Date: February 12th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-104612/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Journal of Cardiothoracic Surgery on March 29th, 2021. See the published version at <https://doi.org/10.1186/s13019-021-01420-2>.

Abstract

Background: Anatomic variation may increase the difficulty and risk of anatomic segmentectomy. The preoperative three-dimensional computed tomography bronchography and angiography (3D-CTBA) can provide a detailed model of the segmental structure, and contribute to precise and safe segmentectomy.

Case presentation: This is a case of anomalous bronchi and pulmonary vessels in the right upper posterior segment (RS²). Under the guidance of 3D-CTBA, anatomic RS² segmentectomy was performed accurately and safely. The postoperative condition was uneventful.

Conclusions: This rare case highlights the importance of 3D-CTBA to guide accurate segmentectomy with anatomic variation.

Background

With the popularization of chest computed tomography (CT), the discovery of small pulmonary nodules is increasingly common, which also promotes the progress of segmentectomy [1, 2]. As reported in the previous cases, the anatomic structure of the pulmonary segment is sometimes variable [3-5]. The public's attention is now shifting to how to achieve an accurate anatomic segmentectomy. The development and application of various 3D imaging softwares have given a strong technical support in terms of the successful resection [6]. Preoperative reconstruction of the segment can clearly show the anatomic structure and judge whether there are variations, so as to make an accurate surgical plan. In this case, a 34-year-old female patient experienced a ground-glass nodule (GGN) at the right upper posterior segment (RS²). The preoperative three-dimensional CT bronchography and angiography (3D-CTBA) revealed multiple anatomic variations. Owing to the detailed planning of the operation, video-assisted thoracoscopic surgery (VATS) RS² segmentectomy was performed successfully.

Case presentation

A 34-year-old female was admitted to us with a GGN in RS², which was discovered by chest CT during a health checkup 1 year ago. A review of CT indicated that the subpleural GGN was slightly larger than before, with a diameter of 7 mm and a CT value of -400HU (Figure 1). She had no positive signs, no history of smoking, no history of malignant tumors, and no family history of lung cancer. Preoperative 3D-CTBA revealed multiple variations in the right upper lobe: (1) The apical subsegmental bronchi B^{1a} and B^{1b} originated from the posterior segmental bronchus (B²) and the anterior segmental bronchus (B³), respectively. (2) The right upper pulmonary arteries shared a trunk without a posterior ascending artery (Asc.A²). (3) The right upper lobe had no central vein, with only 1 posterior intrasegmental vein (V^{2t}); the other 2 veins pointed to intersegmental plane respectively without another posterior intrasegmental vein (V^{2b}) (Figure 2). VATS anatomic RS² segmentectomy and lymph node sampling were conducted accurately With the guidance of 3D-CTBA (Figure 3); all anatomic variations were successfully detected during the operation (Figure 4). Minimally invasive adenocarcinoma (MIA) was the fast-frozen pathology.

The chest radiograph illustrated that the right lung was completely redilated, and on the second postoperative day, the **incisal margin** of the segment displayed no obvious exudation (Figure 5). The drainage tube was removed on the second postoperative day, and the patient was discharged on the third postoperative day. The postoperative pathology revealed MIA with negative surrounding lymph nodes.

Video description

Intravenous general anesthesia combined with double-lumen endotracheal intubation and contralateral one-lung ventilation were performed. The observation port was made in the seventh intercostal space of the posterior axillary line, and the operation port in the fourth intercostal space of the anterior axillary line. The surgical procedure was as follows: (1) Thoracoscopic exploration revealed a subpleural nodule at the RS². (2) The parietal pleura was sectioned by harmonic, and then the Nos.11 lymph nodes were removed. (3) The V²t was dissociated and pulled by silk, and the Nos.12 and 13 lymph nodes around the bronchus were completely removed. (4) After ligating the V²t, the B² was dissected out and transected with a thick tissular (blue cartridge) 45-mm long endostapler (EC45A, JJMC, USA). (5) According to the preoperative 3D-CTBA, the other 2 veins were judged as the intersegmental veins; hence, they were all preserved. (6) We reventilated the right lung with pure oxygen because the posterior segmental artery was very difficult to ligate; when the intersegmental plane was clear after 18 minutes, the segmental hilum was released sufficiently to expose the posterior segmental artery, and then the artery was ligated safely. (7) The intersegmental plane was divided along the inflation-deflation line using the endostaplers; thus, the posterior segment was removed successfully. A sterile glove was used to collect the specimen. Bleeding and air leakage were not observed after the pleural injection of inflation. No. 24 chest drainage tube was placed in the observation port. The incision was sutured after the right lung was completely dilated. The operative time was 75 minutes, and intraoperative blood loss was 50 mL.

Discussion And Conclusions

In recent years, anatomic segmentectomy for early lung cancer is one of the biggest hotspots in thoracic surgery. It can not only completely resect the tumor but also preserve the normal lung tissue to the maximum extent [7]. The distribution of bronchus, artery, and vein in the pulmonary segment exists variations in some patients. The key to the implementation of this operation is to accurately grasp the anatomic structure of the target segment. The 3D-CTBA can provide a precise anatomic structure, identify the intrasegmental and intersegmental veins from different views [8]. Therefore, some complicated pulmonary segmentectomy should be performed under the guidance of 3D-CTBA [9, 10].

The efficacy of 3D reconstruction for thoracic surgery has been previously described[11], Kimihiro et al. [12] first reported the application of 3D-CTBA in VATS segmentectomy. At present, the main softwares for 3D reconstruction of segment include IQQA, DeepInsight, etc[13, 14]. All the reconstructive softwares can offer precise anatomic structure of pulmonary segments, but sometimes, it is difficult for surgeons to skillfully use them. In our institution, with the assistance of the radiologists, we adopt the CT pulmonary angiography based technology to reconstruct the anatomic structure of each patient who undergo

segmentectomy. It shows the precise structure of bronchi, arteries and veins in a natural surgical field view.

In all kinds of segmental resection, RS² segmentectomy is a common procedure, but anatomic variations may increase its difficulty and risk [15]. Xinfeng et al. [16] reported a tracheal bronchus and a variable central vein entering the left atrium dorsal to the right pulmonary artery trunk in a patient who underwent VATS RS² segmentectomy. Tadashi et al. [17] reported anatomic variations in bronchi and pulmonary vessels in a patient who underwent thoracoscopic RS² segmentectomy. However, to the best of our knowledge, bronchial variation associated with variant pulmonary vessels has rarely been reported. The normal anatomic structure of RS² consists of B², Asc.A², the recurrent artery (Rec.A²), intrasegmental veins (V²t and V²b) and intersegmental veins (V²a and V²c), but in this case, the 3D-CTBA revealed multiple variations: (1) The bronchopulmonary trees of the right upper lobe were divided into (B²+B¹a) and (B³+B¹b); the (B²+B¹a) variation might have been mistaken for B² without the guidance of 3D-CTBA. (2) There was only 1 intrasegmental vein (V²t) without another intrasegmental vein (V²b); intersegmental veins might have been transected as intrasegmental veins without the preoperative 3D-CTBA. (3) The posterior segmental arteries originated from the superior trunk without Asc.A². They were very difficult to expose; therefore, we released the hilum of segment first to expose the arteries sufficiently, and then the artery was safely ligated. This novel strategy has rarely been reported. In conclusion, we present a successful strategy for VATS RS² segmentectomy with multiple anatomic variations.

Declarations

Ethics approval and consent to participate

The article was reviewed and approved by the by the research ethics committee of Huzhou Central Hospital, Affiliated Central Hospital of HuZhou University. Written informed consent was signed by all participants.

Consent for publication

Not applicable.

Availability of data and materials

All data generated or analysed during this study are included in this article.

Competing interests

The authors declare that they have no competing interests.

Funding

Not applicable.

Authors' contributions

Jianbin Zhang performed surgery on this patient, and was the major contributor in writing. Yilv Zhu performed the segmental reconstruction, and provided the photos of the 3D-CTBA. Hongwei Li provided the technical guidance. Caihua Yu and Weiwei Min were involved in patient's care and follow-up. All authors read and approved final manuscripts.

Acknowledgements

Not applicable.

Abbreviations

3D-CTBA: three-dimensional computed tomography bronchography and angiography; RS²: right upper posterior segment; VATS: Video-assisted thoracoscopic surgery; CT: computed tomography; GGN: ground-glass nodule; MIA: minimally invasive adenocarcinoma; B¹: apical segmental bronchus; B²: posterior segmental bronchus; B³: anterior segmental bronchus; Asc.A²: ascending artery; Rec.A²: recurrent artery.

References

1. Nakazawa S, Shimizu K, Mogi A, Kuwano H: **VATS segmentectomy: past, present, and future.** *Gen Thorac Cardiovasc Surg* 2018, **66**(2):81-90.
2. Sato Y, Fujimoto D, Morimoto T, Uehara K, Nagata K, Sakanoue I, Hamakawa H, Takahashi Y, Imai Y, Tomii K: **Natural history and clinical characteristics of multiple pulmonary nodules with ground glass opacity.** *Respirology* 2017, **22**(8):1615-1621.
3. Aragaki M, Iimura Y, Yoshida Y, Hasegawa N: **Anomalous V(2) of the left pulmonary vein detected using three-dimensional computed tomography in a patient with lung cancer: A case report.** *Int J Surg Case Rep* 2017, **37**:208-210.
4. Ohtaka K, Iwashiro N, Watanabe K, Mizota T, Takahashi R, Suzuoki M, Komuro K, Ohara M, Kaga K, Matsui Y: **A left lung abscess with a displaced subsegmental bronchus and anomalous pulmonary artery and vein: a case report.** *Surg Case Rep* 2019, **5**(1):66.
5. Qi W, Yang F, Liu H, Sun Y: **Thoracoscopic right apical segmentectomy for lung cancer with tracheal bronchus and variable pulmonary vein: a case report and literature review.** *J Int Med Res* 2020, **48**(8):300060520947935.
6. Chang SS, Okamoto T, Tokunaga Y, Nakano T: **Intraoperative Computed Tomography Navigation During Thoracoscopic Segmentectomy for Small-sized Lung Tumors.** *Semin Thorac Cardiovasc Surg* 2018, **30**(1):96-101.
7. Tane S, Nishio W, Nishioka Y, Tanaka H, Ogawa H, Kitamura Y, Takenaka D, Yoshimura M: **Evaluation of the Residual Lung Function After Thoracoscopic Segmentectomy Compared With Lobectomy.** *Ann Thorac Surg* 2019, **108**(5):1543-1550.

8. Wu WB, Xu XF, Wen W, Xu J, Zhu Q, Chen L: **Thoracoscopic Pulmonary Sub-Subsegmentectomy Based on Three-Dimensional Images.** *Ann Thorac Surg* 2016, **102**(5):e389-e391.
9. Sato M, Murayama T, Nakajima J: **Thoracoscopic stapler-based "bidirectional" segmentectomy for posterior basal segment (S10) and its variants.** *J Thorac Dis* 2018, **10**(Suppl 10):S1179-S1186.
10. Shimizu K, Nagashima T, Yajima T, Ohtaki Y, Obayashi K, Nakazawa S, Kosaka T, Mogi A, Kuwano H: **Thoracoscopic Medial-Basal Segment Segmentectomy.** *Ann Thorac Surg* 2017, **104**(5):e403-e406.
11. Fukuhara K, Akashi A, Nakane S, Tomita E: **Preoperative assessment of the pulmonary artery by three-dimensional computed tomography before video-assisted thoracic surgery lobectomy.** *Eur J Cardiothorac Surg* 2008, **34**(4):875-877.
12. Shimizu K, Nakano T, Kamiyoshihara M, Takeyoshi I: **Segmentectomy guided by three-dimensional computed tomography angiography and bronchography.** *Interact Cardiovasc Thorac Surg* 2012, **15**(2):194-196.
13. Wu WB, Xu XF, Wen W, Xu J, Zhu Q, Pan XL, Xia Y, Chen L: **Three-dimensional computed tomography bronchography and angiography in the preoperative evaluation of thoracoscopic segmentectomy and subsegmentectomy.** *J Thorac Dis* 2016, **8**(Suppl 9):S710-S715.
14. Xu G, Chen C, Zheng W, Zhu Y, Zheng B, Chen H: **IQQA-3D imaging interpretation and analysis system-guided single-port video-assisted thoracic surgery for anatomical sub-segmentectomy (LS(1+2)a+b).** *J Thorac Dis* 2018, **10**(9):5515-5521.
15. Polaczek M, Szaro P, Jakubowska L, Zych J, Religioni J, Orłowski TM: **Pulmonary veins variations with potential impact in thoracic surgery: a computed-tomography-based atlas.** *J Thorac Dis* 2020, **12**(3):383-393.
16. Xu XF, Chen L, Wu WB, Zhu Q: **Thoracoscopic right posterior segmentectomy of a patient with anomalous bronchus and pulmonary vein.** *Ann Thorac Surg* 2014, **98**(6):e127-129.
17. Akiba T, Morikawa T, Marushima H, Nakada T, Inagaki T, Ohki T: **Computed tomography guided thoracoscopic segmentectomy for lung cancer with variant bronchus.** *Ann Thorac Cardiovasc Surg* 2014, **20**(5):407-409.

Figures




Figure 1

The preoperative computed tomography scanning. A 7-mm ground-glass nodule was identified at the posterior segment of the right upper lobe (arrow) , the CT value was -400HU.




Figure 2

The three-dimensional computed tomography bronchography and angiography. The apical subsegmental bronchi B1a and B1b originated from the posterior segmental bronchus (B2) and the anterior segmental bronchus (B3) respectively; The right upper pulmonary artery shared trunk without posterior ascending artery (Asc.A2); The right upper pulmonary vein had no central vein, with only one posterior intrasegmental vein (V2t), the other two veins pointed to the intersegmental plane respectively. (A) Posterior view; (B) Lateral view.



Figure 3

The QR code of Video. VATS right posterior segmentectomy and lymph node sampling.




Figure 4

The intraoperative view of the posterior segment of the right upper lobe. (A) Exposed the posterior segmental artery after releasing the segmental hilum sufficiently; (B) The structure of lung segmental hilum after resection.




Figure 5

The chest radiograph on the 2nd postoperative day. The incisal margin of the segment displayed no obvious exudation(arrow).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CAREchecklistEnglish2013.pdf](#)