

Pancreatic Safety of DPP-4 Inhibitors in Type 2 Diabetes Mellitus—Protocol for a Systematic Review and Network Meta-Analysis of Randomized Controlled Trials

Fan Yang

Shandong University of Traditional Chinese Medicine

Quanlin Zhao (✉ zhaoquanlin65@163.com)

Shandong University of Traditional Chinese Medicine Affiliated Hospital

Youzi Dong

Shandong University of Traditional Chinese Medicine

Baohua Li

Shandong University of Traditional Chinese Medicine

Bobiao Ning

Shandong University of Traditional Chinese Medicine

Protocol

Keywords: Dipeptidyl-peptidase IV inhibitors, Pancreatic safety, Type 2 diabetes mellitus, Systematic review, Network meta-analysis

Posted Date: November 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1047322/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Dipeptidyl-peptidase IV inhibitor (DPP-4i) is a common hypoglycemic medication in treating type 2 diabetes mellitus (T2DM). It has become widely utilized in clinical practice due to its ability to effectively manage blood glucose while posing a low risk of hypoglycemia and weight gain. However, there is no consensus on DPP-4i's pancreatic safety due to a paucity of clinical evidence. The safe event appears to be easily overlooked. This review aims to evaluate the pancreatic safety of DPP-4i in patients with type 2 diabetes mellitus using the standard pairwise and network meta-analysis (NMA) methods.

Methods: MEDLINE, Embase, Pubmed, Web of Science, and the Cochrane Central Register of Controlled Trials will be used to search for published literature and clinical trial registries (such as www.ClinicalTrials.gov) will be used to look for unpublished trials. English-language randomized controlled trials (RCTs) will be included. Two independent reviewers will screen literature for eligibility, extract available data, and assess the risk of bias. All divergences will be resolved after rechecking the source papers and further discussion among the reviewers with a complete consensus before inclusion. Pancreatic safety events involving acute pancreatitis, chronic pancreatitis, and pancreatic cancer will be the primary outcome. The secondary result will be the change in pancreatic enzyme from the baseline. A random-effects model will be used for pairwise meta-analysis with a 95% confidence interval, and Bayesian network meta-analysis will be conducted between the various interventions.

Discussion: This systematic review and network meta-analysis will evaluate the pancreatic safety of DPP-4i in patients with type 2 diabetes mellitus. The findings of this study may supplement the evidence-based information on DPP-4i, improve existing understanding of this issue, and assist patients and clinicians in making better treatment decisions by raising their awareness of the problem.

Systematic review registration: submitted in PROSPERO, 31 August 2021

Background

DPP-4i is an incretin-based pharmacotherapy, a new antidiabetic drug that differs from conventional oral glucose-lowering medications. They decrease the inactivation of incretin hormones such as glucagon-like peptide-1 (GLP-1) and gastric inhibitory polypeptide (GIP), raising the level of endogenously generated incretin hormones by inhibiting the dipeptidyl-peptidase IV, which degrades principal hormones into inactive products. Incretin hormones stimulate insulin secretion while inhibiting glucagon secretion in a glucose-dependent way, lowering blood glucose and reducing the risk of hypoglycemia and weight gain. Apart from the effects mentioned above, Dpp-4i has certain cardiovascular safety advantages that protect the heart and do not raise the risk of severe cardiovascular adverse events or mortality in T2DM patients. As a result, this medication is frequently given to treat T2DM as a monotherapy or in conjunction with other hypoglycemic medicines.

In recent years, more attention has been devoted to the cardiovascular or renal outcomes of DPP-4i. However, other safety events, such as pancreatic events that are uncommon and controversial, are more

likely to be overlooked. Clinically, related pancreatic events mainly refer to pancreatic exocrine diseases, such as pancreatitis and pancreatic cancer. According to the mechanism of action of DPP-4i, which increases GLP-1 in the body, some researchers hypothesized that GLP-1-based therapy could hasten the potential adverse effects of exocrine dysplasia and induce α -cell hyperplasia, which can negatively affect the pancreatic ducts and increase the risk of acute or chronic pancreatitis, as well as pancreatic cancer[1]. An experiment that reveals increased pancreatic ductal turnover, ductal metaplasia, and isolated pancreatitis in rats treated exclusively with DPP-4i found a potential connection between DPP-4i treatment and pancreatic events[2]. These findings could provide evidence for how incretin therapy contributes to pancreas-related disorders. Regrettably, other researchers do not share this viewpoint. The TECOS Study discovered no statistically significant difference between the Sitagliptin and placebo groups, while numerical differences exist[3]. The population-based research of incretin-based treatment in Taiwan found no link between the DPP-4i and acute pancreatitis and the risk of pancreatic cancer in the short term[4]. However, it is challenging to overcome restrictions imposed by technique and objective situations. The safety of DPP-4i medicines on the pancreas is still a worry for diabetic individuals. While DPP-4is have specific benefits for antidiabetic treatment, none of them could offset the slight risk of having unfavorable pancreatic consequences.

Numerous researchers have investigated this topic in vivo and vitro, and some secondary analysis has also been conducted, but no consensus has been reached based on the evidence gathered thus far. Moreover, most of the published meta-analyses were a pairwise comparison of two treatments, which might skew the results' validity. As a result, our research aims to compile all randomized controlled trials that compare the pancreatic endpoints of DPP-4i therapy to other antidiabetic drugs or placebo in T2DM patients to assess the pancreatic safety of DPP-4i in a pairwise and network meta-analysis.

Methods And Design

PRISMA-P (Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols) will be used to guide the protocol for this systematic review and NMA [5]. A completed PRISMA-P checklist was utilized to guarantee the quality of the protocol, as indicated in **Additional file1**. In addition, the review will be published using the PRISMA extension statement for network meta-analysis[6]. Our protocol has been registered with PROSPERO; any amendments and their reasons will be added to the platform.

Eligibility criteria

Types of participants

We will include studies with patients diagnosed with type 2 diabetes using the criteria formulated by WHO[7] and who are 18 years old and taking DPP-4is or other active antidiabetic medications, regardless of gender, race, or nationality.

Types of interventions and comparators

We will take into account the research that evaluated the following therapies involving DPP-4is (mainly including Sitagliptin, Saxagliptin, Vildagliptin, Linagliptin, and Alogliptin) compared with placebo or other hypoglycemic drugs (like Biguanides, Sulfonylureas, Thiazolidinediones, α -glycosidase inhibitors, SGLT-2 inhibitor, and DLP-1 receptor agonist) in patients with T2DM.

Types of studies

Only randomized controlled trials (RCTs) will be enrolled in this review, which compared DPP-4i therapies with other active hypoglycemic medications or placebo in T2DM patients with a minimum intervention length of 4 weeks. Either blinding or open-label studies will be considered. Furthermore, the literature will only be available in English. Other types of studies, such as in vivo or in vitro studies, case reports, reviews, and non-RCTs, will be eliminated.

Outcome measures

The primary outcome of our study is pancreatic safety events, such as pancreatitis and pancreatic cancer. The secondary outcome of interest will include the change of pancreatic enzyme from baseline.

Search strategy

Medline, Embase, Pubmed, Web of Science, the Cochrane Central Register of Controlled Trials (CENTRAL) were searched from their inception dates to August 10th, 2021. Clinical trial registries (such as www.ClinicalTrials.gov) were also searched for unpublished trials. Moreover, we will check the systematic reviews and meta-analyses to search for further relevant studies and contact the authors to obtain incomplete data. Two researchers will perform the search, and all discrepancies will be resolved by discussion with a third reviewer. To find qualifying studies in PubMed, we will use the following search strategy, and it will be tweaked for each database. **Table 1** shows the search strategies and retrieval type which will be employed in this investigation.

Table1 Search strategies for PubMed

NO.	Retrieval type
#1	Diabetes Mellitus, Type 2 [Mesh]
#2	<p>Diabetes Mellitus, Type 2 [Title/Abstract] OR Diabetes Mellitus, Noninsulin-Dependent [Title/Abstract] OR Diabetes Mellitus, Ketosis-Resistant [Title/Abstract] OR Diabetes Mellitus, Ketosis Resistant [Title/Abstract] OR Ketosis-Resistant Diabetes Mellitus [Title/Abstract] OR Diabetes Mellitus, Non Insulin Dependent [Title/Abstract] OR Diabetes Mellitus, Non-Insulin-Dependent [Title/Abstract] OR Non-Insulin-Dependent Diabetes Mellitus [Title/Abstract] OR Diabetes Mellitus, Stable [Title/Abstract] OR Stable Diabetes Mellitus [Title/Abstract] OR Diabetes Mellitus, Type II [Title/Abstract] OR NIDDM [Title/Abstract] OR Diabetes Mellitus, Noninsulin Dependent [Title/Abstract] OR Diabetes Mellitus, Maturity-Onset [Title/Abstract] OR Diabetes Mellitus, Maturity Onset [Title/Abstract] OR Maturity-Onset Diabetes Mellitus [Title/Abstract] OR Maturity Onset Diabetes Mellitus [Title/Abstract] OR MODY [Title/Abstract] OR Diabetes Mellitus, Slow-Onset [Title/Abstract] OR Diabetes Mellitus, Slow Onset [Title/Abstract] OR Slow-Onset Diabetes Mellitus [Title/Abstract] OR Type 2 Diabetes Mellitus [Title/Abstract] OR Noninsulin-Dependent Diabetes Mellitus [Title/Abstract] OR Noninsulin Dependent Diabetes Mellitus [Title/Abstract] OR Maturity-Onset Diabetes [Title/Abstract] OR Diabetes, Maturity-Onset [Title/Abstract] OR Maturity Onset Diabetes [Title/Abstract] OR Type 2 Diabetes [Title/Abstract] OR Diabetes, Type 2 [Title/Abstract] OR Diabetes Mellitus, Adult-Onset [Title/Abstract] OR Adult-Onset Diabetes Mellitus [Title/Abstract] OR Diabetes Mellitus, Adult Onset [Title/Abstract]</p>
	#1 OR #2
	Dipeptidyl-Peptidase IV Inhibitors [Mesh]
#3	<p>Dipeptidyl-Peptidase IV Inhibitors [Title/Abstract] OR Dipeptidyl Peptidase IV Inhibitors [Title/Abstract] OR DPP-4 Inhibitor [Title/Abstract] OR DPP 4 Inhibitor [Title/Abstract] OR Inhibitor, DPP-4 [Title/Abstract] OR DPP-IV Inhibitor [Title/Abstract] OR DPP IV Inhibitor [Title/Abstract] OR Inhibitor, DPP-IV [Title/Abstract] OR DPP-4 Inhibitors [Title/Abstract] OR DPP 4 Inhibitors [Title/Abstract] OR DPP-IV Inhibitors [Title/Abstract] OR DPP IV Inhibitors [Title/Abstract]</p>
#4	<p>OR Gliptin [Title/Abstract] OR Dipeptidyl Peptidase 4 Inhibitor [Title/Abstract] OR Dipeptidyl-Peptidase IV Inhibitor [Title/Abstract] OR Dipeptidyl Peptidase IV Inhibitor [Title/Abstract] OR Inhibitor, Dipeptidyl-Peptidase IV [Title/Abstract] OR Dipeptidyl-Peptidase 4 Inhibitor [Title/Abstract] OR Inhibitor, Dipeptidyl-Peptidase 4 [Title/Abstract] OR Dipeptidyl-Peptidase 4 Inhibitors [Title/Abstract] OR Dipeptidyl Peptidase 4 Inhibitors [Title/Abstract] OR Gliptins [Title/Abstract] OR DPP4 Inhibitor [Title/Abstract] OR Inhibitor, DPP4 [Title/Abstract] OR DPP4 Inhibitors [Title/Abstract] OR Sitagliptin [Title/Abstract] OR Saxagliptin [Title/Abstract] OR Vildagliptin [Title/Abstract] OR Linagliptin [Title/Abstract] OR Alogliptin [Title/Abstract]</p>
#5	<p>OR DPP4 Inhibitors [Title/Abstract] OR Sitagliptin [Title/Abstract] OR Saxagliptin [Title/Abstract] OR Vildagliptin [Title/Abstract] OR Linagliptin [Title/Abstract] OR Alogliptin [Title/Abstract]</p>
	#4 OR #5
	<p>Randomized Controlled Trial [Publication Type] OR RCT [Title/Abstract] OR randomized [Title/Abstract] OR controlled [Title/Abstract] OR placebo [Title/Abstract]</p>
	pancrea* [Title/Abstract] OR safety [Title/Abstract]
	#3 AND #6 AND #7 AND #8
#6	
#7	

#8

#9

Study selection process

Two reviewers will conduct literature screening independently. They will screen titles and abstracts of all the retrieved records to find potentially eligible studies and then examine the entire text to select studies that match the inclusion criteria. After rechecking the source papers, debate among the two reviewers, and adjudicating by the third, disagreements will be resolved with an entire consensus before inclusion. A PRISMA flow diagram will outline the research selection procedure and reasons for exclusions [8], and it was indicated in **Additional file2**. If a study was published in duplicate, we would choose the version with the most detailed content and data.

Data extraction

We will ensure a standardized data extraction template in advance, and two investigators will independently extract the data from all eligible studies in duplicate. Any discrepancies will be resolved by consensus or arbitrated by the third. The following trial information will be extracted: study author, publication year, study design, sample size, types of intervention and control, background therapy, and funding source. Moreover, population characteristics containing mean age, gender proportions, racial, duration of disease, length of the trial, loss of follow-up, and baseline level of HbA1c will also be collected. What is more, the following outcome measures will be gathered: Relative Risk of pancreatic safety event (involving pancreatitis and pancreatic cancer) and post-intervention values or changes from baseline with corresponding standard deviations for the pancreatic enzyme. Additionally, if the supplied data is incomplete, we will contact the authors for more information.

Risk of bias assessment

Two assessors will assess the risk of bias and quality of all included studies, with the third reviewer participating in the discussion as needed. They will pilot numerous samples before the formal evaluation to get to an agreement on assessment standards. Cochrane risk of bias tool (ROB tool) will be used to assess the risk of bias for trials concerning seven points: the judgment of the random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, and other sources of bias, which are graded as low, high or unclear risk of bias[9]. Furthermore, we will try to gather all the information related to this review to control publication bias, and if more than 10 trials are included, we will assess the reporting bias through a funnel plot.

Quality assessment

The collected evidence will be interpreted by the GRADE Working Group approach for rating the quality of treatment effect estimates from NMA. According to GRADE, the assessment requires estimates from direct, indirect, and combined evidence from direct and indirect sources, as well as quality ratings for the direct and indirect comparisons[10], which are primarily used to assess the quality of each piece of evidence in five areas: limitation, imprecision, inconsistency, indirectness, and publication bias.

Data Synthesis and Analysis

A descriptive overview of available data will be compiled and reported, namely trial and population characteristics, interventions, results, and risk of bias evaluations. We will plot a network regarding the evidence structure of direct comparisons, in which the size of nodes will be proportional to the sample size of each intervention, and the thickness of the lines will be proportional to the cumulative number of RCTs for each pairwise comparison. We will also use the contribution plot to present the influence of each direct piece of evidence[11]. Both traditional pairwise meta-analysis and the network meta-analysis will be conducted. All quantitative analyses will be carried out using Stata 16[12] and WinBUGS[13].

Pairwise meta-analyses

A random effects model will be employed for traditional pairwise meta-analyses if the included studies are of heterogeneity. The I^2 value will measure the degree of heterogeneity. Relative Risk (RR) for dichotomous data will be calculated as effect measures with 95% confidence intervals, and continuous data will be presented as weighted mean difference (WMD) with 95% confidence intervals. If different scales are involved in studies, we will use standardized mean differences (SMD) to present continuous data to eliminate its effect on results. Besides, sensitivity analysis will be performed to validate the stability of the results or exclude studies with a high risk of bias.

Network meta-analyses

NMA is an upgrade of classical meta-analysis, which can simultaneously combine direct and indirect analysis, estimate the efficacy of interventions through common comparators even if they have not been investigated head to head in randomized clinical trials, and thus analyze the effects of multiple interventions compared with each other.

Before the analysis, examine the assumptions of consistency, heterogeneity, and similarity of the included studies within and across connections in the entire network of interventions to determine whether direct and indirect evidence is reasonable[14]. An approach of loop inconsistency is used to evaluate the presence of inconsistency in each closed loop in which an intervention effect measured using an indirect comparison is not equivalent to the intervention effect measured using direct comparison. The I^2 statistic will quantify the global heterogeneity, and the Cochran Q test and its P-value will also be used to evaluate the heterogeneity[15]. The predictive interval graph and confidence interval will be plotted to present the influence of heterogeneity on each pairwise comparison. Furthermore, the

similarity will be evaluated by comparing the critical clinical and methodological characteristics that can influence the effects of studies between two sets.

Suppose the above assumptions of consistency, heterogeneity, and similarity are reasonable. In that case, NMA will be performed in a random effects model, by a Bayesian framework using Markov Chains Monte Carlo (MCMC) methods with WinBUGS 1.4.3 and Stata 16 software to synthesize all the available evidence[16]. The results of all pairwise comparisons will be reported as appropriate effect values with 95% confidence intervals. The network geometry of interventions will be plotted to present their concise characteristics, while forest plots and contribution figures will be used to show the combined effect value[17]. Besides, we will display values of the Surface Under the Cumulative Ranking (SUCRA) curve for each intervention as well as rankings of effects[18]. And convergence will be assessed by the Gelman Rubin statistic method and inspection of Monte Carlo errors[16].

Subgroup analyses and meta-regression

We will conduct subgroup analysis and meta-regression to explore the source of heterogeneity if significant heterogeneity and sufficient data exist. Subgroup analyses will combine effect sizes for each subgroup from the perspectives of clinical or methodological heterogeneity, including age, duration of disease, length of the trial, loss of follow-up, the baseline level of HbA1c, and quality of study. And in meta-regression, regression analysis will be used to explore the influence of some covariables on the merger effect in meta-analysis.

Discussion

Dpp-4i is widely used in clinical practice due to its sound hypoglycemic effect and safety, but there are concerns that its use may increase the risk of pancreatic events. Pancreatic events, such as pancreatitis and pancreatic cancer, are rare but real in DPP-4i safety events. Some authorities, such as the European Medicines Agency and Food and Drug Administration, issued a warning on pancreatitis for all available DPP-4is[19], but it has not yet been contraindicated. Some clinical studies suggest a significant association exists between pancreatic events and DPP-4i, and Dpp-4i using is related to increased risks of pancreatic cancer in patients with type 2 diabetes[20-22]. The results of some systematic reviews and meta-analyses also support this viewpoint[23-25]. Nevertheless, most of the present data have not suggested any negative trend[19,26-28], although they are insufficient to draw reliable conclusions. Generally, the correlation observed in trials, whether positive or negative, could be owing to low incidence, limited sample size, chance, or bias but merits further investigation. More importantly, in addition to the hazards of pancreatic events, the health status of the pancreas is strongly related to the development of T2DM, any possibility of increased risk on the pancreas should be taken seriously.

The current studies have not concluded the association between DPP-4i and pancreatic safety. Both the European Medicines Agency and Food and Drug Administration indicate that pancreatic events will continue to be considered a risk associated with DPP-4i until more data are available[29]. Among numerous analytical methods, NMA can synthesize all available data for statistical processing through

quantitative synthesis to further guide clinical decisions. In this way, our study will analyze and compare the pancreatic outcomes of different DPP4i treatments for T2DM, focusing on randomized controlled trials that could ensure the quality of studies to some extent, although it may limit the number of studies included. There are few NMAs in this direction presently, and most of them are typical pairwise meta-analyses. Therefore, it is meaningful to gather various direct and indirect evidence as comprehensively as possible to provide reliable recommendations for pancreatic safety and clinical medication of DPP4i.

Abbreviations

DPP-4i: Dipeptidyl-peptidase IV inhibitor; T2DM: type 2 diabetes mellitus; GLP-1: glucagon-like peptide-1; GIP: gastric inhibitory polypeptide; NMA: Network meta-analysis; RCTs: randomized controlled trials; PRISMA-P: Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols; PRISMA: Preferred Reporting Items for Systematic review and Meta-Analysis; ROB: Risk of bias; RR: Relative Risk; WMD: weighted mean difference; SMD: Standardized mean differences; MCMC: Markov Chains Monte Carlo; SUCRA: Surface Under the Cumulative Ranking

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Not applicable

Competing interests

The authors declare that they have no competing interests.

Funding

This research was supported by the National Natural Science Foundation of China. (NO. 81774254)

Authors' contributions

FY and **QZ** contributed to the design of this study. **FY, QZ, YD, BL,** and **BN** drafted this **protocol**. **FY** and **YD** provided professional advice on evidence-based medicine, and **QZ** provided clinical expertise. **BL** and **BN** will be in charge of searching, screening, collection, and quality assessment. **FY, YD, BL,** and **BN** will be involved in data analysis. All authors read and approved the final manuscript.

Acknowledgements

Not applicable

Author details

¹The First Clinical Medical College, Shandong University of Traditional Chinese Medicine, Jinan, Shandong, China.

²Division of Comprehensive Internal Medicine, Shandong University of Traditional Chinese Medicine Affiliated Hospital, Jinan, Shandong, China.

*Correspondence: zhaoquanlin65@163.com

Fan Yang¹, Email: yangfan_viola@163.com

Quanlin Zhao^{2*}, Email: zhaoquanlin65@163.com

Youzi Dong¹, Email: Dong_youz@163.com

Baohua Li¹, Email: 10728178@qq.com

Bobiao Ning¹, Email: 4936752@qq.com

References

1. Andersen DK, Andren-Sandberg Å, Duell EJ, et al. Pancreatitis-diabetes-pancreatic cancer: summary of an NIDDK-NCI workshop. *Pancreas*. 2013;42:1227-1237.
2. Matveyenko Aleksey V, Dry Sarah, Cox Heather I et al. Beneficial endocrine but adverse exocrine effects of sitagliptin in the human islet amyloid polypeptide transgenic rat model of type 2 diabetes: interactions with metformin. *Diabetes*. 2009;58:1604-15.
3. Buse John B, Bethel M, Angelyn, Green Jennifer B et al. Pancreatic Safety of Sitagliptin in the TECOS Study. *Diabetes Care*. 2017;40:164-170.
4. Tseng Chao-Ming, Liao Wei-Chih, Chang Chi-Yang et al. Incretin-based pharmacotherapy and risk of adverse pancreatic events in the ethnic Chinese with diabetes mellitus: A population-based study in Taiwan. *Pancreatology*. 2017;17:76-82.
5. Shamseer Larissa, Moher David, Clarke Mike et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*. 2015;350:g7647.
6. Hutton Brian, Salanti Georgia, Caldwell Deborah M et al. The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Ann Intern Med*. 2015;162:777-84.

7. World Health Organization. Definition and diagnosis of diabetes mellitus and intermediate hyperglycemia: report of a WHO/IDF consultation. Geneva: WHO Document Production Services; 2006.
8. Moher David, Liberati Alessandro, Tetzlaff Jennifer et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6:e1000097.
9. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.2 (updated February 2021). Cochrane, 2021. Available from www.training.cochrane.org/handbook.
10. A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis. *BMJ*. 2015;350:h3326.
11. Chaimani Anna, Higgins Julian P T, Mavridis Dimitris et al. Graphical tools for network meta-analysis in STATA. *PLoS One*. 2013;8:e76654.
12. Stata Statistical Software: Release 16. 2019.
13. Lunn DJ, Thomas A, Best N, Spiegelhalter D. WinBUGS—a Bayesian modelling framework: concepts, structure, and extensibility. *Stat Comput*. 2000;10
14. Cipriani Andrea, Higgins Julian P T, Geddes John R et al. Conceptual and technical challenges in network meta-analysis. *Ann Intern Med*. 2013;159:130-7.
15. Peter Emanuel L, Deyno Serawit, Mtewa Andrew et al. Safety and efficacy of *Momordica charantia* Linnaeus in pre-diabetes and type 2 diabetes mellitus patients: a systematic review and meta-analysis protocol. *Syst Rev*. 2018;7:192.
16. Dias Sofia, Sutton Alex J, Ades A E et al. Evidence synthesis for decision making 2: a generalized linear modeling framework for pairwise and network meta-analysis of randomized controlled trials. *Med Decis Making*. 2013;33:607-17.
17. Mills Edward J, Thorlund Kristian, Ioannidis John P A, Demystifying trial networks and network meta-analysis. *BMJ*. 2013;346:f2914.
18. Salanti Georgia, Ades A E, Ioannidis John P A, Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *J Clin Epidemiol*. 2011;64:163-71.
19. Dicembrini Ilaria, Monterecci Chiara, Nreu Besmir et al. Pancreatitis and pancreatic cancer in patients treated with Dipeptidyl Peptidase-4 inhibitors: An extensive and updated meta-analysis of randomized controlled trials. *Diabetes Res Clin Pract*. 2020;159:107981.
20. Singh Awadhesh Kumar, Gangopadhyay Kalyan Kumar, Singh Ritu, Risk of acute pancreatitis with incretin-based therapy: a systematic review and updated meta-analysis of cardiovascular outcomes trials. *Expert Rev Clin Pharmacol*. 2020;13:461-468.
21. Zhang Zeqing, Chen Xi, Lu Puhuan et al. Incretin-based agents in type 2 diabetic patients at cardiovascular risk: compare the effect of GLP-1 agonists and DPP-4 inhibitors on cardiovascular and pancreatic outcomes. *Cardiovasc Diabetol*. 2017;16:31.

22. Tkáč Ivan,Raz Itamar,Combined Analysis of Three Large Interventional Trials With Gliptins Indicates Increased Incidence of Acute Pancreatitis in Patients With Type 2 Diabetes. *Diabetes Care*. 2017;40:284-286.
23. Lee Minyoung,Sun Jiyu,Han Minkyung et al. Nationwide Trends in Pancreatitis and Pancreatic Cancer Risk Among Patients With Newly Diagnosed Type 2 Diabetes Receiving Dipeptidyl Peptidase 4 Inhibitors. *Diabetes Care*. 2019;42:2057-2064.
24. Knapen Lotte M,de Jong Roy G P J,Driessen Johanna H M et al. Use of incretin agents and risk of acute and chronic pancreatitis: A population-based cohort study. *Diabetes Obes Metab*. 2017;19:401-411.
25. Lai Yun-Ju,Hu Hsiao-Yun,Chen Hsin-Hua et al. Dipeptidyl Peptidase-4 Inhibitors and the Risk of Acute Pancreatitis in Patients With Type 2 Diabetes in Taiwan: A Population-Based Cohort Study. *Medicine (Baltimore)*. 2015;94:e1906.
26. Abd El Aziz Mirna,Cahyadi Oscar,Meier Juris J et al. Incretin-based glucose-lowering medications and the risk of acute pancreatitis and malignancies: a meta-analysis based on cardiovascular outcomes trials. *Diabetes Obes Metab*. 2020;22:699-704.
27. Kim Young-Gun,Kim Seirhan,Han Seung Jin et al. Dipeptidyl Peptidase-4 Inhibitors and the Risk of Pancreatitis in Patients with Type 2 Diabetes Mellitus: A Population-Based Cohort Study. *J Diabetes Res*. 2018; 2018:5246976.
28. Hong Jin-Liern,Buse John B,Jonsson Funk Michele et al. The Risk of Acute Pancreatitis After Initiation of Dipeptidyl Peptidase 4 Inhibitors: Testing a Hypothesis of Subgroup Differences in Older U.S. Adults. *Diabetes Care*. 2018;41:1196-1203.
29. Egan Amy G,Blind Eberhard,Dunder Kristina et al. Pancreatic safety of incretin-based drugs—FDA and EMA assessment. *N Engl J Med*. 2014;370:794-7.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1PRISMAPchecklist.docx](#)
- [Additionalfile2PRISMAFlowDiagram.pdf](#)