Physiological changes in chicken embryos inoculated with drugs and viruses highlight the need for more standardisation of this animal model

Simone Sommerfeld¹, Antonio Vicente Mundim¹, Rogério Reis Silva¹, Jéssica Santos Queiroz¹, Maisa Paschoal Rios¹, Fabiana Oliveira Notário¹, Alessandra Aparecida Medeiros¹, Marcelo Emílio Beletti², Rodrigo Rodrigues Franco³, Foued Salmen Espindola³, Luiz Ricardo Goulart³, Belchiolina Beatriz Fonseca¹

¹School of Veterinary Medicine, Federal University of Uberlândia, Uberlândia 38402-018, Brazil; antoniomundim@ufu.br (A.V.M.); rogerioreissilva98@gmail.com (R.R.S.); jesk.queiroz@hotmail.com (J.S.Q.); maisapaschoal@hotmail.com (M.P.R.); fabiana0394@gmail.com (F.O.N.); alessandra.medeiros@ufu.br (A.A.M.); biafonseca@ufu.br (B.B.F.). ²Institute of Biomedical Sciences, Federal University of Uberlândia, Uberlândia 38405-319; belettim@gmail.com (M.E.B.). ³Institute of Biotechnology, Federal University of Uberlândia, Uberlândia 38405-319, Brazil; rodrigorf@yahoo.com.br (R.R.F.); foued@ufu.br (F.S.E.); lrgoulart@ufu.br (L.R.G.)

* Correspondence: s_sommerfeld@hotmail.com

Abstract

Background: Several works have been developed using the *Gallus gallus* embryo as an experimental model to study the toxicity of drugs and infections. Studies that seek to standardise the evaluated parameters are needed to better understand and identify the viability of chicken embryos (CE) as an experimental model. Therefore, we sought to verify whether macroscopic, histopathological, blood count, metabolites and/or enzymes changes and oxidative stress in CE of different ages are peculiar to the model. To achieve this goal, *in ovo* assays were performed injecting a virus (*Gammacoronavirus*) and two drugs (filgrastim and dexamethasone) that cause known changes in adult chickens and other animals. **Results:** Embryo response to virus and drug challenges may not occur as expected for adult chickens and even different species. While macroscopic and microscopic damage was visible in the case of viral

infections, the white blood cell count and inflammation biomarkers did not change. Filgrastim (FG) testing did not result in the expected effects for CE. On the other hand, with dexamethasone (DX), changes in blood parameters and biomarkers were inherent to the model and depended on the stage of CE development. **Conclusions:** Our work reinforces the importance of standardization and correct use of the model so that the results of infection, toxicity and pharmacokinetics are recorded.

Keywords: chorioallantoic membrane, *gammacoronavirus*, toxicity

1. Background

The embryonated egg is self-sufficient, and its natural development at 37 °C and 60% humidity guarantees the maintenance of these animals without more complex means of support. Furthermore, within the egg, the embryo is a highly controlled, accessible and relatively transparent model in which normal physiology, disease pathology and the consequences of experimental manipulation can be easily visualized(1), allowing high reproducibility and simplified and economical experiments(2).

Numerous works have been carried out using the *Gallus gallus* embryo as an experimental model for evaluating the toxic effects of drugs and infections. These studies usually evaluated blood, allantoic and amniotic fluid biochemical parameters, erythrocyte morphology, oxidative stress and histopathological lesions(3–7). As well as in the evaluation methods and collected materials, there is much variation in the ages and inoculation routes, among which the shell membrane, the allantoic fluid, the chorioallantoic membrane (CAM), the yolk sac and the amnion are widely used(7–11).

Chicken embryo (CE) development is completed in just 21 days, being the fastest among the most used animal models and equal only to the mouse embryo(12). Thus, on each day of incubation age, many physiological changes can significantly

influence the studies and responses being evaluated, making the definition of the day of inoculation and collection crucial for realising an excellent experimental study.

The standardisation of parameters, such as the day of inoculation and collection, the inoculation route used, the biological materials collected and the types of analyses performed are crucial for an experiment's success. Studies that seek the standardisation of such parameters are needed to better understand and identify the viability of CE and *in ovo* studies as models for preclinical tests. Thus, this study aims to verify if the results of viability, weight, pathological and histopathological changes, blood count, dosage of metabolites and/or enzymes and oxidative stress in CE of different ages are peculiar to the model, proposing and discussing the need for standardisation for embryos, using as a reference a virus and two drugs that cause known changes in adults and/or other species.

2. Methods

This research was performed in the following laboratories of the Uberlândia Federal University: Poultry Egg Incubation, Nanobiotechnology, Biochemistry, Molecular Biology, Animal Pathology, Veterinary Clinical, and Rodent Vivarium Network. The studies were divided into different phases.

2.1. Challenge of CE with *Gammacoronavirus*

The first part of the experiment was carried out in specific pathogen-free (SPF) eggs of *Gallus gallus* challenged with a *gammacoronavirus*, the infectious bronchitis virus (IBV) that causes embryonic chicken lesions. The CE were incubated in an artificial incubator (Premium Ecológica®) at 37 °C, 58% humidity, with the eggs being turned at a two-hour interval. The eggs were weighed and inoculated at 10 embryonic incubation days (EID). This age was used as between 9–10 EID, lesions in CE can be

distinguished(13). A total of 4.35 log viral particles/CE/of IBV (strain Massachusetts, H52), diluted in sterile Phosphate-buffered saline (PBS), were inoculated in allantoic liquid from 10 CE. In parallel, four embryos were inoculated as negative controls (NC) with sterile PBS (diluent of the virus).

After 24 hours of incubation, the CE were evaluated to remove the non-specific mortalities. At 17 EID, 0.5mL of allantoic liquid was collected. Blood was collected by sectioning the umbilical vessels and divided into two microtubes: one with EDTA K3 for blood cell analysis and the other with clot activator for serum analysis. In the allantoic fluid and serum, the minerals, metabolites and enzymes there were quantified. After collecting the blood, the CE were immediately euthanised, weighed and evaluated for macroscopic changes. The livers were collected, stored in formalin for histopathological tests and in liquid nitrogen for oxidative stress evaluation.

2.2. Inoculation with dexamethasone and filgrastim

2.2.1. Drug dosages: pilot test

Prior to commencing the study, we carried out a pilot test to determine the dose of each drug. In this part of the experiment, we used a commercial line of CE, Hy-Line W36, as SPF birds only are mandatory for research with viruses. The eggs were incubated from zero EID in conditions identical to those mentioned in item 2.1. There is no standardised allometric extrapolation calculation for CE. Therefore, we usually test the adult chicken dose in 10 EID CE with an average weight of 20 g (CE and embryonic annexes). The intention was to determine the dose that would not kill the embryo.

In the pilot test, we tested two routes: shell membrane (SM) and chorioallantoic membrane (CAM). We used an approximate dose for a hatched animal

as the basis. The 2–4 mg/Kg doses were used for DX(14). According to the drug manufacturer, there is no indication of FG for birds, so we based it on human doses (5 mg/Kg). Therefore, the following doses were tested: 4, 0.4 and 0.08 µg/CE of DX and 150, 15 and 1.5 µg/CE of FG, with three CE per dose via SM and CAM plus the NC for each route, totaling 42 embryos. After seven days, at 17 EID, the CE were evaluated for macroscopic changes. The dose chosen was based on the group with no expressive mortality or severe lesions in the embryos. For DX, the dose of 0.08 µg/CE did not cause gross lesions in any CE, but the 0.4 and 4 µg/CE resulted in green allantoic fluid in CE inoculated via SM and CAM. Doses of 1.5, 15 and 150 µg/CE of FG did not alter the CE. Therefore, the used dose of DX was 0.08 µg/CE (~4 µg/Kg) and FG was 150 µg/CE (~7.5 µg/Kg). As the results for embryos inoculated via CAM and SM were identical, in the first moment, the SM route was used as it was the easiest, with a lower risk of death or lesions, and is an essential route for young or old embryos (which were used in this work).

2.2.2. Drug tests in CETest of the drugs in CE

We started the experiment using CE of 12 EID (Hy-Line W36), as they are older CE and their organs and physiologies more mature than those of younger CE. The egg incubation conditions were identical to those mentioned in item 2.1. A total of 18 CE were used with six CE in the following groups inoculated via SM: (i) treated with 0.08 µg/CE of DX; (ii) treated with 150 µg/CE of FG; and (iii) NC, inoculated with water (diluent of drugs) only. The CE were treated at intervals of 24 hours for three days. At 12 EID, the eggs were weighed. After seven days post-inoculation (pi), at 19 EID, we carried out blood and liver collection. We evaluated the weight and macroscopic

1 lesions similar to those described in item 2.1, except for allantoic fluid, as the amount
2 of this fluid was very low at this age.

3 We also performed inoculations with FG (150 µg/CE) and DX (0.08 µg/CE)
4 and a NC group at intervals of 24 hours for three days, with six or eight eggs per group,
5 at ages zero, three and seven EID via SM. After nine, 11 and 10 days of incubation,
6 the CE inoculated at zero, three and seven EID, respectively, were euthanised,
7 weighed and evaluated regarding mortality and macroscopic lesions. We did not
8 conduct blood, metabolites or microscopy analysis of the CE inoculated at zero, three
9 and seven EID.

10 Another assay was performed in CE at 10 EID. In this part of the experiment,
11 the route of inoculation for DX was SM. As there were no changes, as expected, in
12 blood granulocyte counts in CE inoculated at 12 EID with FG, and there were no
13 changes in young CE, the CAM route was used for FG in CE of 10 EID. As the FG
14 pathway in humans is subcutaneous and the CAM pathway in CE has direct access to
15 the vessels, we hypothesised that this could be an alternative to test this drug. Thus,
16 it could be possible to assess whether the route could interfere with the results for FG.
17 This part of the experiment was carried out twice, with a total of 18 CE each time,
18 totalling 36 CE. The groups were divided as follows: (i) CE inoculated with 0.08 µg/CE
19 of DX via SM; (ii) NC of DX – CE inoculated with water via SM; (iii) CE inoculated with
20 150 µg/CE via CAM; (iv) NC of FG – CE inoculated with water via CAM. The CE were
21 treated at intervals of 24 hours for 3 days. After seven days pi, at 17 EID, blood,
22 allantoic fluid, liver weight and macroscopic lesions were collected, similar to that
23 described in item 2.1. The experiment was repeated at an interval of 15 days, identical
24 to the previous experiment.

2.3. Weight of the CE and annexes

During the experiments, the eggs were numbered and weighed on the first day of inoculation and their weights recorded. Then, the embryo and yolk were weighed immediately after collecting blood and allantoic fluid on the day of collection. As the embryo weight is related to the initial egg weight, we performed an adjustment for an initial egg weight of 50 grams, according to Ribeiro et al. (2020)(7)

2.4. Macroscopic evaluations

We checked and counted the CE that died and identified the death date based on embryo development degree. For live animals, we observed their annexes for the presence of circulatory changes, malformation and colour changes. We also checked the embryos by external evaluation and evaluated the internal organs for circulatory changes, malformation and colour changes and compared the treated groups with the animals in their respective control group.

2.5. Blood cell count, haematocrit, haemoglobin and erythrocyte indices

To determine the haematocrit value (Ht), we filled capillary tubes up to 2/3 with a blood sample. We centrifuged them at 12,000 g for 5 minutes for later reading on a microhaematocrit scale.

Haemoglobin (Hb) concentration was measured using the cyanmethemoglobin method based in Collier (1944)(15) with modifications using Drabkin's solution and reading by spectrophotometry at an absorbance of 540 nm. The resulting colour is of an intensity proportional to the haemoglobin content in the blood. In parallel, we measured the haemoglobin by calculating 1/3 of the haematocrit value.

The total red blood cell (RBC) count was obtained using the Natt and Herrick (1952)(16) solution. Therefore, we performed a manual count in a Neubauer chamber for the haemocytometer method using blood dilution of 1:200. Counting was done in the five diagonal squares of the central reticulum of the chamber. The result was multiplied by 10,000 to obtain the value of erythrocytes per microliter of blood(17).

The total white blood cell count and thrombocytes were also determined using the haemocytometer method by diluting the blood with the Natt and Herrick (1952)(16) solution. The leucocytes and thrombocytes were counted simultaneously in the four external reticula of the Neubauer chamber. The result was multiplied by 500 to obtain the value of leucocytes and thrombocytes per microliter of blood(17). Two different people counted the cells to confirm the precision of the result. The characteristics of the cells (thrombocytes and eosinophils) were confirmed by cytochemistry.

We calculated the mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH) and mean corpuscular haemoglobin concentration (MCHC) using the formulas(18):

$$MCV = \frac{(Hc \times 10)}{RBC} \quad MCH = \frac{(Hb \times 10)}{RBC} \quad MCHC = \frac{(Hb \times 100)}{Hc}$$

2.6. Characterisation of blood cells

The blood smear slides were prepared for staining with fast panoptic dye for differential leucocyte counts under an optical microscope (Olympus CX31, 100x immersion oil). Slides were also stained using the Periodic Acid Schiff (PAS) cytochemical technique for identifying thrombocytes and Sudan Black B (SBB) for identifying eosinophils(18). Finally, the cells were counted twice at different moments under an optical microscope.

Knowing that serotonin binds to gaseous formaldehyde, we carried out the marking of thrombocytes. We based this on Swayne et al. (1986)(19) with some

modifications. First, blood smear slides were placed in a box containing formalin and kept for 24 hours at 50 °C. Next, we read the slides under a UV light microscope (EVOS FL Cell Imaging System, Life Technologies Corporation, USA). Thrombocyte diameter was measured using the ImageJ morphometry program. Slides from healthy and adult chickens (24 weeks of age) of the same line were prepared in parallel as a positive control.

2.7. Biochemical analyses of the serum and allantoic liquid

As the collection of blood from the embryo is not simple and some researchers perform analysis of metabolites and enzymes in allantois, we performed the analysis in blood and allantois to determine if the results were similar. Biochemical analysis of serum and allantoic fluid was performed in an automatic biochemical analyser (ChemWell® 2910, Awareness Technology). The analytes aspartate aminotransferase (AST), alanine aminotransferase (ALT), gamma-glutamyl transferase (GGT), alkaline phosphatase (ALP), creatinine (Creat), uric acid (UA), calcium (Ca), phosphorus (P) and c-reactive protein (CRP) (Bioclin®, Minas Gerais, Brazil) were measured in both serum and allantoic fluid.

2.8. Oxidative stress

Immediately after euthanasia, the livers of the CE were removed and stored at -80 °C. The samples were homogenised with 10 mM sodium phosphate buffer (pH

7.4) and centrifuged at $800 \times g$, at 4°C for 10 min. The supernatant was used to quantify the biomarkers of oxidative stress.

2.8.1. *Reactive oxygen species (ROS)*

The samples were incubated with dichloro-dihydro fluorescein diacetate ($10\text{ }\mu\text{M}$) and 5 mM Tris-HCl buffer (pH 7.4) for 3 min. Subsequently, the fluorescence was measured at 530 nm (excitation in 474 nm).

2.8.2. *Lipid peroxidation*

The liver homogenates were incubated with 0.67% thiobarbituric acid (TBA) and 10% trichloroacetic acid (TCA), for 120 min. Then, n-butanol was added to the samples to remove the organic phase and the fluorescence was measured at 553 nm, after excitation at 515 nm. Lipid peroxidation was determined using the malondialdehyde (MDA) analytical curve(20).

2.8.3. *Sulphydryl group*

The sulphydryl group was detected using ditionitrobenzoic acid (DTNB) diluted in 0.2 mM potassium phosphate buffer (pH 8.0). The liver homogenates were incubated for 30 min with 1 mM phosphate buffer (pH 7.4) and 10 mM DTNB solution. The presence of sulphydryl groups was spectrophotometrically detected at 412 nm(21).

2.8.4. *Total antioxidant capacity*

The liver homogenates were incubated with 300 mM sodium acetate buffer (pH 3.6), 10 mM 2,4,6-tri (2pyridyl) -stiazine (TPTZ) and 20 mM ferric chloride at 37°C for

6 min, at 593 nm. A trolox analytical curve determined the antioxidant capacity and sodium acetate buffer was used as a blank(22).

2.9. Histopathology

The fragments of liver fixed in 10% buffered formalin were processed for the preparation of histological slides stained with Haematoxylin and Eosin (HE)(23). Histopathologic morphology was qualitatively assessed and classified for the presence of inflammation, degeneration, necrosis and circulatory change.

Intracellular fat deposits (lipidosis) were evaluated semiquantitatively and classified by assigning a score relative to the level of lipid deposits in the sample, according to the histological classification of Brunt et al. (1999)(24) modified by Angulo (2009)(25) with adaptations.

According to Fáncsi (1982)(26), the cytoplasm of hepatocytes from CE has drops of lipids, and the amount of lipid inclusions increases mainly around the twenty-seventh day. Thus, the presence of small vacuoles in the cytoplasm of hepatocytes was considered physiological in the assessment of lipidosis. The scores were evaluated according to the NC and divided into: 0 corresponding to normal, with discrete lipid deposits; 1 with slight lipid deposits; 2 with moderate lipid deposits; 3 with a marked level of lipid deposition.

In the analysis of the inflammatory process, the number of inflammatory foci and the number of inflammatory cells in the foci were microscopically evaluated in all the livers of the embryos according to the NC. The inflammatory process was classified as: 0 corresponding to normal, with no inflammatory process; 1 – mild, when the number of inflammatory foci was less than three with mild inflammatory infiltrate; 2 – moderate, when the number of inflammatory foci ranged from three to five with

moderate inflammatory infiltrate; and 3 – marked, when the number of inflammatory foci was greater than five with marked inflammatory infiltrate.

Regarding circulatory alterations, haemorrhage and congestion, they were evaluated semiquantitatively in relation to the NC as: 0 corresponding to normal, with no circulatory alterations; 1 – light; 2 – moderate; 3 – accentuated.

2.10. Angiogenesis

While no injury was observed in CE treated with FG at any dose or route and we noticed an increase in blood vessels in some eggs treated with FG. Therefore, we performed another experiment adding five and 25 times the dose of FG used in the previous investigation. The dose increase was to assess whether doses higher than those used in humans were capable of causing visible damage to embryos and test the hypothesis that FG induces angiogenesis. We tested 34 CE (Hy-Line W36) at three EID and divided into the following groups: (i) treated with 0.08 µg/CE of DX; (ii) treated with 150 µg/CE of FG; (iii) treated with 750 µg/CE of FG; (iv) treated with 3.75 mg/CE of FG; (v) NC, inoculated with water (diluent of drugs) only, all by the SM route. The CE were treated at intervals of 24 hours for days. At 12 EID, we evaluated the viability and macroscopic lesions. We evaluated the angiogenesis in CAM using ImageJ software with the Vessels Analysis(27) plugin.

2.11. Statistical analysis

We assessed whether the data followed normality. For parametric data, we used Analysis of variance (ANOVA) followed by Tukey's test and for non-parametric data, we used the Kruskal–Wallis test. For evaluation of two groups, we used the t test for parametric data and Kruskal–Wallis or Wilcoxon tests for non-parametric data.

Person's correlation test was performed. We considered a 95% confidence interval using the program, GraphPad Prism 9.2.

3. Results

3.1. The virus can change blood cell count, calcium and lipid peroxidation in the liver

Of the 10 CE inoculated with *gammacoronavirus*, three died, six had lesions and one survived without lesions. The injuries, as expected, were dwarf or curled embryos with a green liver in one CE, five tiny embryos (being 2 CE with a haemorrhagic or enlarged liver, and 1 CE with milky allantoic fluid). The presence of the virus led to a change in the weight of live embryos, as expected. Embryos inoculated with the virus had an average weight of 13.41 g (± 2.86), while the NC weighed 18.17 g (± 1.62) with $p = 0.0173$. We could not weigh the yolk sac of embryos inoculated with the virus because they disintegrated during manipulation. We performed a blood cell count, and the results are described in Table 1.

Table 1. Blood cell counts and quantification of metabolites, minerals and enzymes in CE treated with *Gammacoronavirus* inoculated at 10 EID.

	NC (blood or serum)*	Virus (blood or serum)*	NC (AL)	Virus (AL)
Ht (%)	21.00 (± 1.63) ^a	15.60 (± 2.30) ^b	-	-
Hg (g/dL)	7.00 (± 0.55) ^a	5.20 (± 0.77) ^b	-	-
(1/3 Ht)				
Hg (g/dL)	8.05 (± 0.16) ^a	6.23 (± 0.99) ^b	-	-
(Cyanmethemoglobin)				
Erythrocytes $\times 10^6/\text{mm}^3$	2.05 (± 0.15) ^a	1.59 (± 0.23) ^b	-	-
MCH (pg)	35.15 (± 3.47) ^a	40.08 (± 9.95) ^a		

MCV (fL)	38.15 (+/-3.47) ^a	40.08 (+/-9.95) ^a		
MCHC (g/dL)	96.96 (+/-17.07) ^a	100.6 (+/-27.12) ^a		
Thrombocytes	2.21 (+/-0.51) ^a	2.75 (+/-2.05) ^a	-	-
x10 ⁴ /mm ³				
Leucocytes x10 ³ /mm ³	8 (+/-2.67) ^a	20.4(+/-22.5) ^a	-	-
Monocytes/mm ³	28 (+/-55) ^a	254 (+/-465.5) ^a	-	-
Lymphocytes/mm ³	650 (+/-730) ^a	2170 (+/-2943) ^a	-	-
Heterophiles x10 ³ /mm ³	7.29 (+/-2.27) ^a	17.6 (+/-18.8) ^a	-	-
Heterophile/lymphocyte	24.09 (+/-18.53) ^a	28.32 (+/-20.51) ^a		
Eosinophils/mm ³	00.00(+/-00.00) ^a	00.00(+/-00.00) ^a	-	-
Unidentified	23.75(+/-47.5) ^a	259(+/-459.1) ^a	-	-
granulocytes/mm ³				
Basophils/mm ³	00.00(+/-00.00) ^a	00.00(+/-00.00) ^a	-	-
UA (mg/dL)	21.76 (+/-23.13) ^a	22.63 (+/-15.17) ^a	18.53 (+/-21.53) ^a	54.30 (+/-29.89) ^a
Creat (mg/dL)	1.42 (+/-1.21) ^{ab}	0.50 (+/-0.35) ^a	3.21 (+/-1.41) ^b	1.55 (+/-1.11) ^{ab}
ALP (U/L)	1857 (+/-1108) ^a	2258 (+/-1540) ^a	80.40 (+/-26.37) ^b	55.74 (+/-49.27) ^b
GGT (U/L)	81.53 (+/-55.34) ^a	252.4 (+/-121.5) ^a	210.00	184.00 (119.40) ^a
			(+/-172.2) ^a	
AST (U/L)	264.00 (+/-40.84) ^a	451.00 (+/-281.1) ^a	200.00	592.00 (+/-283.10) ^a
			(+/-123.3) ^a	
ALT (U/L)	57.50 (+/-233.6) ^a	92.00(+/-85.73) ^a	63.00 (+/-38.1) ^a	48.00 (+/-22.80) ^a
CRP (mg/L)	24.55 (+/-10.31) ^a	24.75 (+/-9.91) ^a	59.50 (+/-32.55) ^a	84.00 (+/-45.72) ^a
Ca (mg/dL)	95.23 (+/-66.18) ^a	10.23 (+/-8.08) ^b	16.05 (+/-10.48) ^b	16.60 (+/-13.77) ^b
P (mg/dL)	6.82 (+/-3.70) ^a	5.50 (+/-2.96) ^a	19.19 (+/-8.57) ^b	10.76 (+/-6.32) ^{ab}

1

2 **Cell count: blood; Quantification of metabolites, enzymes and minerals: Serum and allantoinic liquid.** The value in

3 parentheses is the standard deviation. Different letters on the same line show statistical difference. Monocytes, MCV: non-

4 parametric test. Unidentified granulocytes: cells not tagged in Sudan Black B or PAS but not having a standard format. NC:

5 Negative control. AL: Allantoinic liquid. Ht: Hematocrit. Hg: Hemoglobin. MCH: corpuscular volume. MCH: mean corpuscular

6 haemoglobin. MCHC: mean corpuscular haemoglobin concentration. UA: Uric Acid. Creat: Creatinine. ALP: Alkaline Phosphatase.

7 GGT: gamma-glutamyl transferase AST: aspartate aminotransferase. ALT: alanine aminotransferase. Ca: calcium. P: phosphorus

8 and CRP: c-reactive protein.

Figure 1. Levels of oxidative stress biomarkers in terms of ROS production (A), FRAP (B), lipid peroxidation (C) and sulphydryl groups (D) in liver of CE after infection with *Gammacoronavirus*. FRAP: ferric reducing antioxidant power. ROS: reactive oxygen species. NC: Negative control.

In the histopathological analysis of the liver, congestion and haemorrhage were observed in CE infected with the virus. There was also a moderate inflammatory process predominantly composed of heterophils (Table 2, Figure 2).

Table 2. Histopathological analysis of liver from CE infected with *Gammacoronavirus*.

	NC	Virus
Inflammation	0.00 ^a (Mi: 0.00; Ma: 0.00)	2.00 ^b (Mi: 0.00; Ma: 2.00)
Degeneration	0.00 ^a (Mi: 0.00; Ma: 0.00)	0.00 ^a (Mi: 0.00; Ma: 0.00)
Necrosis	00.00 ^a (Mi: 0.00; Ma: 0.00)	0.00 ^a (Mi: 0.00; Ma: 3.00)
Circulatory change	0.00 ^a (Mi: 0.00; Ma: 0.00)	1.00 ^b (Mi: 0.00; Ma: 3.00)

Mi: Minimum; Ma: Maximum. NC: Negative Control.

Figure 2. Histopathological analysis of CE liver infected with *Gammacoronavirus* at 10 EID. A. NC sample showing sinusoids with a minimal number of erythrocytes (arrow). B. NC sample without perivascular inflammatory infiltrates (arrow). C. Liver of virus-infected CE with congestion characterised by erythrocyte-filled sinusoids (arrows). D. Liver of virus-infected CE with inflammatory infiltrate (arrow). The bars in images A and C represent 200 μm and the bars in images B and D represent 60 μm .

3.2. The age of inoculation and type of drug result in different effects on injury, mortality or embryo viability

The CE were inoculated with the same dose of drugs at 0, 3, 7, 10 and 12 EID for three days. After inoculation, the CE were evaluated daily by light candling, and the dead CE necropsied and removed. Between 7–12 days of inoculation, the eggs were opened and necropsied (Table 3).

Table 3. Evaluation of CE's viability, lesions and embryonic mortality when treated with filgrastim and dexamethasone at different ages.

	Zero EID			3 EID			7 EID			10 EID		10 EID		12 EID		
										CAM		SM				
	NC	FG	DX	NC	FG	DX	NC	FG	DX	NC	FG	CN	DX	NC	FG	DX
Alive (normal)	6	5	0	6	6	1	6	6	0	5	6	6	3	6	6	3
Injured	0	0	2	0	0	5	0	0	3	1	1	0	6	0	0	3
Dead	1	1	4	0	0	0	0	0	3	0	0	0	0	0	0	0
Total	6	6	6	6	6	6	6	6	6	6	7	6	9	6	6	6

Note: eggs broken or killed by the inoculation process were removed from the analysis. CE inoculated at 0, 3, 7, 10 and 12 EID

were evaluated at 9, 11, 10, 17 and 19 EID, respectively. EID: Embryonic incubation days. CAM: Chorioallantoic membrane. SM:

Shell membrane. NC: Negative control. FG: Filgrastim. DX: Dexamethasone.

Of the dead embryos inoculated at zero EID with DX, two were haemorrhagic and died between three to four EID; two died at nine EID and had malformed heads, and two survived but had malformed heads. The only dead CE treated with FG died at nine EID without a specific injury. At three EID, the CE inoculated with DX had the CAM not wholly formed around the CE, and the yolk sac leaked very quickly. Dead CE inoculated at seven EID with DX presented a smaller size, gelatinous albumen,

1 irregular warping and a green liver, and died between 14–15 EID. Injured live CE
2 showed decreased size and curling, plus petechiae on top of the head. The injured CE
3 in NC and FG inoculated at 10 EID showed a delicate increase in the liver. The injured
4 CE inoculated at 10 EID treated with DX had an enlarged liver which was dark or light
5 green. The lesions observed in live injured CE inoculated at 12 EID with DX were white
6 material over the heart (with UA aspect) and beige material within the allantois. This
7 material was different in appearance and colour from the UA commonly found in the
8 allantois in normal embryos.

9 10 3.3. The weight of CE can be a drug analysis tool depending on age

11 As shown above, most CE inoculated with DX at zero EID did not survive, so
12 the weights of the embryos from the control group and the FG-treated group were
13 evaluated, and there was no difference in weights. CE inoculated with DX at three EID
14 showed decreased weight compared to NC and FG groups. The same occurred in CE
15 inoculated at seven EID. The yolk of embryos inoculated with DX presented a lower
16 weight in relation to the NC and FG groups inoculated at three and seven EID. In CE
17 inoculated at 10 EID and 12 EID, a lower weight of embryos treated with DX compared
18 to embryos from NC and FG groups was observed, but this difference was not
19 observed in the yolk weights of these animals (Figure 3).

Figure 3. Adjusted embryo and yolk weight in grams at different ages and treatments. Embryo weight inoculated at 0 EID (A), 3 EID (B), 7 EID (C) inoculated via SM, 10 EID inoculated with FG via CAM (E), 10 EID inoculated with DX via SM (G), 12 EID inoculated at SH (I). Yolk sac weight at 7 EID (D), 10 EID inoculated with FG via CAM (F), 10 EID inoculated with DX (H) via SM, 12 EID (J) inoculated at SH. In CE inoculated at 0 EID, the CE of the DX group died. We could not weigh the yolk sac of embryos inoculated with the DX at 0 and 3 EID because they disintegrated during manipulation. NC: Negative control. FG: Filgrastim. DX: Dexamethasone.

3.4. The blood cell count can change according to the age of inoculation

The blood samples collected from CE treated with FG did not show a statistical difference compared to the NC group for inoculations made at 10 EID and 12 EID, but the same did not occur for CE treated with DX. Compared to the NC group, the samples

from embryos inoculated with DX at 10 EID had a smaller number of erythrocytes, lower values of haematocrit and haemoglobin, and a smaller number of thrombocytes. No statistically relevant differences were observed in the amounts of total leucocytes. In CE inoculated at 12 EID, embryos treated with DX showed a lower number of erythrocytes. The number of thrombocytes was more significant than the NC group, contrary to the CE treated at 10 EID. In the group inoculated at 12 EID, there was a statistical difference for the amounts of total leucocytes, which was higher for the group treated with DX, heterophiles which was higher, and lymphocytes which were lower compared to the NC group (Table 4).

3.5. Not all metabolites, enzymes and/or minerals change even in liver-damaged embryos

Among the serum and allantoic samples analysed, only AST from serum samples of CE treated with DX at 10 EID showed a statistical difference compared to the NC group (Table 5).

3.6. The amount of metabolites, enzymes and minerals in serum and allantois are not always identical.

The UA, Creat, ALP, GGT, ALT, AST, Ca and phosphorus (P) of serum and allantoic samples from embryos treated with FG and DX at 10 EID were evaluated. Only AST and calcium in CE treated with FG and Crea, ALP and P in CE treated with DX showed similar values for the two analysed samples (Table 5). Although the values of the metabolites and enzymes were not equivalent in all analyses, the results were identical since there were no statistical differences except for AST, which increased in the group treated with DX at 10 EID in serum but not in the allantoic fluid.

1

Table 4. Blood cell count in CE treated with filgrastim and dexamethasone at different ages.

	10 EID				12 EID		
	NC (CAM)	FG (CAM)	NC (SM)	DX (SM)	CN (SM)	FG (SM)	DX (SM)
Ht (%)	21.64 (+/-3.38) [#]	18.43 (+/-3.60) [#]	20.33 (+/-1.86) ^a	17.13 (+/-2.35) ^b	27.80 (+/-3.56) ^A	28.67 (+/-3.50) ^A	23.40 (+/-3.71) ^A
Hg (g/dL) (1/3 ht)	7.20 (+/-1.13) [#]	6.14 (+/-1.20) [#]	6.78 (+/-0.62) ^a	5.71 (+/-0.78) ^b	8.93 (+/-1.03) ^A	9.55 (+/-1.17) ^A	9.26 (+/-1.48) ^A
Hg (g/dL) (Cyanmethemoglobin)	8.12 (+/-0.66) [#]	7.52 (+/-1.29) [#]	7.83 (+/-0.65) ^a	7.22(+/-1.12) ^b	10.61 (+/-3.04) ^A	10.55 (+/-1.48) ^A	7.90 (+/-1.13) ^A
Erythrocytesx10 ⁶ /mm ³	1.76 (+/-0.53) [#]	1.73 (+/-0.28) [#]	2.05 (+/-0.27) ^a	1.58 (+/-0.26) ^b	2.59 (+/-0.48) ^A	2.66 (+/-0.38) ^A	1.90 (+/-0.26) ^B
MCH (pg)	50.34 (+/-14.07) [#]	47.35(+/-4.95) [#]	38.35 (+/-2.04) ^a	45.58 (+/-6.69) ^b	41.13 (+/-9.12) ^A	39.65(+/-2.12) ^A	41.09(+/-1.39) ^A
MCV (fL)	107.1 (+/-13.82) [#]	135.0(+/-43.47) [#]	99.73 (+/-8.53) ^a	109.10 (+/-13.39) ^a	102.8(+/-25.82) ^A	108.4(+/-10.98) ^A	128.6(+/-10.29) ^A
MCHC (g/dL)	38.33(+/-3.3) [#]	41.00(+/-3.48) [#]	38.56 (+/-1.85) ^a	42.12 (+/-2.65) ^b	37.75(+/-7.09) ^A	36.92(+/-4.40) ^A	33.28(+/-3.18) ^A
Thrombocytesx10 ³ /mm ³	15.8(+/-0.97) [#]	13.3 (+/-13.1) [#]	16.3 (+/-5.1) ^a	8.75 (+/-5.75) ^b	5.30 (+/-1.35) ^A	5.66 (+/-1.08) ^A	14.5 (+/-7.32) ^B
Leucocytesx10 ³ /mm ³	3.68(+/-2.06) [#]	4.25(+/-2.85) [#]	4.15(+/-0.74) ^a	3.37(+/-2.58) ^a	4.00 (+/-1.05) ^A	4.96 (+/-1.78) ^A	16.3 (+/-8.30) ^B
Monocytes/mm ³	0.00 (+/-0.00) [#]	14.17(+/-33.09) [#]	0.00(+/-0.00) ^a	0.00(+/-0.00) ^a	45.83(+/-40.79) ^{AB}	61.67(+/-34.86) ^A	40.00(+/-132.70) ^B
Lymphocytes*/mm ³	159.01(+/-97.72) [#]	220.80(+/-216.80) [#]	318.5(+/-326.3) ^a	357(+/-396.6) ^a	457.60(+/-114.80) ^A	516.70(+/-139.00) ^A	271.90(+/-139.80) ^B
Heterophiles/mm ³	3523(+/-2020) [#]	3973 (+/-2565) [#]	3104(+/-1251) ^a	2453(+/-1695) ^a	3498 (+/-940.50) ^A	4132(+/-1549) ^A	16900 (+/-8200) ^B
Heterophile/lymphocyte	26.26(+/-34.75) [#]	32.85 (+/-31.77) [#]	38.51 (+/-38.11) ^a	36.80 (+/-43.29) ^a	8.00 (+/-1.55) ^A	7.00 (+/-3.30) ^A	65.77(+/-29.40) ^B
Eosinophils/mm ³	5.00 (+/-15.00) [#]	25.53(+/-69.48) [#]	0.00(+/-0.00) ^a	1.18(+/-6.03) ^a	0.00 (+/-0.00) ^A	0.00 (+/-0.00) ^A	0.00 (+/-0.00) ^A
Basophils/mm ³	0.00 (+/-0.00) [#]	0.00 (+/-0.00) [#]	0.00 (+/-0.00) ^a	0.00 (+/-0.00) ^a	0.00 (+/-0.00) ^A	0.00 (+/-0.00) ^A	0.00 (+/-0.00) ^A

2

1 Statistical comparisons are between FG CAM and NC CAM (test t) or DX SM and NC SM (test t) inoculated at 10 EID (Test t) or between NC, FG, DX inoculated at 12 EID via SM (ANOVA). Different
2 symbols on the same line indicate a statistical difference between FG and NC inoculated at 10 EID via CAM. Different lowercase letters on the same line indicate a statistical difference between DX
3 and NC at inoculated at 10 EID via SM. Different uppercase letters on the same line indicate a statistical difference in CE inoculated at 19 EID. MCH, eosinophils, monocytes DX 10 EID; leucocytes,
4 lymphocytes and monocytes FG 12 EID via CAM: non-parametric test. Ht: Hematocrit. Hg: Hemoglobin. MCH: corpuscular volume. MCH: mean corpuscular haemoglobin. MCHC: mean corpuscular
5 haemoglobin concentration.

1 **Table 5.** Quantification of metabolites, minerals and enzymes in serum and allantois in CE treated with filgrastim and dexamethasone
 2 at different ages.

	10 EID						12 EID				
	CAM				SM		SM				
	NC (serum)	FG (serum)	NC (AL)	FG (AL)	NC (serum)	DX (serum)	NC (AL)	DX (AL)	CN (serum)	FG (serum)	DX (serum)
UA (mg/dL)	11.00 (+/-7.60) [#]	16.49 (+/-14.55) [#]	97.57 (+/-31.16) [*]	87.88 (+/-51.40) [*]	13.86 (+/-10.37) ^{ab}	12.04 (+/-9.84) ^a	60.00 (+/-38.63) ^{ab}	70.30 (+/-45.93) ^b	8.28 (+/-1.88) ^A	12.53 (+/-10.17) ^A	9.50 (+/-10.82) ^A
Creat (mg/dL)	1.31 (+/-0.68) [#]	0.80 (+/-0.63) [#]	2.49 (+/-0.85) [*]	3.19 (+/-0.77) [*]	0.53 (+/-0.35) ^a	1.12 (+/-1.03) ^a	3.57 (+/-0.53) ^b	2.62 (+/-1.29) ^b	1.37 (+/-1.04) ^A	0.70 (+/-0.53) ^A	1.39 (+/-0.53) ^A
APL (U/L)	1836 (+/-815.30) [#]	2380 (+/-298.40) [#]	47.05 (+/-32.58) [*]	61.26 (+/-53.26) [*]	2125 (+/-1365) ^a	4233 (+/-5830) ^a	45.94 (+/-39.65) ^b	77.81 (+/-54.46) ^b	2194 (+/-322.90) ^A	2439 (+/-437.70) ^A	1928 (+/-356.40) ^A
GGT (U/L)	63.17 (+/-85.13) [#]	67.00 (+/-81.56) [#]	14.53 (+/-5.50) [#]	12.49 (+/-6.98) [#]	81.20 (+/-94.30) ^a	49.41 (+/-82.50) ^a	12.72 (+/-11.31) ^a	19.51 (+/-9.54) ^a	47.25 (+/-40.85) ^A	59.84 (+/-64.05) ^A	69.08 (+/-47.69) ^A
AST (U/L)	48.12 (+/-29.53) [#]	49.18 (+/-15.01) [#]	15.14 (+/-11.55) [*]	9.33 (+/-6.02) [*]	37.55 (+/-20.84) ^a	99.89 (+/-68.10) ^b	39.87 (+/-55.81) ^a	13.27 (+/-4.25) ^a	123.1 (+/-81.82) ^A	108.2 (+/-19.21) ^A	556.1 (+/-27.61) ^A
ALT (U/L)	54.02 (+/-51.67) ^{#*}	158.00 (+/-150.10) [#]	8.28 (+/-4.07) [*]	11.71 (+/-8.51) [*]	142.6 (+/-161.3) ^a	142.8 (+/-138.7) ^a	8.00 (+/-1.58) ^a	13.00 (+/-8.69) ^a	56.68 (+/-59.33) ^A	130.5 (+/-226.9) ^A	73.70 (+/-25.15) ^{Av}
Ca (mg/dL)	18.80 (+/-17.03) [#]	21.80 (+/-21.75) [#]	12.50 (+/-7.05) [#]	8.34 (+/-4.62) [#]	9.15 (+/-10.65) ^a	30.53 (30.50) ^a	6.50 (2.80) ^a	15.91 (13.46) ^a	17.98 (+/-9.35) ^A	17.70 (+/-12.28) ^A	13.63 (+/-10.32) ^A

P (mg/dL)	7.48	4.52	15.37	22.13	4.68	5.80	23.65	26.90	5.22	4.76	9.52
	(+/-4.52)#	(+/-2.58)#	(+/-9.04)##*	(+/-9.63)*	(+/-3.33) ^a	(+/-4.21) ^a	(+/-6.32) ^b	(+/-11.91) ^b	(+/-3.89) ^A	(+/-2.85) ^A	(+/-3.28) ^A

- 1 Statistical comparisons are between FG CAM and NC CAM (t test) serum and AL or DX SM and NC SM (t test) inoculated at 10 EID (t test) serum and AL or between NC, FG, DX inoculated at 12 EID
- 2 via SM (ANOVA) in serum because at 19 EID (age at colect) there are not allantoic fluid. Different symbols on the same line indicate a statistical difference between FG and NC inoculated at 10 EID
- 3 via CAM. Different lowercase letters on the same line indicate a statistical difference between DX and NC inoculated at 10 EID via SM. Different uppercase letters on the same line indicate a statistical
- 4 difference in CE inoculated at 12 EID. GGT in DX group inoculated at 10 EID via SM: non-parametric test. GGT and ALP in DX group inoculated at 10 EID via SM; Creat in FG group inoculated at 10
- 5 EID; UA, ALT, group inoculated at 12 EID: non-parametric test.. UA: Uric Acid. Creat: Creatinine. ALP: Alkaline Phosphatase. GGT: gamma-glutamyl transferase AST: aspartate aminotransferase.
- 6 ALT: alanine aminotransferase. Ca: calcium. P: phosphorus and CRP: c-reactive protein.

3.7. The age of inoculation is an essential factor for changing oxidative stress parameters

Figure 4 shows the levels of sulphydryl groups, FRAP, lipid peroxidation and ROS in the liver of CE inoculated with FG and DX at 10 and 12 EID. In comparison to the NC, FG increased the levels of sulphydryl groups ($p < 0.05$) of CE treated at 10 EID. At the same EID, embryos treated with DX had increased levels of ROS production. In CE treated at 12 EID with FG, there was no difference between the NC group but the CE treated with DX showed a decrease in the sulphydryl group and FRAP levels and an increase in the levels of lipid peroxidation and ROS production.

Figure 4. Levels of oxidative stress biomarkers in terms of sulphydryl groups (A), FRAP (B), lipid peroxidation (C) and ROS production (D) in chicken embryos' livers after inoculation with FG via CAM at 10 EID; sulphydryl groups (E), FRAP (F), lipid peroxidation (G) and ROS production (H) in chicken embryos' livers after inoculation with DX via SM at 10 EID; sulphydryl groups (I), FRAP (J), lipid peroxidation (K) and ROS production (L) in chicken embryos' livers after inoculation with FG and DX via SM in 12 EID.

12 EID. FRAP: ferric reducing antioxidant power. ROS: Reactive oxygen species. NC: Negative control. FG: Filgrastim. DX: Dexamethasone.

3.8. There is a correlation between the results for haemoglobin using a Drabkin solution or performing the calculation of one third of the haematocrit

There was a correlation between the assessment of haemoglobin by calculating one third of the haematocrit and the method using the Drabkin solution. The r value in the group of CE inoculated at 12 EID was 0.61, representing a moderate correlation(28). In the groups treated at 10 EID with FG, DX and virus, the r value was 0.86, 0.85 and 0.86, respectively, showing a strong correlation.

3.9. Characterisation of granulocytes and thrombocytes by cytochemistry

Thrombocytes were labelled by the formaldehyde method (Figure 5) and the analysis showed that the thrombocytes' diameters in CE were lower than in adult animals. Besides, the FG and DX increased the diameter of the thrombocytes.

Figure 5. Diameter of formalin-labelled thrombocytes evaluated. A: diameter of thrombocytes in SPF CE inoculated with *Gammacoronavirus*. A1: thrombocytes stained in group treated with virus. B. Diameter of thrombocytes in CE treated with filgrastim at 10 EID. B1: thrombocytes stained in group treated with FG inoculated at 10 EID via CAM. C. Diameter of thrombocytes in CE treated with DX inoculated at 10 EID via SM. B1: thrombocytes stained in the group treated with DX. D. Diameter of thrombocytes in CE treated with FG and DX inoculated at 12 EID via SM. B1: thrombocytes stained in the group treated with FG. E. Thrombocytes stained in the NC group in CE inoculated at 10 EID. E1: trans, E2: UV, E3: over. NC: Negative control

Blood smear slides were stained by PAS cytochemical methods for identification of thrombocytes by staining of cytoplasmic glycogen granules and SBB for identification of eosinophils by staining of cytoplasmic granules. These two methods were used to differentiate lymphocytes that were negative for PAS and heterophils that were negative for SBB (Figure 6).

Figure 6. Blood smear slides stained by cytochemical methods. (A) Sudan Black B (SBB) positive eosinophils (arrow) and SBB negative heterophils (arrowhead). (B) Periodic Acid Schiff (PAS) positive thrombocytes (arrow) and PAS negative lymphocytes (arrowhead).

3.10. Unidentified granulocytes were found in several groups

Despite the various methods of analysis used to identify the cells, some cells with a rounded nucleus and basophilic round cytoplasmic granules were found on some slides in groups NC, FG, DX, and in the group infected with the virus (Figure 7). These cells were not positive by cytochemistry used in our work.

Figure 7. Unidentified granulocytes in blood smear stained with fast Panotico® (arrow 100x).

3.11. Liver histopathological analysis

The degenerative change observed in the CE livers was lipidosis. The hepatocytes were vacuolated, ranging from few and small light vacuoles to large light vacuoles, shifting the nucleus to the periphery, with the hepatocyte resembling an adipocyte. As in virus-infected embryos, the inflammatory process had a periportal distribution and was predominantly composed of heterophils (Figure 8).

Figure 8. Histopathological analysis of the livers of CE treated with DX and FG at 10 EID. A. CE liver sample from NC with a small amount of lipids in vacuoles in hepatocytes considered normal for comparison with the other groups (arrow). B. Sample from the NC without inflammatory infiltrate (arrow). C. DX-treated CE sample showing large cytoplasmic vacuoles in hepatocytes containing lipid characterising lipid degeneration (arrow). D. DX-treated CE sample with sinusoids filled with erythrocytes characterising congestion (arrows). E. FG-treated CE sample with perivascular inflammatory infiltrate (white arrow) and congestion (black arrows). The bars in images A, B and C represent 60 μm and the bars in images D and E represent 200 μm .

The animals inoculated with FG at 10 EID showed an inflammatory process with infiltration of heterophils. Embryos inoculated with DX at 10 EID showed a mild inflammatory process and circulatory alteration and animals inoculated at 12 EID showed a mild inflammatory process and degeneration (lipidosis) (Table 6).

Table 6. Histopathological analysis of liver from embryos inoculated with FG and DX at 10 and 12 EID.

	10 EID				12 EID		
	NC (CAM)	FG (CAM)	NC (SM)	DX (SM)	NC	FG	DX
Inflammation	0.00 (Mi: 0.00; Ma: 0.00) [#]	1.00 (Mi: 0.00; Ma: 2.00) [*]	0.00 (Mi: 0.00; Ma: 0.00) ^a	0.00 (Mi: 0.00; Ma: 2.00) ^b	0.00 (Mi: 0.00; Ma: 0.00) ^A	0.00 (Mi: 0.00; Ma: 1.00) ^{AB}	1.00 (Mi: 0.00; Ma: 2.00) ^B
Degeneration	0.00 (Mi: 0.00; Ma: 0.00) [#]	0.00 (Mi: 0.00; Ma: 0.00) [#]	0.00 (Mi: 0.00; Ma: 0.00) ^a	0.00 (Mi: 0.00; Ma: 3.00) ^a	0.00 (Mi: 0.00; Ma: 0.00) ^A	0.00 (Mi: 1.00; Ma: 1.00) ^{AB}	1.00 (Mi: 1.00; Ma: 3.00) ^B
Necrosis	00.00 (Mi: 0.00; Ma: 0.00) [#]	0.00 (Mi: 0.00; Ma: 0.00) [#]	0.00 (Mi: 0.00; Ma: 0.00) ^a	0.00 (Mi: 0.00; Ma: 2.00) ^a	0.00 (Mi: 0.00; Ma: 0.00) ^A	0.00 (Mi: 0.00; Ma: 0.00) ^A	0.00 (Mi: 0.00; Ma: 0.00) ^A
Circulatory change	0.00 (Mi: 0.00; Ma: 0.00) [#]	0.00 (Mi: 0.00; Ma: 1.00) [#]	0.00 (Mi: 0.00; Ma: 0.00) ^a	1.00 (Mi: 0.00; Ma: 3.00) ^b	0.00 (Mi: 0.00; Ma: 0.00) ^A	0.00 (Mi: 0.00; Ma: 1.00) ^A	0.00 (Mi: 0.00; Ma: 0.00) ^A

Mi: Minimum, Ma: Maximum. NC: Negative control. CAM: Chorioallantoic membrane. FG: Filgrastim. SM: Shell Membrane. DX:

Dexamethasone. EID: Embryonated incubation days.

3.12. FG in high doses does not cause injury to angiogenesis in CE

We tested the hypothesis that FG could cause angiogenesis in CE using several doses. Unlike that believed, FG did not induce angiogenesis (measured by vascular

density and vessel length density), just as DX did not change the number of vessels (Supplementary Figure 1). As already reported in the previous experiment, DX led to incomplete CAM formation. The high dose of FG also did not cause visible changes in the embryo.

4. Discussion

CE provide an ideal model for investigating the development of pathogens such as viruses and testing drug effects, evaluating both toxicity and pharmacokinetics. Thus, CE as an animal model can be an important ally for carrying out experiments that require quick and less complex execution and with limited resources.

The CE and its annexes constitute a favourable environment for replicating several viruses, being widely used for the isolation or production of vaccines(9). IBV causes dwarfism, haemorrhage and death when inoculated into embryonated eggs. Some strains of the *gammacoronavirus* can cause nephropathies and others haemagglutination(29). However, IBV has a minimal agglutination capacity for erythrocytes in chickens(30). Our study showed a decrease in haemoglobin concentration, haematocrit value and the number of erythrocytes in infected embryos compared to the NC group, indicating anaemia. Haematimetric indices did not differ in relation to the NC group. Thus, we can classify anaemia as normocytic and normochromic. This means that the cells were of standard size and haemoglobin concentration, so there was a decrease in RBC, but there was no response from the bone marrow to release cells into the bloodstream. Normocytic and normochromic anaemia can indicate decreased erythrocyte production, which can develop rapidly in birds with diseases involving infectious agents(18).

Although there was no statistical difference in the number of leucocytes in the infected animals compared to embryos in the NC group, some animals had a high leucocyte count ($> 30,000/\text{mm}^3$). In contrast, others had a low count ($< 4,000/\text{mm}^3$) (Supplementary Table 1). These results contributed to the increase in the standard deviation in this analysis. The response of the white blood cell count during an infection can vary. In the initial moment of disease, leucocytosis may occur as a response by the body. However, as the leucocytes are consumed, leucopenia may occur when the demand for leucocytes is above the production capacity of these cells by the bone marrow(18). From 12 EID, the haematopoiesis in CE is more active and occurs mainly in bone marrow (31). Therefore, in this work, haematopoiesis in the CE was intense since we used animals between 10–17 EID. However, although the CE used in this work were from the same flock and the mother had high consanguinity, there may be an inherent variation in the model itself.

There was no change in the Creat and UA levels in serum and allantois samples from infected CE compared to the NC group, and macroscopic kidney lesions were not observed either. On the other hand, the liver of infected animals was macroscopically altered, with an increase in volume and a greenish colour. In addition, we observed a moderate inflammatory reaction by histopathologic analyses (Figure 2), although liver enzymes were not increased in the serum or allantois compared to the control group(32). There was also a decrease in lipid peroxidation rather than an increase. The absence of a rise in GGT, ALT, AST and the reduction in lipid peroxidation indicate that the stress was very intense, reducing liver responsiveness.

Serum calcium levels from virus-infected CE decreased compared to the NC group. Possibly the viruses can hijack the host cell's machinery and utilise the host cell's calcium to create an environment adapted to meet their own demands for

1 replication(33). This hypothesis is supported in the work of Cao et al. (2011)(34). These
2 authors observed that the expression level of some calcium-binding proteins was
3 increased after *in ovo* IBV infection. These proteins facilitate the transcellular transport
4 of calcium, suggesting that IBV may disrupt cellular Ca_2 homeostasis for its own
5 benefit.

6 In addition to virus experiments, drug studies using CE as a model provide a
7 technically simple way to study complex biological systems for *in vivo* drug toxicity and
8 pharmacokinetic assessment. For example, DX is a widely used glucocorticoid. Its
9 administration in CE can cause immunosuppressive effects increasing embryonic
10 catecholamines that alter development and cause death(35).

11 There was high macroscopic lesion and mortality rate after the DX inoculation,
12 similar to other studies(35–37). As expected, the mortality was age-dependent. In
13 addition, in our study, CE treated with DX reduced the weight in all CE that lived. The
14 reduced weight in CE treated with DX can be related to muscle and bone development
15 inhibition(36,37). Furthermore, high doses of glucocorticoids can generate
16 suppression of growth hormone activity in the pituitary gland which is fully established
17 at the beginning of the last week of the embryonic development of CE(37). Thus, we
18 can suggest that DX promotes a delay in embryonic development, negatively
19 influencing embryo weight gain, which may explain the delay in the complete formation
20 of CAM found by us.

21 Unlike DX, FG did not cause significant changes and deaths. The only death
22 observed in an embryo inoculated at zero EID was not accompanied by changes and
23 occurred in an embryo from the NC group, probably caused by a natural process.
24 Similarly, the injury in the CE inoculated via CAM possibly resulted from the inoculation
25 process since haemorrhage may occur using this route.

CE treated with DX at 10 and 12 EID were anaemic. The CE treated with DX showed a decreased erythrocyte count and decreased haematocrit and at 10 EID, haemoglobin concentration compared to CE in the NC group. There were also decreases in haematocrit, haemoglobin and erythrocyte values in animals treated with DX at 12 EID, but only erythrocytes showed a statistical difference for the NC group. Despite that, there was a high correlation between haemoglobin and haematocrit (r value = 0.78), haematocrit and erythrocyte (r value = 0.77) and haemoglobin and erythrocytes (r value = 0.81). Therefore, the erythrocyte reduction accompanied the reduction in haemoglobin and haematocrit. Although the CE treated with DX had anaemia, the haematimetric indices were not statistically different between the groups treated and the NC, defining anaemia as normocytic and normochromic. Chickens treated with corticosteroids show increased energy expenditure(38). Embryos treated with DX at 10 EID had higher yolk consumption, indicated by the lower yolk weight of these embryos compared to animals in the NC group. The nutritional deficit associated with liver damage caused by the drug may have impaired the production of erythrocytes, causing normocytic normochromic anaemia.

In CE inoculated with DX at 10 EID, there was a decrease in the number of thrombocytes compared to embryos from the NC group. Perhaps the production of thrombocytes in CE treated with DX at 10 EID was impaired by the damage caused in the liver. Thrombopoietin (TPO) is the regulator of megakaryocyte development and thrombocyte production and its expression in chickens occurs mainly in the liver(39). Hemopoietic activity in the liver starts at seven EID with a peak at 14 EID(40), but is more active from 12 EID(31). In animals treated with DX at 12 EID, the opposite was observed. There was an increase in the number of thrombocytes compared to CE from the NC group. Thrombocytosis may reflect a rebound response after recovery from

1 other conditions associated with excessive use of thrombocytes. Considering that CE
2 from 12 EID had a more mature liver, active bone marrow and hemopoietic activity(31),
3 thrombocytosis can be explained in CE treated at 12 EID. Additionally, it should also
4 be considered that in birds, thrombocytes have a phagocytic function(17)and the
5 influence of glucocorticoids on these cells is unknown(41).

6 In embryos treated with DX at 12 EID, there was an increase in the number of
7 leucocytes, different to that observed in CE inoculated at 10 EID. This can be explained
8 by the onset of lymphoid activity in the Bursa of Fabricius at 12 EID(42), with a greater
9 capacity to respond to the stimulus caused by the drug. In mammals treated with DX,
10 an initial leucocytosis may occur, mainly due to neutrophilia(30). In fact, CE inoculated
11 at 12 EID showed an increase of H/L ratio due to the increase in heterophils and a
12 decrease in lymphocytes. This corroborates several studies in born animals(43–45).
13 During CE development, granulopoiesis is more predominant; however, at hatching,
14 the granulocytes begin to be replaced by lymphocytes by the first three days(18).

15 Taken together, the results show that embryos with a small difference in embryo
16 development stage can completely alter the cell count response. We do not know if
17 this occurs with other drugs. However, knowing that the embryo is an ascending animal
18 model, further work must be carried out to consider the best age for using the model
19 depending on the expected objective.

20 As a granulocyte colony-stimulating factor (G-CSF), FG is used in human
21 medicine to increase levels of neutrophils in the bloodstream. Therefore, we expected
22 that the same effect would be observed in the CE in this experiment, increasing
23 heterophils that have characteristics and performance corresponding to human
24 neutrophils. However, no increase in granulocytes was observed in animals treated
25 with FG at 10 EID or 12 EID either by inoculation in CAM or SM. Perhaps this did not

1 occur because FG is a synthetic compound for human use and may not have the same
2 results in other species.

3 The analysis of biochemical parameters provides essential data for the
4 assessment of the clinical status of the animal. However, blood samples collected in
5 research with CE do not always allow this analysis because it is not easy and takes
6 time. Therefore, alternatives samples have been used to assess these parameters,
7 such as amniotic fluid and allantoic fluid(5,7,46). In this work, we compared the
8 biochemical analyses of serum and allantoic fluid and observed that the values found
9 for the two samples were not always similar. However, the differences observed
10 between groups in serum samples were also observed in allantoic samples, except for
11 AST, which may indicate that this enzyme does not have a good analysis from
12 allantois.

13 It is known that proteins found in serum have different physical and biochemical
14 properties and change in various physiological and pathological conditions. One of the
15 problems with enzymatic analysis methods is that the reagents were designed to
16 provide the substrate and its optimal concentrations for human plasma, but these
17 variables can change depending on the species. For example, birds have deficient
18 levels of activity of the enzyme ALT in the liver tissue, so in cases of severe liver
19 damage, this enzyme may present normal values. Aspartate aminotransferase (AST)
20 activity occurs in multiple tissues, but the main ones are liver and muscle, being
21 considered sensitive but not very specific in cases of liver problems. GGT activity is
22 increased in all conditions in which hepatocellular damage is present. Alkaline
23 phosphatase (ALP) is associated with the regulation of bird growth, participating in
24 chondrogenic and osteoblastic activities. Thus, physiological variations can be
25 observed, with higher activity levels resulting from bone growth in young birds.

Elevations in ALP may be associated with liver disease even if its activity in this organ is minimal²⁶.

In our study, only the AST enzyme of CE treated with DX at 10 EID showed a statistical difference compared to the CN group. However, histopathological findings and results of oxidative stress biomarkers indicate that there was liver damage. At 12 EID, there was no increase of AST in CE treated with DX. At this age, just 50% of the CE had a macroscopic injury (Table 2) and maybe because of this, the AST level did not increase. However, the AST maximum value of the group treated of DX at 19 EID was greater than the NC (Supplementary Table 2). Thus, AST may be the best parameter for analysis of liver function in embryo serum.

High concentrations (up to five times) of UA in plasma can lead to precipitation of this acid in the form of crystals, which accumulate in tissues. Situations of hypouricemia are rare and may be related to severe liver damage with a consequent decrease in UA production(47). The excretion of Crea occurs via the kidneys, but in birds, most creatine is excreted before being converted to creatinine(18). Thus, increased Crea concentrations are rare and may occur in severe renal impairment, significantly if filtration is affected(47). In our study, no changes in urea or Creat were observed in any of the treatments. Macroscopic lesions were also not observed. This can mean that there was no damage to the CE kidneys. However, as we did not perform histopathological analysis of the kidneys, we cannot rule out the possibility that the high damage caused by the treatment does not lead to changes in biochemical parameters.

Drugs can cause increased production of oxidants and the formation of free radicals, which, by exceeding the body's ability to neutralise and scavenge these radicals, can cause organ damage(48). Our study showed an increase in oxidative

stress biomarkers ROS in the livers of embryos treated with DX at 10 EID and 12 EID and an increase in lipid peroxidation in embryos treated at 12 EID. In adult animals, corticosteroids can increase oxidative stress(49), but a similar approach was never studied in CE. The increased energy expenditure triggered by high circulating corticosteroid levels may be responsible for the augmented formation of ROS as reflected by increased lipid peroxidation(49). The body uses enzymatic and non-enzymatic antioxidants to neutralise damage caused by free radicals and minimise excessive oxidative stress. Sulphydryl groups are an example of essential antioxidants in controlling these processes and protecting against damage. Embryos that were treated at 12 EID showed decreased sulphydryl groups and FRAP values, indicating that these embryos had difficulty matching the damage caused by free radicals in the induction of oxidative stress by DX.

Embryos treated with FG at 12 EID did not show changes in oxidative stress biomarkers compared to CE from the NC group. However, in CE inoculated at 10 EID via CAM, FG increased the sulphydryl group. Filgrastim is a granulocytic colony-stimulating factor (G-CSF). It has biological activity identical to that of endogenous human G-CSF with a free cysteine at position 17 with an ionised sulphydryl group that is very reactive to free radical oxidation(50). From this, we can conclude that FG may have shown an antioxidant effect, with possible protection of the embryo against ROS.

To measure haemoglobin in this experiment, we used Drabkin's solution and compared it with the values obtained using the Drabkin solution method. Our study showed a moderate correlation in embryos treated at 12 EID and a strong correlation in embryos treated at 10 EID(28). Thus, we can conclude that the calculation based on haematocrit can be used to approximate the haemoglobin value in situations where measurement by spectrophotometry is not possible. However, this replacement is only

possible if there is no suspicion of haemolysis, since haemolysis due to collection or pathological problems, promotes a decrease in haematocrit without a proportional reduction in haemoglobin.

To perform the differential count of leucocytes in birds, there is a great difficulty in differentiating between thrombocytes and lymphocytes. Though not identical, the nuclei of thrombocytes and small lymphocytes are too similar to serve as a basis for distinguishing between these two cell types. In the present study, cells in fast panoptic stained CE blood smears with small, round or oval nuclei with dense chromatin clumps were categorised as thrombocytes if they had cytoplasmic vacuoles and colourless cytoplasm. Cells classified as small lymphocytes had similar nuclei but scant amounts of blue or dark blue cytoplasm without vacuoles. To establish a basis for the categorisation of these cells, some cytochemical properties of these cells were compared. The typical thrombocytes were visualised by UV after being exposed to gaseous formaldehyde. Swayne et al. (1986)(19) observed that most of the thrombocytes (99%) had been fluorescent after gaseous formaldehyde treatment and all of the small lymphocytes had been non-fluorescent. This fluorescence resulted from serotonin condensation products. We measured the thrombocytes found in the slides of animals inoculated in this study and compared them with cells found in the blood smear slides of adult chickens. Thrombocytes from adult chickens had a larger size compared to the NC group, which averaged 8.5 μm . We could not determine why FG increased the size of thrombocytes.

Using the PAS, we characterised the thrombocytes as PAS positive while lymphocytes and erythroblasts were PAS negative (Figure 6). The eosinophils were SBB positive, while the heterophils were SBB negative (Figure 6). The use of PAS and SBB is important for better cell classification.

Despite the different forms of differentiation used to identify cells in the blood smears of embryos in this experiment, some cells remained without conclusive identification. The authors identified these cells as being a type of granulocyte, but they could be confused with eosinophils, basophils or granulocyte precursors(17,18,51). These cells were found in samples from embryos treated with FG (20%) and DX (25%) inoculated at 10 EID, animals infected with virus (100%), and in animals from the NC group (23.07%) inoculated at 10 EID.

Another response observed in the use of FG in humans is the stimulation of endothelial cells with consequent angiogenesis(52–54). In this study, we analysed the blood vessels of the CAM of animals inoculated with FG to identify a possible increase in vessel density characterising the occurrence of angiogenesis. However, there was no statistical difference between CE inoculated with FG and the NC group.

5. Conclusion

Overall, the results of this work provide vital information on the use of CE as an experimental model. The response of CE to challenges from viruses and drugs does not always go as expected. Although macro and microscopic damage was visible in viruses, white blood cell counts and inflammation biomarkers such as CRP did not change. It is important to mention that some drugs can be innocuous and not result in expected effects on CE, which was the case with FG. In the case of DX, changes in blood parameters and biomarkers seem to be inherent to the model and are highly dependent on the developmental stage of the CE.

This article reiterates the wonderful value of the CE as an animal model. However, our work sheds light on the importance of standardisation and the correct use of the model (considering the laboratory analysis, drug, age, route) so that the infection, toxicity and pharmacokinetic results are reliable.

1

2 6. List of abbreviations

3 CE: Chicken embryo

4 FG: Filgrastim

5 DX: Dexamethasone

6 CAM: Chorionallantoic membrane

7 SPF: Specific pathogen-free

8 IBV: Infectious bronchitis virus

9 EID: Embryonic incubation days

10 PBS: Phosphate-buffered saline

11 NC: Negative control

12 SM: Shell membrane

13 Ht: Haematocrit value

14 Hb: Haemoglobin concentration

15 RBC: Red blood cell

16 MCV: Mean corpuscular volume

17 MCH: Mean corpuscular haemoglobin

18 MCHC: Mean corpuscular haemoglobin concentration

19 PAS: Periodic acid Schiff

20 SBB: Sudan Black B

21 AST: Aspartate aminotransferase

22 ALT: Alanine aminotransferase

23 GGT: Gamma-glutamyl transferase

24 ALP: Alkaline phosphatase

25 Creat: Creatinine

- 1 UA: Uric acid
- 2 Ca: Calcium
- 3 P: Phosphorus
- 4 CRP: C-reactive protein
- 5 ROS: Reactive oxygen species
- 6 TBA: Thiobarbituric acid
- 7 TCA: Trichloroacetic acid
- 8 MDA: Malondialdehyde
- 9 DTNB: Ditionitrobenzoic acid
- 10 HE: Haematoxylin and eosin
- 11 FRAP: Reducing antioxidante power
- 12 G-CSF: Granulocyte colony-stimulating factor

13

14 **Ethics approval and consent to participate:** The project was evaluated and
 15 approved by the Ethics and Research with Animals Committee of the Universidade
 16 Federal de Uberlândia (certificate A011/20 and nº 008/21).

17

18 **Authors' Contributions:** B.B.F. idealized and coordinated the project. S.S., B.B.F.,
 19 A.V.M. and L.R.G. designed all the experiments. S.S., R.R.S. and B.B.F. carried out
 20 drugs and virus inoculation in chicken embryos. S.S., B.B.F., R.R.S, M.P.R., J.S.Q.,
 21 F.O.N., collected the biological material and macroscopic analysis. F.S.E. and R.R.F.
 22 performed the biochemical studies of the liver chicken embryos. F.O.N and A.A.M.
 23 performed the histopathological analyses of the liver chicken embryos. S.S., M.P.R.,
 24 J.S.Q. and M.E.B. performed the blood analyses of the chicken embryos. S.S., A.V.M.
 25 and B.B.F. wrote the manuscript. All the authors revised the manuscript.

Acknowledgements: To Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Amparo a Pesquisa do Estado de Minas Gerais (FAPEMIG) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) that support our work. To Felipe Cesar Gonçalves and Igor Paula de Castro who support us with technical experience. To Hy Line Company that support us with eggs in our research.

Funding: This research was funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001, Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG).

Data Availability Statement: Data are contained within the article or supplementary material. Full data will be made available if the publisher requests it.

Conflicts of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Consent for publication: All authors have read and agreed to the published version of the manuscript.

SUPPLEMENTARY MATERIAL

Supplementary Figure 1. Angiogenesis was measured by vascular density and vessel length density using ImageJ software with the Vessels Analysis 58 plugin. The CE were inoculated at 3 EID and evaluated at 12 EID. A. vascular density, B. vessel length density, C. Photos CAM (NC), D. Photo CAM treated with DX, E. Photo CAM treated with FG (150 ug/CE), F. Photo CAM treated with FG (750 ug/CE), G. Photo CAM treated with FG (3.75 mg/CE)

- 1 **Supplementary Table 1.** Maximum and minimum values and median of
 2 haematological analysis of CE infected with *Gammacoronavirus* and NC.

		Virus	NC
Hc	Minimum	13	19
	Maximum	18	23
	Median	15	21
Hb 1/3Hc	Minimum	4.33	6.33
	Maximum	6	7.67
	Median	5	7
Hb Drabkin	Minimum	5.02	7.7
	Maximum	7.39	8.23
	Median	6.605	8.085
Erythrocytes	Minimum	1220000	1900000
	Maximum	1790000	2240000
	Median	1680000	2020000
MCV	Minimum	80.81	83.33
	Maximum	121.1	147.5
	Median	92.99	87.21
MCH	Minimum	32.37	29.88
	Maximum	42.74	58.28
	Median	37.7	38.4
MCHC	Minimum	35	35.86
	Maximum	47.06	44.13
	Median	39.53	39.6
Leucocytes	Minimum	3500	5500
	Maximum	54000	11000
	Median	6000	7750
Heterophils	Minimum	3325	5280
	Maximum	45360	10340

	Median	5640	6788
Lymphocytes	Minimum	70	120
	Maximum	6480	1710
	Median	180	385
Monocytes	Minimum	0	0
	Maximum	1080	110
	Median	70	0
Eosinophils	Minimum	0	0
	Maximum	0	0
	Median	0	0
Thrombocytes	Minimum	9000	18000
	Maximum	51500	29500
	Median	19000	20500

Ht: Hematocrit. Hg: Hemoglobin. MCH: corpuscular volume. MCH: mean corpuscular haemoglobin. MCHC: mean corpuscular haemoglobin concentration.

Supplementary Table 2. Maximum and minimum values and median of metabolites, enzymes and minerals from the serum and allantoic fluid of CE infected with *Gammacoronavirus* and their NCs.

		Serum		Allantoic	
		Virus	NC	Virus	NC
UA	Minimum	4.24	1.27	7.15	1.27
	Maximum	36.94	50.07	85.39	50.07
	Median	24.67	17.85	56.93	11.39
Crea	Minimum	0.19	0.26	0.33	1.63
	Maximum	0.96	2.94	2.75	4.92
	Median	0.435	1.235	1.05	3.155

ALP	Minimum	235.7	736.5	18.2	58.2
	Maximum	3845	3388	123.1	82.9
	Median	2476	1651	24,1	68.1
GGT	Minimum	10.2	77.4	40	58
	Maximum	136	356	434	346
	Median	89.6	288	183	166
ALT	Minimum	24	40	20	20
	Maximum	100	220	120	80
	Median	53	54	60	40
AST	Minimum	54	120	120	220
	Maximum	560	780	380	920
	Median	221	452	150	640
Ca	Minimum	2.5	20.5	4.4	10.4
	Maximum	21.5	146.4	39	34.2
	Median	8.45	118.8	14.2	15.8
P	Minimum	2.7	1.4	2.5	11.6
	Maximum	8.4	9.7	18.3	29.4
	Median	5.45	8.1	8.7	17.7
CRP	Minimum	10.8	15	26	40
	Maximum	35.4	38	100	152
	Median	26	23	56	86

NC: negative control; AL: allantoic liquid; UA: Uric acid; Crea: Creatinine; ALP: alkaline phosphatase; AST: Aspartate transaminase; ALT: Alanine Aminotransferase; GGT: gamma-glutamyl transferase; CRP: C-reactive Protein.

Supplementary Table 3. Maximum and minimum values and median of haematological analysis of CE treated with FG and DX at 10 EID and their respective NCs.

		NC	DX	NC	FG
Hc	Minimum	17	13	19	14
	Maximum	22	21	28	25
	Median	20.5	17.5	21	17
Hb 1/3Hc	Minimum	5.67	4.33	6.33	4.67
	Maximum	7.33	7	9.33	8.33
	Median	6.835	5.835	7	5.67
Hb Drabkin	Minimum	6.48	5.16	7.08	5.58
	Maximum	8.75	8.75	9.17	9.76
	Median	7.86	7.375	8.085	7.755
Erythrocytes	Minimum	1730000	1205000	1125000	1135000
	Maximum	2530000	1940000	2600000	1995000
	Median	2012500	1537500	1772500	1760000
MCV	Minimum	86.96	95.74	92.39	76.92
	Maximum	112.5	135.3	125.3	195.6
	Median	98.78	108.1	104.2	138.8
MCH	Minimum	34.58	40.98	31.24	36.19
	Maximum	40.91	64	75.29	50.04
	Median	38.56	45.85	48.04	44.19
MCHC	Minimum	35	37.56	32.75	38.15
	Maximum	41.63	47.45	42.7	49.21
	Median	39.04	42.71	37.63	39.63
Leucocytes	Minimum	1000	500	2000	500
	Maximum	5000	7500	8500	8500
	Median	4000	2500	3000	3250
Heterophils	Minimum	980	495	1780	475
	Maximum	4500	6300	8160	7735
	Median	3600	2300	2880	3218
Lymphocytes	Minimum	20	5	60	25
	Maximum	810	1200	340	640

	Median	210	200	120	150
	Minimum	0	0	0	0
Monocytes	Maximum	0	0	0	85
	Median	0	0	0	0
	Minimum	0	0	0	0
Eosinophils	Maximum	45	210	0	0
	Median	0	0	0	0
	Minimum	11500	1000	5750	2500
Thrombocytes	Maximum	22500	15500	34000	39500
	Median	14000	8500	14250	7000

Ht: Hematocrit. Hg: Hemoglobin. MCH: corpuscular volume. MCH: mean corpuscular haemoglobin. MCHC: mean corpuscular haemoglobin concentration.

Supplementary Table 4. Maximum and minimum values and median of metabolites, enzymes and/or minerals from the serum and allantoinic fluid of CE treated with FG and DX inoculated at 10 EID and their respective NCs.

		Serum		Allantoic		Serum		Allantoic	
		NC	DX	NC	DX	NC	FG	NC	FG
UA	Minimum	1.83	0.63	17.52	0.76	2.53	2.64	33.65	17.46
	Maximum	32.13	22.8	115.4	129.9	22.15	38.71	127.8	174.7
	Median	10.82	12.21	53.24	76.43	9.835	7.98	109.4	82.23
Crea	Minimum	0.15	0.12	2.7	1.1	0.41	0.28	1.81	2.16
	Maximum	1.08	3.06	4.26	5.67	2.34	1.98	4.29	4.65
	Median	0.51	0.695	3.735	2.18	1.14	0.51	2.16	3.01
ALP	Minimum	562.6	904.1	9.1	21.5	1114	1830	8.5	9.8
	Maximum	4157	17399	130.9	150.8	3110	2661	98	126.3

	Median	2460	2263	31.9	75	1404	2473	39.1	35.8
GGT	Minimum	5.1	2.4	1.3	4.3	3.1	1.1	7.4	4.7
	Maximum	229	232	27	31	228	204	22.9	21
	Median	14.4	16.4	8.35	17.8	33.3	28	13.95	12.9
ALT	Minimum	14.8	25.8	6	2	5.6	6.2	2	2
	Maximum	437.9	428	10	21	118	364	15	29
	Median	71.4	123	8	14.5	41.95	145.8	8	10
AST	Minimum	11.7	39	5.9	6	15	39.8	3.2	4
	Maximum	63	200.4	31	20.8	74.4	71.5	31.1	17.9
	Median	38.8	79.2	16	14	61.8	42.7	12.6	6
Ca	Minimum	3.1	5.6	3.4	0.8	2.6	3.6	6.3	2.6
	Maximum	30.6	94.2	10.2	41.6	40	60	25.3	14.7
	Median	5.3	20.9	5.8	12.1	16.9	9.8	10.04	6.3
P	Minimum	1.79	0.28	15.6	16.1	2.34	2	4.7	11.1
	Maximum	10.8	13.2	32.5	47.6	15.8	9	29.5	37.2
	Median	3.94	4.03	23.4	20.8	7.8	4.525	15.4	22

1 UA: Uric Acid. Creat: Creatinine. ALP: Alkaline Phosphatase. GGT: gamma-glutamyl transferase AST: aspartate

2 aminotransferase. ALT: alanine aminotransferase. Ca: calcium. P: phosphorus and CRP: c-reactive protein.

3
4
5 **Supplementary Table 5.** Maximum and minimum values and median of metabolites,
6 enzymes and minerals in serum of CE treated with FG and DX at 12 EID and the NC.

		NC	FG	DX
UA	Minimum	5.95	5.72	3.42
	Maximum	10.5	30.18	28.8
	Median	8.7	7.93	5.34
Crea	Minimum	0.32	0.19	0.71

	Maximum	2.76	1.3	1.92
	Median	1.62	0.42	1.47
ALP	Minimum	1754	2071	1528
	Maximum	2566	2980	2391
	Median	2200	2180	1896
GGT	Minimum	9.6	5.6	2.7
	Maximum	104.4	164	108
	Median	37.5	40.2	82.8
ALT	Minimum	4.3	10.6	49.4
	Maximum	150	534	95.4
	Median	58.2	22.5	75
AST	Minimum	57.9	65.9	124.3
	Maximum	259.2	175.8	1319
	Median	112.2	104.4	390.3
Ca	Minimum	9.7	4.1	5.9
	Maximum	32.4	31.5	28.8
	Median	13.8	21.3	9.9
P	Minimum	1.1	1.5	5.9
	Maximum	11.4	9	13.2
	Median	4.4	3.8	9.5

1 UA: Uric Acid. Creat: Creatinine. ALP: Alkaline Phosphatase. GGT: gamma-glutamyl transferase AST: aspartate
2 aminotransferase. ALT: alanine aminotransferase. Ca: calcium. P: phosphorus and CRP: c-reactive protein.

3

4

5

6

7

8

9

Supplementary Table 6. Maximum and minimum values and blood count treated with FG and DX inoculated at 12 EID and the NC.

		NC	FG	DX
Hc	Minimum	23	24	18
	Maximum	31	34	26
	Median	30	29	26
Hb 1/3Hc	Minimum	7.67	8	6.67
	Maximum	10.33	11.33	8.67
	Median	10	9.665	8.67
Hb Drabkin	Minimum	8	8.58	6.9
	Maximum	14.08	12.13	9.76
	Median	9.34	10.62	7.84
Erythrocytes	Minimum	1900000	2020000	1600000
	Maximum	3210000	3020000	2290000
	Median	2550000	2745000	1860000
MCV	Minimum	80.99	89.66	113
	Maximum	131.6	118.8	139.8
	Median	117.7	112.5	125.8
MCH	Minimum	28.23	36.07	39.37
	Maximum	53.32	42.45	42.62
	Median	42.11	39.74	41.33
MCHC	Minimum	31.14	31.14	30.17
	Maximum	45.43	43.58	37.54
	Median	34.86	35.55	32.87
Leucocytes	Minimum	2500	2500	6000
	Maximum	5000	8000	30000
	Median	4250	4500	14000
Heterophils	Minimum	2175	1800	5820

	Maximum	4500	7280	29700
	Median	3725	3893	15593
Lymphocytes	Minimum	325	315	140
	Maximum	650	720	500
	Median	427.5	571	220
Monocytes	Minimum	0	0	0
	Maximum	100	100	0
	Median	47.5	75	0
Eosinophils	Minimum	0	0	0
	Maximum	0	0	0
	Median	0	0	0
Thrombocytes	Minimum	4000	4000	6000
	Maximum	7500	7000	23500
	Median	5000	5750	15000

- 1 Ht: Hematocrit. Hg: Hemoglobin. MCH: corpuscular volume. MCH: mean corpuscular haemoglobin. MCHC: mean corpuscular
- 2 haemoglobin concentration.
- 3

6. REFERENCES

1. Kain KH, Miller JWI, Jones-Paris CR, Thomason RT, Lewis JD, Bader DM, et al. The chick embryo as an expanding experimental model for cancer and cardiovascular research. *Dev Dyn*. 2014;243(2):216–28.
2. Lokman NA, Elder ASF, Ricciardelli C, Oehler MK. Chick Chorioallantoic Membrane (CAM) Assay as an In Vivo Model to Study the Effect of Newly Identified Molecules on Ovarian Cancer Invasion and Metastasis. 2012;9959–70.
3. Halik G, Strojny B, Hotowy A, Grodzik M, Piast R. Toxicity studies of six types of carbon nanoparticles in a chicken-embryo model. 2017;2887–98.
4. Hrubá H, Abdelsalam EEE, Anisimov N, Bandouchova H, Havelkova B, Heger T, et al. Reproductive toxicity of fluoroquinolones in birds. *BMC Vet Res*. 2019 Dec;15(1):209.
5. Khosravi A, Sharifi I, Tavakkoli H, Derakhshanfar A, Keyhani AR, Salari Z, et al. Embryonic toxico-pathological effects of meglumine antimoniate using a chick embryo model. Yildirim A, editor. *PLoS One*. 2018 May;13(5):e0196424.
6. Ping, Liu; Deng, Guangfu; Guo, XiaoQuan; Kuang, Jun; Zhang, Caiying; Cao, Huabin and Hu G. Clinicopathology of Gout in Growing Layers Induced by Avian Nephrotrophic Strains of Infectious Bronchitis Virus. 2015;8318.
7. Ribeiro LN de M, de Paula E, Rossi DA, Monteiro GP, Júnior ECV, Silva RR, et al. Hybrid Pectin-Liposome Formulation against Multi-Resistant Bacterial Strains. *Pharmaceutics*. 2020 Aug;12(8):769.
8. Haselgrübler R, Stübl F, Stadlbauer V, Lanzerstorfer P, Weghuber J. An In Ovo Model for Testing Insulin-mimetic Compounds. *J Vis Exp*. 2018 Apr;(134).

9. Guy JS. Isolation and Propagation of Coronaviruses in Embryonated Eggs. *Methods Mol Biol.* 2015;454:109–17.
10. Fauzia E, Barbhuyan TK, Shrivastava AK, Kumar M, Garg P, Khan MA, et al. Chick Embryo: A Preclinical Model for Understanding Ischemia-Reperfusion Mechanism. *Front Pharmacol.* 2018 Sep;9.
11. Shi L, Ko ML, Huang CC-Y, Park S-Y, Hong M-P, Wu C, et al. Chicken Embryos as a Potential New Model for Early Onset Type I Diabetes. *J Diabetes Res.* 2014;2014:1–10.
12. Bronner-Fraser M. *Methods in Cell Biology.* 2017. 409 p.
13. Maier HJ, Bickerton E, Britton P. Coronaviruses: Methods and protocols. *Coronaviruses Methods Protoc.* 2015;1282:1–282.
14. Viana FAB. *Guia terapêutico veterinário.* CEM. 2019;
15. Collier HB. Standardization of Blood Haemoglobin Determinations. *Can Med Assoc J* [Internet]. 1944;50(6):550–2. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20323122><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC1581573>
16. Natt MP, Herrick CA. A New Blood Diluent for Counting the Erythrocytes and Leucocytes of the Chicken. *Poult Sci* [Internet]. 1952;31(4):735–8. Available from: <http://dx.doi.org/10.3382/ps.0310735>
17. Campbell TW. Hematology. In: *Avian Medicine: Principles and Application.* 1997.
18. Thrall MA. *Veterinary Hematology and Clinical Chemistry.* Veterinary Hematology and Clinical Chemistry. 2012. 776 p.
19. Swayne DE, Stockham SL, S.Johnson G. Cytochemical Properties of Chicken Blood Cells Resembling Both Thrombocytes and Lymphocytes. *Vet Clin Pathol.* 1986;15(2):17–24.

20. Justino AB, Pereira MN, Peixoto LG, Vilela DD, Caixeta DC, De Souza AV, et al. Hepatoprotective Properties of a Polyphenol-Enriched Fraction from *Annona crassiflora* Mart. Fruit Peel against Diabetes-Induced Oxidative and Nitrosative Stress. *J Agric Food Chem*. 2017;65(22):4428–38.
21. Aksenov MY, Markesbery WR. Changes in thiol content and expression of glutathione redox system genes in the hippocampus and cerebellum in Alzheimer's disease. *Neurosci Lett*. 2001;302(2–3):141–5.
22. Franco RR, Justino AB, Martins MM, Silva CG, Campana PRV, Lopes JCD, et al. Phytoscreening of Vochysiaceae species: Molecular identification by HPLC-ESI-MS/MS and evaluating of their antioxidant activity and inhibitory potential against human α -amylase and protein glycation. *Bioorg Chem* [Internet]. 2019;91:103122. Available from: <https://doi.org/10.1016/j.bioorg.2019.103122>
23. Tolosa, E. M. C.; Rodrigues, C. J.; Behmer OA., Freitas-Neto AG. Manual de técnicas para histologia normal e patológica. 2003;2. ed.:331p.
24. Brunt EM, Janney CG, Bisceglie AM Di, Neuschwander-tetri BA, Bacon BR. Nonalcoholic steatohepatitis: A proposal for grading and staging the histological lesions. 1999;94(9).
25. Angulo P. Nonalcoholic fatty liver disease. *Encycl Mol Mech Dis*. 2009;346(16):1429–1429.
26. FánCSI T. Ultrastructural Studies of the Goose Embryo Liver. *Anat Histol Embryol*. 1982;11(2):138–46.
27. Elfarnawany MH. Signal Processing Methods for Quantitative Power Doppler Microvascular Angiography. *Electron Thesis Diss Repos* [Internet]. 2015;(August). Available from: <https://ir.lib.uwo.ca/etd/3106>
28. Mukaka MM. Statistics corner: A guide to appropriate use of correlation

- coefficient in medical research. *Malawi Med J*. 2012;24(3):69–71.
29. Decaro N. Gammacoronavirus: Coronaviridae. *The Springer Index of Viruses*. 2011;100:403.
 30. Schultze B, Cavanagh D, Herrler G. Neuraminidase treatment of avian infectious bronchitis coronavirus reveals a hemagglutinating activity that is dependent on sialic acid-containing receptors on erythrocytes. *Virology*. 1992;189(2):792–4.
 31. Sheng G. Primitive and definitive erythropoiesis in the yolk sac: A bird's eye view. *Int J Dev Biol*. 2010;54(6–7):1033–43.
 32. Zhang Z, Sun M, Deng J, Yu J, Yang X, Zhao W, et al. Zika virus induced more severe inflammatory response than dengue virus in chicken embryonic livers. *Front Microbiol*. 2019;10(MAY):1–9.
 33. Zhou Y, Frey TK, Yang JJ. Viral calciomics: Interplays between Ca²⁺ and virus. *Cell Calcium*. 2009;46(1):1–17.
 34. Cao Z, Han Z, Shao Y, Geng H, Kong X, Liu S. Proteomic analysis of chicken embryonic trachea and kidney tissues after infection in ovo by avian infectious bronchitis coronavirus. *Proteome Sci* [Internet]. 2011;9(1):11. Available from: <http://www.proteomesci.com/content/9/1/11>
 35. Heiblum R, Arnon E, Chazan G, Robinzon B, Gvaryahu G, Snapir N. Glucocorticoid Administration During Incubation: Embryo Mortality and Posthatch Growth in Chickens. *Poult Sci* [Internet]. 2001;80(9):1357–63. Available from: <http://dx.doi.org/10.1093/ps/80.9.1357>
 36. Eriksen MS, Haug A, Torjesen PA, Bakken M. Prenatal exposure to corticosterone impairs embryonic development and increases fluctuating asymmetry in chickens (*Gallus gallus domesticus*). *Br Poult Sci*. 2003;44(5):690–7.

37. Mashaly MM. Effect of Exogenous Corticosterone on Chicken Embryonic Development. Poult Sci [Internet]. 1990;70(2):371–4. Available from: <http://dx.doi.org/10.3382/ps.0700371>
38. Lin H, Decuypere E, Buyse J. Oxidative stress induced by corticosterone administration in broiler chickens (*Gallus gallus domesticus*): 2. Short-term effect. Comp Biochem Physiol - B Biochem Mol Biol. 2004;139(4):745–51.
39. Bartunek P, Karafiat V, Bartunkova J, Pajer P, Dvorakova M, Kralova J, et al. Impact of chicken thrombopoietin and its receptor c-Mpl on hematopoietic cell development. Exp Hematol. 2008;36(4):495–505.
40. Wong GK, Cavey MJ. Development of the liver in the chicken embryo. I. Hepatic cords and sinusoids. Anat Rec. 1992;234(4):555–67.
41. Maxwell MH. Avian blood leucocyte responses to stress. Worlds Poult Sci J. 1993;49(1):34–43.
42. Bellairs R, Osmond M. The Atlas of Chick Development Third Edition. 2014. 225 p.
43. Vicuña, E. A.; Kuttappan VA, Latorre JD, Faulkner OB, Hargis BM, Tellez G, Bielke LR. Effect of dexamethasone in feed on intestinal permeability , differential white blood cell counts , and immune organs in broiler chicks. 2012;2075–80.
44. Jain NC. Essentials of veterinary hematology. 1993;
45. Aengwanich W. Effects of dexamethasone on physiological changes and productive performance in broilers. 2007.
46. Elsayed M, Mohamed N, Hatab M, Elaroussi M. Oxidative Stress of in-Ovo Ochratoxin A Administered during Chick Embryonic Development. Brazilian J Poult Sci. 2019;21(1).

47. Ritchie, J. Branson; harrison, J. Greg; Harrison LR. Avian Medicine: Principles and application. 2009. 2009 p.
48. Mandelker L, Vajdovich P. Oxidative stress in applied basic research and clinical practice - Studies on experimental models. Vol. 56, European Journal of Histochemistry. 2012. 9 p.
49. Lin H, Decuypere E, Buyse J. Oxidative stress induced by corticosterone administration in broiler chickens (*Gallus gallus domesticus*): 1. Chronic exposure. *Comp Biochem Physiol - B Biochem Mol Biol*. 2004;139(4):737–44.
50. Hasegawa M. A thermodynamic model for denaturation of granulocyte colony-stimulating factor: O-linked sugar chain suppresses not the triggering deprotonation but the succeeding denaturation. *Biochim Biophys Acta (BBA)/Protein Struct Mol*. 1993;1203(2):295–7.
51. Weiss, Douglas J.; Wardrop KJ. Schalm's Veterinary Hematology. 2010.
52. Bussolino F, Ziche M, Wang JM, Alessi D, Morbidelli L, Cremona O, et al. In Vitro and In Vivo Activation of Endothelial Cells by Colony-stimulating Factors. 1991;87(March):986–95.
53. Bussolino, Federico; Wang, Ming Ji; Defilippi, Paola; Turrini, F.; Sanavio, Florella; Edgell, S. J. C.; Aglietta, Massimo; Arese, Paolo; Mantovani A. Granulocyte and granulocyte-macrophage colony stimulating factors induce human endothelial cells to migrate and proliferate. 1989.
54. Bocchietto E, Cucielmetti A, Silvagno F, Bussolino F, Cenetica D, Medica C, et al. Proliferative and Migratory Responses of Murine Microvascular Endothelial Cells to Granulocyte-Colony-Stimulating Factor. 1993;95:89–95.