Comparison of Bacterial And Archaeal Microbiome In Two Bioreactors Fed With Cattle Sewage And Corn Biomass

Caterina Agrimonti

University of Parma

Giovanna Visoli (✉ giovanna.visoli@unipr.it)

University of Parma <https://orcid.org/0000-0001-9712-3223>

Graziano Ferrari

AVG Consulting Srl via Farini 1

Anna Maria Sanangelantoni

University of Parma

Research Article

Keywords: microbial communities, bioreactors, cattle manure, molecular analysis

Posted Date: December 14th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1050429/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

The bacterial and archaeal communities of two biogas producing plants (P1 and P2), associated with a 999 kW cogeneration unit, both located in North Italy, were analyzed at start up and fully operating phases, by means of various molecular approaches: i) Automated Ribosomal Intergenic Spacer Analysis; ii) cloning and sequencing of PCR amplicons of archaeal genes *16Srrna* and *mcrA*; iii) 16S rDNA high throughput next generation sequencing. P1 and P2 use the same technology and both were fed with cattle manure and corn silage. During the study of P1 also the post digester (fed with pig manure) was analyzed. The aim of this research was to characterize the bacterial and archaeal community in two very similar plants to profile the core microbiome. The results of this analysis highlighted that the two plants (producing comparable quantities of volatile fatty acids, biogas, and energy) differed in anaerobic microbiota (Bacteria and Archaea). Notably the methanogenic community of P1 was dominated by the strict acetoclastic *Methanosaeta* (*Methanotherix*) (up to 23.05%) and the unculturable *Candidatus Methanofastidiosum* (up to 32.70%), while P2 was dominated by the acetoclastic, but more substrate-versatile, *Methanosarcina* archaeal genus (49.19%). The data demonstrated that the performances of plants with identical design, in similar operating conditions, yielding comparable amount of biogas (average of 8662 m³/day and 7916 m³/day respectively for P1 and P2), VFA (1643 mg/L and 1634 mg/L) and energy recovery (23.90-24 MWh/d) depends on the stabilization of an effective and functionally optimized methanogenic community rather than on the species composition

Statement Of Novelty

Even though many papers have been published about utilization of organic waste for biogas production, the relationships between microbial communities operating in different plants and efficiency of biogas production, is a still opened question. In this work, we have found that bacterial and archaeal species are different also in plants located in the same geographical area, fed with the same substrate (cow manure and corn silage) and operating at the same conditions. However, these differences do not affect the efficiency of biogas production. Interestingly, our results evidenced the role of not well characterized species (i.e. *Candidatus methanofastidiosum*) in the process, suggesting that other pathways than those so far identified, are involved in biogas production.

Introduction

In the last years, the biogas production, starting from organic wastes, has been considerably increased to reduce the consumption of fossil fuels and the environmental impact of methane emissions from zoogenic fluids [1]. The biogas produced by anaerobic digestion from such renewable sources is transformed in fuels that can be used for energy production and transport.

Despite the high operating costs of these plants and the necessity of significant financial incentives, the anaerobic digestion (AD) is a well-established technology and remains a fundamental energy source in the emerging market for renewable energy. This is the reason why, in the circular economy perspective, it is considered a key technology on the path to independence from fossil fuels [2,3]. Italy is one of the leading countries in the biogas production where agricultural, landfill, sewage and manure substrates are converted into biogas using AD and into electricity and heat as illustrated by Benato and Macor in 2019 [4].

As many commercial AD facilities suffer from several problems like long start up times or system instability, the optimization of the process is an important deal. The attention, in this regard, had formerly been focused on the evaluation of substrates and the implementation of plants characteristics, but it was soon hypothesized that the efficiency of the plants was more likely dictated by the composition and activity of the microbial communities from the incoming substrates [5].

The overall process of biogas production is due to the complex, synergistic, interaction among four functional guilds performing the different reactions (hydrolysis, acidogenesis, acetogenesis, methanogenesis). Early in the process, microbial enzymes hydrolyse polysaccharides, proteins and fats into sugars, amino acids and fatty acids. These compounds are then converted into volatile fatty acids (VFAs) and alcohols that are further converted into acetic acid, CO₂ and hydrogen (Fig.1). Methanogenic microorganisms belonging to the Archaea domain carry out the final step, the production of methane [6, 7]. Acetoclastic and hydrogenotrophic methanogens, transforming acetic acid, CO₂ and hydrogen into methane are particularly sensitive to process conditions, like temperature, pH, redox and inhibitors like sulfidric acid produced through the reduction of sulphates [8, 9, 10]. The methanogenic Archaea, mostly represented in biodigesters, belong to *Methanomicrobiales*, *Methanosarcinales* and *Methanococcales* orders [11,12]. The process of biogas formation has been thoroughly dissected, however a little number of prokaryotes involved in anaerobic digestion have been so far isolated because many microbial species are not culturable and little is known about their dynamics and interactions. Methods based on DNA analysis demonstrated to be the best way to study AD [13, 14, 15]. The gene encoding ribosomal 16S RNA in Bacteria and Archaea is the most used target sequence for many molecular techniques, together with intergenic spacer (IGS) between the small (16S) and large (23S) rRNA genes in the *rna* operon. In addition, methanogens can be identified by targeting the gene *mcrA*, encoding α -subunit of methyl-coenzyme M reductase (MCR), the key enzyme involved in final steps of methanogenesis. Unlike other enzymes involved in methanogenic metabolism, MCR is highly specific for methanogens [16, 17].

The molecular techniques used for microbiome analysis include ARISA (Automated Ribosomal Intergenic Spacer Analysis) analyzing IGS, clone library sequencing and NGS (Next Generation Sequencing). ARISA, based on the different length of ribosomal intergenic spacers, is a molecular technique for characterizing non-cultivable microbial communities directly from environmental DNA (the whole DNA extracted from samples). It has been widely

applied for a rapid analysis of biodiversity in large samples, but suffers the limitation of not allowing the recognition, at a taxonomic level, of the species present in the sample [18]. The sequencing of 16S *rma* or *mcrA* genes clone libraries offers a rather precise identification of microorganism but it does not cover the whole microbiome diversity. 16S rDNA NGS is much more exhaustive about the microbiome diversity but is less precise in the identification of microorganisms at species level.

In this study, in order to describe the microbiome evolution in two full-scale biodigesters, managed in the same way, in the different steps of their operation, an integrated approach was followed consisting in *i*) ARISA analysis, to generally characterize the bacterial diversity among all the samples collected; *ii*) amplification, cloning and sequencing of archaeal 16S *rma* and *mcrA* genes in one sample for each digester; *iii*) 16S profiling with the NGS approach to describe the relative abundance of phyla/genera of Bacteria and Archaea. The anaerobic biodigesters described (P1 and P2) are operating in two sites in Lombardia (Italy) separated by 200 km distance. Both plants were fed with cattle sewage and corn silages as substrates and operated in mesophilic regimen. P1 was followed from the startup phase up to the phase of full operation (included), while P2 was analyzed when already operative at full capacity. As for P1, a sample of the post-digester was included in the analyses.

Although several studies have studied and compared microbial communities of laboratory-scale biodigesters or full-scale digesters [5, 12, 14, 15, 19] the novelty of this study is the use of a simplified system in which two full-scale digesters, with identical technical characteristics, conducted in the same way and both fed with cattle sewage and corn biomass, derived from the respective local farms, in different seasons, were compared. It was interesting to discover that the two plants performed in very similar way despite a different phylogenetic composition of the bacterial and archaeal communities. The important finding is that an effective microbial succession developed, starting from the inoculum, and reached in both cases an optimized (although different) equilibrium of functions.

Materials And Methods

Plant characteristics and management, sample collection and DNA extraction

The biodigesters location was the Po Valley: P1 N 44° 57' 55.30" E 10° 27' 57.63"; P2 N 45° 10' 15.52", E 8° 38' 45.01" (DMS). Both plants, were CSTR (Continuous Stirred Tank Reactor) reactors with 7200 m³ reaction volume and hydraulic retention time of 120 days. Operational parameters were monitored with online instruments: for the gas composition analysis an Awite series 7 analyzer (Await Bioenergia S.r.l Bolzano, Italy) was used; the Ultrasonic flow measuring system was Prosonic Flow 200 (Endress+Hauser Instruments International AG, Switzerland); a PT100 sensor was used to measure temperature and the portable pH 3110 (WTW-Xylem Analytics-Germany) for measuring pH. Chemical analyses were done by external laboratories (Accredia certificated) following the standard procedures: Water & Life Lab <https://www.waterlifelab.it/> and Studio Alfa-
<https://www.studioalfa.it/>.

Plant P1 was monitored since the start-up phase in the autumn-winter. Four phases were identified (Fig.2): I) 03 Oct-03 Nov; II) 04 Nov-5 Dec; III) 6 -18 Dec; IV) 19-31 Dec. In the phase I) the plant was loaded with a total of 1120 t of bovine sewages. In addition, an inoculum of 540 t of a substrate coming from a similar plant, already operative, was added on 28-29 Oct to speed up the anaerobic digestion. In the phase II) the manure amount added to biodigester was reduced and dispensed more regularly, although not daily (from 11.60 t /day up to 84.20 t day⁻¹). From 6 Dec the average daily amount of sewage added was 20.24 t.

The corn biomass was regularly loaded into biodigester from 7 Nov, starting with 1 t, then gradually increasing until 15 Dec when 44.50 t were loaded. The daily load of plant biomass was then regulated around 39-40 t/day. In addition, in the phase III) a vertical axis mixer was used to integrate two smaller submerged mixers. Finally, in the phase IV), the primary digester was integrated with a supporting post digester where pig manure was added. The whole process was daily monitored for temperature, pH, VFAs, biogas production and energy cogeneration.

The P2 biodigester was started on 15 Mar, and became fully operative by 13 Apr. Since then, the average load of cattle sewage was 37.84 t/day with corn silage varying from 19.90 to 49.20 t/day. The collection of samples started 8 May.

Five aliquots of digestate were collected in both plants, at each specific point and stored at -20 °C until DNA extraction. In Table 1 are reported the data concerning VFA production (mg/L), T (°C), pH, Alkalinity (mg/L), ratio VFA/Alk, Ammonia nitrogen concentration (mg/Kg), the amount of corn silage (t /day) and manure (t/day) added to the biodigesters, biogas production (m³/day), energy production (MWh/day) and are highlighted the samples used for microbial analysis.

All the samples were analyzed with ARISA (see afterwards) and only some representative samples were selected for clone libraries construction and 16S rDNA NGS profiling (Table 1).

Samples were centrifuged at 14000 x g for 10 min, supernatant was discarded and DNA was extracted from the pellets with FastDNA SPIN Kit for soil and FasPrep® Instrument (MP Biomedicals, Santa Ana, CA). DNA samples were checked with electrophoresis on agarose gel 0.9%.

PCR conditions

All PCR were conducted with primers pairs listed in Table 2, in a mix of 25 µl containing 5 µl of 1:1000 diluted DNA, 1X PCR Buffer, 200 µM dNTPs, 100nM primers and 0.8 U µl⁻¹ of Taq. A thermal cycler TECHNE TC-512 (Bibby Scientific Ltd, Staffordshire, UK) was used as follows: hotstart 94 °C for

3 min, 35 cycles of denaturation (94 °C for 45 sec), annealing (58 °C for 1 min) and extension (72 °C for 2 min) followed by a final extension (72 °C for 7 min).

The primer ReubR, for the amplification of bacterial the ITSReub, was 5'-6 FAM labelled to detect amplicons through capillary electrophoresis.

ARISA (Automated Ribosomal Intergenic Spacer Analysis)

Amplicons obtained from primers ITSF/ITSReub (Table 2) were purified with Wizard®SV Gel and PCR Clean-Up System (Promega, Madison WI, US). An aliquot of 4 µl of purified DNA was mixed with 9 µl of Hi-Di™ formamide (Applied Biosystems, Foster City, CA, US) and 1 µl of GeneScan-2500 ROX™ Size Standard (Applied Biosystems). After denaturation at 95 °C for 3 min and fast cooling, DNA was injected in an ABI Prism 310 (Applied Biosystems, CA, US) capillary filled with POP-4™ Performance Optimized Polymer for electrophoresis. ARISA electropherograms were analyzed using the GeneScan 3.1 software program and elaborated with Microsoft Excel. The raw data were processed excluding the peaks whose heights and areas, proportional to the intensity of the fluorescence, were less than 1% of the sum of the same. Then the peaks were grouped according to differences of 4 bp for the fragments up to 700 bp and 5 bp for those up to a length of 1000 bp [20, 21]. For each sample the presence (1) or absence (0) of a certain peak was determined obtaining a matrix utilised for calculation of Jaccard similarity coefficient and construction of dendrograms using the programme DendroUPGMA, available at the site <http://genomes.urv.cat/UPGMA>.

Amplification and cloning of 16S*rma* and *mcrA* (methyl coenzyme M reductase A) genes from the archaeal community

The DNA extracted from selected biodigesters samples (see Table 1) was diluted 1:100 and amplified with the primers, specific for Archaea, reported in Table 2 (69F and ARC934R for archaeal 16S rRNA gene and the pair *mcrAF*/*mcrAR* for methyl coenzyme M reductase A gene for methanogenic Archaea). The amplicons obtained were purified by GFX PCR and a Gel Band Purification Kit (GE Healthcare UK Limited –HP7 9NA, UK), ligated into pGEM®-T Easy vector System I (Promega) and transformed into the strain JM109 of *Escherichia coli*. The recombinant colonies were subjected to colony PCR with primers SP6/T7 (Table 2) to check the presence of inserts.

ARDRA (Amplified Ribosomal DNA Restriction Analysis) and sequencing

In order to select, for sequencing, the recombinant plasmids containing fragment of 16S*rma* and *mcrA* genes from different Archaea, the ARDRA analysis was performed respectively on 100 and 50 clones for each clone library. Restriction analysis of the cloned inserts was conducted with the enzyme *Hae*III for 16S*rma*, and with enzymes *Rsa*I and *Alu*I for *mcrA*. 5 µl of amplified DNA was digested with 5 U of each enzyme and 1X of the respective buffer at 37 °C for 4 hr.

The restriction fragments were loaded on 2.5% NuSieve® low melting agarose gel (Lonza Group Ltd, Basel, Switzerland) on TBE buffer 0.5% (45 mM Tris borate, 1 mM EDTA pH 8.0) and subjected to electrophoresis at 60 V for 4 hr. Images were acquired with Biorad Gel Doc 2000 and the software Quantity One (Bio-Rad, Hercules, CA, US).

One representative clone for the most represented ARDRA profiles was selected for the partial sequencing of 16S*rma* gene and *mcrA* at BMR Genomics srl in Padova (Italy). The obtained sequences were then compared with those stored in the GenBank database at the NCBI (National Center for Biotechnology Information) by using the BLAST (Basic Local Alignment Search Tool) program (<http://www.ncbi.nlm.nih.gov/blast>). The sequences were deposited in GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) and their accession number are reported in Table 3

Biodigesters microbiome profiling by 16S rRNA gene amplification and NGS sequencing

The bacterial and archaeal community profiles of the samples were generated by NGS technologies at the Genprobio S.r.l. Laboratory. Partial 16S rRNA gene sequences of duplicate samples were obtained from the extracted DNA by Polymerase Chain Reaction (PCR), using the primer pair Probio_Uni and Probio_Rev, targeting the V3 region of the 16S rRNA gene sequence [22] for bacterial community and the primers pair ArchV56 [23] for archaeal community. Amplifications were carried out using a Verity Thermocycler (Applied Biosystems) and PCR products were purified by the magnetic purification step involving the Agencourt AMPure XP DNA purification beads (Beckman Coulter Genomics GmbH, Bernried, Germany) in order to remove primer dimers. Sequencing was performed using an Illumina MiSeq sequencer with MiSeq Reagent Kit v3 chemicals. The fastq files were processed using a custom script based on the QIIME software suite [24]. Paired-end read pairs were assembled to reconstruct the complete amplicons.

To calculate downstream diversity measures, operational taxonomic units (OTUs) were defined at 100% sequence homology using DADA2 [25] OTUs not encompassing at least two sequences of the same sample were removed. All reads were classified to the lowest possible taxonomic rank using QIIME2 and a reference dataset from the SILVA database v132 [26]. The biodiversity of the samples (alpha-diversity) was calculated with Chao1 and Shannon indices. The 16S rDNA reads were sent to GenBank to obtain their under-accession numbers. They are available as bioproject PRJNA534196.

Results

Performance of the reactors

In the plant P1, the biogas production started during phase II, after the addition of corn silage and of an inoculum coming from a similar, fully operating, plant (Fig.2). In the primary digester, the biogas production increased from 4329 m³/day in 24 Nov up to 10.919 m³/day in 22 Dec, with an average daily production of 7916 m³. During the entire process, the pH was around 7.4-7.5, optimal values for substrate degradation and methanogenic activity and the temperature was maintained between 42.2 °C and 44.2°C (average 42.8°C). The concentration of volatile fatty acids (VFA) was, on average, 1643 mg/L (Table 1). At the end of December, 2 month and 20 days from startup, P1 was supported with a secondary biodigester that operated in the same condition of pH and temperature, but was fed with pig slurry instead of cattle slurry. The VFA concentration detected in the secondary biodigester, in 29 Dec, was lower than in primary digester (1152 mg/L). This can be explained by the dilution effect or considering that cows and pigs have different diets and the dry matter content of pigs zootechnical effluent varies from 3 to 6% (versus 8%-15% of cows), and has a faster biological degradability due to the lower amount of fiber.

While the P1 digester was followed and sampled from the start-up phases up to the stabilization phase, the sampling in P2 biodigester was done when the production of biogas had already stabilized.

In P2 the biogas production started at 24 Apr and reached 8662 m³/day by 22 May with an average value of 7796 m³/day (comparable with P1 biodigester). The concentration of VFA was highly comparable to that of P1, with an average value of 1634 mg/L, and also the pH values were similar, while the average temperature was higher (44.2°C). In both plant the daily energy production was comparable in the full operational phase as reported in Table 1 (23.90-24 MWh/ day).

Bacterial community analysis by ARISA

The Automated rRNA Intergenic Spacer Analysis (ARISA) is a molecular method for analyzing bacterial community diversity without the bias imposed by culture-based approaches. With the DNA extracted from all the samples reported in Table 1 in P1 and P2 biodigesters, ARISA profiles were obtained and compared by using a pairwise similarity coefficient, the Jaccard index. The dendrogram resulting from analysis with the programme DendroUPGMA showed that bacterial microbiome of P1 and P2 clustered in two distinct groups and, for each plant, samples were similarly grouped (Fig.3). In P1 digester one cluster encompassed samples CM2 and CM3, collected when the VFA production was almost stabilized, but the biogas production was still low; in the second cluster the communities of CM4, CM5, CM6 and CM7, collected in the phase of temperature and biogas production stabilization, were more similar and separated from CM8 when the biogas production decreased and, as expected, from CMP (the post digester). In P2, similarly, the samples clustered in two main groups one encompassing VL1 to VL6 the other encompassing VL8, VL10 and VL12, characterized by a lower VFA production and higher temperatures. ARISA does not allow description of the community in terms of bacterial genera or species, therefore, to have more detailed information, the bacterial community profiling by NGS of 16S rDNA was performed.

Bacterial community 16S profiling by NGS

The bacterial community profile by NGS of 16S rDNA was determined on duplicate samples CM3, CM8 and CMP (for P1) and VL5 (for P2). MiSeq runs produced an average of 75566 sequences for the samples collected in P1 and 83931 from that collected in P2. The rarefaction curves obtained with Chao 1 and Shannon indices (a measure used to estimate the alpha diversity in samples and evaluate whether sequencing efforts allowed to capture all the microbial diversity) highlighted the lowest index of diversity in VL5. In Fig.4 (panel a) are shown only results with Shannon index. As expected, the most represented phyla in both P1 and P2 biodigesters are the Firmicutes, with similar percentages in all samples (from 62.64% to 74.54%), followed by Bacteroidetes (14.55% to 19.43%) and Cloacimonetes (from 1.57% to 8.72%) (Fig.5).

The most represented class is that of Clostridia (between 50.50 and 61.98%), that includes the uncultured group MBA03 (10.79% to 21.27%), the genus *Sedimentibacter* (7.93% to 13.94%), and *Clostridium sensu stricto*, present at 12.29% in CM3 and in lower percentages (< 2%) in the other samples (Fig.6). The main differences between the plants P1 and P2 concern the presence of *Clostridium sensu strictu*, very low in VL5 (1.16%), *Hydrogenispora*, (8.93% in P2 vs < 1.5% in all samples of P1), uncultured bacterium of *Lentimicrobiaceae* family (9.47% in P2 vs <2% in all samples of P1), *Proteiniphilum* (5.29% in P2 vs <1.5% in all samples of P1) and a bacterium belonging to the *Bacteroidales* family UCG-001 (3.10-3.58% in P1 samples vs < 0.28% in P2). In the plant P1 is remarkable the presence of the family W27, (class *Cloacimonetes*) that increases from 1.15% of CM3 to 7.39% in CMP and is absent in the plant P2 (Fig.6).

Archaeal community analysis by cloning and sequencing of 16S rRNA genes and *mcrA* genes

The first approach for archaeal community characterization was carried out through classical molecular methods, namely the cloning and sequencing of amplified target genes. The sequences encoding partial 16S rRNA (about 870 bp) and part of the gene encoding methyl CoM reductase alpha subunit (*mcrA*) gene (about 500 bp) were amplified from the DNA extracted from samples CM8 (in P1) and VL10 (in P2) and cloned into the vector pGEM®-T Easy. The cloned fragment (100 clones/gene library) were analyzed by ARDRA and generated eight reproducible profiles (A-E for P1 and F-I for P2) (Table 3). Sequence analysis of the phylotype A, representing 36% of the analyzed clones, showed 98.20% similarity with *Methanotrix soehngenii* (*Methanosaeta*) an acetoclastic archaeum. Other methanogenic species belonging to genus *Methanoculleus* (hydrogenotrophic and similar to *Methanobacterium*) as well as uncultured bacterial species were identified in P1 (Table 3) while in P2 the dominant archaeal genera were the acetoclastic, but more versatile, *Methanosarcina* (98.80% similarity with the specie *thermophila*) and *Methanoculleus*. Within these samples, several *Clostridiales* and other uncultured bacteria were found. This may be due to the utilization of degenerate primers amplifying also bacterial sequences very abundant in biodigesters.

Archaeal community 16S profiling by NGS

The MiSeq runs produced an average of 10912 sequences for the samples collected in P1 and 11184 from those collected in P2. Rarefaction curves obtained with Chao1 and Shannon biodiversity indices highlighted, as for bacteria, the lowest index of diversity in sample VL5 (P2) and the highest in sample CM3 (P1). In Fig.4 (panel b) are shown only the rarefaction curves obtained with Shannon index, but identical results were obtained with Chao1 biodiversity index (data not shown).

The most represented genera in both plants, are *Methanobacterium*, *Methanosarcina*, *Methanosaeta*, *Candidatus Methanofastidiosum*, *Candidatus Methanoplasma*, *Methanobrevibacter*, uncultured microorganisms of *Bathyarchaeia* class, and *Methanoculleus* (Fig.7). *Methanobacterium* is present in both plants, at percentages variable between 14.37 – 22.87% in P1, and 31% in P2.

P2 has a prevalence of *Methanosarcina* (49.19%), which is present, in P1, in the samples CM3 and CMP (8.38% and 4.62% respectively), and in very low amount in CM8 (0.73%). This was highlighted also with the approach of cloning and sequencing the 16S*rrnA* and *mcrA* genes, previously described. P1 has a significant abundance of *Methanosaeta* in all samples (20%), almost absent in P2 (< 0.05%). P1 is also characterized by the presence of *Ca. Methanofastidiosum*, that increases from 10.30% in sample CM3 to 24.70% in CMP and 32.70% in CM8; while it is almost absent in P2 (0.44%). Samples CM3 and CMP of P1 contains *Methanobrevibacter* (10.80% and 6.42%, respectively), a genus strongly reduced in samples CM8 (1.45%) and VL5 (0.39%) of P2. *Bathyarchaeia* are present in samples of both plants, but in variable percentages: 19.98% in CM3, 5.44% in CM8, 2.22% in CMP and 8.88% in VL5. *Methanoculleus* is rather abundant in sample CM8 (4.87%) while is less than 2.5% in the other samples. CMP is characterized by *Ca. Methanoplasma* (10.05%), absent in P2 sample and very low (< 2%) in CM3 and CM8. The sample CMP is characterized by the presence of *Ca. Methanoplasma* (10.05%) absent P2 sample, and very low (< 2%) in CM3 and CM8.

Discussion

This work highlighted that two full-scale, structurally identical, biogas producing plants (P1 and P2), located in two different sites in North Italy, associated with a 999 kW cogeneration unit, both fed with bovine manure and corn silage, and analyzed in two different seasons, highly differ in the archaeal microbiome and, even at less extent, in bacterial microbiome, but do not differ in biogas and energy productivity.

Bacterial biodiversity

The Automated Ribosomal Intergenic Spacer Analysis (ARISA), a molecular method for analyzing bacterial community diversity without the bias imposed by the cultivation steps, showed that bacteria of P1 and P2 cluster in two distinct groups which are consistent with the operational phases (Fig.3).

The deeper analysis with NGS 16S profiling showed that bacterial populations of all samples are dominated by the phylum Firmicutes followed by Bacteroidetes. The members of these phyla have a wide metabolic capacity, including degradation of lignocellulose of corn silage and residual of bovine digestion [27]. Firmicutes are often present in fermentative plants, and in some cases, they are the most abundant bacteria [28, 29, 30]. Indeed, they possess the ability to metabolize a large variety of molecules into acetate and butyrate, VFA intermediate, and therefore they play an important role in the hydrolysis of primary substrates in anaerobic digesters. P1 was characterized by an increase in the CM8 and CMP samples of the phylum Cloacimonetes, very low in P2. Some authors suggest that members of this phylum are involved in lignocellulose degradation and play a role in the syntrophic oxidation of propionate [30, 31]. The genome reconstruction of two candidates of Cloacimonetes, *Candidatus Cloacimonas acidaminovorans* [32] and *Candidatus Syntrophosphaera thermopropionivorans* [33] indicate that they possess the genes of the methylmalonylCoA pathway responsible of the oxidation of propionate. The oxidation of propionate and other short-chain fatty acids is a key step for methanogenesis to occur, as it produces acetate and H₂ utilizable by both acetoclastic and hydrogenotrophic methanogens.

Archaeal biodiversity

Either cloning and sequencing of archaeal target genes and NGS archaeal 16S profiling pointed out that biogas plants P1 and P2 are characterized by the presence of *Methanobacterium* (14.37 to 22.87% in samples from P1 and 31% in P2). *Methanosaeta* is abundant in P1 (up to 23.05%), while *Methanosarcina* is the predominant genus in P2 (49.19%), in which *Methanosaeta* is almost absent. This means that, although P1 and P2 have a similar trend in biogas and VFA production and operate in the same conditions of pH and temperature, they must have a substantial difference in the metabolic fermentative pathways. *Methanosarcina* and *Methanosaeta* are both acetoclastic methanogens, but operate, respectively, at high and low concentration of acetate [34]. Moreover, differently from members of *Methanosaeta* genus, that are strictly acetoclastic, most *Methanosarcina* are mixotrophic, they utilize not only acetate, but also hydrogen derived from acetate oxidation conducted by anaerobic bacteria (in general *Clostridia*) and are able to produce methane even from methanol and methylamines. Archaea belonging to *Methanosarcina* genus, in addition, are able themselves of conducting acetate oxidation, bypassing the bacterial action [35]. Differently from *Methanosarcina*, *Methanosaeta* are rather sensitive to VFA and ammonia but in this work VFA concentration does not significantly differ in P1 and P2 and ammonium concentration, even though is different in P1 and P2, never exceeds 2000 mgL⁻¹. Therefore, the difference may be related to environmental condition of manure storage, considering that P1 started in autumn-winter, while P2 operates in spring-summer. Powell et al. (2008) observed that ammonia production and emission were significantly higher in spring than in fall-winter, not only because the temperature increases the evaporation, but also because the excreted N in urine is higher and urease

activity is stimulated, producing ammonia [36, 37]. In this case the concentration of ammonium was lower in the spring/summer bioreactor (P2) where *Methanosarcina* was the most abundant methanogen.

Karakashev et al. [35] suggest that, in the absence of *Methanosaeta*, the hydrogenotrophic methanogenic pathway dominated, also in presence of *Methanosarcina* that shifted their metabolism from acetoclastic to hydrogenotrophic methanogenesis. The same authors showed that acetate oxidation is reduced in presence of *Methanosaeta*. The P1 microbiome has an evolution characterized by a decrease of *Methanosarcina* (from 8.38 of CM3 to 0.73 of CM8) and by a parallel increase of *Candidatus Methanofastidiosum*, (formerly called WSA2 group) while *Methanosaeta* remains substantially unchanged. The genome analysis of *Ca. Methanofastidiosum* conducted by Nobu et al. [38] revealed that this group has the peculiar capacity of producing methane through methylated thiol reduction. Moreover, members of this group may utilize acetate (as well as malonate or propionate) with CO₂ as carbon source. It is possible that, in this way, they maintain low the acetate concentration favoring the growth of *Methanosaeta*. Our data showed a certain competition between *Methanosarcina* and *Methanosaeta*, but also between *Methanosarcina* and *Ca. Methanofastidiosum*. In general, *Methanosarcina* prevails over *Methanosaeta* because of the greater tolerance to NH₄⁺, acetate and the metabolic versatility; but it is possible that *Ca. Methanofastidiosum* prevails on *Methanosarcina*, favoring the development of *Methanosaeta*. The group of WAS2 was detected in different environments, like freshwater and marine sediments, contaminated groundwater and sludge bioreactor [38, 39]. Recently this genus has been detected also in municipal sludge bioreactor [40] and in bioreactors fed with pig manure [41] but, being uncultivable microorganism, their metabolic features were deduced only by genome analysis. However, a possible competition of WAS2 with *Methanosarcinales* for acetate has been hypothesized by Rivière et al. [5].

Methanobrevibacter is a typical microorganism of rumen and manure, operating a hydrogenotrophic metabolism [42]. It is present in the initial phases of bioreactor activity, with fresh manure inoculum, but it is strongly reduced in the advanced phases, where it is replaced by *Methanosaeta*. It is almost absent in VL5 where *Methanosarcina* predominates. A shifting of *Methanobrevibacter* to *Methanosarcina* has been observed in another research by Ciotola et al. [43]. However, the global percentage of hydrogenotrophic Archaea remains substantially unchanged, in CM3 and CM8, as the reduction of *Methanobrevibacter* is compensated by an increase of *Methanobacterium* and *Methanoculleus*. The percentage of hydrogenotrophic (around 30%) is similar also in CMP and VL5, however, in VL5 *Methanobacterium* is the prevalent hydrogenotrophic microorganism. The presence of acetoclastic Archaea is constant in all samples, being the increase of *Methanosaeta* compensated by the decrease of *Methanosarcina*.

In the ARISA analysis of bacterial community, the post digester of P1 (CMP) clusters separately from the samples CM4,5,6,7, as expected since it was fed with pig manure and had lower VFA concentration. Nevertheless, it seems to be closer to CM8 and this can be explained with the massive inoculum from the primary digester to the secondary digester. The lowering of VFA concentration can be the result of dilution, and the similarity of microbial communities can be ascribed to the inoculum. The NGS analysis of archaeal community indicated a similar trend, with less differences between CM8 and CMP where a predominance of *Candidatus Methanofastidiosum*, *Methanosaeta* and about 30% of hydrogenotrophic Archaea were detected. It is rather surprising the increase of *Methanosarcina* to 4.62% in comparison to 0.72 % of CM8, and a slight reduction of *Methanosaeta*. These results are somewhat in contrast with data reported in the literature, but according to the observation that low concentrations of VFA favor the development of *Methanosaeta* and not of *Methanosarcina* [44]: The slight reduction of *Candidatus Methanofastidiosum*, seems to confirm its competition with *Methanosarcina*. Another peculiarity of CMP is the presence of *Candidatus Methanoplasma*, a group of obligate hydrogen-dependent methylotrophs, but being *Candidatus Methanoplasma termitum* the only deeply characterised species of this group [45] it is difficult to infer its role in CMP and in methanogenesis in general.

Conclusions

In this study an integrated molecular approach was applied to analyze and compare the bacterial and archaeal communities of two full-scale Anaerobic Digestors (7200 m³ reaction volume). The two plants, identical in structure, are used for production of biogas and energy from corn silage and bovine manure and are located in Po Valley at 200 km distance. The plants are managed from the same company, in a comparable way. The molecular analyses were conducted from the phase of start-up to the phase of full operation in autumn-winter for P1, while P2 was analyzed when already in full activity during the spring.

It was possible to observe that, in the plant P1, there was, along four phases, a progressive stabilization of the biogas production, that reached values comparable to that of P2. This was accompanied by a reduction of biodiversity of bacterial and archaeal communities, thus reflecting a competition and selection of microbial populations and microbial functions. Although the plant technology, the operating characteristic, the biogas and energy production were comparable in both plants, the microbiome, in particular Archaea, resulted to be notably different. This may be linked to the different seasons in which the plants were monitored and the samples collected, but also highlights that biogas production efficiency does not necessarily depend on the phylogenetic structure of microbial community acting in AD, but rather on their optimal synergic activity. Indeed, despite the differences observed, the percentage of hydrogenotrophic methanogens remained substantially unchanged in the two plants, suggesting that this pathway is necessary. Was also interesting to observe the relevant presence of *Candidatus Methanofastidiosum*, an unculturable Archaeum found in extreme environment and, more recently, in some bioreactors, probably involved in methanogenesis through methylated thiol reduction. This underlines that different pathways are relevant in methane production, other than acetoclastic and hydrogenotrophic.

From the data obtained, it is possible to conclude that, in the two biogas plants, two different effective microbial succession developed independently, starting from the inoculum and reached, in both cases, an optimized (although different) equilibrium of the community, ensuring the necessary

metabolic functions. Although the phylogenetic composition of the two communities showed important differences it appears that these microbial communities carry out similar functional processes, regardless of differences in their structure (functional similarity).

Langer et al (2015) already reached these conclusions describing the functional redundancy and structural changes of microbial communities in four lab-scale (12 L), continuously stirred tank reactors. The diverse microbial communities optimized their metabolism in a way that ensured efficient biogas production [44]. With the present study, conducted in full scale dimension, the same conclusion can be drawn.

Declarations

The authors declare that they have no Conflict of interest.

Data Availability

Data are available within the article. NGS and conventional sequences are available in GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>).

Funding

This research was supported by funds to Prof. Anna Maria Sanangelantoni from FIL, of the University of Parma Local Funding for Research

Author contributions

Caterina Agrimonti: Formal analysis, Data curation, Writing- Original draft preparation

Giovanna Visioli: Validation, Writing - Review & Editing

Anna Maria Sanangelantoni: Conceptualization, methodology, investigation, writing and reviewing.

Graziano Ferrari: Sample and data collection

References

1. Marañón, E., Salter, A.M., Castrillón, L., Heaven, S., Fernández-Nava Y.: Reducing the environmental impact of methane emissions from dairy farms by anaerobic digestion of cattle waste. *Waste Manag.* 31, 1745-1751 (2011)
2. Stahel, W.R.: The circular economy. *Nature*, 531, 435-438 (2016). doi: 10.1038/531435a.
3. Blades, L., Morgan, K., Douglas, R., Glover, S., De Rosa, M., Cromie, T., Smyth, B.: Circular Biogas-Based Economy in a Rural Agricultural Setting. *Energy Procedia* 123, 89–96 (2017). <https://doi.org/10.1016/j.egypro.2017.07.255>
4. Benato, A., Macor, A.: Italian Biogas Plants: Trend, Subsidies, Cost, Biogas Composition and Engine Emissions. *Energies* 12(6), 979 (2019). <https://doi.org/10.3390/en12060979>.
5. Rivière, D., Desvignes, V., Pelletier, E., Chaussonnerie, S., Guermazi, S., Weissenbach, J., Li, T., Camacho, P., Sghir, A.: Towards the definition of a core of microorganisms involved in anaerobic digestion of sludge. *ISME J.* 3, 700–714 (2009). <https://doi.org/10.1038/ismej.2009.2>.
6. Enzmann, F. Mayer, F., Rother, M., Holtmann. D.: Methanogens: biochemical background and biotechnological applications. *AMB Express* 4, 8 (2018), <https://doi.org/10.1186/s13568-017-0531-x>.
7. Conrad, R.: Importance of hydrogenotrophic, acetoclastic and methylotrophic methanogenesis for methane production in terrestrial, aquatic and other anoxic environments: A mini review. *Pedosphere* 30, 25-39 (2020). [https://doi.org/10.1016/S1002-0160\(18\)60052-9](https://doi.org/10.1016/S1002-0160(18)60052-9).
8. Yekta, S.S., Ziels, R.M., Björn, A., Skjellberg, U., Ejlertsson, J., Karlsson, A., Svedlund, M., Willén, M., Svensson, B.H.: Importance of sulfide interaction with iron as regulator of the microbial community in biogas reactors and its effect on methanogenesis, volatile fatty acids turnover, and syntrophic long-chain fatty acids degradation. *J.Biosci. Bioeng.* 123(5), 597-605 (2017). <https://doi.org/10.1016/j.jbiosc.2016.12.003>.
9. Dai, X.H., Hu, C.L., Zhang, D., Dai, L.L., Duan, N.N.: Impact of a high ammonia- ammonium-pH system on methane-producing archaea and sulfate-reducing bacteria in mesophilic anaerobic digestion. *Bioresour. Technol.* 245, 598-605 (2017), doi: [10.1016/j.biortech.2017.08.208](https://doi.org/10.1016/j.biortech.2017.08.208).
10. Ziyi, Y., Wen, W., Yanfeng, H., Ruihong, Z., Guangqing, L.: Effect of ammonia on methane production, methanogenesis pathway, microbial community and reactor performance under mesophilic and thermophilic conditions. *Renew. Energy* 125, 915-925 (2018) <https://doi.org/10.1016/j.renene.2018.03.032>.
11. Raskin, L., Zheng, D., Griffin, M.E., Stroot, P.G., Misra, P.: Characterization of microbial communities in anaerobic bioreactors using molecular probes. *Antonie Van Leeuwenhoek* 68, 297–308 (1995). <https://doi.org/10.1007/BF00874140>.

12. Krause, L., Diaz, N.N., Edwards, R.A., Gartemann, K.H., Krömeke, H., Neuweget, H., Pühler, A., Runte, K.J., Schlüter, A., Stoye, J., Szczepanowski, R., Tauch, A., Goesmann, A.: Taxonomic composition and gene content of a methane-producing microbial community isolated from a biogas reactor. *J. Biotechnol.* 136, 91–101 (2008).
13. Yu, Y., Lee, C., Kim, J., Hwang, S.: Group-specific primer and probe sets to detect methanogenic communities using quantitative real-time polymerase chain reaction. *Biotechnol. Bioeng.* 89, 670–679 (2005). <https://doi.org/10.1002/bit.20347>.
14. Sundberg, C., Al-Soud, W.A., Larsson, M., Alm, E., Yekta, S.S., Svensson, B.H., Sørensen, S.J., Karlsson, A.: 454 pyrosequencing analyses of bacterial and archaeal richness in 21 full-scale biogas digesters. *FEMS Microbiol. Ecol.*, 85, 612–626 (2013). <https://doi.org/10.1111/1574-6941.12148>.
15. Ma, S., Jiang, F., Huang, Y., Zhang, Y., Wang, S., Fan, H., Liu, B., Li, Q., Yin, L., Wang, H., Liu, H., Ren, Y., Li, S., Cheng, L., Fan, W., Deng, Y.: A microbial gene catalog of anaerobic digestion from full-scale biogas plants. *Gigascience*, 10 gaa164 (2021). <https://doi.org/10.1093/gigascience/giaa164>.
16. Luton, P.E., Wayne, J.M., Sharp, R.J., Riley, P.W.: The *mcrA* gene as an alternative to 16S rRNA in the phylogenetic analysis of methanogen populations in landfill. *Microbiology*, 148 3521–3530 (2002). <https://doi.org/10.1099/00221287-148-11-3521.14> S.
17. Rastogi, G., Ranade, D.R., Yeole, T.Y., Patole, M.S. Shouche, Y.S.: Investigation of methanogen population structure in biogas reactor by molecular characterization of methyl-coenzyme M reductase A (*mcrA*) genes, *Bioresour. Technol.* 99 5317–5326 (2008). <https://doi.org/10.1016/j.biortech.2007.11.024>
18. Ranjard, L., Poly, F., Lata, J.C., Mougél, C., Thioulouse, J., Nazaret, S.: Characterization of bacterial and fungal soil communities by automated ribosomal intergenic spacer analysis fingerprints. *Appl. Environ. Microbiol.* 67, 4479–87 (2001). <https://doi.org/10.1128/aem.67.10.4479-4487.2001>
19. Tao, Y., Ersahin, M.E., Ghasimi, D.S.M., Ozgun, H., Wang, H., Zhang, X., Guo, M., Yang, Y., Stuckey, D.C., van Lier, J.B.: Biogas productivity of anaerobic digestion process is governed by a core bacterial microbiota, *Chem. Eng. J.* 380, 122425 (2020). <https://doi.org/10.1016/j.cej.2019.122425>.
20. Hewson, I., Fuhrman, J.A.: Richness and diversity of bacterioplankton species along an estuarine gradient in Moreton Bay, Australia. *Appl. Environ. Microbiol.* 70, 3425–3433 (2004). <https://doi.org/10.1128/AEM.70.6.3425>.
21. Kovacs, A., Yacoby, K., Gophna, U.: A systematic assessment of automated ribosomal intergenic spacer analysis (ARISA) as a tool for estimating bacterial richness. *Res. Microbiol.* 161, 192–197(2010). <https://doi.org/10.1016/j.resmic.2010.01.006>
22. Milani, C., Hevia, A., Foroni, E., Duranti, S., Turrone, F., Lugli, G.A., Sanchez, B., Martín, R., Gueimonde, M., van Sinderen, D., et al.: Assessing the fecal microbiota: An optimized ion torrent 16S rRNA gene-based analysis protocol. *PLoS ONE*, 8, e68739, (2013), <https://doi.org/10.1371/journal.pone.0068739>
23. Fischer, M.A., Güllert, S., Neulinger, S.C., Streitz, W.R., Schmitz, R.A.: Evaluation of 16S rRNA gene primer pairs for monitoring microbial community structures showed high reproducibility within and low comparability between datasets generated with multiple archaeal and bacterial primer pairs. *Front. Microbiol.* 7, 1297–1311 (2016). <https://doi.org/10.3389/fmicb.2016.01297>
24. Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Gonzalez Peña, A., Goodrich, J.K., Gordon J.I., et al.: QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods*, 7, 335–336 (2010). <https://doi.org/10.1038/nmeth.f.303>.
25. Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J., Holmes, S.P.: DADA2: High-resolution sample inference from Illumina amplicon data, *Nat. Methods* 13, 581–583 (2016). <https://doi.org/10.1038/nmeth.3869>.
26. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O.: The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools, *Nucleic Acids Res.* 41, D590–D596 (2013). <https://doi.org/10.1093/nar/gks1219>.
27. Liu, T., Sun, L., Müller, B., Schnürer, A.: Importance of inoculum source and initial community structure for biogas production from agricultural substrates. *Bioresour. Technol.* 245, 768–777 (2017). <https://doi.org/10.1016/j.biortech.2017.08.213>.
28. Cardinali-Rezende, J., Colturato, L.F.D.B., Colturato, T.D.B., Chartone-Souza, E., Nascimento, A.M.A., Sanz, J.L.: Prokaryotic diversity and dynamics in a full-scale municipal solid waste anaerobic reactor from start-up to steady-state conditions. *Bioresour. Technol.* 119, 373–383 (2012). <https://doi.org/10.1016/j.biortech.2012.05.136>.
29. Kröber, M., Bekel, T., Diaz, N.N., Goesmann, A., Jaenicke, S., Krause, L., Miller, D., Runte, K.J., Viehöver, P., Pühler, A., Schlüter, A.: Phylogenetic characterization of a biogas plant microbial community integrating clone library 16S-rDNA sequences and metagenome sequence data obtained by 454-pyrosequencing. *J. Biotechnol.* 142, 38–49 (2009). <https://doi.org/10.1016/j.jbiotec.2009.02.010>.
30. Wei, J., Ge, T., Wah, Y.: Monitoring of microbial communities in anaerobic digestion sludge for biogas optimisation. *Waste Manag.* 71, 334–341 (2018). <https://doi.org/10.1016/j.wasman.2017.10.007>.

31. Calusinska, M., Goux, X., Fossépré, M., Muller, E.E.L., Wilmes, P., Delfosse, A.: A year of monitoring 20 mesophilic full-scale bioreactors reveals the existence of stable but different core microbiomes in bio-waste and wastewater anaerobic digestion systems, *Biotechnol Biofuels*. 111–19 (2018). <https://doi.org/10.1186/s13068-018-1195-8>.
32. Pelletier, E., Kreimeyer, A., Bocs, S., Rouy, Z., Gyapay, G., Chouari, R., Rivière, D., Ganesan, A., Daegelen, P., Sghir, A., Cohen, G.N., Médigue, C., Weissenbach, J., Le Paslier, D.: "*Candidatus* Cloacamonas acidaminovorans": Genome sequence reconstruction provides a first glimpse of a new bacterial division, *J. Bacteriol.* 190, 2572–2579 (2008). <https://doi.org/10.1128/JB.01248-07>.
33. Dykstra, S., Gallert, C.: *Candidatus* Syntrophosphaera thermopropionivorans: a novel player in syntrophic propionate oxidation during anaerobic digestion, *Environ. Microbiol. Rep.* 11, 558–570 (2019). <https://doi.org/10.1111/1758-2229.12759>.
34. Conklin, A.S., Stensel, H.D., Ferguson, J.F.: The growth kinetics and competition between *Methanosarcina* and *Methanosaeta* in mesophilic anaerobic digestion. *Proc. Water Environ. Fed.* 2005, 100–121(2012). <https://doi.org/10.2175/193864705783867792>.
35. Karakashev, D., Batstone, D.J., Trably, E., Angelidaki, I.: Acetate oxidation is the dominant methanogenic pathway from acetate in the absence of *Methanosaetaceae*. *Appl. Environ. Microbiol.* 72, 5138–5141 (2006). <https://doi.org/10.1128/AEM.00489-06>.
36. Powell, J.M., Broderick, G.A., Misselbrook, T.H.: Seasonal diet affects ammonia emissions from tie-stall dairy barns, *J. Dairy Sci.* 91857–869 (2008). <https://doi.org/10.3168/jds.2007-0588>.
37. Minato, K., Kouda, Y., Yamakawa, M., Hara, S., Tamura, T., Osada, T.: Determination of GHG and ammonia emissions from stored dairy cattle slurry by using a floating dynamic chamber, *Anim. Sci. J.* 84, 165–177(2013). <https://doi.org/10.1111/j.1740-0929.2012.01053.x>.
38. Nobu, M.K., Narihiro, T., Kuroda, K., Mei, R., Liu, W.T.: Chasing the elusive Euryarchaeota class WSA2: Genomes reveal a uniquely fastidious methyl-reducing methanogen, *ISME J.* 10, 2478–2487 (2016). <https://doi.org/10.1038/ismej.2016.33>.
39. D. Wilkins, X-Y. Lu, Z. Shen, J. Chen, P.K.H. Lee, Pyrosequencing of mcrA and archaeal 16S rRNA genes reveals diversity and substrate preferences of methanogen communities in anaerobic digesters, *Appl. Environ. Microbiol.* 81, 604–613 (2015). <https://doi.org/10.1128/aem.02566-14>
40. Shakeri Yekta, S., Liu, T., Axelsson Bjerg, M., Safaric, L., Karlsson, A., Bjorg, A., Schnurer, A.: Sulfide level in municipal sludge digesters affects microbial community response to long-chain fatty acid loads, *Biotechnol. Biofuels* 12, 259-273 (2019). <https://doi.org/10.1186/s13068-019-1598-1>
41. Yang, J., Wang, D., Luo, Z., Zeng, W., Huang, H.: The role of reflux time in a leach bed reactor coupled with a methanogenic reactor for anaerobic digestion of pig manure: Reactor performance and microbial community, *J. Clean Prod.* 242, 118367(2020). <https://doi.org/10.1016/j.jclepro.2019.118367>
42. Ozbayram, E., Ince, O., Ince, B., Harms, H., Kleinstuber, S.: Comparison of rumen and manure microbiomes and implications for the inoculation of anaerobic digesters, *Microorganisms* 6, 15 (2018). <https://doi.org/10.3390/microorganisms6010015>.
43. Ciotola, R.J., Martin, J.F., Castaño, J.M., Lee, J., Michel, F.: Microbial community response to seasonal temperature variation in a small-scale anaerobic digester. *Energies* 6, 5182–5199 (2013). <https://doi.org/10.3390/en6105182>.
44. Wahid, R., Mulat, D.G., Gaby, J.C., Horn, S.J.: Effects of H₂:CO₂ ratio and H₂ supply fluctuation on methane content and microbial community composition during in-situ biological biogas upgrading. *Biotechnol Biofuels* 12, 104 (2019), <https://doi.org/10.1186/s13068-019-1443-6>.
45. Lang, K., Schuldes, J., Klingl, A., Poehlein, A., Daniel, R., Brune, A.: New mode of energy metabolism in the seventh order of methanogens as revealed by comparative genome analysis of "*Candidatus* Methanoplasma termitum", *Appl. Environ. Microbiol.* 81, 1338–1352 (2015). <https://doi.org/10.1128/AEM.03389-14>
46. Langer, S.G., Ahmed, S., Einfalt, D., Bengelsdorf, F.R., Kazda, M.: Functionally redundant but dissimilar microbial communities within biogas reactors treating maize silage in co-fermentation with sugar beet silage. *Microb. Biotechnol.* 8, 828-836 (2015)
47. Cardinale, M., Brusetti, L., Quatrini, P., Borin, S., Puglia, A.M., Rizzi, A., Zanardini, E., Sorlini, C., Corselli, C., Daffonchio, D.: Comparison of different primer sets for use in automated ribosomal intergenic spacer analysis of complex bacterial communities. *Appl. Environ. Microbiol.* 70, 6147-6156 (2004). <https://doi.org/10.1128/AEM.70.10.6147-6156.2004>
48. Klocke, M., Nettmann, E., Bergmann, I., Mundt, K., Souidi, K., Mumme, J., Linke, B.: Characterization of the methanogenic Archaea within two-phase biogasreactor systems operated with plant biomass. *Syst. Appl. Microbiol.* 31, 190–205 (2008). <https://doi.org/10.1016/j.syapm.2008.02.003>

Tables

Table 1. Samples collected in the biodigesters P1 (CM2 to CMP) and P2 (VL1 to VL12), respective operating conditions, VFA, biogas and energy production. Samples CM3, CM8, CMP and VL5 (in bold characters) were used for NGS analysis and *CM8** and *VL10* (bold italic) for cloning and

sequencing of archaeal sequences. n.r. Not reported.

Sample	Sampling site	Time of collection	VFA (mgL ⁻¹)	T °C	pH	Alkalinity (mgL ⁻¹)	Ratio VFA/Alk	Ammonia nitrogen concentration (mg Kg ⁻¹)	Feeding		Production	
									Com silage (t day ⁻¹)	Manure (t day ⁻¹)	Biogas (m ³ day ⁻¹)	Energy (MWh day ⁻¹)
CM2	Primary digester	17 Nov	1784	42.2	7.51	15840	0.11		8.90	54.80	2486	0
CM3	Primary digester	24 Nov	1689	42.5	7.46	14325	0.12	1875	21.70	0	4329	0
CM4	Primary digester	01 Dec	1677	43.1	7.46	13699	0.12		33.80	0	6703	0.7
CM5	Primary digester	08 Dec	1766	43.3	7.47	13598	0.13		41.40	0	8259	17.70
CM6	Primary digester	15 Dec	1834	43.1	7.48	13334	0.14	1951	44.50	20.20	9140	24
CM7	Primary digester	22 Dec	1463	44.2	7.44	12663	0.12		40.70	16.90	10919	24
CM8	Primary digester	26 Dec	1804	42.5	7.49	12464	0.14		39.10	17.80	8031	24
CM8*	Primary digester	26 Dec	1804	42.5	7.49	12464	0.14		39.10	17.80	8031	24
CMP	Post digester	29 dec	1152	41.5	7.67	14145	0.13		n.r.	n.r.	n.r.	
VL1	Primary digester	08 May	1790	40.9	7.33	10968	0.16	1100	39.70	0	7920	22
VL2	Primary digester	15 May	1783	42.2	7.43	9855	0.18		41.00	0	8180	23.90
VL3	Primary digester	22 May	1379	42.5	7.35	9423	0.15		41.80	24.90	8662	23.20
VL4	Primary digester	29 May	1751	41.7	7.45	8499	0.21		32.20	49.20	6737	23.90
VL5	Primary digester	5 June	1844	42.8	7.51	8974	0.21	1315	36.70	19.90	7637	23
VL6	Primary digester	12 June	1776	44.3	7.50	8835	0.20		37.30	20.30	8159	23.90
VL8	Primary digester	19 June	1367	46.9	7.50	8711	0.16		36.00	27.00	7538	23.90
VL10	Primary digester	26 June	1434	47.9	7.51	9021	0.16		36.00	27.00	7690	24
VL12	Primary digester	03 July	1403	47.7	7.53	9504	0.15	1320	39.90	27.10	7642	24

Table 2. Primers utilised in this study. "Y", "R" "M" and "W" mean degenerate bases.

Primer	Sequence	Ta (°C)	Ref.
ITSF	5'-GTCGTAACAAGGTAGCCGTA-3'	58	[47]
ITSReub	5'-GCCAAGGCATCCACC-3'		
	5'-YGAYTAAGCCATGCRA GT-3'	47	[48]
69F	5'-TGCTCCCCCGCCAATTCCT-3'		
ARC934R			
	5'-TATTTAGGTGACACTATAG-3'	50	
SP6	5'-TAATACGACTCACTATAGGG-3'		
T7			
	5'-GGTGGTGTGTTGGATTACARTATGCW-3'	55	[16]
mcrAF	5'-TTCATTGCRTAGTTWGGRTAGTT-3'		
mcrAR			

Table 3. ARDRA profiles detected after restriction analysis of amplicons derived from rRNA 16S of Archaea, their relative abundance in the analysed clones and similarity with bacteria species established with BLAST analysis.

Profiles P1 (Accession number)	Relative abundance in P1	Similar microorganism (Accession number)	Sequence similarity	Functional group
16S A (CM8) (MW381770)	36%	<i>Methanothrix soehngenii</i> (NR_104707.1)	98.20%	Acetoclastic
16S B (CM8) (MW381771)	17%	<i>Methanoculleus bourgensis</i> (NR_114489.1)	98.07%	Hydrogenotrophic
16S C (CM8)	6%	Uncultured bacterium (FJ205838.1)	73%	
16S D (CM8)	8%	Uncultured bacterium (<i>JX102010.1</i>)	96.85%	
16S E (CM8) (MW381772)	6%	<i>Methanoculleus bourgensis</i> (AB065298.1)	98.68%	Hydrogenotrophic
16S mix (CM8)	27%	<i>Unidentified miscellaneous clones</i>		
16S F (VL10) (MW381773)	32%	<i>Methanosarcina thermophila</i> (NR_044725.1)	98.8%	Acetoclastic
16S G (VL10) (MW381774)	25%	<i>Methanoculleus hydrogenitrophicus</i> (NR_116881.1)	99.52%	Hydrogenotrophic
16S I (VL10) (MW381775)	10%	<i>Hydrogenispora ethanolica</i> (NR_125455.1)	90.22%	
16S mix (VL10)	33%	<i>Unidentified miscellaneous clones</i>		

Profiles P1 (Accession number)	Similar microorganism (Accession number)	Sequence similarity	Functional group
Mcr1 (CM8) (MW390769)	Uncultured methanogenic archeon (JQ686784.1)	98.92%	
Mcr2 (CM8) (MW390770)	Uncultured methanogenic archeon (JQ686770.1)	99%	
Mcr3 (CM8) (MW390771)	Uncultured methanogenic archeon (AB615638.1)	99.15%	
Mcr4 (CM8) (MW390772)	Uncultured <i>Methanobacterium</i> sp (KJ487752.1)	98%	Hydrogenotrophic
Profiles P2 (Accession number)	Similar microorganism (Accession number)	Sequence similarity	Functional group
Mcr5 (VL10) (MW390773)	<i>Methanoculleus bourgensis</i> (LT549891.1)	98.76%	Hydrogenotrophic
Mcr6 (VL10) (MW390774)	<i>Methanosarcina flavescens</i> (CP032683.1)	97.54%	Acetoclastic
Mcr7 (VL10) (MW390775)	Uncultured methanogenic archeon (EF628139.1)	99%	
Mcr8 (VL10) (MW390776)	<i>Methanosarcina thermophila</i> (AB353225.1)	98.30%	Acetoclastic

Figures

Figure 1

The key stages of anaerobic digestion process, schematically represented.

Figure 2

Synoptic diagram of the start-up phases of P1.

Figure 3

UPGMA dendrogram of P1 and P2 obtained from ARISA profiles.

Figure 4
Rarefaction curves derived by Shannon diversity index of Bacteria (a) and Archaea (b) of samples CM3, CM8, CMP and VL5 calculated on the basis of 16S rDNA NGS taxonomy profiling.

Figure 5

Relative abundance of bacterial phyla in the samples CM3, CM8, CMP and VL5 calculated through 16S rDNA NGS taxonomy profiling (average of duplicated biological samples).

Figure 6

Relative abundance of bacterial genera (> 1%) in the samples CM3, CM8, CMP and VL5 calculated through 16S rDNA NGS taxonomy profiling (average of duplicated biological samples).

Figure 7

Relative abundance of archaeal genera (> 1%) in the samples CM3, CM8, CMP and VL5 calculated through 16S rDNA NGS taxonomy profiling (average of duplicated biological samples).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Graphicalabstract.pptx](#)