

Modified Commando Procedure in Complicated Infective Endocarditis - A Case Series.

Po Sung Chen

Cheng Hsin General Hospital <https://orcid.org/0000-0002-5707-2414>

Chungyi Chang

Cheng Hsin General Hospital

Yicheng Chuang

Cheng Hsin General Hospital

Ichen Chen

Cheng Hsin General Hospital

Tingchao Lin (✉ tingchaolin@gmail.com)

Cheng Hsin General Hospital

Case report

Keywords: Infective endocarditis, Intervalvular fibrous body, Prosthetic valve endocarditis, Valvular heart disease, Surgical reconstruction

Posted Date: November 19th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-105185/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Journal of Cardiothoracic Surgery on April 13th, 2021. See the published version at <https://doi.org/10.1186/s13019-021-01480-4>.

Title page

Modified Commando procedure in complicated infective endocarditis — a case series.

Author:

Posung Chen^{1, 2}, Chungyi Chang¹, Yicheng Chuang¹, Ichen Chen¹, Tingchao Lin^{1, 3}

¹ Division of Cardiovascular Surgery, Heart Center, Cheng Hsin General Hospital, Taipei, Taiwan

² Department of Surgery, Kinmen Hospital, Kinmen, Taiwan

³ Department of Medicine, School of Medicine, National Yang-Ming University, Taipei, Taiwan

Corresponding author:

Posung Chen even.sinfonia@gmail.com

Tingchao Lin tingchaolin@gmail.com

Abstract

Background: Complicated infective endocarditis (IE) with perivalvular abscess and destruction of intervalvular fibrous body (IFB) has high mortality risk and requires emergent or urgent surgery mostly.

Case presentation: We presented four patients with complicated infective endocarditis combined with perivalvular abscess and IFB destruction. Three patients had prosthetic valve endocarditis and one patient had native valve endocarditis. They all received modified Commando procedure successfully. No surgical mortality or re-exploration for bleeding.

Conclusions: We suggest that modified Commando procedure may have some benefit in improving survival rate of patients with complicated IE and reducing complications.

Keyword: Infective endocarditis; Intervalvular fibrous body; Prosthetic valve endocarditis; Valvular heart disease; Surgical reconstruction

Introduction

Complicated infective endocarditis (IE) with perivalvular abscess and destruction of the intervalvular fibrous body (IFB) has a high mortality risk and mostly requires emergent or urgent surgery. David et al. [1] reported the technique of aortic valve replacement (AVR), mitral valve replacement (MVR) combined with reconstruction of the IFB, which is known as the Commando procedure. However, a high mortality rate and re-exploration for bleeding were noted in early series, which might be related to the emergent surgical procedure, fragile infected tissue, and difficulty of achieving hemostasis due to inaccessible posterior suture lines. The complicated condition is sometimes inevitable, but we would like to introduce a modified Commando procedure, which may improve surgical exposure, solidify the infected tissue, decrease tension, and transfer the suture line to the anterior aspect for better hemostasis. Here, we describe the cases of 4 patients who underwent reoperation due to prosthetic IE with IFB involvement and the surgical technical modifications we introduced to the origin procedure.

Case

Patient 1

A 46-year-old woman was admitted to our hospital due to intermittent dyspnea on exertion for 2 months and fever for 1 week. She had undergone mitral valve repair and tricuspid valve repair (TVr) 8 years prior due to IE with symptomatic severe mitral regurgitation (MR) and tricuspid regurgitation. Transthoracic echocardiography (TTE) revealed moderate to severe aortic regurgitation (AR), severe MR and vegetation over the anterior mitral leaflet that extended to the left ventricular outflow tract (LVOT). Recurrent IE was diagnosed. Antibiotic therapy was initiated, but intermittent fever persisted, so urgent surgery was arranged.

The modified Commando procedure was performed during the reoperation. Cardiopulmonary bypass was established via the ascending aorta and bicaval cannulation. Custodiol HTK cardioplegic solution was used. Surgical exposure was achieved through oblique aortotomy and an extended transeptal approach. The aortotomy was extended from the noncoronary cusp into the IFB and onto the left atrial roof. In the aortic valve area, left-coronary-cusp and noncoronary-cusp perforation with perivalvular abscess formation over the aortic root was observed (Fig. 1a). In the mitral valve area, vegetation was observed on the anterior mitral leaflet extending to the IFB and LVOT (Fig. 1b).

All infected tissues were removed (Fig. 1c), and the remaining margin was treated with glutaraldehyde solution carefully. Reconstruction of the IFB was performed with one diamond-shaped patch of bovine pericardium folded into two triangular shapes (Fig. 1d), one for reconstruction of the interatrial septum and the other for reconstruction of the aortic root. MVR with a 27-mm Medtronic ATS mechanical prosthetic valve and AVR with a 20-mm Medtronic ATS mechanical prosthetic valve were performed (Fig. 1e). After reconstruction of the interatrial septum and aortic root, another bovine pericardium patch was used for reconstruction of the left atrial roof and right atrium to close the opening of the extended transeptal approach (Fig. 1f).

The postoperative course was uneventful except that a permanent pacemaker (PPM) was implanted for complete heart block. There were no other complications and no restrictions on physical activity after discharge. Follow-up TTE examinations at 1 year and 3 years revealed good valve function, and no paravalvular leakage was noted.

Patient 2

A 41-year-old man was admitted to our hospital for decompensated heart failure. He had undergone MVR and TVr 3 months prior and anal fistulectomy 1 month prior.

TTE revealed dislocation of the mitral prosthetic valve and severe right ventricular dysfunction with an ejection fraction of 28%. At surgery, vegetation on the aortic valve, mitral prosthetic valve and IFB was detected. Nearly total dehiscence of the mitral prosthetic valve and posterior atrial-ventricular groove containing rupture was also detected (Fig. 2a). The ruptured atrial-ventricular groove was repaired with one bovine pericardium patch, and the modified Commando procedure was performed. A PPM was implanted postoperatively, and the residual course was uneventful.

Patient 3

A 60-year-old man was admitted to our hospital for heart failure and intracardiac shunt (LVOT to left atrium). He had IE and left medial frontal cerebral infarction and undergone AVR at another hospital 2 weeks prior. In addition, he had been intubated preoperatively due to respiratory failure caused by influenza A infection. During surgery, dehiscence of the aortic prosthetic valve with an annular abscess and vegetation over the aortic root to the mitral valve and one large perforation of the IFB were detected (Fig. 2b). The modified Commando procedure was performed, along with aortic root reconstruction with a metallic valve and gelatin vascular graft and the Cabrol method for coronary ostia reimplantation (Fig. 2c). The postoperative course was prolonged due to preoperative respiratory failure and poor brain

infarction-related mobility. A PPM was implanted for heart block. Multiple debridement and muscular flap reconstruction procedures were performed for sternal infection. He was discharged in a wheelchair and has undergone continuous rehabilitation.

Patient 4

An 81-year-old woman was admitted to our hospital for fever and dyspnea for 3 days. TTE revealed aortic valve vegetation with severe AR and severe MR. IE complicated with valvular insufficiency was diagnosed. At surgery, vegetation on the aortic noncoronary cusp, LVOT and IFB, an aortic annular abscess with perforation and nearly total dehiscence of the mitral prosthesis were detected. The modified Commando procedure was performed with bioprosthetic valves due to the advanced age of the patient. The postoperative course was smooth, except PPM implantation was performed for heart block.

Discussion

Surgical treatment for infective endocarditis involving the IFB is challenging.

Among the limited studies of IFB repair, the incidence of mortality and re-exploration for bleeding is worth noting. David et al. [1] reported that 43 patients (14 with endocarditis) underwent double valve replacement and IFB reconstruction. Of the 14 patients with endocarditis, eleven had prosthetic valve endocarditis. There were three operative deaths (21.4%). Davierwala et al. [2] reported that 25 patients underwent the Commando procedure. Twenty of them had prosthetic valve endocarditis (80%). Eight patients (32%) died within 30 days of surgery, and the major cause of mortality (3 patients) was uncontrolled bleeding due to poor patch anchoring in infection-ridden and devitalized tissue. Additionally, Forteza et al. [3] reported that 26 patients with IE underwent the Commando procedure, resulting in an in-hospital mortality rate of 15.4% and a rate of re-exploration for bleeding of 15.4%.

There were no cases of in-hospital mortality or re-exploration for bleeding in our series. The modification of the original Commando procedure may have contributed to these results. First, we used an extended transeptal approach, with wide opening of the right atrium, interatrial septum, left atrium, and aortic root. This approach provides better exposure than an extended aortotomy incision in the LA

roof only. It also provides excellent exposure and an unobstructed approach for complete debridement and valve replacement. Second, an extra patch is used to repair the right atrium to further reduce the tension. The folded bitriangular bovine pericardium patch, which was used for left atrial roof and aortic reconstruction in the abovementioned studies, was used to reconstruct the interatrial septum and aortic root. This method may lessen the tension on the IFB and transfer the suture line from the posterior aspect of the heart to the anterior aspect, which may help to identify bleeding and facilitate hemostasis. Third, we used glutaraldehyde solution to treat the remaining tissue surface. Glutaraldehyde treatment has the benefit of eliminating bacteria and reinforcing the remaining tissue, making further sutures more stable and reducing the risk of tissue tearing and bleeding; it may also decrease the postoperative infection rate. Nakamura et al. [4] reported on 35 IE patients who underwent mitral valve surgery with glutaraldehyde treatment and concluded the safety of this glutaraldehyde treatment. However, AV nodes may be affected during the procedure, which may contribute to the high PPM implantation rate in our series. Last, the rocking motion of the mitral valve during systole and diastole is a problem.

[1] We sew the noncoronary cusp part of the aortic prosthesis and the anterior part of the mitral prosthesis together to enhance the new IFB and reduce the risk of patch dehiscence or valve dehiscence. This procedure does not increase the risk of

bleeding, and follow-up echocardiography does not reveal any sign of structural destruction or paravalvular leakage.

The choice of prosthesis in patients with infective endocarditis is controversial. Few studies have suggested that homografts have the benefit of preventing recurrent endocarditis. [5, 6] However, Moon et al. reported that 306 patients with IE (97 patients with prosthetic endocarditis) underwent valve replacement and that survival was independent of valve type. [7] As previous studies have suggested, [1, 2, 3] we believe that the radical debridement of infected tissue is more important than the type of valve. An aortic valve homograft with a preserved anterior leaflet of the mitral valve may be ideal in this procedure, but the uncertain availability in most urgent or emergent cases is a disadvantage. In addition, the durability is doubtful and may result in patients facing a high-risk reoperation. Therefore, we chose to use mechanical prostheses in most patients to reduce the possibility of a high-risk reoperation.

Conclusion

Infective endocarditis involving the IFB is an urgent or emergent situation that requires immediate treatment with high mortality and morbidity rates. The modified Commando procedure as presented may improve the exposure of the surgical field,

reduce the 'rocking motion' of the prosthetic valve, reduce tension via the extra patch, and help to achieve hemostasis by transferring the suture line from the posterior aspect to the anterior aspect. The results are optimal, with no cases of in-hospital mortality, re-exploration for bleeding, or paravalvular leakage. More extended experience and long-term surveillance are required to confirm its efficacy and long-term durability.

Abbreviations

IE: Infective endocarditis; IFB: Interventricular fibrous body; AVR: Aortic valve replacement; MVR: Mitral valve replacement; TVr: Tricuspid valve repair; MR: Mitral regurgitation; TTE: Transthoracic echocardiography; AR: Aortic regurgitation; LVOT: Left ventricular outflow tract; LA: Left atrial; PPM: Permanent pacemaker

Declarations**Acknowledgements**

None.

Authors' contributions

PC wrote the original manuscript and prepared the figures. CC, YC and IC revised the paper. TL drafted the final manuscript. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

We announce that the patient involved is consent to participate, not applicable on ethics approval.

Consent for publication

We announce that all presentations of this case report have consent to publish.

Competing interests

The authors declare that they have no competing interests.

References

1. David TE, Kuo J, Armstrong S. Aortic and mitral valve replacement with reconstruction of the intervalvular fibrous body. *J Thorac Cardiovasc Surg.* 1997;114(5):766-71.
2. Davierwala PM, Binner C, Subramanian S, et al. Double valve replacement and reconstruction of the intervalvular fibrous body in patients with active infective endocarditis. *Eur J Cardiothorac Surg.* 2014;45(1):146-52.
3. Forteza A, Centeno J, Ospina V, et al. Outcomes in aortic and mitral valve replacement with intervalvular fibrous body reconstruction. *Ann Thorac Surg.* 2015;99(3):838-45.
4. Nakamura K, Hashimoto K, Sakamoto Y, et al. Safe Use of Glutaraldehyde to Repair the Destroyed Valve in Active Infective Mitral Valve Endocarditis. *Circ J.* 2018;82(10):2530-2534.
5. Sabik JF, Lytle BW, Blackstone EH, et al. Aortic root replacement with cryopreserved allograft for prosthetic valve endocarditis. *Ann Thorac Surg.* 2002;74(3):650-9.
6. Yankah AC, Pasic M, Klose H, et al. Homograft reconstruction of the aortic root for endocarditis with periannular abscess: a 17-year study. *Eur J Cardiothorac Surg.* 2005;28(1):69-75.
7. Moon MR, Miller DC, Moore KA, et al. Treatment of endocarditis with valve replacement: the question of tissue versus mechanical prosthesis. *Ann Thorac Surg.* 2001;71(4):1164-71.

Figures




Figure 1

a Aortic valve perforation: non-coronary-cusp & left-coronary-cusp. b Mitral valve post annuloplasty with vegetable formation and perforation. c Post removal of infective tissue (aortic valve, mitral valve, partial aortic annulus and IFB). d Bovine pericardium (two triangular shape) for reconstruction of IFB, interatrial septum and aortic root. e Post AVR and MVR. f After reconstruction of interatrial septum and aortic root, proceed to reconstruction of left atrial roof and right atrium.




Figure 2

a Patient 2, dehiscence of mitral prosthetic valve. b Patient 3, vegetations over aortic perivalvular area and IFB perforation. c Patient 3, post modified Commando procedure and aortic root reconstruction with Cabrol method.