Genotype by Environment Interaction and Grain Yield Stability of Drought Tolerant Cowpea landraces in Ethiopia

Tesfaye Walle Mekonnen (✉ tesfaye.walle@gmail.com)

University of the Free State Faculty of Natural and Agricultural Sciences <https://orcid.org/0000-0002-1943-9991>

Firew Mekbib

Haramaya University

Berhanu Amsalu

Ethiopian Institute of Agricultural Research

Melaku Gedil

IITA: International Institute of Tropical Agriculture

Maryke Labuschagne

University of the Free State Faculty of Natural and Agricultural Sciences

Research Article

Keywords: ASV, Environment, GGE biplot, YSI, Yield

Posted Date: November 23rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1054648/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Euphytica on April 12th, 2022. See the published version at <https://doi.org/10.1007/s10681-022-03011-1>.

Abstract

Cowpea is one of the most important indigenous food and forage legumes in Africa. It serves as a primary source of protein for poor farmers in drought-prone areas of Ethiopia. The crop is used as a source of food, and insurance crop during the dry season. Cowpea is adaptable to a wide range of climatic conditions. Despite this, the productivity of the crop is generally low due to lack of stable and drought tolerant varieties. In this study, 25 cowpea genotypes were evaluated in five environments using a triple lattice design during the 2017 and 2018 main cropping seasons. The objectives of this study were to estimate the magnitude of genotype by environment interaction (GEI) and grain yield stability of selected drought tolerant cowpea genotypes across different environments. The additive main effect and multiplicative interaction (AMMI) model indicated the contribution of environment, genotype and GEI as 63.98 6%, 2.66% and 16.30% of the total variation for grain yield, respectively. The magnitudes of the GEI sum of squares were 6.12 times that of the genotypes for grain yield. The IPCA1, IPCA2 and IPCA3 were all significant and explained 45.47%, 28.05% and 16.59% of the GEI variation, respectively. The results from AMMI, cultivar superior measure (Pi), genotype plus genotype-by-environment (GGE) biplot yield stability index (YSI), and AMMI stability value (ASV) analyses identified NLLP-CPC-07-145-21, NLLP-CPC-103-B and NLLP_CPC-07-54 as stable and high yielding genotypes across environments. Thus, these genotypes should be recommended for release for production for drought prone areas. NLLP-CPC-07-143, Kanketi and CP-EXTERETIS were the least stable. The AMMI1 biplot showed that Jinka was a high potential and favorable environment while Babile was an unfavorable environment for cowpea production.

Introduction

Cowpea [*Vigna unguiculata* (L.) Walp.] is one of the most important food and forage legumes grown in the semi-arid tropics and some temperate regions of the world (Timko and Singh, 2008). Cowpea, at all stages of growth, serves as food source (Ahenkora et al., 1998; Singh et al., 2003). Poor people in developing countries of the tropics derive their protein, animal feed and cash income from the production of the crop (Diouf 2011). The young leaves, green pods and green grains are used as vegetables, and the dry grains are used in various preparations for both human food and livestock feed (Filho et al., 2017; Owade et al., 2020). The cowpea grain is highly nutritious and contains about 22.8 - 28.9% protein with an average of 25.6% (Weng et al. 2019). The portion of the cowpea crop above ground (except for pods) serves as a useful source of nutrient-rich fodder for livestock in many areas of the world (Singh et al., 2003; Weng et al., 2019). Cowpea can withstand harsh growing conditions, particularly temperature and moisture stress in comparison to other crops (Agbicodo et al., 2009; Goufo et al., 2017; Fatokun et al., 2012).

Environment and growing season, and rainfall distribution and intensity may have positive or negative impacts on cowpea genotypes (Marina et al., 2017). Plant breeders evaluate genotypes in multi-environments, representing favorable and unfavorable growing conditions, to estimate and understand the complexity of the genotype across environments (Mohammadi and Amri 2008).

Cowpea is greatly influenced by seasonal environmental fluctuations and shows large genotype by environment interaction (GEI), which is a major challenge in obtaining a full understanding of genetic control of varieties when compared across a series of environments (Kuruma et al., 2019a; Olajide & Ilori, 2017; Shegro et al., 2020; Simion, 2018; Simion et al., 2018). Studying and understanding of GEI is important to plant breeding programs for improving yield and yield components (Alberts, 2004; Yan & Tinker, 2006) and is also used for identifying the basic causes of differences between genotypes for yield stability (Yan & Tinker, 2006). GEI exists when the ranks of genotypes show an obvious shift from one environment to another (Leflon et al. 2008). Measuring GEI is important to determine the optimum breeding strategy for releasing genotypes with adequate adaptation to target environments (Arpita et al., 2019; Yan, 2016).

The methods commonly used for quantifying GEI and stability include; principal component analysis (Zobel et al., 1988), additive main effect and multiplicative interaction (AMMI) (Zobel et al., 1988), GGE biplot analysis (Yan and Tinker 2006), cultivar superiority measure (Lin and Binns 1988), AMMI stability value (Purchase et al. 2000) and yield stability index (Tumuhimbise et al. 2014). Understanding the magnitude of effects of the environment, genotype and their interaction on yield and stability performance of cowpea genotype across environments is important because it reduces the efficiency of the genetic gain through the development of high yielding genotypes with desirable agronomic traits (Hall et al., 2003; Simion et al., 2018). Ethiopia is a victim of repeated droughts that cause partial or total crop failure, and subsequently, famine in the country. Cowpea can be used to reduce the consequence of drought.

Knowledge on the effect of genotype, environment, and their interactions on drought tolerant cowpea grain yield is limited in Ethiopia. The objectives of the present study, therefore, were to estimate the magnitude of GEI and grain yield stability of selected drought tolerant cowpea genotypes across drought prone environments.

Materials And Methods

Study Area

The field experiments were conducted during the 2017 and 2018 main cropping season at five environments namely Babile, Melkassa, Mieso, Jinka and Sirnka (Table 1).

Table 1. Description of experimental sites

		Geographical location			Temperature (°C)		
Experimental sites	Soil type	Altitude			Rainfall	Min T	Max T
		(m.a.s.l)	Latitude (N)	Longitude (E)	(mm)		
Babile	Sandy	1647	9° 13' 09"	42° 19'	432	13.85	28.9
Melkassa	Andosols	1550	8°3 0'	39° 24'	763	15.73	27.31
Mieso	Vestisol	1470	9°14'	40°45'	787	18.4	30.70
Jinka	Vestisol	1383	5° 52'	36° 38'	1247.7	17.60	30.00
Sirinka (Kobo)	Euric Fluvsol	1470	12° 09'	39° 28'	637.3		

m.a.s.l= meters above sea level, E= east, N= North, Min= minimum, Max= maximum temperature, °C= degree centigrade, mm= millimeter. Weather data accessed from meteorological stations at each site

Plant Materials

A total of 25 genotypes were used for this study. Of these, two released varieties were used as standard checks and 23 genotypes were selected for drought tolerance from a drought stress experiment which included 324 genotypes (Table 2).

Table 2 List of genotypes used for genotypes by environment interaction

Genotypes	Genotype code	Source	Breeding status of the genotypes
NLLP-CPC-07-10	G1	MARC	Landrace
CP-EXTERETIS	G2	MARC	Landrace
NLLP-CPC-07-169	G3	MARC	Landrace
Dass 001	G4	MARC	Landrace
ACC-215-821	G5	MARC	Landrace
ACC-216-747	G6	MARC	Landrace
NLLP-CPC-07-139	G7	MARC	Landrace
NLLP-CPC-07-145-21	G8	MARC	Landrace
NLLP-CPC-07-143	G9	MARC	Landrace
ACC-215-762	G10	MARC	Landrace
NLLP-CPC-103-B	G11	MARC	Landrace
NLLP-CPC-07-156	G12	MARC	Landrace
NLLP-CPC-07-54	G13	MARC	Landrace
NLLP-CPC-07-28	G14	MARC	Landrace
NLLP-CPC-07-36	G15	MARC	Landrace
NLLP-CPC-07-19	G16	MARC	Landrace
NLLP-CPC-07-03-B	G17	MARC	Landrace
NLLP-CPC-07-167	G18	MARC	Landrace
NLLP-CPC-07-166	G19	MARC	Landrace
NLLP-CPC-07-157	G20	MARC	Landrace
NLLP-CPC-07-140	G21	MARC	Landrace
NLLP-CPC-07-57	G22	MARC	Landrace
ACC-211-490	G23	MARC	Landrace
Bole (Standard check)	G24	MARC	Released variety
Kanketi (Standard check)	G25	MARC	Released variety

MARC= Melkassa Agricultural Research Center

Experimental Design and Procedures

The experiments were laid out using a 5×5 triple lattice design. The plot size was 4 m long, 0.75 m between rows and 0.2 m between plants. It consisted of four rows with 20 plants per row. The distance between plots, intra blocks and replications was 1 m, 1.5 m and 2 m, respectively. The data were collected from the middle two rows.

Data Collection and Analyses

According to the descriptors of cowpea (IBPGR, 1983), yield on a plot basis was collected and converted to grain yield per hectare (kg ha⁻¹). SAS (SAS 2013) software was used for combined analysis of variance (ANOVA) over environments and seasons. A mixed linear model was used (Gomez & Gomez 1984). Environment and season were considered as random and genotypes as fixed effects (Hartley, 1950). F-Max ratio was used to test the homogeneity of error variances before analyzing the combined data. GEA-R was used for AMMI, GGE biplot, and cultivar superiority measure (Pi) analysis (Pacheco et al. 2016).

The Additive Main effect and Multiplicative Interaction (AMMI) model analysis proposed by Zobel et al. (1988) was used for analyzing GEI. AMMI partitions the sum of squares into interaction principal component (IPC) axes.

GGE biplots were constructed from the data (Yan et al., 2000; Yan & Rajcan, 2002; Yan et al., 2007; Yan, 2011). The GGE biplot has many visual interpretations that AMMI does not have. It allows visualization of crossover GEI (Yan et al., 2007). Moreover, the GGE biplot is more logical for biological objectives in terms of explaining the first principal component score, which represents genotypic level rather than additive level (Yan et al., 2000). The GGE biplot is built on the first two major components of a principal component analysis (PCA) using the Site Regression (SREG) model. When the first component is highly correlated with the genotype main effect, the proportion of the yield is considered to be due only to the characteristics of the genotype. The second component represents the variation in the yield due to the GEI (Yan, 2011). The GGE biplots were generated using a singular value decomposition model of the first two principal components (Yan, 2002; Yan & Rajcan, 2002).

GEA-R was used for AMMI, GGE biplot analysis and cultivar superiority measure (Pi) (Pacheco et al. 2016). AMMI stability Value (ASV) (Purchase *et al.*, 2000) would be essential to quantify and rank genotypes according to their yield stability (Farshadfar et al. 2011).

$$ASV = \sqrt{\left[\frac{[IPCA\ 1\ Sum\ of\ squares]}{[IPCA\ 2\ Sum\ of\ squares]} (IPCA\ 1\ scores)^2 + [IPCA\ 2\ scores]^2 \right]}$$

Where ASV = AMMI's stability value, SS = sum of squares, IPCA1 = interaction of first principal component, IPCA2 = interaction of second principal component.

Yield stability index (YSI) incorporates both mean yield and stability in a single criterion. Low values of both parameters show desirable genotypes with high mean yield and stability (Tumuhimbise et al. 2014). The yield stability index was calculated using the following formula:

$$YSI = RASV + R$$

Where: RASV is the ranking of the AMMI stability value and R the ranking of genotypes in all environments.

Cultivar superiority (Pi) measures the deviation from the yield of a given genotype in relation to the maximum in each environment. The significant difference of cultivar superiority (Pi) was compared by computing a cutoff point for each value. Even though distributional properties of the cultivar superiority measure is not exactly known, the cut-off point was calculated by multiplying the 5% or 1% significant F-values for Pi at environment (E) and genotype (E-2) degrees of freedom by the deviation from regression mean squares, where G and E denote the number of genotypes and environments respectively (Lin and Binns 1988). The Cultivar superiority (Pi) measures was calculated using the following formula:

$$P_i = \frac{n(X_i - \bar{M})^2 + \sum_j (X_{ij} - \bar{X}_i - \bar{M}_j + \bar{M})^2}{2n}$$

Where X_{ij} is the response of the i th genotype in the j th environment, M_j is the genotype with maximum response among all genotype in the j th environment, $\bar{X}_i$ is the mean of genotype i in the environment, $\bar{M}$ is the genotype with maximum response among all genotype in the j th environment.

Results

Analysis of variance for grain yield across environments

The results of the combined ANOVA across the tested genotypes showed that environment (E) and season (Y) main effects, $G \times E$, $G \times Y$, $E \times Y$, $G \times E \times Y$ were all highly significant ($p < 0.0001$) for grain yield, and the genotype main effect (G) was significant ($p < 0.05$) for grain yield (Table 3).

Table 3 Sum of squares from combined ANOVA for grain yield of 25 cowpea genotypes

Source of variation	DF	Mean square
Replication	2	71291.20
Genotype	24	208390.70*
Environment	4	30026430.10**
Season	1	57369811.20**
Genotype* Environment	96	318675.40**
Genotype* Season	24	334214.70**
Environment* Season	4	25866297.40**
Genotype* Environment* Season	96	301461.90**
Error	498	121728.80

*, **= significant at 5% and 1% probability level, respectively, DF= degree of freedom

Grain yield performance across the environments

G2, G8, G10, G3 and G 18 had high yield performance at specific environments Sirnka, Jinka, Melkassa, Miesso, and Babile, respectively. However, G20, G9, G12, G15, G11, and G9 were the poorest performers at Sirnka, Jinka, Melkassa, Miesso, and Babile, respectively (Table 4). The highest grain yielding was recorded at Jinka (2445.7 kg ha⁻¹), while the poorest grain yield was recorded at Babile (555.81 kg ha⁻¹). The highest yielding genotype (G8) across the environments and seasons was G8 with a yield of 1641.25 kg ha⁻¹ while the poorest yielding genotype was G9 with a yield of 1267.62 kg ha⁻¹.

Table 4 Grain yield performance across the environments

N/S	G	Simka	G	Jinka	G	Melkassa	G	Mieso	G	Babile	G	GM
1	G2	1804.7A	G8	2445.7A	G10	1905.4A	G3	2252.2A	G18	909.77A	G8	1641.25
2	G11	1757.7BA	G6	2333.3BA	G13	1769.3BA	G19	2192.7A	G5	898.4BA	G17	1595.47
3	G24	1693BAC	G12	2282.7BAC	G7	1748.4BA	G12	2166.6BA	G20	896.94BA	G13	1582.33
4	G4	1602.8BDAC	G13	2240.2BDAC	G8	1696.6BAC	G8	2163.4BA	G9	852.86BAC	G6	1566.2
5	G16	1597.4BDAC	G5	2150.6BDAC	G17	1694.9BAC	G23	2156.2BAC	G22	834.36BAC	G14	1551.03
6	G13	1584BDAC	G1	2139.4EBDAC	G14	1691BAC	G21	2049.4BDAC	G25	824.45BAC	G12	1529.22
7	G10	1555.6EBDAC	G15	2109.4EBDACF	G21	1656.3BDAC	G25	2015.5EBDAC	G23	821.78BAC	G11	1523.12
8	G17	1549.7EBDAC	G22	2106.9EBDACF	G16	1544.8BDEC	G11	1979.5EBDACF	G15	821.02BAC	G21	1521.99
9	G7	1536.5EBDAC	G14	1973.4EBDAGCF	G1	1541BDEC	G14	1975.4EBDACF	G17	817.86BAC	G1	1515.8
10	G15	1517.2EBDAC	G3	1972EBDAGCF	G11	1516.2BDEC	G17	1966.1EBDACF	G21	787.9BDAC	G19	1513.18
11	G6	1503.3EBDAC	G19	1971.1EBDAGCF	G20	1489.4FBDEC	G20	1936.9EBDACF	G4	783.2BDAC	G3	1489.99
12	G12	1449.7EBDACF	G17	1948.8EBDAGCF	G24	1476.9FBDEC	G6	1886.6EBDACF	G2	770.67BDAC	G15	1486.16
13	G18	1430EBDACF	G11	1806.3EBDHGCF	G15	1466FBDEC	G4	1832.8EBDACF	G13	770.05BDAC	G7	1480.54
14	G3	1404.3EBDCF	G4	1786.5EDHGCF	G6	1393.1FGDEC	G1	1815.9EBDACF	G7	756.3BDEC	G4	1471.45
15	G21	1401EBDCF	G16	1778.2EDHGCF	G4	1352FGDE	G10	1804.5EBDACF	G12	746.35BDEC	G10	1468.25
16	G25	1382.8EBDCF	G7	1746.7EDHGF	G22	1342.1FGDE	G5	1794.5EBDACF	G14	734.93DEC	G5	1459.13
17	G14	1380.5EBDCF	G21	1715.4EDHGF	G18	1329.1FGDEH	G18	1704.2EBDCF	G1	734.75DEC	G16	1450.65
18	G19	1349.2EDGCF	G18	1615.8EHGF	G19	1321.7FGEH	G2	1671.3EDCF	G19	731.19DEC	G23	1432.33
19	G1	1348EDGCF	G23	1610.8EHGF	G9	1281.1FGEH	G9	1655.4EDF	G10	728.84DEC	G18	1397.76
20	G5	1347.3EDGCF	G20	1594.5HGF	G2	1278.9FGEH	G16	1615.2EDF	G16	717.69DEC	G2	1397.63
21	G23	1316.2EDGCF	G24	1594.3HGF	G23	1256.7FGEH	G7	1614.8EDF	G8	717.22DEC	G22	1396.82
22	G9	1223.7EDGF	G25	1527.7HG	G25	1178.8FGH	G22	1586.8EDF	G6	714.69DEC	G24	1390.28
23	G8	1183.4EGF	G2	1462.6HG	G3	1168.7FGH	G24	1577.2EDF	G3	652.89FDE	G25	1385.84
24	G22	1114GF	G10	1346.9H	G5	1104.9GH	G13	1548.2EF	G24	610.04FE	G20	1379.77
25	G20	981.2G	G9	1324.9H	G12	1000.8H	G15	1517.2F	G11	555.81F	G9	1267.62
	Mean	1440.52		1863.36		1448.16		1858.13		767.6		1475.75
	LSD	380.87		529.94		333.77		488.26		15248		176.99
	R2	59.72		55.53		79.99		88.33		93.34		
	CV	23.08		24.82		20.12		22.92		17.34		

G= genotype, GM= grand mean, CV coefficient of variation

Additive main effect and multiplicative interaction analysis (AMMI)

AMMI analysis (Table 5) showed that environment and GEI effects were highly significant ($p<0.0001$), and genotype effect was significant ($p<0.05$) for grain yield. The test environments contributed 63.98% of the total variation in yield. Genotype and GEI accounted for 2.66% and 16.30% of the total variation for grain yield, respectively. The ratio of genotype effect to genotype + genotype $\times$ environment (G+G $\times$ E) was 0.14. The magnitude of the GEI sum of squares was 6.12 times that of the genotype sum of squares for grain yield. The AMMI model extracted three highly significant ($p<0.0001$) IPCA's from the interaction PC axes (Table 5). Those three IPCA's accounted for a total of 90.11% of the observed variation due to GEI. IPCA1, IPCA2, and IPCA3 captured 45.47%, 28.05%, and 16.59% of the sum of squares, respectively.

Table 5 AMMI analysis of variance for grain yield (kg ha⁻¹) of 25 cowpea genotypes evaluated at five environments

Source of variation	DF	SS	M.S.	Sum of squares explained (%)		
				Total VE	GEI E	GEI cum
Treatments	124	77849952	627822			
Genotypes	24	2500687	104195*	2.66		
Environments	4	60052810	15013202**	63.98		
Block	10	559100	55910			
Interactions	96	15296455	159338**	16.3		
IPCA 1	27	6710300	248530**		45.47	45.47
IPCA 2	25	4336683	173467**		28.05	73.52
IPCA 3	23	2668520	116023*		16.59	90.11
Residuals	21	1580952	75283			
Error	240	15453907	64391			
Total	374	93862959	250970			

**, **= significant at 5% and 1% probability level, respectively, Total VE=total variation explained, GEI E= GEI explained, GEI cum= GEI cumulative, SS= sum of squares, MS= mean squares, DF= degree of freedom*

Lin and Binns Cultivar superiority measure (Pi)

According to this stability model, the three most stable genotypes with the lowest Pi values were G8, G17 and G13, which ranked 1st, 2nd and 3rd for grain yield (Table 6). The most unstable genotypes were G9, G2, and G25 which ranked 25th, 20th and 24th for grain yield.

Table 6 Grain yield and cultivar superiority value (Pi)

Genotypes	$\bar{y}$ Mean	R^2	y Rank	Pi	P_i Rank
G1	1515.80	0.94	9	65612.64	20
G2	1397.63	0.61	20	171567.2	2
G3	1489.99	0.90	11	99346.36	14
G4	1471.45	0.95	14	97339.92	15
G5	1459.13	0.78	16	114667.1	10
G6	1566.2	0.91	4	53755.99	23
G7	1480.54	0.79	13	101497.7	12
G8	1641.25	0.88	1	47458.89	24
G9	1267.61	0.83	25	234259.9	1
G10	1468.25	0.50	15	150238.8	5
G11	1523.12	0.90	7	76205.07	19
G12	1529.22	0.82	6	100484	13
G13	1582.331	0.73	3	62452.33	21
G14	1551.03	0.96	5	55618.29	22
G15	1486.16	0.79	12	93679.66	16
G16	1450.65	0.88	17	106111.3	11
G17	1595.47	0.98	2	44651.05	25
G18	1397.76	0.97	19	146153.6	7
G19	1513.18	0.93	10	80879.75	18
G20	1379.77	0.69	24	167535.7	3
G21	1521.99	0.90	8	81427.34	17
G22	1396.83	0.76	21	135756	9
G23	1432.33	0.80	18	137337.8	8
G24	1390.28	0.76	22	146642	6
G25	1385.85	0.79	23	161189.5	4

$\bar{y}$ = grain yield

GGE biplots

Which-won-where and what

The polygon view of the genotypes in the GGE biplot for 25 genotypes is presented in Figure 1. The cumulative variation contributed by PC1 (AXIS 1) and PC2 (AXIS2) was 69.16%, both which were highly significant. The biplot showed that two environments (Babile and Sirnka) grouped in the same mega environments. The other three environments (Jinka, Melkassa, and Miesso) each fell in a different mega environment. The plot showed that G8, G13, G3, G16 ANG G4 recorded the highest grain yield in Jinka, Melkassa, Miesso, Sirnka, and Babile. On the other hand, G9, G25, and G23 did not fall in a specific environment, and were poor yielders.

Ideal genotypes (ranking genotypes)

The first two principal components (PC1 and PC2) were highly significant ($p < 0.0001$) and explained 42.31% and 26.85% of the yield variation among the genotypes, respectively (Figure 2). The GGE biplot which identifies ideal genotypes which are high yielding and stable across the test environments, identified G8, G6 and G1, which fell close to the center of the concentric circle as ideal genotypes. Based on the average environment coordination (AEC) method, G9, G25 and G23 were the most unstable and low yielding genotypes across the environments (Figure 2)

Ideal environment (ranking environment)

Jinka was close to the concentric circle and provided the most ideal environment and the most powerful to discriminate performance of the tested genotypes (Figure 3). In contrast, Babile was located far from the center of the concentric circle, indicating poor discriminating power.

Mean vs. stability

The GGE biplot visualizes performance and effectively identifies the best performing genotypes across environments with the help of the Average Environment Coordinate (AEC). The mean of PC1 and PC2 scores of the tested environments is represented by the arrowhead, and the AEC ordinate is the line that passes through the biplot origin and it is perpendicular to the AEC abscissa (Figure 4).

The length of the abscissa discriminates the grain yield of genotypes that are above and below average yield if right and left of the biplot origin, respectively (Figure 4). The length of the ordinate approximates the GEI associated with the genotype stability, and a longer ordinate corresponds to higher variability and lower stability and vice-versa. NLLP-CPC-07-10, NLLP-CPC-07-57, ACC-216-747, NLLP-CPC-07-145-21, NLLP-CPC-07-28, NLLP-CPC-07-36, and NLLP-CPC-103-B were above average yielders with higher stability, however, NLLP-CPC-07-169, ACC-215-821, NLLP-CPC-07-156, NLLP-CPC-07-54, and NLLP-CPC-07-03-B had above average yield but with lower stability. Furthermore, CP-EXETERETIS, Dass 001, NLLP-CPC-07-143, NLLP-CPC-07-166, NLLP-CPC-07-167, NLLP-CPC-07-157, and NLLP-CPC-07-140 were stable but their yield below average, and NLLP-CPC-07-139, ACC-215-762, NLLP-CPC-07-19, ACC-211-490 Bole and Kanketi were not stable and their yield also below the average.

AMMI stability value (ASV) and yield stability index (YSI)

Utilizing yield stability index (YSI), the combination of AMMI stability values and average grain yield was estimated to quantify and classify the genotype (Table 7). According to the YSI model, G14, G17, and G21 were the most stable genotypes across environments and high grain yielders. On the other hand, genotypes G10, G2, and one check (Bole) were unstable as indicated by high YSI values of 40, 41, and 45 respectively, and those genotypes had poor productivity and lower stability.

Table 7 AMMI stability value (ASV), yield stability index, ranks and IPCA scores

Genotypes	IPCA1	IPCA2	ASV	rASV(A)	YLD	r YLD(B)	YSI (A+B)
G1	-0.41	-0.36	0.73	13	1515.80	9	22
G2	1.00	0.01	0.92	21	1397.63	20	41
G3	0.33	0.13	0.84	19	1489.99	11	30
G4	-0.67	0.46	0.34	6	1471.45	14	20
G5	-0.05	-0.84	0.80	16	1459.13	16	32
G6	-0.09	-0.11	0.81	17	1566.20	4	21
G7	-0.18	-0.62	0.74	14	1480.54	13	27
G8	0.16	-0.41	0.96	22	1641.25	1	23
G9	0.19	-0.03	0.68	11	1267.62	25	36
G10	0.16	0.03	1.55	25	1468.25	15	40
G11	-0.45	0.35	0.53	8	1523.12	7	15
G12	0.59	0.10	1.13	24	1529.22	6	30
G13	-0.10	0.23	0.84	20	1582.33	3	23
G14	0.13	0.15	0.17	1	1551.03	5	6
G15	-0.48	-0.39	0.68	10	1486.16	12	22
G16	0.06	0.70	0.48	7	1450.65	17	24
G17	0.66	-0.15	0.30	5	1595.47	2	7
G18	0.15	0.61	0.25	3	1397.76	19	22
G19	-0.39	0.59	0.77	15	1513.18	10	25
G20	0.21	0.11	0.27	4	1379.77	24	28
G21	-0.52	0.01	0.24	2	1521.99	8	10
G22	-0.50	-0.24	0.83	18	1396.82	21	39
G23	0.38	-0.46	0.70	12	1432.33	18	30
G24	-0.62	-0.09	1.03	23	1390.28	22	45
G25	0.41	0.25	0.65	9	1385.84	23	32

Discussion

The majority of cowpea growing environments are extremely vulnerable to moisture stress, and the farmers use this crop as an insurance crop as they experience prolonged drought in this area. The higher yield variation contributed by the environment over genotype and GEI, indicates that the test

environments were highly variable and had a great impact on genotypes. The significant GEI necessitates the need to identify adaptable genotypes with consistent high grain yield (Yan & Tinker, 2006).

In addition, the response of genotypes varied considerably for grain yield due to the genetic makeup of the materials and the interaction between genetic constitution and environmental influences. This is in agreement with Shegro et al. (2020) who reported that genotype and environment directly affect the yield potential of cowpea. From this study, the effect of environment, season, and environment $\times$ season was high as it was responsible for 33.98%, 16.23%, and 29.27% of the total variation for grain yield, respectively. Thus, the environment and the season effects were very high, contributing to diverse cowpea grain yield. Cowpea grain yield will therefore also be largely affected by climate change. A previous study also reported variation in responses of genotype across environments in different seasons (Kuruma et al. 2019).

Generally, the existing variation due to environment, season, genotype performance and GEI in relation to genotype effect suggested that there was the possibility of mega environment effects for different genotypes. Therefore, based on the variable response of genotypes, it would help to map the mega environments (MEs) suitable for the improvement of grain yield to combat the rapid climate change.

Grain yield increment is the goal of cowpea for any stressed environment because yield is governed by multi traits with different levels of expression for various environments and their interaction. In the present study, 52% of the tested genotypes recorded a higher yield than the standard checks (Bole and Kanketi). Genotype NLL-CPC-07-145-21 had a yield advantage of 15.26% and 22.25% compared to the worst genotype (NLLP-CPC-07-143), and the standard checks, respectively. Therefore identification of the highest yielding and adaptable genotypes for the specific range of environments is important for the selection and evaluation of superior genotypes in multi-environment studies and are the main targets of cowpea breeding programs. Kuruma et al. (2019) and Muranaka et al. (2016) reported that high grain yield variation could be due to greater differences between the genotypes.

AMMI analysis showed that the environmental effects accounted for the most (63.98%) of the total variation compared to the other components, implying that differential cowpea yield performance was typically caused by environmental changes. This is in agreement with the findings of Shegro et al. (2020) and Simion (2018) that environment made the largest contribution to grain yield variation in cowpea. The magnitude of the GEI sum of squares was 6.12 times that of the genotype sum of squares for grain yield of cowpea, indicating that there were considerable differences in genotypic responses across environments, which was contrary with the findings of Shegro et al. (2020).

In the present investigation, the three IPCA's accounted for 90.11% of the interaction sum of squares. Zobel et al. (1988) stated that AMMI with the first two multiplicative terms was the best predictive model. In this study, the high (45.47%) and significant contribution of IPCA1 to the total variation across the tested environments implies that IPCA1 could identify stable and unstable genotypes based on the value scores or nearest or furthest to zero, which is in line with the findings of previous investigations (Muranaka et al., 2016; Shegro et al., 2020; Simion, 2018; Yaw et al., 2020). The positive and negative IPCA scores of genotypes in AMMI analysis are the best indicators of stability or adaptation over environments. High positive interaction of the genotypes like G2 (NLLP-CPC-07-169) in IPCA1 in an environment can exploit the agro-ecological conditions of the specific environment (Simka). Therefore, it would be possible to identify adaptable and suitable genotype/s for the specific environment. Kandus et al. (2010) and Yan et al. (2007) reported that the different high yielding genotypes fall in a specific environment, and it shows crossover GEI, suggesting that the test environment could be classified into mega environments.

Using Pythagorean Theorem the distance from the origin (0:0) in a two-dimensional scattergram indicates the most stable genotypes (Purchase et al. 2000). In the ASV method, a genotype with the lowest ASV score is the most stable; accordingly, G14, G17, and G21 were stable and these genotypes were the highest yielding among the tested genotypes, indicating that the yield performance and stability showed a similar trend. Oliveira et al. (2014) noted that the dynamics of stable genotype and yield response is always parallel to the mean response of the tested environments. Unstable genotypes like Bole, CP-EXTERETIS, and ACC-215-762 had high ASV values, and they were adapted to a specific favorable environment. Likewise, Oladosu et al. (2017) reported that the higher the IPCA score, and AMMI stability value, the more specifically adapted a genotype is to a certain environment. In this study, crossover stability and yield did not have the same trend for all genotypes across season and environment. Jadhav et al. (2019) and Mohammadi & Amri (2008) suggested that principle stability *per se* should, however, not be the only selection parameter because the most stable genotypes would not necessarily give the best yield performance. Therefore, there is a need for approaches that incorporate both mean yield and stability in a single index.

The lowest YSI value is considered as the most stable, with high grain yield (Bose et al. 2014). NLLP-CPC-07-28, NLLP-CPC-07-145-21, and NLLP-CPC-0754 were the most stable genotypes with good yield performance. Thus, according to the YSI method, the most desirable genotypes can be considered as widely adapted and with grain yield above the grand mean among 25 genotypes. Similarly Zali et al. (2012) also indicated that both yield and stability of performance should be considered simultaneously to exploit the useful effect of GEI and to select genotypes for diverse environments.

The polygon view of the "which-won-where and what" GGE-biplot (Figure 1) showed that genotypes NLLP-CPC-07-143, ACC-215-762, NLLP-CPC-07-156, NLLP-CPC-07-54, NLLP-CPC-07-145-21, and ACC-211-490 were genotype markers located farthest from the biplot origin in various directions and it shows that the genotypes were well adapted to specific environments. Shegro et al. (2020) reported similar results on cowpea. If the environment markers fall in different sectors it shows that different cultivars won in different environments (Oladosu et al., 2017; Shegro et al., 2020). In addition, because of long environment vectors, the rest of the genotypes scattered around the biplot origin. This clearly explained that the environment effect was higher than the genotype effect. This implies that the genotype had less response for genotype by environment interaction because of high environment exertion to the GEI. The GGE-biplot showed that the tested environments occurred in different sectors, indicating that the particular environment had different high yielding genotypes for those sectors, this indicating the existence of crossover GEI. It also indicated the possibility to classify the environments into mega-environments for cowpea production. Yan & Rajcan (2002) stated that the presence or absence of crossover GEI indicates the existence of different mega-environments. Generally, the GGE biplot effectively identified the best performing genotypes across environments and best genotypes for specific environments, whereby specific genotypes can be recommended for specific environments and can be used to evaluate the yield and stability of genotypes, which is not possible with AMMI analysis (Kaya et al., 2006; Yan & Tinker, 2006).

GGE biplots help visualize and compare the distance between each genotype and the ideal genotype located at the center of the concentric circle (Yan and Rajcan 2002). The ideal genotypes, based on proximity to the center of the concentric circle of the GGE biplot were ACC-216-747 and NLLP-CPC-07-145-21, with high yield and stability (Figure 2). In addition, NLLP-CPC-07-10 was located on the next homocentric circle and might be considered as a desirable genotype. In principle, the ideal genotype should have the longest vector, highest mean performance and with zero GEI, and/or it should perform consistently in all environments. Because of the genetic background or nature of the traits (yield) and level of expression, the ideal genotype does not always exist in reality. Therefore, such like genotypes the breeders can be used as a reference for genotype for further study. Genotypes which were high yielding but were not stable across environments could be recommended for a particular environment.

Yan & Rajcan (2002) and Yan et al. (2007) specified that the environments with long vectors (PC1 scores) and relatively small angles or absolute with the AEC abscissa are valuable for greater discriminatory capacity (in terms of the genotype main effect) and is representative of the other environments. Therefore, Jinka was in the epicenter of the concentric circle, and it was identified as a highly discriminating environment for these genotypes, thus this environment is considered as an ideal environment for developing high yielding genotypes, or for identify ideal genotypes. Hence, Jinka allowed the genotype to express genetic potential, minimizing population development expenses by discriminating the worst genotypes at an early stage.

Conclusions

The present study showed that genotype, environment, and genotype $\times$ environment, genotype $\times$ season and environment $\times$ season interaction effects were significant for grain yield. The performance of tested genotypes was largely affected by the environment. Jinka was the most ideal environment and it had discriminatory power for genotype performance in all seasons. The AMMI showed that the environmental effects accounted for 63.98% of the total yield variation. The GGE biplot showed that the tested environments occurred in different sectors which could be used for classifying the environments into mega-environments for cowpea production. In terms of ideal genotypes (having a long vector), good yielding capacity and stability, NLLP-CPC-07-145-21, NLLP-CPC-103-B and NLLP-CPC-07-54 were identified as ideal genotypes for drought-prone environments for the country and could be proposed for release for production. The results of this study confirmed that the ideal genotype should have a long vector, high yield performance, but such genotypes do not always exist in reality.

Declarations

Acknowledgements

The authors sincerely acknowledge the contribution of Wolkite University, Melkassa Agricultural Research Center (EIAR), and The McKnight Foundation USA, for providing the research funds to complete this project. The author acknowledges the staff of national lowland pulse improvement program of Melkassa Agricultural Research Center (MARC) is highly appreciated for their comprehensive support.

Conflict of interest: The authors declare that they have no conflict of interest.

References

- Agbicodo EM, Fatokun CA, Muranaka S, et al (2009) Breeding drought tolerant cowpea: Constraints, accomplishments, and future prospects. *Euphytica* 167:353–370. <https://doi.org/10.1007/s10681-009-9893-8>
- Ahenkora K, Adu Dapaah HK, Agyemang A (1998) Selected nutritional components and sensory attributes of cowpea (*Vigna unguiculata* [L.] Walp) leaves. *Plant Foods Hum Nutr* 52:221–229. <https://doi.org/10.1023/A:1008019113245>
- Arpita D, Ashok K, Parihar D, et al (2019) Deciphering Genotype-by- Environment Interaction for Targeting Test Environments and Rust Resistant Genotypes in Field Pea (*Pisum sativum* L .). 10: <https://doi.org/10.3389/fpls.2019.00825>
- Bose LK, Jambhulkar NN, Pande K, Singh ON (2014) Use of AMMI and other stability statistics in the simultaneous selection of rice genotypes for yield and stability under direct-seeded conditions. *Chil J Agric Res* 74:1–9. <https://doi.org/10.4067/S0718-58392014000100001>
- Diouf D (2011) Recent advances in cowpea [*Vigna unguiculata* (L .) Walp .] " omics " research for genetic improvement. 10:2803–2810. <https://doi.org/10.5897/AJBx10.015>
- Farshadfar E, Mahmodi N, Yaghotipoor A (2011) AMMI stability value and simultaneous estimation of yield and yield stability in bread wheat (*Triticum aestivum* L.). *Aust J Crop Sci* 5:1837–1844
- Fatokun CA, Boukar O, Muranaka S (2012) Evaluation of cowpea (*Vigna unguiculata* (L .) Walp .) germplasm lines for tolerance to drought. *Plant Genet Resour Charact Util* 10:171–176. <https://doi.org/10.1017/S1479262112000214>
- Filho JT, Noronha C, Dos G, et al (2017) GENOTYPE BY ENVIRONMENT INTERACTION IN GREEN COWPEA ANALYZED VIA MIXED MODELS 1. 30:687–697
- Gomez KA, Gomez AA (1984) Statistical procedures for agricultural research, 2nd edn. John Wiley and Sons Inc., New York, USA
- Goufo P, Moutinho-pereira JM, Jorge TF, Correia CM (2017) Cowpea (*Vigna unguiculata* L . Walp .) Metabolomics: Osmoprotection as a Physiological Strategy for Drought Stress Resistance and Improved Yield. *Front Plant Sci* 8:1–22. <https://doi.org/10.3389/fpls.2017.00586>

- Hall AE, Cisse N, Thiaw S, et al (2003) Development of cowpea cultivars and germplasm by the Bean / Cowpea CRSP. *F Crop Res* 82:103–134. [https://doi.org/10.1016/S0378-4290\(03\)00033-9](https://doi.org/10.1016/S0378-4290(03)00033-9)
- Hartley O. (1950) The Maximum F-Ratio as a Short-Cut Test for Heterogeneity of Variance. *Biometrika* 37:308–312
- Jadhav S, Balakrishnan D, V GS, et al (2019) Genotype by Environment (G × E) Interaction Study on Yield Traits in Different Maturity Groups of Rice. 22:425–449. <https://doi.org/10.1007/s12892-018-0082-0>
- Kandus, M., D. Almora, R.B. Ronceros and JCS (2010) Statistical models for evaluating the genotype-environment interaction in Statistical models for evaluating the genotype-environment interaction in maize (*Zea mays* L .) Modelos estadísticos para evaluar la interacción genotipo-ambiente en maíz (*Zea mays*. *Phyton* (B Aires) 79:39–46. <https://doi.org/10.32604/phyton.2010.79.039>
- Kaya Y, Akçura M, Taner S (2006) GGE-Biplot analysis of multi-environment yield trials in bread wheat. *Turkish J Agric For* 30:325–337
- Kuruma RW, Sheunda P, Muriuki C (2019) YIELD STABILITY AND FARMER PREFERENCE OF COWPEA (*VIGNA UNGUICULATA*) LINES IN SEMI-ARID EASTERN KENYA. *afrika Focus* 32:65–82
- Leflon M, Lecomte C, Barbottin A, et al (2008) Characterization of Environments and Genotypes for Analyzing Genotype × Environment Interaction Some Recent Advances in Winter Wheat and Prospects for QTL Detection Characterization of Environments and Genotypes for Analyzing Genotype × Environment Intera. *J Crop Improv* \ 14:249–298. <https://doi.org/10.1300/J411v14n01>
- Lin CS, Binns MR (1988) A SUPERIORITY MEASURE OF CULTIVAR PERFORMANCE FOR CULTIVAR X LOCATION DATA. *Can J Plant Sci* 68:193–198
- Marina M, Fernández J, Ochoa J, et al (2017) Genotype by environment interactions in cowpea (*Vigna unguiculata* L . Walp .) grown in the Iberian Peninsula. *Crop Pasture Sci* 68:924–931
- Mohammadi R, Amri ÆA (2008) Comparison of parametric and non-parametric methods for selecting stable and adapted durum wheat genotypes in variable environments. *Euphytica* 159:419–432. <https://doi.org/10.1007/s10681-007-9600-6>
- Muranaka S, Shono M, Myoda T, Takeuchi J (2016) Genetic diversity of physical , nutritional and functional properties of cowpea grain and relationships among the traits. *Plant Genet Resour Charact Util* 14:67–76. <https://doi.org/10.1017/S147926211500009X>
- Oladosu Y, Rafii MY, Abdullah N, et al (2017) Genotype × Environment interaction and stability analyses of yield and yield components of established and mutant rice genotypes tested in multiple locations in Malaysia. 67:590–606. <https://doi.org/10.1080/09064710.2017.1321138>
- Olajide AA, Ilori CO (2017) Effects of Drought on Morphological Traits in Some Cowpea Genotypes by Evaluating Their Combining Abilities. *Adv Agric* 2017:1–10
- Oliveira EJ De, Paulo J, Freitas X De, Jesus ON De (2014) AMMI analysis of the adaptability and yield stability of yellow passion fruit varieties. *Sci Agrcola* 71:139–145
- Owade JO, Abong G, Okoth M, Mwang AW (2020) A review of the contribution of cowpea leaves to food and nutrition security in East Africa. *Food Sci Nutr* 8:36–47. <https://doi.org/10.1002/fsn3.1337>
- Pacheco A, Vargas M, Alvarado G, et al (2016) USER ' S MANUAL GEAR (Genotype by Environment Analysis with R). *GEAR* 1–42
- Purchase JL, Hatting H, Deventer CS Van (2000) Genotype × environment interaction of winter wheat (*Triticum aestivum* L .) in South Africa : I . AMMI analysis of yield performance. *South African J Plant Soil* 17:95–100. <https://doi.org/10.1080/02571862.2000.10634877>
- SAS II (2013) SAS 9.4 Statements ®. 476
- Shegro A, Willem G, Jansen S, et al (2020) Genotype and genotype 3 environment interaction effects on the grain yield performance of cowpea genotypes in dryland farming system in South Africa. *Euphytica* 216:1–11. <https://doi.org/10.1007/s10681-020-02611-z>
- Simion T (2018) Adaptability Performances of Cowpea [*Vigna Unguiculata* (L .) Walp] Genotypes in Ethiopia. *Food Sci Qual Manag* 72:43–47
- Simion T, Mohammed W, Fenta BA (2018) Genotype by environment interaction and stability analysis of cowpea [*Vigna unguiculata* (L .) Walp] genotypes for yield in Ethiopia. *J Plant Breed Crop Sci* 10:. <https://doi.org/10.5897/JPBCS2018.0753>
- Singh BB, Ajeigbe HA, Tarawali SA, et al (2003) Improving the production and utilization of cowpea as food and fodder. *F Crop Res* 84:169–177. [https://doi.org/10.1016/S0378-4290\(03\)00148-5](https://doi.org/10.1016/S0378-4290(03)00148-5)
- Timko, M.P. and Singh BB (2008) Cowpea, a Multifunctional Legume. In: Paul, H. Moore and Ray M (ed) *Genomics of Tropical Crop Plants*, 1st edn. Springer Science+Business Media, LLC, pp 227–258
- Tumuhimbise R, Melis R, Shanahan P, Kawuki R (2014) ScienceDirect Genotype × environment interaction effects on early fresh storage root yield and related traits in cassava. *Crop J* 2:329–337. <https://doi.org/10.1016/j.cj.2014.04.008>

Weng Y, Jun Q, Eaton S, et al (2019) Evaluation of Seed Protein Content in USDA Cowpea Germplasm. 54:814–817. <https://doi.org/10.21273/HORTSCI13929-19>

Yan W (2016) Analysis and Handling of G × E in a Practical Breeding Program. 56:2106–2118. <https://doi.org/10.2135/cropsci2015.06.0336>

Yan W (2011) GGE Biplot vs. AMMI Graphs for Genotype-by-Environment Data Analysis. *tJournal Indian Soc Agric Stat* 65:183–193

Yan W (2002) Singular-value partitioning in biplot analysis of multienvironment trial data. *Agron J* 94:990–996. <https://doi.org/10.2134/agronj2002.0990>

Yan W, Hunt LA, Sheng Q, Szlavics Z (2000) Cultivar evaluation and mega-environment investigation based on the GGE biplot. *Crop Sci* 40:597–605. <https://doi.org/10.2135/cropsci2000.403597x>

Yan W, Kang MS, Ma B, et al (2007) GGE biplot vs. AMMI analysis of genotype-by-environment data. *Crop Sci* 47:643–655. <https://doi.org/10.2135/cropsci2006.06.0374>

Yan W, Rajcan I (2002) Biplot Analysis of Test Sites and Trait Relations of Soybean in Ontario. *Crop Sci* 42:11–20. <https://doi.org/10.2135/cropsci2002.0011>

Yan W, Tinker NA (2006) Biplot analysis of multi-environment trial data: Principles and applications. *Can J Plant Sci* 86:623–645. <https://doi.org/10.4141/P05-169>

Yaw E, Isaac O, Amegbor K, et al (2020) Genotype × environment interactions of yield of cowpea (*Vigna unguiculata* (L.) Walp.) inbred lines in the Guinea and Sudan Savanna ecologies of Ghana. *J Crop Sci Biotechnol* 23:453–460. <https://doi.org/10.1007/s12892-020-00054-5>

Zali H, Farshadfar E, Sabaghpour SH (2012) Evaluation of genotype × environment interaction in chickpea using measures of stability from AMMI model. *Ann Biol Res* 3:3126–3136

Zobel RW, Wright MJ, Gauch HG (1988) Statistical Analysis of a Yield Trial. *Agron J* 80:388–393. <https://doi.org/10.2134/agronj1988.00021962008000030002x>

Figures

Figure 1

Polygon view of GGE-biplot for 25 genotypes evaluated across five environments

Figure 2

GGE-biplot showing the ideal genotypes based on mean grain yield performance across environments

Figure 3

GGE-biplot showing the ideal environments for the 2017 and 2018 seasons

Figure 4

Mean vs. Stability view of GGE biplot showing the mean performance and stability of 25 genotypes