

Health-Related Quality of Life and Associated Factors in Chinese Menstrual Migraine Patients

Wenxiu Luo

Affiliated Hospital of North Sichuan Medical College

Xing Cao

Affiliated Hospital of North Sichuan Medical College

Jiayu Zhao

Affiliated Hospital of North Sichuan Medical College

Jiaming Yang

Affiliated Hospital of North Sichuan Medical College

Yu Cen

North Sichuan Medical College [Search North Sichuan Medical College]: North Sichuan Medical University

Jinlong He

North Sichuan Medical College [Search North Sichuan Medical College]: North Sichuan Medical University

Jing Luo

North Sichuan Medical College [Search North Sichuan Medical College]: North Sichuan Medical University

Yunling Zhong

North Sichuan Medical College [Search North Sichuan Medical College]: North Sichuan Medical University

Ying Luo

North Sichuan Medical College [Search North Sichuan Medical College]: North Sichuan Medical University

Xinyue Wang

North Sichuan Medical College [Search North Sichuan Medical College]: North Sichuan Medical University

Liqu Yang

North Sichuan Medical College [Search North Sichuan Medical College]: North Sichuan Medical University

Xingyu Feng

North Sichuan Medical College [Search North Sichuan Medical College]: North Sichuan Medical University

Xiaoqing Pang

North Sichuan Medical College [Search North Sichuan Medical College]: North Sichuan Medical University

Jiaming Luo (✉ jiamingluo@nsmc.edu.cn)

North Sichuan Medical College [Search North Sichuan Medical College]: North Sichuan Medical University <https://orcid.org/0000-0003-0025-0379>

Research Article

Keywords: Menstrual migraine, Health related quality of life, Associated factors, SF-36

Posted Date: November 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1055240/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Abstract

Background

Menstrual migraine is a special form of migraine with significant impact on quality of life for those women afflicted. Presently, there is no study reported on quality of life in menstrual migraine patients. The study reported here aimed to assess the health-related quality of life and identify its associated factors among Chinese menstrual migraine patients.

Methods

The cross-sectional study group consisted of 109 patients with menstrual migraine and the control group consisted of 397 female patients with non-menstrual migraine. In total, 506 patients completed questionnaires for demographic and clinical information, Self-rating Idea of Suicide Scale, Hamilton Depression Scale, Hamilton Anxiety Scale, Headache Impact Test-6, Perceived Social Support Scale, Pittsburgh Sleep Quality Index. Health-related quality of life was measured using the 36-Item Short Form Survey.

Results

Compared with non-menstrual migraine patients, five dimensions of health-related quality of life were all found to be significantly impaired in menstrual migraine patients. Headache frequency ($\beta=-0.218$, $P=0.014$), the impact of headache on daily life ($\beta=-0.270$, $P=0.002$), depression symptoms ($\beta=-0.345$, $P<0.001$) were significantly associated with physical component summary, depression symptoms ($\beta=-0.379$, $P<0.001$), social support ($\beta=0.270$, $P<0.001$), suicidal ideation ($\beta=-0.344$, $P<0.001$) were closely related to mental component summary.

Conclusion

Menstrual migraine patients had significantly poorer health-related quality of life in many domains than non-menstrual migraine patients. Headache frequency, the impact of headache on daily life, depression symptoms, social support and suicidal ideation were significantly associated with health-related quality of life in menstrual migraine patients.

Trial registration:

ChiCTR1800014343. Registered 01 July 2018 at Chinese Clinical Trial registry. <http://www.chictr.org.cn>

Background

Migraine is a common debilitating headache disorder that was ranked as the second highest cause of disability in the 2016 Global Burden of Disease study[1]. Menstrual migraine (MM) is a subclass of migraine that can be classified into two types: pure menstrual migraine and menstrually-related migraine[2], and the prevalence of MM in female migraine patients is nearly 60%[3]. Compared with non-menstrual Migraine (NMM), MM is characterised by longer duration, greater frequency, less responsive to treatment, and may cause more severe disability and heavier life burden to patients[4, 5].

Health-related quality of life is defined as an individual's subjective perception of the impact of disease and treatment on physical, psychological, social and somatic domains of functioning and well-being[6]. HRQoL assessments could provide patients, researchers and policy makers with information about the status of a patient's health in an efficient manner[7]. Limited number of studies indicate a negative effect of migraine on quality of life[8, 9]. In clinical practice, it has also been observed that a considerable proportion of MM patients suffer from anxiety and depression symptoms, sleep disorders and other unpleasant mental and physical experience, leading to decreased quality of life. Therefore, it is of great significance to actively search for and devote to improving controllable risk factors for MM patients. Clinicians have found the contribution of that demographic factors such as age, clinical factors such as frequency and severity of pain, and psychological factors such as anxiety and depression symptom play an important role in decreasing the migraine patients' quality of life. However, studies on the quality of life and its associated factors in patients with MM are scarce. Based on the particularities of MM, further study of factors related to quality of life with MM is warranted.

The purpose of this cross-sectional study was to explore the HRQoL of MM versus NMM and to identify its associated factors among Chinese MM patients. It was hypothesized that (1) MM patients have lower HRQoL when compared with NMM; (2) Some clinical and psychological factors could affect the HRQoL of MM patients. Our research would provide a more comprehensive understanding of the HRQoL and associated factors among Chinese MM patients. Understanding these factors may help the development of individualized strategies for patients, thereby raising the HRQoL and reducing the financial, social and psychological burden.

Method

Participants and study design

A cross-sectional study was conducted from February 2019 to February 2020 at the neurology department of a hospital in western China. A total of 545 female migraine patients who met the diagnostic criteria were screened, and 39 persons who did not complete the HRQoL scale were excluded (3 with MM, 36 with NMM). A total of 506 female patients diagnosed with MM (n = 109) and NMM (n = 397) were included in this study, the inclusion criteria are in line with the International Classification of Headache Disorders, Third Edition (ICHD-3) developed by the Headache Classification Committee of the International Headache Society (Two experienced neurologists confirmed all diagnoses using ICHD-3 diagnostic criteria). Exclusion criteria: 1. Patients who had been diagnosed with psychiatric illness before

the headache occurred, including anxiety disorder, depressive disorder, sleep disturbance, etc; 2. Patients with secondary headaches; 3. patients suffered from other disease that affect the quality of life; 4. patients who could not understand the content of questionnaire or did not complete the questionnaire; 5. Patients who refuse to sign informed consent. All participants signed an informed consent, and this study was approved by the Survey Ethics Committee of the first author's affiliated institution. All authors had full access to all data.

Data Collection

Demographic and headache information

Demographic variables include age, ethnicity (Han/minority), height (cm), weight (kg), body mass index (BMI), BMI is calculated as $BMI = \text{body mass (kg)} / (\text{height (m)})^2$. Headache related information include headache frequency (the average number of days with headache per month over the last 3 months), duration (<24h/24-72h/>72h), severity (mild/moderate/severe), with or without family History and with or without aura.

Health-related quality of life

HRQoL was measured using the Mandarin version of 36-Item Short Form Survey (SF-36) which has been tested with satisfactory reliability and validity[10, 11]. The SF-36 is a widely used survey of self-reported physical and mental health with 36 questions that measures 8 dimensions of health: physical functioning (PF), role-physical (RP), bodily pain (BP), general health (GH), vitality (VT), social functioning (SF), role-emotional (RE), and mental health (MH). The eight dimensions can also be summarized as physical component summary (PCS) and mental component summary (MCS), which are respectively average value of physical health dimensions (PF, RP, BP, GH) and mental health dimensions (VT, SF, RE, MH)[7].

Impact of Headache on daily life

We performed the Headache Impact Test-6 (HIT-6) to assess the impact of headache on daily life. The HIT-6 is a concise and reliable tool for measuring headache burden based on six domains with total score ranging from 36 to 78. The higher the score, the greater impact of headaches on daily activities[12].

Anxiety and depression symptoms

The severity of anxiety symptoms and depression symptoms of patients were assessed using the Hamilton Anxiety Scale (HAMA) and the Hamilton Depression Scale (HAMD) respectively. The higher the score, the more severe the anxiety or depression symptoms are[13, 14].

Social support

Perceived Social Support Scale (PSSS) consists of 12 items, assessing perceived support from family, friends, and significant others. The total score ranges from 12-84, with higher scores representing higher perceived social support[15].

Sleep quality

We measured sleep quality of patients using the Pittsburgh Sleep Quality Index (PSQI), a questionnaire that assesses subjectively perceived sleep quality over the last month. The PSQI is a self-report instrument that consists of 7 components: subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleeping medications, and daytime dysfunction. The total score ranges from 0-21, with higher scores indicating poorer sleep quality[16].

Suicidal ideation

Self-rating Idea of Suicide Scale (SIOSS) is a self-report tool in Chinese with 26 questions that evaluate the suicide ideation of patients. We use the sum of three factors (despair, sleep and optimism) to assess the level of suicidal ideation. The SIOSS has been shown to have good reliability and validity[17]. A higher score reflects a higher level of suicidal ideation.

Data quality control

All the data collectors including psychiatrists and neurologists have received special training about data collection and management to ensure uniformity in criteria evaluation.

Statistical methods

Statistical analyses were performed with IBM SPSS version 26.0 software. Measurement data was expressed as Mean (standard deviation (SD)) and t test was used to compare between two groups. Count data was presented as percentages (N, %), and chi-square test was used to compare between two groups. Possible associated factors of PCS and MCS were analyzed using univariable logistic regression, stepwise multiple linear regression was performed on the factors with statistical significance in the univariate regression analyses to identify the independent factors that influenced PCS and MCS. $P < 0.05$ was considered statistically significant.

Results

Demographic and clinical characteristics

A total of 506 patients consented to participate in the study, including 109 MM patients and 397 NMM patients, with mean ages of 23.62 years (SD=7.59) and 21.48 years (SD=7.08), respectively. Table 1 presents the demographic and clinical characteristics about the participants and illustrates the differences between MM and NMM.

Health-related quality of life

Figure 1 (Figure 1: Mean scores in SF-36 for menstrual migraine (MM) patients vs. non-menstrual migraine (NMM). PF, physical functioning; RP, role-physical; BP, bodily pain; GH, general health; VT: vitality; SF: social functioning; RE: role-emotional; MH, mental health; PCS, physical component summary; MCS:

mental component summary. * $P < 0.05$) presents the mean scores of different dimensions of the HRQoL among the MM and NMM groups. MM patients had significantly lower mean scores for the five dimensions of HRQoL (PF, RF, GH, SF, MH) compared with NMM patients. The average PCS and MCS scores of MM patients were 67.82(14.08) and 67.82(14.08), respectively, which were significantly lower than those of NMM patients. Detailed data are presented in Table 1.

Table 1 Demographic and clinical data of our study sample

Variable	MM (N=109)	NMM (N=397)	P value
	Mean (SD)/ N (%)		
Age, years *	23.62 (7.59)	21.48 (7.09)	0.009
BMI	20.19 (2.50)	20.14 (2.38)	0.848
Headache frequency, days/month	3.97 (4.57)	3.83 (4.58)	0.750
PF*	89.13 (12.07)	91.52 (10.18)	0.037
RF*	59.17 (38.29)	69.40 (36.99)	0.012
BP	70.28 (14.28)	72.90 (14.95)	0.102
GH*	53.11 (17.21)	57.84 (17.74)	0.013
VT	61.24 (15.43)	62.01 (15.29)	0.641
SF*	79.68 (15.90)	93.78 (13.18)	<0.001
RE	42.20 (39.19)	46.26 (40.77)	0.353
MH*	58.80 (15.08)	62.21 (14.80)	0.034
PCS*	67.82 (14.08)	72.91 (14.25)	0.001
MCS*	60.31 (16.45)	66.07 (15.54)	0.001
Ethnicity			0.131
Han	105 (96.3)	366 (92.2)	
Minority	4 (3.7)	31 (7.8)	
Family history			0.058
Yes	78 (71.6)	245 (61.7)	
No	31 (28.4)	152 (38.3)	
Aura			0.547
Yes	17 (15.6)	53 (13.4)	
No	92 (84.4)	344 (86.6)	
Degree of headache			0.141
Mild	32 (29.4)	148 (37.3)	
Moderate	60 (55.0)	209 (52.6)	
Severe	17 (15.6)	40 (10.1)	
Headache duration			0.539

<24h	99 (90.8)	372 (93.7)
24-72h	7 (6.4)	16 (4.0)
>72h	3 (2.8)	9 (2.3)

BMI, body mass index; PF, physical functioning; RP, role-physical; BP, bodily pain; GH, general health; VT: vitality; SF: social functioning; RE: role-emotional; MH, mental health; PCS, physical component summary; MCS: mental component summary.

Univariate analysis of factors associated with PCS and MCS

Univariate analysis showed that the PCS and MCS scores were both correlated with headache frequency, anxiety and depression symptoms, suicidal ideation, sleep quality and social support, while the impact of headache on daily life was only related to PCS scores. Detailed data are presented in Table 2.

Table 2
Univariate linear of factors associated with SF-36

Variable	β	t Value	95%CI	P Value
PCS				
Headache frequency	-0.364	-0.405	(-1.671,-0.572)	<0.001
HIT-6	-0.359	-3.977	(-0.719,-0.241)	<0.001
HAMD	-0.399	-4.507	(-1.058,-0.412)	<0.001
HAMA	-0.400	-4.513	(-1.073,-0.418)	<0.001
PSSS	0.260	2.786	(0.091,0.543)	0.006
PSQI	-0.335	-3.658	(-2.279,-0.677)	<0.001
SIOSS	-0.360	-3.989	(-1.869,-0.628)	<0.001
MCS				
Headache frequency	-0.244	-2.607	(-1.547,-0.211)	0.010
HIT-6	-0.151	-1.585	(-0.533,-0.059)	0.116
HAMD	-0.600	-7.760	(-1.619,-0.960)	<0.001
HAMA	-0.459	-5.351	(-1.371,-0.630)	<0.001
PSSS	0.473	5.553	(0.433,0.914)	<0.001
PSQI	-0.306	-3.312	(2.517,-0.632)	0.001
SIOSS	-0.630	-8.382	(-3.155,-1.948)	<0.001
PCS, physical component summary; HIT-6, Headache Impact Text-6; HAMD, Hamilton Rating Scale for Depression; HAMA, Hamilton Rating Scale for Anxiety; PSSS, Perceived social support scale; PSQI, Pittsburgh Sleep Quality Index; SIOSS, Self-rating Idea of Suicide Scale.MCS: mental component summary.				

Step-wise multiple linear regression of factors associated with PCS and MCS

Headache frequency ($\beta=-0.218$, $P=0.014$), depression symptoms ($\beta=-0.345$, $P<0.001$), the impact of headache on daily life ($\beta=-0.270$, $P=0.002$) were negatively associated with PCS, depression symptoms ($\beta=-0.379$, $P<0.001$), suicidal ideation ($\beta=-0.344$, $P<0.001$) were negatively associated with MCS, while social support ($\beta=0.270$, $P<0.001$) was positively associated with MCS. Detailed data are presented in Table3.

Table 3: Step-wise multiple linear regression of factors associated with SF-36

Variable	β	t Value	95%CI	P Value
PCS				
Headache frequency	-0.218	-2.503	(-1.198,-0.139)	0.014
HIT-6	-0.270	-3.151	(-0.599,-0.136)	0.002
HAMD	-0.345	-4.164	(-0.936,-0.332)	<0.001
MCS				
HAMD	-0.379	-5.134	(-1.124,-0.498)	<0.001
PSSS	0.270	3.912	(0.190, 0.581)	<0.001
SIOSS	-0.344	-4.434	(-2.018,-0.771)	<0.001
PCS, physical component summary; HIT-6, Headache Impact Text-6; HAMD, Hamilton Rating Scale for Depression; MCS: mental component summary; PSSS, Perceived social support scale;SIOSS, Self-rating Idea of Suicide Scale.				

Disussion

This study provides, for the first time, a comprehensive assessment of the HRQoL and its associated factors in Chinese patients with MM. We found that the HRQoL in MM patients is lower in five dimensions of HRQoL (PF, RF, GH, SF, MH) compared with NMM patients. In addition, in the stepwise linear regression model, we found that PCS is related to headache frequency, depression symptoms, the impact of headache on daily life. MCS is related to depression symptoms, suicidal ideation, social support in MM patients. Such findings point the way to scientific strategies to improve HRQoL for MM patients.

Migraine increases the burden of patients[18], and seriously affects the life quality of patients[8]. Nicodemo et al. used SF-36 to evaluate the life quality of MM, and found the scores of MM in the six dimensions of HRQoL (PF, RP, BP, GH, VT, SF) were lower than those of healthy women, and there is no significant difference in the scores of the two dimensions (RE, MH)[19]. Based on the severity and refractory of MM, we performed subgroup analysis of migraine to compare the HRQoL in patients with MM and NMM. We found MM patients had lower mean scores for the five dimensions (PF, RF, GH, SF, MH) compared with NMM patients, and there were no statistically significant differences in scores for the three dimensions (BP, VT, RE). In addition, patients with MM scored lower than NMM for both PCS and MCS.

Our study demonstrated that depression symptoms are independently associated with PCS and MCS, after adjusting the confounding factors. Depression is a common comorbidity in migraine patients, and HRQoL was reduced in patients who had both migraine and depression relative to migraine patients who were not depressed[20]. A recent study in the United States found that depression symptoms are a predictor of headache frequency and migraine-related disability[21]. Hence, it is not difficult to understand that depression symptoms are closely associated with physical and mental health. In agreement with the

study finding in Brazil, we demonstrated that the severity of depression symptoms is a predictor of HRQoL among MM[22]. Pradeep et al. reported the presence of depression was noted to add to the magnitude of migraine-related disability and diminished the quality of life in migraine patients, which is similar to the results of our study[23]. This study[23] also found that anxiety had a negative impact on the quality of life of patients who suffer from migraine. In our study, although univariate analysis showed that anxiety symptoms may be a significant risk factor for HRQoL in MM patients, no significant difference was detected by multivariate analysis. At present it cannot be certain that anxiety symptoms are independently associated with HRQoL among MM patients.

MM patients who reported more frequent migraine attacks and greater impact of headaches on daily life could impair HRQoL by affecting their physical health. A downward trend in the quality of life of migraine was noted with the increased headache frequency[20]. Previous study has proved that the more frequent migraine attacks and greater impact of headaches on daily life were predictors of detrimental effects on quality of life in migraine patients[23], which is similar to our results. In a recent study, Irimia found that a positive linear association between headache frequency and the risk of anxiety, depression in migraine patients[24]. Patients with monthly headache days ≥ 3 days are at higher risk of anxiety, while those with ≥ 19 days are at risk of depression. Moreover, patients with monthly headache days ≥ 10 days are often accompanied by severe disability[24]. It is interesting to note that patients experiencing only one to six headaches per year still show a reduction in quality of life, it might be due to the unpredictability of attacks that magnifies the effect of the few headache days on quality of life in a remarkable way[20]. Richard et al[25] examined the association between headache-free days and the disease burden of migraine, and found headache-related disability shows a decrease tendency with the headache-free days increasing[25].

Our research showed that the perception of social support is positively associated with MCS. An Italian study of chronic migraine patients with medication overuse found that social support is a predictor of the quality of life to some extent[26]. Moreover, a French study also found that the higher perceived social support was, the higher the probability of being an active consulter for migraine[27, 28]. Such headache counseling allowed patients to actively take prevention and treatment strategies to minimize the burden of migraine and relieve adverse emotions, and thus engaged more in social activities and improved HRQoL, especially in mental health related HRQoL.

As the limitations of daily social and work-related activities caused by migraine, the mental health of patients was impacted, and severe cases may lead to suicidal ideation. Our research found that suicidal ideation can affect MCS and could consequently predict HRQoL of MM patients. Many scholars agreed that migraine patients are associated with a poor quality of life and a higher likelihood of suicidal ideation[8, 9, 29, 30]. This study demonstrated, for the first time to our knowledge, the correlation between HRQoL and suicidal ideation among MM patients. Providing psychological treatment to MM patients with suicidal ideation may help to reduce suicide risk, as can the application of active treatment, improving their mental health and HRQoL.

Sleep disturbance, as a clinical triggers for migraine, is associated with migraine, creating a vicious cycle[31]. Although sleep quality is associated with the PCS and MCS of patients with MM in the univariate analysis, it is no longer a significant associated factor after adjusting for the potential confounders in the multivariate model. Thus, we cannot consider sleep quality as a predictor for HRQoL in MM patients. In the current study, the evidence describing the relationship between sleep quality and HRQoL was still insufficient among migraine patients[32],and this result was in line with the previous study.

Conclusion

Considering HRQoL by migraine subtype, we found that MM patients had a worse HRQoL with statistically significantly lower mean scores in many domains than NMM patients. Depression symptoms are closely linked to physical and mental health in MM patients. For MM patients, PCS is related to the headache frequency and the impact of headaches on daily life, and MCS is related to the social support and suicidal ideation. Moreover, sleep quality did not have a significant impact on the HRQoL among MM patients. The early application of individualized treatment may help improve the HRQoL in patients with MM.

There were some limitations to this study. The sample representativeness was limited because the sample was derived from the outpatient population of a hospital in southwest China. A multicenter study would be needed with a larger sample size and more associated factors to determine the factors that had an association with HRQoL of MM patients.

List Of Abbreviations

HRQoL: Health-related quality of life; MM: Menstrual migraine; NMM: non-menstrual migraine; SIOSS: Self-rating Idea of Suicide Scale; HAMD: Hamilton Depression Scale; HAMA: Hamilton Anxiety Scale; HIT-6: Headache Impact Test-6; PSSS: Perceived Social Support Scale; PSQI: Pittsburgh Sleep Quality Index; SF-36: 36-Item Short Form Survey; ICHD-3: International Classification of Headache Disorders, Third Edition; BMI: Body mass index; PF: Physical functioning; RP: Role-physical; BP: Bodily pain; GH:General health; VT:Vitality; SF:Social functioning; RE: Role-emotional; MH: Mental health; PCS: Physical component summary; MCS: Mental component summary; SD: Standard deviation.

Declarations

Ethics approval and consent to participate

Approval was obtained from the ethics committee of Affiliated Hospital of North Sichuan Medical College.The procedures used in this study adhere to the tenets of the Declaration of Helsinki. An informed consent was obtained from each participant.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used or analysed during the current study are available from the corresponding author on reasonable request.

Conflicts of interest

The authors declare that they have no competing interests

Funding

The research received funding from the Bureau of Science and Technology and Intellectual Property of Nanchong City (No. NSMC20170420).

Authors' contributions

Material preparation, data collection and analysis were performed by Wenxiu Luo, Xing Cao and Jiaming Luo. The first draft of the manuscript was written by Wenxiu Luo and Xing Cao, and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript. All authors certify responsibility of the paper.

Acknowledgements

We thank all the patients for their support in this research.

References

1. Disease GBD, Injury I, Prevalence C: Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017;**390**:1211–1259. doi: 10.1016/S0140-6736(17)32154-2.
2. Headache Classification Committee of the International Headache Society (IHS) The International Classification of Headache Disorders, 3rd edition. *Cephalalgia*. 2018;**38**:1-211. doi: 10.1177/0333102417738202.
3. Broner SW, Bobker S, Klebanoff L: Migraine in Women. *Semin Neurol*. 2017;**37**:601–610. doi: 10.1055/s-0037-1607393.
4. Pavlovic JM, Stewart WF, Bruce CA, Gorman JA, Sun H, Buse DC, Lipton RB: **Burden of migraine related to menses: results from the AMPP study**. *J Headache Pain*. 2015;**16**:24. doi: 10.1186/s10194-015-0503-y.
5. MacGregor EA, Victor TW, Hu X, Xiang Q, Puenpatom RA, Chen W, Campbell JC: **Characteristics of menstrual vs nonmenstrual migraine: a post hoc, within-woman analysis of the usual-care phase of**

- a nonrandomized menstrual migraine clinical trial. *Headache*. 2010;**50**:528–538. doi: 10.1111/j.1526-4610.2010.01625.x.
6. Goh KKK, Lai PSM, Lim SK: **Cross cultural adaptation and validation of the Malay Kidney Disease Quality of Life (KDQOL-36)**. *BMC Nephrol*. 2019;**20**:226. doi: 10.1186/s12882-019-1397-8.
 7. Liu S, Wang X, Xu Y, Xu Q, Zhi Y: **Health-related quality of life and its risk factors in Chinese hereditary angioedema patients**. *Orphanet J Rare Dis*. 2019;**14**:191. doi: 10.1186/s13023-019-1159-5.
 8. Peters GL: **Migraine overview and summary of current and emerging treatment options**. *Am J Manag Care*. 2019;**25**:S23-S34.
 9. Silberstein SD: Migraine. *Lancet*. 2004;**363**:381–391. doi: 10.1016/S0140-6736(04)15440-8.
 10. Li L, Wang HM, Shen Y: **Chinese SF-36 Health Survey: translation, cultural adaptation, validation, and normalisation**. *J Epidemiol Community Health*. 2003;**57**:259–263. doi: 10.1136/jech.57.4.259.
 11. Wang R, Wu C, Zhao Y, Yan X, Ma X, Wu M, Liu W, Gu Z, Zhao J, He J: **Health related quality of life measured by SF-36: a population-based study in Shanghai, China**. *BMC Public Health*. 2008;**8**:292. doi: 10.1186/1471-2458-8-292.
 12. Houts CR, Wirth RJ, McGinley JS, Cady R, Lipton RB: **Determining Thresholds for Meaningful Change for the Headache Impact Test (HIT-6) Total and Item-Specific Scores in Chronic Migraine**. *Headache*. 2020;**60**:2003–2013. doi: 10.1111/head.13946.
 13. Zimmerman M, Martinez JH, Young D, Chelminski I, Dalrymple K: **Severity classification on the Hamilton Depression Rating Scale**. *J Affect Disord*. 2013;**150**:384–388. doi: 10.1016/j.jad.2013.04.028.
 14. Bhamre R, Sawrav S, Adarkar S, Sakaria R, S JB: **Psychiatric comorbidities in patients with inflammatory bowel disease**. *Indian J Gastroenterol*. 2018;**37**:307–312. doi: 10.1007/s12664-018-0870-9.
 15. Li Y, Long Z, Cao D, Cao F: **Social support and depression across the perinatal period: A longitudinal study**. *J Clin Nurs*. 2017;**26**:2776–2783. doi: 10.1111/jocn.13817.
 16. Huang Y, Zhu M: **Increased Global PSQI Score Is Associated with Depressive Symptoms in an Adult Population from the United States**. *Nat Sci Sleep*. 2020;**12**:487–495. doi: 10.2147/NSS.S256625.
 17. Xie P, Wu K, Zheng Y, Guo Y, Yang Y, He J, Ding Y, Peng H: **Prevalence of childhood trauma and correlations between childhood trauma, suicidal ideation, and social support in patients with depression, bipolar disorder, and schizophrenia in southern China**. *J Affect Disord*. 2018;**228**:41–48. doi: 10.1016/j.jad.2017.11.011.
 18. Leonardi M, Raggi A: **A narrative review on the burden of migraine: when the burden is the impact on people's life**. *J Headache Pain*. 2019;**20**:41. doi: 10.1186/s10194-019-0993-0.
 19. Nicodemo M, Vignatelli L, Grimaldi D, Sancisi E, Fares JE, Zanigni S, Pierangeli G, Cortelli P, Montagna P, Cevoli S: **Quality of life, eating and mood disorders in menstrual migraine: a case-control study**. *Neurol Sci*. 2008;**29 Suppl 1**:S155-157. doi: 10.1007/s10072-008-0911-8.

20. Becker WJ: **Assessing health-related quality of life in patients with migraine.** *Can J Neurol Sci.* 2002;**29 Suppl 2**:S16-22. doi: 10.1017/s031716710000189x.
21. Rettig EK, Ergun G, Warfield JR, Slater SK, LeCates SL, Kabbouche MA, Kacperski J, Hershey AD, Powers SW: **Predictors of Improvement in Pediatric Chronic Migraine: Results from the Cognitive-Behavioral Therapy and Amitriptyline Trial.** *J Clin Psychol Med Settings.* 2021. doi: 10.1007/s10880-021-09782-4.
22. Vieira RV, Vieira DC, Gomes WB, Gauer G: **Alexithymia and its impact on quality of life in a group of Brazilian women with migraine without aura.** *J Headache Pain.* 2013;**14**:18. doi: 10.1186/1129-2377-14-18.
23. R P, S CN, S H, K R: **Migraine Disability, Quality of Life, and Its Predictors.** *Ann Neurosci.* 2020;**27**:18–23. doi: 10.1177/0972753120929563.
24. Irimia P, Garrido-Cumbrera M, Santos-Lasaosa S, Aguirre-Vazquez M, Correa-Fernandez J, Colomina I, Pozo-Rosich P: **Impact of monthly headache days on anxiety, depression and disability in migraine patients: results from the Spanish Atlas.** *Sci Rep.* 2021;**11**:8286. doi: 10.1038/s41598-021-87352-2.
25. Lipton RB, Lee L, Saikali NP, Bell J, Cohen JM: **Effect of Headache-Free Days on Disability, Productivity, Quality of Life, and Costs Among Individuals with Migraine.** *J Manag Care Spec Pharm.* 2020;**26**:1344–1352. doi: 10.18553/jmcp.2020.20103.
26. D'Amico D, Grazzi L, Bussone G, Curone M, Di Fiore P, Usai S, Leonardi M, Giovannetti AM, Schiavolin S, Raggi A: **Are depressive symptomatology, self-efficacy, and perceived social support related to disability and quality of life in patients with chronic migraine associated to medication overuse? Data from a cross-sectional study.** *Headache.* 2015;**55**:636–645. doi: 10.1111/head.12534.
27. Lanteri-Minet M, Massiou H, Nachit-Ouinekh F, Lucas C, Pradalier A, Radat F, Mercier F, El Hasnaoui A: **The GRIM2005 study of migraine consultation in France I. Determinants of consultation for migraine headache in France.** *Cephalalgia.* 2007;**27**:1386–1397. doi: 10.1111/j.1468-2982.2007.01426.x.
28. Radat F, Lanteri-Minet M, Nachit-Ouinekh F, Massiou H, Lucas C, Pradalier A, Mercier F, El Hasnaoui A: **The GRIM2005 study of migraine consultation in France. III: Psychological features of subjects with migraine.** *Cephalalgia.* 2009;**29**:338–350. doi: 10.1111/j.1468-2982.2008.01718.x.
29. Ehi Y, Sahin L, Mutlu MF: **Migraine management in pregnancy.** *Clin Exp Obstet Gynecol.* 2016;**43**:661–665.
30. Pei JH, Wang XL, Yu Y, Zhang YB, Gou L, Nan RL, Chen HX, Dou XM, Han ZJ: **Prevalence of suicidal ideation and suicide attempt in patients with migraine: A systematic review and meta-analysis.** *J Affect Disord.* 2020;**277**:253–259. doi: 10.1016/j.jad.2020.08.019.
31. Ferini-Strambi L, Galbiati A, Combi R: **Sleep disorder-related headaches.** *Neurol Sci.* 2019;**40**:107–113. doi: 10.1007/s10072-019-03837-z.
32. Wang, XM, Zhang YH, Shao GF. **Analysis of the factors affecting migraine patients' quality of life. (article in Chinese)** *Journal of Neuroscience and Mental Health.* 2018;**18**:28–31.

Figures

Mean scores in SF-36 for MM vs NMM




Figure 1

Mean scores in SF-36 for menstrual migraine (MM) patients vs. non-menstrual migraine (NMM). PF, physical functioning; RP, role-physical; BP, bodily pain; GH, general health; VT: vitality; SF: social functioning; RE: role-emotional; MH, mental health; PCS, physical component summary; MCS: mental component summary. * P<0.05