

Stunting and Its Associated Factors Among Children Age 6-59 Months at Debre Tabor Town, Northwest Ethiopia, 2019; a Community Based Cross-sectional Study

Bisrat Dessie Getu (✉ edombsr10@gmail.com)

Debre Tabor Health science college

Kendalem Asmare Atalell

University of Gondar

Destaye Guadie Kassie

University of Gondar

Kefyalew Amogne Azanaw

Debre Tabor Health Science College

Nigusie Selomon Tibebu

Debre Tabor University

Research

Keywords: Children, Stunting, Debre Tabor town, Ethiopia

Posted Date: February 9th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1055501/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Introduction: Globally in U5 year children, greater than half of all deaths are attributable to under-nutrition. Stunting is one of the major under-nutrition problems in children, especially in developing countries like Ethiopia. Therefore, this study was aimed to assess the prevalence of stunting and its associated factors among children age 6-59 months at Debre tabor Town, Northwest Ethiopia, 2019.

Methods: Community-based cross-sectional study design was conducted among children age 6-59 months at Debre Tabor town, from February to March 2019. Data were collected using a pretested, semi-structured an interviewer-administered questionnaire adapted from WHO stepwise surveillance (WHO STEPS). A systematic random sampling technique was used to engage 620 parent-child pairs. Data was entered into Epi-info version 7 and exported to SPSS version 22 for analysis. Height for age Z score was computed using WHO Anthro plus software. Both the bivariable and a multivariable logistic regression analyses were computed and adjusted odds ratio (AOR) at p-value <0.05 was used to determine the statistically significant association between factors and the outcome variable.

Result: The prevalence of stunting in this study was 46% (95% CI: 41.9%-50.3%). Age of the child [AOR=1.98, 95%CI: (1.08, 3.67)], occupation of the mother [AOR=2.13, 95% CI: (1.16, 3.92)], educational status of the father [AOR=2.88, 95% CI: (1.45, 5.70)] and occupation of the father [AOR=5.05, 95% CI: (2.46, 10.36)] were variables significantly associated with stunting.

Conclusion: The prevalence of stunting was found to be high among children age 6-59 months in the study area as compared to the national average. The age of the child, occupation of the mother, the educational status of the father, and occupation of the father were factors associated with stunting? Therefore, attention should be targeted to improve the occupational and educational status of the parents.

Introduction

Globally, in under five children greater than half of all deaths are attributable to under-nutrition [1]. Besides, stunting is one of the most common markers of under-nutrition, which is a linear growth failure or inability to gain a potential height for a particular age [2, 3].

According to UNICEF, and World Bank 2019 reports, more than one in five children under the age of 5 years had stunted growth [1]. Even though stunting decreases globally from in the last decades, it is increased in alarming rates in Africa[2, 4]. Furthermore, in Ethiopia, according to the Ethiopian Demographic health survey (EDHS) 2016, stunting was 38% [5].

According to WHO conceptual frame work on childhood stunting, the household and family incomes, breastfeeding techniques of children, infection, and complementary feeding practices were some of the contributing factors of stunting [6]. The global review of stunting in low- and middle-income countries also identified that growth restriction in-utero and lack of access to sanitation as the main driving factors

for stunting [7]. Consequently, stunting in children results in susceptibility to infectious disease, diminish intellectual ability, and poor school performance [4, 8-10]. Moreover, stunted children may never regain their height loss again in their lifetime [11].

Despite the global burden of stunting, it is not often recognize in the communities, especially in low and middle-income countries like Ethiopia [12, 13]. Even though, stunting in Ethiopia, reduced from 58% according to Ethiopian Demographic Health Survey (EDHS) 2000 [14], to 38% EDHS 2016 [5], in under-five children. However, this progress is not sufficient to meet the global target [15]. The risk factors of stunting in a different setting is different for under-five children [16]. Thus, to address this gap, this study was aimed to assess the prevalence of stunting and its associated factors among children age 6-59 months at Debre tabor Town, Northwest Ethiopia, 2019.

Methods

Study design and setting

Community-based cross-sectional study design was conducted to assess stunting among children age 6-59 months at Debre tabor town, Northwest Ethiopia, from February to March 2019.

The study was conducted at Debre tabor town, which is found in the Northwest part of Ethiopia. The general population of Debre tabor town is estimated to be 84,382 of this 10, 868 were children, age 6-59 months [17].

Study participants and sampling

All children age 6-59 months living at Debre Tabor town were eligible for this study. The town has six kebeles; each kebele was considered as a cluster. Then, three clusters (50% of the cluster) were selected randomly by the lottery method. Households with children age 6-59 months in the three clusters were obtained from health extension workers in each cluster. Using health extension workers registrations as a frame of reference studied households were selected by systematic random sampling technique. For more than one eligible child in one household, the lottery method was used.

Dependent Variable

- **Stunting** (below -2 SD) children age from 6-59 months

Independent Variables

- Socio- Demographic, and Economic Characteristics
- Maternal Characteristics
- Childs' Characteristics
- Environmental Characteristics

Operational Definition

- **Stunting:** A child is defined as stunted if the height-for-age Z-score is found to be below -2 SD of the median of the WHO standard curve.

Data collection tools and procedures

Data were collected by using a pretested, semi-structured interviewer-administered questionnaire, which is adopted from WHO stepwise surveillance (WHO STEPS) for child malnutrition. The questionnaire comprised of socio-demographic, economic, environmental conditions, and child health-related characteristics.

The data were collected by five BSc nurses. Mothers or caregivers were interviewed whereas; anthropometry measurements were taken from children. The height of infants aged six to 23 months was measured in a recumbent position to the nearest 0.1 cm, using a board with an upright wooden base and movable headpieces. Children aged 24 to 59 months were measured in a standing up position to the nearest 0.1 cm. additionally, child weight was measured by an electronic digital weight scale to the nearest 0.1 kg. The calipers of the measurements were checked before and after the measurement of the individual child.

Data processing and analysis

Data were entered into EPI-info version 7 and then exported to SPSS version 12 for analysis. WHO Anthro-Plus software was used to convert nutritional data into Z scores of the indices HAZ using the WHO standard. In addition, principal component analysis (PCA) was used to compute family wealth status.

Both bivariate and multivariable binary logistic regression models were computed to identify factors associated with stunting among children age 6-59 months. A bivariate binary logistic regression model was first computed, and variables less than 0.2 in the bivariate analysis were entered into the multivariable logistic regression model. In the final multivariable analysis, variables having a p-value less than 0.05 at 95% CI were considered as significantly associated with stunting.

Result

Socio-demographic and economic characteristics of the study participants

A total of 620 children age 6-59 months were included in the analysis. The majority (91.0%) of the mothers/caregivers were married and (90.6%) Orthodox Christian in Religion. Nearly three fourth (72.4%) of the fathers had high school and above educational status (Table1).

Table 1
Socio-demographic characteristics of children age 6-59 months at Debre tabor town,
Northwest Ethiopia, 2019 (n=620).

Characteristics		Frequency	Percentage
Marital status	Married	564	91.0
	Single	56	9.0
Religion	Orthodox	562	90.6
	Muslim	44	7.1
	Protestant	14	2.3
Ethnicity	Amhara	601	96.9
	Oromo	17	2.7
	Gurage	2	0.3
The educational status of the mother	No formal education	127	20.5
	Primary education	143	23.1
	Highschooland above	350	56.5
The educational status of the father	No formal education	97	15.6
	Primary education	74	11.9
	Highschooland above	449	72.4
Occupation of mother	Housewife	284	46.5
	Daily laborer	41	6.7
	Private organization	21	3.4
	Merchant	95	15.5
	Government employee	170	27.8
Occupation of father	Daily laborer	95	15.5
	Private organization	96	15.7
	Merchant	158	25.8
	Government employee	264	43.1
Family size	Less than 5	503	81.1
	Greater than or equal to 5	117	18.9
Totalnumberof	≤2	466	75.2

under-five children		Frequency	Percentage
Characteristics			
	>2	154	24.8
Total number of children	Less than four	576	92.9
	Greater than or equal to four	44	7.1
Wealth status of the family	Poor	204	32.9
	Medium	208	33.5
	Rich	208	33.5

Child characteristics

The mean age of the study participants was 27 (± 14 SD) months. Slightly higher than half (52.3%) of the children were males. Six hundred twelve (98.7%) of children were ever immunized (Table 2).

Table 2
Child characteristics of children age 6-59 months at Debre tabor town, Northwest Ethiopia, 2019(n=620).

Characteristics		Frequency	Percentage
Sex of the child	Male	324	52.3
	Female	296	47.7
Age of the child 27 ±14 SD	6-11 months	110	17.7
	12-23 months	146	23.5
	24-35months	159	25.6
	36- 47 months	138	22.3
	48-59 moths	67	10.8
Ever breastfeeding	Yes	615	99.0
	No	5	1.1
Ever immunized	Yes	612	98.7
	No	8	1.3
Breastfeeding initiations time	Before 1 hour	500	81.0
	Greater than 1 hours	120	19.0
Pre-lacteal feeding use	Yes	47	7.6
	No	573	92.4
Types of pre-lacteal feeding	Water	16	2.6
	Butter	14	2.3
	Milk	15	2.4
	Others (Habish, meqamesha)	2	0.3
SD: Standard Deviation			

Maternal characteristics

Five hundred ninety-one (95.3%) of the mothers/caregivers utilize family planning methods. Nearly three fourth (72.6%) of the mothers were utilized Depo-Provera (Tabe3).

Table 3
Maternal characteristics of children age 6-59 months at Debre tabor
Town, Northwest Ethiopia 2019 (n=620).

Characteristics		Frequency	Percentage
Age of the mothers/caregivers	<25	69	11.1
	25-30	303	48.9
	>30	248	40.0
Family planning utilizations	Yes	591	95.3
	No	29	4.7
Pills	Yes	119	19.2
	No	501	80.8
Depo-Provera	Yes	450	72.6
	No	170	27.4
Norplant	Yes	104	16.8
	No	516	83.2
Is the pregnancy planned	Yes	548	88.4
	No	72	11.6

Environmental characteristics

Five hundred and fifty-six (89.7%) of the respondents use private tap water as their main water source. The majority of the respondents 615 (99.2%) have latrine.

Eighty-five (13.7%) of the respondents don't have a separate kitchen for their food preparations (Table 4).

Table 4

Environmental characteristics of children age 6-59 months at Debre tabor town, Northwest Ethiopia, 2019 (n=620).

Characteristics		Frequency	Percentage
Source of drinking water			
Private well	Yes	129	20.8
	No	491	79.2
Private tap	Yes	556	89.7
	No	64	10.3
Public tap	Yes	26	4.2
	No	594	95.8
Spring water	Yes	92	14.8
	No	528	85.2
Amount of water used in the household	<50 liters	332	53.5
	50-75 liters	282	45.5
	>75litrs	6	1.0
Water treatment utilizations	Yes	98	15.8
	No	522	84.2
Latrine	Yes	615	99.2
	No	5	0.8
Type of latrine	Private pit/wooden slab	503	81.2
	Privateslab/cement Slab	77	12.5
	Sharedlatrine/wooden Slab	7	1.2
	Shared VIP latrine	13	2.1
	Others like an open field	20	3.0
Waste disposal	Open field	190	30.6
	Private pit	49	7.9
	Common pit	21	3.4
	Composting	10	1.6
	Burning	302	48.7

Characteristics		Frequency	Percentage
	Other	48	7.7
Kitchen	Yes	535	86.3
	No	85	13.7

Prevalence of stunting among children age 6-59 months

The overall prevalence of stunting among children age 6-59 months at Debre tabor town was 46% at 95% CI (41.9-50.3) (Figure (1)).

Factors associated with stunting among children age 6-59 months

In the bivariate logistic regression analysis, age of the child, educational level of the mother, educational level of the father, occupation of the father, occupation of the mother, total number of children, family size, family planning use, and family wealthy status were significantly associated with stunting. However, in the multivariable logistic regression analysis; the age of the child, the educational status of the father, occupation of the father, and occupation of the mother remained significantly associated with stunting.

The odds of having stunting among children with the age group of 12-23 months were nearly 2 times higher as compared to children with age groups of 6-11 months age (AOR=1.98; 95%CI: 1.08,3.67). The odds of having stunting among children who had a daily laborer fathers were 5 (AOR= 5.03; 95%CI: 2.45,10.36) times and children with father who works in private organizations were 2.4 (AOR=2.36; 95%CI: 1.29, 4.33) times higher as compared with children whose fathers were governmental employees respectively. Similarly, the odds of having stunting among children who had a father with no formal education were nearly 3 (AOR=2.88; 95%CI: 1.45, 5.70) times and children with fathers who attend primary school were 2 (AOR=1.98; 95%CI: 1.03, 3.79) times higher as compared to children with fathers who attended high school and above educational level respectively. The odds of having stunting among children who had housewife mothers were 2 times higher than children of their mothers who were a governmental employee (AOR=2.13; 95%CI: 1.16, 3.92) (Table 5).

Table 5

Factors associated with stunting among children age 6-59 months at Debre tabor Town Northwest, Ethiopia, 2019 (n=620).

Variable		Stunted			
		Yes	No	COR at 95% CI	AOR at 95% CI
Age	6-11 months	35	75	1	1
	12-23 months	70	76	1.97(1.18,3.31)	1.98(1.08,3.67)*
	24-35 months	73	86	1.81(1.09,3.02)	1.54(0.84,2.82)
	35-47 months	69	69	2.14(1.27,3.61)	1.45(0.78,2.71)
	48-59 months	38	29	2.81(1.49,5.26)	1.45(0.68,3.10)
Maternal education	No formal education	85	42	4.25(2.76,6.54)	0.89(0.46,1.73)
	Primary education	87	56	3.26(2.18,4.88)	1.22(0.73,2.06)
	High school and above	113	237	1	1
Educational status of the father	No formal education	71	26	4.84(2.97,7.89)	2.88(1.45,5.70)**
	Primary education	52	22	4.19(2.45,7.15)	1.98(1.03,3.79)*
	High school and above	162	287	1	1
Occupation of the mothers	Housewife	169	115	5.88(3.77,9.17)	2.13(1.16,3.92)*
	Daily laborer	28	13	8.62(4.04,18.38)	1.47(0.57,3.81)
	Private organizations	6	15	1.60(0.58,4.43)	0.65(0.19,2.10)
	Merchant	43	52	3.31(1.91,5.74)	1.3(0.68,2.67)
	Gov't employee	34	136	1	1
Father occupation	Daily laborer	76	19	13.03(7.31,23.2)	5.05(2.46,10.36)*
	Private organization	56	40	4.56(2.78,7.49)	2.36(1.29,4.33)**
	Merchant	86	72	3.89(2.55,5.94)	1.71(0.99,2.96)
	Gov't employee	62	202	1	1
Total number of family	Less than 5	214	289	1	1

1=reference, *(p<0.05) significant, ** (P<0.001) highly significant

	Greater than or equal to 5	71	46	2.08(1.38,3.14)	0.57(0.32,1.01)
Total number of children	Less than four	258	318	1	1
	Greater than or equal to four	27	17	1.96(1.04,3.67)	0.65(0.27,1.58)
Total number of under-five children	≤ 2	206	260	1	1
	>2	79	75	0.75(0.52,1.04)	1.13(0.72,1.78)
Current family planning use	Yes	202	268	1	1
	No	83	67	1.64(1.34,2.4)	1.52(0.95,2.42)
Family wealth	Poor	125	79	3.05(2.04,4.56)	1.38(0.84,2.26)
	Medium	89	119	1.44(0.97,2.15)	1.11(0.70,1.77)
	Rich	71	137	1	1
1=reference, *(p<0.05) significant, ** (P<0.001) highly significant					

Discussion

Stunting is a major public health problem globally. Therefore, the aim of this study was to determine the prevalence of stunting and its associated factors among children age 6-59 months in Debre Tabor town, Northwest Ethiopia.

The overall prevalence of stunting in this study was 46% (95%CI; 41.9, 50.3), which is in line with a cross-sectional study conducted in Labella Town, Ethiopia 47.3% [18], Wukro town, Tigray region, Ethiopia 49.2% [19], Arba Minch Health and Demographic Surveillance Site (HDSS), Ethiopia 47.9% [20], Pastoral Communities of Afar Regional State, Ethiopia 43.1% [21], Amhara regional state, Ethiopia 46.3% [22] and Bule Hora, Ethiopia 47.6% [23].

However, the finding in this study is higher than the national average reported by EDHS 2016 38% [5] and other studies conducted at West Gojjam, Ethiopia 24.9% [24], Aykel town, Northwest Ethiopia 28.4% [25], Sodo Zuria District, South Ethiopia 24.9% [26], Evidence from 2016 Demographic and Health Survey, Ethiopia 38.3% [27], Bishoftu Town, Oromia Region, Ethiopia 16.1% [28], Damot Gale district, Southern Ethiopia 41.7% [29].

This might be due to dietary diversity differences in different areas of Ethiopia. Ethiopia is a multicultural country, which had different child feeding cultures [30, 31]

The result in our study is also higher than study conducted in Zambia 40% [32], Nigeria 38.7% [33], Southwestern Nigeria 18.6% [8], Thailand 28.8% [34], and china 8.4% [35]. This might be directly related to the difference in the economic status of the countries.

The underline cause of stunting in children is poverty and lack of educations. Therefore, children lived-in low-income countries like Ethiopia lead to the inadequacy of foods, there is also a problem with the preparations of clean and diversified foods, which directly may cause stunting [36].

On the other hand, the prevalence of stunting in this study was lower than the study done in Belesa, Ethiopia 57.7% [37], and Kayin State, Myanmar 59.4% [36]. The higher result observed in Belesa might be due to the fact that the Belesa district is mostly affected by drought, and the study was conducted among rural households [38, 39].

The odds of having stunting among children with the age group of 12-23 months were nearly 2 times higher as compared to children with age groups of 6-11 months age (AOR=1.98; 95%CI: 1.08,3.67), which is supported by a study conducted in Labella Town, Ethiopia [18], Damot Gale district, southern Ethiopia 41.7% [29], SodoZuria District, South Ethiopia 24.9% [26] and Zambia [32].

This is due to the fact that, in most studies, breastfeeding is universal in Ethiopia, and mostly it continues up to 1 year of age. As we all now breastfeeding greatly reduces the occurrence of stunting. Therefore, after one year of age when most children cease breastfeeding, they might be exposed to stunting.

The other possible explanation might be, in most cases, children are exposed to family food at the age of one to two years. This might again expose the child to stunting [32]. Researchers suggested that children accumulate growth delay during the first 2 years of life with stunting peaking around 2 and 3 years after which they stabilize [40].

The odds of having stunting among children who had a daily laborer father were 5 (AOR= 5.03; 95%CI: 2.45,10.36) times and children with a father who works in private organizations were 2.4 (AOR=2.36; 95%CI: 1.29, 4.33) times higher as compared with children whose fathers were governmental employees respectively. This might be related to the family income and feeding status of the children, children with government employee fathers would have better income as compared to their counterparts.

The odds of having stunting among children who had a father with no formal education were nearly 3 (AOR=2.88; 95%CI: (1.45-5.70) times and children with fathers who attend primary school were 2 (AOR=1.98; 95%CI: (1.03-3.79) times higher as compared to children with fathers who attended high school and above educational level respectively.

This is supported by a study conducted at West Gojjam, Ethiopia 24.9% [24].

The possible reason might be educated fathers had better awareness regarding the kinds and amounts of food appropriate for their children. As the level of education increase, the knowledge of fathers towards child feeding practices, common childhood disorders, child nutritional problems will increase, therefore, stunting will be decreased.

The odds of having stunting among children who had housewife mothers were 2 times higher than children of their mothers who were a governmental employee (AOR=2.13; 95%CI: 1.16, 3.92). This might

be due to related to the income and educational level of the mothers. If the mothers were employed, they should have formal education which implies they will have better awareness regarding child feeding practice. On the other hand, if the mothers were employed, they will have a good income as compared to housewife mothers.

Limitation of the study

Even though we tried to minimize, there might be a sort of recall bias. Since the study is on an interview base. Furthermore, the study is cross-sectional it does not show a cause-effect relationship between outcome and associated factors

Conclusion

The prevalence of stunting was found to be higher among children age 6-59 months in Debre tabor town as compared to the national average. The age of the child, the educational status of the father, occupation of the father, and occupation of the mother were variables significantly associated with stunting in this study. Therefore, attention should be targeted to improve the occupational status and the educational levels of the parents.

Declarations

Ethical Approval and Consent to Participate

Ethical clearance was obtained from the school of the nursing ethical review committee on behalf of the University of Gondar review board. The verbal informed consent was acceptable and approved by Ethical review board on the behalf of University of Gondar. A permission letter was obtained from the Debre Tabor town department of health. Participants were informed about voluntarism and that they can withdraw at any time of the study if they want not respond. For those who were a volunteer to participate, verbal informed consent was obtained from the mother/caregivers/ for the children involved in this study. At the end of the interview, participants were informed about stunting and associated potential effects.

Competing Interest

The authors report no conflicts of interest in this work.

Authors' contribution

BD and KA wrote the proposal, participated in data collections, analyzed the data, and draft the paper. DG, MC, DA, KA and NS approves the proposal with some revisions, participated in data collections and analysis. All authors were participated in the preparation of the manuscript and approve the final manuscript.

Consent for publication

All authors were taken consent to participate in a study and having their data published in a journal article. Authors were give verbal consent to publish their data prior to submitting their paper to a journal.

Acknowledgments

The authors would like to thank the University of Gondar for financial support that made this study possible. They would also like to thank the data collectors for their tolerance and collaboration during the work.

Funding statement

This research didn't receive any grant from any funding agency in the public, commercial, or not-for-profit sectors.

Data availability statement

Data will be available upon request from the corresponding author.

References

1. UNICEF. Malnutrition prevalence remains alarming: stunting is declining too slowly while wasting still impacts the lives of far too many young children. 2020.
2. Demissie S, Worku AJSJPH. Magnitude and factors associated with malnutrition in children 6-59 months of age in the pastoral community of Dollo Ado district, Somali region, Ethiopia. 2013;1(4):175-83.
3. Reinhardt K, Fanzo J. Addressing chronic malnutrition through multi-sectoral, sustainable approaches: a review of the causes and consequences. *Frontiers in nutrition*. 2014;1:13.
4. WonDiMAgegn ZTJCRiN, Journal FS. Magnitude and determinants of stunting among children in Africa: a systematic review. 2014;2(2):88-93.
5. [Ethiopia] CSA. Ethiopia Demographic and Health Survey (EDHS); 2016 2016.
6. Stewart C, Iannotti L, Dewey K, Michaelsen K, Onyango A. WHO conceptual framework on Childhood Stunting: Context, Causes, and Consequences. Geneva: World Health Organization. 2013.
7. Danaei G, Andrews KG, Sudfeld CR, Fink G, McCoy DC, Peet E, et al. Risk Factors for Childhood Stunting in 137 Developing Countries: A Comparative Risk Assessment Analysis at Global, Regional, and Country Levels. *PLoS Med*. 2016;13(11):e1002164-e.
8. Olodu MD, Adeyemi AG, Olowookere SA, Esimai OA. Nutritional status of under-five children born to teenage mothers in an urban setting, south-western Nigeria. *BMC Research Notes*. 2019;12(1):116.
9. Asmare B, Taddele M, Berihun S, Wagnev F. Nutritional status and correlation with academic performance among primary school children, northwest Ethiopia. *BMC research notes*. 2018;11(1):805.

10. Seyoum D, Tsegaye R, Tesfaye A. Undernutrition as a predictor of poor academic performance; the case of Nekemte primary school students, Western Ethiopia. *BMC research notes*. 2019;12(1):727.
11. Caulfield LE, Richard SA, Rivera JA, Musgrove P, Black RE. Stunting, wasting, and micronutrient deficiency disorders. *Disease Control Priorities in Developing Countries 2nd edition: The International Bank for Reconstruction and Development/The World Bank*; 2006.
12. De Onis M, Branca F. Childhood stunting: a global perspective. *Maternal & child nutrition*. 2016;12:12-26.
13. Bloss E, Wainaina F, Bailey RCJJotp. Prevalence and predictors of underweight, stunting and wasting among children aged 5 and under in western Kenya. 2004;50(5):260-70.
14. [Ethiopia] CSA. Ethiopia Demographic and Health Survey (EDHS); 2000. Central Statistical Agency [Ethiopia] 2000.
15. De Onis M, Dewey KG, Borghi E, Onyango AW, Blössner M, Daelmans B, et al. The World Health Organization's global target for reducing childhood stunting by 2025: rationale and proposed actions. 2013;9:6-26.
16. Medhin G, Hanlon C, Dewey M, Alem A, Tesfaye F, Worku B, et al. Prevalence and predictors of undernutrition among infants aged six and twelve months in Butajira, Ethiopia: the P-MaMiE Birth Cohort. *BMC public health*. 2010;10(1):27.
17. office Zh. south Gondar zone population statistics. 2018.
18. Yalew B, Amsalu F, Bikes DJJNDT. Prevalence and factors associated with stunting, underweight, and wasting: A community-based cross-sectional study among children age 6-59 months at Lalibela Town, Northern Ethiopia. 2014;4(147):2161-0509.1000147.
19. Gebru TT, Tesfamichael YA, Bitow MT, Assefa NE, Abady GG, Mengesha MB, et al. Stunting and associated factors among under-five children in Wukro town, Tigray region, Ethiopia: a cross-sectional study. *BMC research notes*. 2019;12(1):504.
20. Bogale B, Gutema BT, Chisha Y. Prevalence of Stunting and Its Associated Factors among Children of 6–59 Months in Arba Minch Health and Demographic Surveillance Site (HDSS), Southern Ethiopia: A Community-Based Cross-Sectional Study. *Journal of Environmental and Public Health*. 2020;2020.
21. Gebre A, Reddy PS, Mulugeta A, Sedik Y, Kahssay M. Prevalence of malnutrition and associated factors among under-five children in pastoral communities of the afar regional state, Northeast Ethiopia: a community-based cross-sectional study. *Journal of nutrition and metabolism*. 2019;2019.
22. Mengesha DK, Merkeb Y. Prevalence of malnutrition and associated risk factors among under-five children in Amhara Region, Ethiopia: evidence from 2016 Ethiopian Demographic and Health Survey. 2020.
23. Asfaw M, Wondaferash M, Taha M, Dube LJBPh. Prevalence of undernutrition and associated factors among children aged between six to fifty-nine months in Bule Hora district, South Ethiopia. 2015;15(1):41.
24. Amare D, Negesse A, Tsegaye B, Assefa B, Ayenie B. Prevalence of undernutrition and its associated factors among children below five years of age in Bure Town, West Gojjam Zone, Amhara National

- Regional State, Northwest Ethiopia. *Advances in Public Health*. 2016;2016.
25. Girma A, Woldie H, Mekonnen FA, Gonete KA, Sisay M. Undernutrition and associated factors among urban children aged 24–59 months in Northwest Ethiopia: a community-based cross-sectional study. *BMC pediatrics*. 2019;19(1):214.
 26. Dake SK, Solomon FB, Bobe TM, Tekle HA, Tufa EG. Predictors of stunting among children 6–59 months of age in Sodo Zuria District, South Ethiopia: a community based cross-sectional study. *BMC nutrition*. 2019;5(1):23.
 27. Kasaye HK, Bobo FT, Yilma MT, Woldie M. Poor nutrition for under-five children from poor households in Ethiopia: Evidence from 2016 Demographic and Health Survey. *PLoS one*. 2019;14(12):e0225996.
 28. Haile A. Prevalence of Undernutrition and Associated Factors among Children Aged 6-59 Months in Bishoftu Town, Oromia Region, Ethiopia. *Food Proc Nutr Sci*. 2020;1(1):8-24.
 29. Abera L, Dejene T, Laelago T. Magnitude of stunting and its determinants in children aged 6–59 months among rural residents of Damot Gale district; southern Ethiopia. *BMC research notes*. 2018;11(1):557.
 30. Mehta NM, Corkins MR, Lyman B, Malone A, Goday PS, Carney L, et al. Defining pediatric malnutrition: a paradigm shift toward etiology-related definitions. *Journal of Parenteral and Enteral Nutrition*. 2013;37(4):460-81.
 31. Wuneh MT, Guracho YD. Nutritional status and feeding practice of children 6–59 months old, Metekele zone of Benishangul-Gumuz region, Northwest Ethiopia. *Science Journal of Clinical Medicine*. 2017;6(6):120-9.
 32. Mzumara B, Bwembya P, Halwiindi H, Mugode R, Banda JJBN. Factors associated with stunting among children below five years of age in Zambia: evidence from the 2014 Zambia demographic and health survey. 2018;4(1):51.
 33. Ekpo UF, Omotayo AM, Dipeolu MA. Prevalence of malnutrition among settled pastoral Fulani children in Southwest Nigeria. *BMC Research Notes*. 2008;1(1):7.
 34. Praditsorn P, Churak P, Wimonpeerapattana W, Moore T, Bovill M. Prevalence of undernutrition and associated factors among children 6 to 59 months of age in refugee camps along Thailand-Myanmar border. *Southeast Asian Journal of Tropical Medicine and Public Health*. 2019;50(2):372-82.
 35. Bhutia DT. Protein-energy malnutrition in India: the plight of our under-five children. *Journal of family medicine and primary care*. 2014;3(1):63.
 36. Hein AK, Hong SA, Puckpinyo A, Tejativaddhana P. Dietary diversity, social support, and stunting among children aged 6–59 months in an internally displaced person camp in Kayin state, Myanmar. *Clinical Nutrition Research*. 2019;8(4):307-17.
 37. Fentahun W, Wubshet M, Tariku A. Undernutrition and associated factors among children aged 6-59 months in East Belesa District, northwest Ethiopia: a community based cross-sectional study. *BMC public health*. 2016;16(1):506.

38. Bachewe FN, Yimer F, Minten B. Agricultural price evolution in drought versus non-drought affected areas in Ethiopia: An updated assessment using national producer data (January 2014 to January 2017). International Food Policy Research Institute (IFPRI), 2017.
39. Irenso AA, Dessie Y, Berhane Y, Assefa N, Canavan CR, Fawzi WW. Prevalence and predictors of adolescent linear growth and stunting across the urban-rural gradient in eastern Ethiopia. Tropical Medicine & International Health. 2020;25(1):101-10.
40. Song Y, Agardh A, Ma J, Li L, Lei Y, Stafford RS, et al. National trends in stunting, thinness, and overweight among Chinese school-aged children, 1985–2014. International Journal of Obesity. 2019;43(2):402-11.

Figures




Figure 1

Prevalence of stunting among children age 6-59 months at Debre tabor town Northwest Ethiopia, 2019(n=620)