Altered Cytokine and BDNF Levels in Individuals with Autism Spectrum Disorders

Yvonne M.Y. Han (✉ yvonne.han@polyu.edu.hk)

The Hong Kong Polytechnic University

Suk Yu Yau

The Hong Kong Polytechnic University

Melody M.Y. Chan

The Hong Kong Polytechnic University

Chun Kwok Wong

The Chinese University of Hong Kong

Agnes S. Chan

The Chinese University of Hong Kong

Research Article

Keywords: Immunologic function, Autism, Cognitive function, BDNF, Biomarkers

Posted Date: November 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1057100/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Brain Sciences on March 29th, 2022. See the published version at <https://doi.org/10.3390/brainsci12040460>.

Abstract

Previous studies have shown that immunological factors are involved in the pathogenesis of autism spectrum disorders (ASDs). The present study examined whether immunological abnormalities are associated with cognitive deficits and problem behaviors in children with ASD and whether children with ASD show different immunological biomarkers and brain-derived neurotrophic factor BDNF levels than typically developing (TD) children. Sixteen TD and 18 children with ASD, aged 6–18 years, voluntarily participated in the study. Participants' executive functions were measured using neuropsychological tests, and behavioral measures were measured using parent ratings. Immunological measures were assessed by measuring the levels of chemokine ligand 2 (CCL2) and chemokine ligand 5 (CCL5) in participants' blood serum. Children with ASD showed greater deficits in cognitive functions as well as altered levels of immunological measures when compared to TD children, and their cognitive functions and behavior deficits were significantly associated with increased CCL5 levels and decreased BDNF levels. These results provide evidence to support the notion that altered immune functions are associated with neuronal deficits, are indexed by decreased levels of BDNF and manifest as selective cognitive and behavioral symptoms in individuals with ASD.

1. Introduction

Autism spectrum disorder (ASD) is a pervasive and lifelong developmental disorder. The prevalence of ASD is high, with recent statistics of 1 in 54 individuals being reported, making ASD one of the most prevalent of all childhood developmental disorders¹. There is great phenotypic heterogeneity in ASD, ranging from severe mental retardation to isolated cognitive problems, such as stereotyped behavior and difficulty in understanding others' feelings^{2–4}. Although the causes of ASD are not well understood, increasing evidence has suggested that dysregulation of the immune system could play a role in the pathogenesis of ASD. Epidemiological studies have reported a greater family history of autoimmune disease in persons with autism than in healthy controls⁵. Within individuals with ASD, a growing number of studies have reported altered profiles in various cellular and humoral immune subsets, as measured in both postmortem brains of ASD patients and peripheral blood circulation in ASD patients⁶. The suggested autotoxic role of cytokines has been supported through recent clinical studies^{7,8}, in which active neuroinflammatory processes with cytokine profiling and activation of microglia and astrocytes were found in the brains of patients with ASD, suggesting neuroimmune dysfunction in the brains of autistic individuals.

Among the various cytokines, subsets of proinflammatory chemokines appear to play a crucial role in mediating neuroinflammation found in individuals with ASD⁹. For example, postmortem brain and spinal cord samples from individuals with ASD showed heightened proinflammatory cytokines, including chemokine macrophage chemoattractant protein 1/chemokine ligand 2 (MCP-1/CCL2), compared to control specimens⁷. Marked increases in CCL2 and eotaxin were also found in the cerebrospinal fluid (CSF) of children with ASD¹⁰. In addition to the elevated neuroinflammatory response in the brains of individuals with ASD, ASD patients showed altered levels of circulating chemokines in the blood. For example, altered gene expression, elevated plasma CCL2 levels and elevated RANTES/CCL5 levels were observed in ASD patients. Notably, chemokine levels have been shown to correlate with the risk^{11,12} and severity of behavioral

abnormalities^{13–15} of ASD. These findings suggest that the levels of chemokine production of CCL2 and CCL5 could be related to the level of severity of cognitive abnormalities in individuals with ASD, and these biomarkers can potentially serve as important tools for recognizing cognitive impairments in ASD patients.

Although research supports immunological involvement in the pathogenesis of ASD, there is relatively limited research investigating the relationship among altered immune systems and cognitive-behavioral deficits in individuals with ASD. Some research suggests a mechanism through which the excess activation of cytokines can lead to brain dysfunction related to disrupted production of growth factors and neurogenesis¹⁶. Brain-derived neurotrophic factor (BDNF) is a member of the “neurotrophin” family of growth factors and is known to have important functions in the formation, branching and connectivity of synaptic connections during development^{17,18}. It also plays a pivotal role in synaptic plasticity and long-term potentiation and is critically involved in learning, memory, and higher-order thinking^{19,20}. Pertinently, BDNF is found throughout the brain and in the periphery²¹. Circulating levels of BDNF have been shown to reflect BDNF levels in the brain^{22,23}. Moreover, low levels of BDNF in the brain and serum have been reported in various neuropsychiatric disorders, including schizophrenia^{24,25} and Alzheimer’s disease^{26,27}.

Additionally, recent studies have provided evidence to support the involvement of BDNF in autism through its neurotrophic effects on the developing brain^{16,28}. This notion is supported by previous studies that have demonstrated aberrant synaptic development and disordered connectivity as an underlying cause of the behavioral characteristics observed in individuals with ASD^{29,30}. However, the role of circulating BDNF remains unclear, and previous studies have shown inconsistent results. While some studies reported increased circulating levels of BDNF in patients with ASD^{16,31,32}, others showed lower circulating levels of BDNF in individuals in the ASD groups than in the typically developing (TD) controls³³. Nevertheless, due to its essential role in neuronal development, altered levels of BDNF could explain the abnormal cortical connectivity reported in individuals with ASD³⁴.

Although it is largely unknown how immunologic factors specifically affect neural networks in the brains of individuals with ASD, a better understanding of the relationship between immunologic function, neurophysiology and cognitive function in people with ASD is highly significant. Establishing these relationships would provide a foundation for the causal mechanism and developmental course of neurodevelopmental disorders in ASD patients that could be used in future research. Given the evidence linking ASD with abnormal immunologic function, altered levels of BDNF, and impaired synaptic connectivity, we postulate that the differing severities of autistic features would be associated with impaired synaptic connectivity in the brains of autistic individuals; moreover, dysregulation of neurotrophic factor function and increased proinflammatory responses would be the potential mechanisms. The present study investigated the role of immune dysfunction in children with ASD by examining the associations among immune responses, serum levels of BDNF, and severity of ASD symptoms. For this purpose, we investigated whether proinflammatory chemokines and altered BDNF levels play a role in the selective cognitive and behavioral symptoms of ASD by (1) examining the levels of the proinflammatory chemokines CCL2 and CCL5 and the levels of BDNF in the blood serum of children with ASD, then comparing these levels with those of TD children; (2) determining whether there was a correlation between chemokine (CCL2 and CCL5) and BDNF

levels; and (3) determining whether proinflammatory chemokines and altered BDNF levels play a role in the selective cognitive and behavioral symptoms of people with ASD.

2. Material And Methods

2.1 Participants

Twenty-one children with high-functioning ASD and 19 TD children aged between 6 and 17 years participated voluntarily in the study with their parents' written informed consent. All of the children had an intelligence quotient (IQ) of 90, as measured by the short form of the Wechsler Intelligence Scale for Children–Fourth Edition, Hong Kong (WISC–IV HK³⁵). Participants in the study were recruited by advertisements that were posted on our websites and invitation emails that were sent to parents in the contact database at the Neuropsychology Laboratory of the Chinese University of Hong Kong. All children with ASD were diagnosed based on the diagnostic criteria of the Diagnostic and Statistical Manual of Mental Disorders fifth edition (DSM-5)³⁶. The diagnosis was confirmed by a clinical psychologist using the Autism Diagnostic Interview-Revised (ADI-R)³⁷, which is a standard clinical interview conducted with parents on the developmental history and current functioning of the children. TD children had no history of delay in developmental milestones or any neurological or psychiatric disorders as reported by their parents, did not meet the DSM–5 diagnostic criteria of ASD and did not have ADI-R subscale scores over the specific cutoffs. Three children were excluded from the ASD group because two of them suffered from flu and had fever within 5 days of the blood test, and 1 child with ASD was prescribed immunosuppressive drugs. In addition, 16 children from the TD group were selected from a pool of 19 TD children to match the age (ASD: $M = 10.28$, $SD = 3.39$; TD: $M = 11.36$, $SD = 3.61$) and IQ (ASD: $M = 102.56$, $SD = 7.82$; TD: $M = 105.06$, $SD = 10.24$) of the individuals in the ASD group. Thus, 18 children with ASD and 16 TD children who were medication free and in good physical health at the time of blood sample assays were included for final analysis.

2.2 Procedures

All of the children and parents provided written informed consent before participating in the assessment and interviews. All children were individually administered the WISC-IV HK so that their IQ could be assessed and a neuropsychological assessment of executive functioning was conducted that comprised five standardized tests: the D2 test of attention (D2)³⁸, the Children's Color Trails Test (CCTT)³⁹, the Five Point Test (FPT)⁴⁰, the modified Wisconsin Card Sorting Test (WCST)⁴¹, and the Rey–Osterrieth Complex Figure Test (Rey-O)⁴². Information on the children's developmental and medical history was collected through structured and individual interviews with parents. All of the assessments and interviews were administered by trained research assistants who were blind to the rationale of the study and the group assignment.

Peripheral blood samples were collected from the children on separate days. A registered nurse drew 3 ml of blood mixed with ethylenediaminetetraacetic acid (EDTA) and 3 ml of clotted blood from each child using venipuncture at a medical clinic. The blood samples were kept in a thermally insulated bag and transported to the clinical laboratory where blood assays were performed. Blood sample processing and assays were performed by an experienced laboratory technician who was blind to the clinical characteristics and group assignments of the participants. Each blood sample was centrifuged at 3,000 rpm for 15 minutes, and the

harvested serum was then stored at -80°C for later processing. Serum levels of CCL2, CCL5 and BDNF were measured by enzyme-linked immunosorbent assay (ELISA) according to the manufacturer's instructions. This study was conducted in accordance with the Helsinki Declaration of the World Medical Association Assembly. The research protocol was approved by the Joint Chinese University of Hong Kong–New Territories East Cluster Clinical Research Ethics Committee (Ref. No. 2013.520) and the Human Subjects Ethics Sub-Committee of the Hong Kong Polytechnic University (Ref. No. HSEARS20170316006).

2.3 Measures

2.3.1 Immunological measures

The concentrations of the chemokines CCL2 and CCL5 were assessed using BD™ human chemokine cytometric bead array (CBA) reagent (Becton Dickinson Biosciences Pharmingen, CA, USA). Samples were analyzed on a multifluorescence BD FACSCalibur™ flow cytometer using BD CellQuest™ and BD™ CBA software. Serum BDNF levels were measured using commercial ELISA kits purchased from Abcam (ab99978, Cambridge, UK).

2.3.2 Behavioral measures

ADI-R. The ADI-R³⁷ is an investigator-based interview for parents of children whose behaviors are relevant to the diagnosis of ASD or pervasive developmental disorder. The test results in separate scores in three domains: (1) reciprocal social interaction; (2) communication; and (3) restricted, repetitive, and stereotyped patterns of behaviors. All of these domains correspond to the DSM-V diagnostic criteria of ASD and were chosen to reflect the severity of their children's core ASD symptoms, with higher scores indicating greater autistic features. Questions were asked concerning both the early developmental and current functioning of the child.

2.3.3 Executive functioning measures

WISC-IV HK short form. The short form of the WISC-IV (HK)³⁵ is a psychometric measure that can be used to evaluate the general intellectual ability of children between 6 and 16 years old. The short form comprises two verbal subtests, Digit Span and Similarities, and two performance subtests, Matrix Reasoning and Coding. The measure yields a standard score of IQ that has a mean of 100 and a standard deviation of 15.

D2. The D2³⁸ is a test of selective and sustained attention in which participants have to cancel out all target characters (a “d” that has two dashes placed above and/or below) and ignore other nontarget characters (a “d” or a “p” with more or less than two dashes above and/or below) within 20 seconds for each of the 14 trials. The concentration performance is calculated as the number of correctly cancelled target characters minus the total number of commission and omission errors demonstrated.

CCTT. The CCTT³⁹, which has a similar but simpler format compared to the traditional Trail Making Test, provides an evaluation of children's attention, cognitive flexibility, and speeded visuomotor tracking ability without the influence of language. Color Trail 1 requires participants to quickly connect scattered numbers from 1 to 8 in ascending order, while in Color Trail 2, participants have to correctly connect scattered numbers 1 to 15 that are concurrently embedded in alternate pink and yellow circles.

FPT. The FPT ⁴⁰ measures figural fluency capacity and provides an evaluation of children's cognitive flexibility. Participants are required to generate as many novel designs as possible by drawing straight lines to connect identically arranged dots presented in squares within 3 minutes.

Rey-O. The Rey-O ⁴² tests children's visuospatial constructional functions, visuographic memory, and planning. Participants are presented with a geometric figure and some colored pencils and are asked to copy the design on a piece of paper, after which they are asked to recall the figure from memory following a delay period of 3 minutes.

Modified WCST. In the modified version ⁴¹, participants have to sort a series of cards according to two rules (i.e., the color or shape of each card). The participants received no information about the sorting rule when the cards were presented, but they were explicitly informed whether their choice was correct after each response. In each trial, three cards arranged in a pyramid were presented by E-prime (Psychology Software Tools, Inc. Pittsburgh, PA) on a computer screen. The card at the top was the target card, and the two cards at the bottom were the choice cards. For half of the trials, the correct choice card was on the left, and for the other half of the trials, it was on the right. The participants had to match the target card with the correct choice card, based on the sorting rule at that moment, by pressing the left or right button on a button box that corresponded to the side of the choice card. There was no time limit for the participant to respond, and the stimuli stayed on the screen until the participant gave a response. There were a total of 5 test blocks with alternating sorting rules. A new block began after 7, 8, or 9 consecutive correct responses were made. The task was terminated if the criterion number of the consecutive correct responses was not met within 50 trials. A practice block that required 8 consecutive correct responses had to be completed first before the 5 test blocks begin.

2.4 Data Analyses

Independent sample t-tests were performed to compare the cognitive, behavioral, and immunological functioning of individuals in the ASD and TD groups. To reduce the number of statistical comparisons, one executive composite score was computed by summing and averaging the Z scores of the main dependent variables (DVs) of the five standardized neuropsychological tests. This method comprised D2 results, in which the main DV was the concentration performance; CCTT, in which the main DVs were the time to completion for Part 1 and Part 2; FPT, in which the main DV was the number of unique designs found in 3 minutes; and Rey-O, in which the main DV was immediate recall. For the modified WCST, the main DV was the mean number of total errors, which included both the number of perseverative errors and the failure to maintain sets. A low executive function composite score indicated poor performance. The relationship between the severity of autistic symptoms and executive functioning and between the concentrations of chemokines and BDNF was examined using Pearson's correlations (two-tailed). Between-group comparisons of categorical variables were analyzed using chi-square tests. The likelihood ratio was reported for distributions that had more than 20% of cells with an expected count of less than 5. Because specific hypotheses were tested, the alpha level was unadjusted and set at .05 for all of the tests. All statistical analyses were performed using SPSS software (SPSS, Inc., Chicago, IL, USA).

3. Results

3.1 Demographic and clinical characteristics

Table 1 shows the demographic and clinical characteristics of the children. The ASD and TD groups were matched for age ($t = 0.90$, $p = .38$), sex (likelihood ratio = 3.37, $p = .066$) and intellectual functioning ($t = 0.81$, $p = .43$) as measured by WISC-IV HK. The ASD group demonstrated significantly more ASD-related symptoms than the TD group (t ranged from 6.93 to 11.05, $ps < .001$).

Table 1

Demographic and clinical characteristics of children with autism (ASD) and the typically developing (TD) controls.

Characteristics	ASD (n = 18)	TD (n = 16)	t or χ^2	p
Age	10.28 (3.39)	11.36 (3.61)	.90	.38
IQ	102.56 (7.82)	105.06 (10.24)	.81	.43
Gender (Male %) [^]	61.5%	38.5%	3.37	.066
ADI-R social interaction	19.17 (7.22)	4.19 (3.43)	7.86	<.001***
ADI-R communication	13.67 (4.61)	1.25 (1.13)	11.05	<.001***
ADI-R stereotyped behavior	5.17 (2.28)	0.25 (0.77)	8.60	<.001***
<i>Note.</i>				
Data are presented as means (SD); IQ, intelligence quotient as assessed by the Chinese version of Wechsler Intelligence Scale for Children Forth Edition; ADI-R, Autism Diagnostic Interview-Revised; SRS-2, Social Responsiveness Scale Second Edition; [^] Likelihood ratio Chi-square test were performed for distribution violating the sample size assumption of Chi-square test.				
*** $p < .001$.				

3.2 Deficient executive functioning in children with ASD

Table 2 presents the means and standard deviations of the executive function composite score and the individual results in each of the five neuropsychological tests that were conducted. An independent sample t-test showed that individuals in the ASD group demonstrated significantly lower executive function composite scores than individuals in the TD group. Specifically, individuals in the ASD group produced a significantly less unique design in the FPT ($t = 3.70$; $p < .001$) and committed a larger number of total errors (i.e., perseverative errors and failures to maintain sets) in the modified WCST ($t = 2.51$; $p = .021$) than individuals in the TD group. Individuals in the ASD group also tended to have worse concentration performance ($t = 1.84$; $p = .075$), take longer to complete the first trial of the CCTT ($t = 1.88$; $p = .069$), and have lower accuracy in the immediate recall of the Rey-O ($t = 2.00$; $p = .054$), although the group differences were only marginally significant. In sum, individuals in the ASD group had significantly poorer performance across the executive functioning tasks compared to the TD counterparts.

Table 2
Comparison of mean scores in executive function tasks between children with autism (ASD) and typically developing (TD) controls.

	ASD (n = 18)	TD (n = 16)	<i>t</i>	<i>p</i>
Executive Function Composite	-5.65 (7.32)	0.34 (4.68)	2.80	.009**
D2 Test of Attention				
Concentration performance	124.50 (43.04)	151.75 (43.88)	1.84	.075
Children's Color Trails Test				
Trial 1 – Time to completion (second)	29.50 (12.34)	22.31 (9.59)	1.88	.069
Trial 2 – Time to completion (second)	58.72 (24.17)	49.56 (18.98)	1.22	.23
Five Point Test				
Number of unique designs in 3-minute	20.78 (6.66)	31.31 (9.82)	3.70	.001**
Rey–Osterrieth Complex Figure Test				
Immediate recall	10.36 (5.16)	14.66 (7.31)	2.00	.054
Modified Wisconsin Card Sorting Test				
Mean number of total errors	3.00 (3.13)	1.10 (0.68)	2.51	.021*
<i>Note.</i>				
Data are presented as means (SD).				
** $p < .01$; * $p < .05$.				

3.3 Deviated serum concentrations of chemokines and BDNF in children with ASD

The results of an independent sample t-test showed that compared to individuals in the TD group, individuals in the ASD group had a significantly higher level of the inflammatory chemokine CCL5 ($t = 2.86$; $p = .008$) and a marginally significant trend of higher CCL2 levels ($t = 2.00$; $p = .056$) (Table 3). In addition, BDNF levels were lower in individuals in the ASD group than in individuals in the TD group ($t = 2.31$; $p = .028$).

Table 3
Comparison of mean concentrations of chemokines and neurotrophin in children with autism (ASD) and typically developing (TD) controls.

	ASD (n = 18)	TD (n = 15 [^])	<i>t</i>	<i>p</i>
Chemokines				
CCL2 (pg/mL)	181.10 (68.18)	143.17 (37.69)	2.00	.056
CCL5 (pg/mL)	43867.64 (15124.57)	31070.68 (8103.60)	2.86	.008**
Neurotrophin				
BDNF (pg/mL)	12.08 (5.02)	15.94 (4.21)	2.31	.028*
<i>Note.</i>				
Data are presented as means (SD).				
[^] One datum is missing from the TD group.				
** $p < .01$; * $p < .05$.				

3.4 Associations between the severity of ASD symptoms, the executive function composite score and between the levels of chemokines and BDNF

Pearson's correlation analysis was performed to assess the possible correlations between the behavioral symptoms of ASD as indicated by the three subscale scores of ADI-R, the executive function composite, the mean levels of the inflammatory chemokines CCL2 and CCL5, and BDNF. The results showed a significant correlation between the level of CCL5 and the social interaction subscale of the ADI-R and between the level of CCL5 and the communication subscale of the ADI-R; more specifically, a higher level of CCL5 was correlated with a higher score on the two subscales ($r = .35$, $p = .047$; $r = .36$; $p = .041$, respectively) (Table 4). Similarly, a significant correlation was found between the level of BDNF and each of the social interaction, communication and stereotyped behavior subscales of the ADI-R (r ranges from $-.39$ to $-.49$; p from $.004$ to $.027$) individually. More specifically, the lower the level of the BDNF was, the higher the three subscale scores of the ADI-R were. In addition, it was found that the executive function composite score was negatively correlated with the scores on the social interaction and communication subscales of the ADI-R ($r = -.36$, $p = .037$; $r = -.58$; $p < .001$, respectively).

Table 4

Pearson association analysis of the three subscales of ADI-R with executive function composite, mean concentrations of CCL2, CCL5, and BDNF in all participants enrolled (n = 33[^]).

	ADI-R social interaction	ADI-R communication	ADI-R stereotyped behavior	Executive Function Composite	CCL2	CCL5	BDNF
ADI-R social interaction	1.00						
ADI-R communication	.90***	1.00					
ADI-R stereotyped behavior	.75***	.78***	1.00				
Executive Function Composite	-.36*	-.58***	-.28	1.00			
CCL2	.144	.20	.10	-.09	1.00		
CCL5	.35*	.36*	.28	-.26	.29	1.00	
BDNF	-.49**	-.47**	-.39*	.33	.08	-.16	1.00
<i>Note.</i>							
ADI-R, Autism Diagnostic Interview-Revised							
[^] One datum is missing from the TD group.							
*** $p < .001$; ** $p < .01$; * $p < .05$							

4. Discussion

The present study examined cognitive dysfunction and behavioral deficits in high-functioning children with ASD and investigated whether these abnormalities were associated with altered proinflammatory chemokine and BDNF levels. When compared to age-, IQ- and sex-matched TD individuals, individuals with ASD showed significantly more severe social communication and interaction deficits with more restrictive repetitive behaviors, weaker executive functioning performance, higher mean level of pro-inflammatory chemokine CCL5, and lower mean level of blood serum BDNF. Importantly, a higher CCL5 level was found to be associated with greater deficits in social interaction and communication, and a lower serum BDNF level was found to be associated with a higher severity across all ASD core behavioral symptoms.

Consistent with previous studies that included individuals with ASD that were of a younger^{11,13} or older⁴³ age when compared to this study sample, we revealed an elevated pro-inflammatory chemokine profile in individuals with ASD. Specifically, CCL5 levels were found to be significantly higher, and the trend of higher levels of CCL2 was found to be marginally significant in our sample. Importantly, this research pioneered the establishment of a significant positive association between CCL5 and ASD social deficits measured by the

ADI-R, the 'gold standard' ^{44,45} in ASD diagnostic evaluation. This association implies that immune dysfunction could be a possible mechanism underlying the manifestation of ASD core behavioral symptoms. Indeed, our current results are supported by previous animal studies showing that reducing proinflammatory chemokines with the administration of resveratrol ⁴⁶ could result in a reduction in ASD behavioral phenotypes ^{47,48}. Notably, the mechanisms of how immune dysfunction results in ASD behavioral deficits remain to be investigated, and abnormal spinogenesis associated with cytokine and chemokine dysregulation ⁴⁹ might be a possible candidate for further studies. For instance, converging data from postmortem human brain tissues ^{50,51} and animal studies using ASD-related models ^{52,53} have suggested an increase in the number of immature dendritic spines in individuals with ASD. In addition, abnormal turnover and structure of dendritic spines have been observed in individuals with ASD that could consequently contribute to delayed spine maturation [see Phillips and Pozzo-Miller ⁵⁴ for a review]. Furthermore, abnormalities in spine dynamics have been shown to result in selective cognitive and behavioral symptoms with neural circuit dysfunction (i.e., disordered synaptic connectivity) in individuals with ASD ⁵⁵. Future research that investigates the causal relationship between proinflammatory cytokines and immature spinogenesis in ASD animal models, as well as the association between aberrant immune function and functional connectivity in human ASD subjects, would be beneficial to improve our understanding of the pathophysiology of this neurodevelopmental disorder.

A recent review concluded that significantly reduced BDNF expression is commonly found in patients with neuropsychiatric disorders ⁵⁶. In line with this observation, as well as being consistent with some previous studies ^{57,58}, our data showed that children with ASD were found to have a lower level of serum BDNF than TD individuals. Furthermore, a lower BDNF level was found to be correlated with a greater severity in all core ASD behavioral symptoms (i.e., social communication/interaction deficits and restricted repetitive behaviors) reflected in the ADI-R scores, suggesting the importance of BDNF abnormalities in the manifestation of ASD symptomatology. BDNF is known to play a crucial role in modulating structural and synaptic plasticity in the brain ^{59,60}. Notably, enhancing BDNF concentration has been found to promote dendritic ⁶¹ and axonal ⁶² growth in an activity-dependent manner ⁶³. Also, in ASD human and animal studies, a generalized reduction in the size and number of dendrites has been observed ⁶⁴; moreover, reduced axonal diameter and white matter in brain regions related to social cognition and information processing hubs has been found using diffusion tensor imaging techniques in ASD patients ^{65,66}. Based on these findings, together with our data, we postulate that BDNF reduction in individuals with ASD might be associated with the dendritic and axonal abnormalities in these individuals, though this concept has to be further verified in future studies. It is worth pointing out that our results are in contrast with two studies with a Chinese population showing that BDNF is higher, instead of lower, in individuals with ASD when compared to TD individuals ^{67,68}. This outcome could be attributed to the fact that the other studies recruited ASD participants that were younger in age (mean age = approximately 4 years) when compared to our sample (mean age = 10.28 years). Also, previous studies have shown that BDNF levels decrease with increasing age in healthy humans (Erickson et al., 2010). A higher BDNF level in younger ASD participants and a lower BDNF level in older ASD participants might imply that these individuals differ in the developmental trajectory of BDNF levels that might modulate cognitive development, which warrants further research.

Social communication is a highly complex construct that requires an individual to process and integrate relevant information from the environment to produce adaptive behaviors for social interaction ⁶⁹. Successful integration and the manipulation of social information require a person to have multiple intact executive functions, such as cognitive flexibility and attention, involving coordination between primarily the frontal lobe and other brain regions ⁷⁰. Importantly, both cognitive flexibility ⁷¹ and attention ⁷² have been found to be associated with social communication skills. Impairments in these cognitive processes have been hypothesized to underlie social communication dysfunction, which is a core symptom in people with ASD regardless of their level of functioning ⁷³. Indeed, our results showed that ASD individuals had weaker executive functioning, specifically in cognitive flexibility (indicated by statistically poorer performance on the FPT and modified WCST), which is consistent with results from a meta-analysis ⁷⁴. We further showed that poorer executive functioning was associated with poorer social interaction/communication skills (indexed by higher ADI-R subscores in these domains), which reiterated the importance of intact executive functions in social information processing. Increasing evidence supports the notion that the executive functioning deficits in individuals with ASD are due to underlying brain abnormalities that affect neural networks across these individuals' brains. For instance, previous human studies have reported disordered long-distance corticocortical connectivity and localized connectivity in the brains of individuals with ASD when they perform various cognitive tasks, including cognitive flexibility ⁷⁵; in addition, disordered connectivity has been shown to be correlated with the severity of impairment in social communication and of restrictive/repetitive behaviors ⁷⁶. Given that connections of the neuronal circuit have been shown to be mediated by the remodeling of dendrites and that this remodeling process has been associated with changes in BDNF and cytokine levels in the brain, as discussed above, we hypothesized that BDNF expression, chemokine concentration and executive functioning are interrelated. Surprisingly, although BDNF and chemokine levels, as well as executive functioning, were indeed abnormal in individuals with ASD, as shown and discussed above, the correlations of BDNF levels and executive functioning composite scores ($p = .057$), CCL5 levels and executive functioning composite scores ($p = .138$), and between CCL5 and BDNF levels ($p = .366$) did not reach statistical significance. However, our results could be limited by a relatively small sample size, or they could also be explained by the inherent heterogeneous clinical and neurobiological profiles of individuals with ASD ⁷⁷. Future research with a larger sample size and a more homogenous sample might be helpful in exploring these relationships to strengthen our understanding of neuroinflammation, neuronal modeling and executive functioning in individuals with ASD.

5. Conclusion

This study aimed to investigate the relationships between immunological abnormalities, cognitive deficits and behavioral manifestations in children with ASD. When comparing the results from 16 TD and 18 ASD individuals aged 6–18 years, ASD individuals exhibited greater deficits in cognitive functions along with altered levels of immunological measures. Importantly, aberrant cognitive functions and associated behavioral deficits in individuals with ASD were significantly associated with heightened CCL5 levels and lower BDNF levels. Our results support the notion that abnormal immune functions are associated with neuronal deficits, are indexed by decreased levels of BDNF and manifest as selective cognitive and behavioral symptoms in individuals with ASD.

Declarations

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgements

The authors would like to especially thank all the children and parents who participated in the research. This study was supported by the research grant (YBYQ) from The Hong Kong Polytechnic University, and donations from Mr. Sau-wang Law and Mr. Sau-hung Li to The Chinese University of Hong Kong.

References

1. Maenner, M. J., Shaw, K. A. & Baio, J. Prevalence of autism spectrum disorder among children aged 8 years—autism and developmental disabilities monitoring network, 11 sites, United States, 2016. *MMWR Surveillance Summaries*, **69**, 1 (2020).
2. Chan, A. S. *et al.* Disordered connectivity associated with memory deficits in children with autism spectrum disorders. *Research in Autism Spectrum Disorders*, **5**, 237–245 (2011).
3. Cheung, M., Chan, A. S., Sze, S. L., Leung, W. W. & To, C. Y. Verbal memory deficits in relation to organization strategy in high-and low-functioning autistic children. *Research in Autism Spectrum Disorders*, **4**, 764–771 (2010).
4. Happé, F. Autism: cognitive deficit or cognitive style? *Trends in cognitive sciences*, **3**, 216–222 (1999).
5. Sweeten, T. L., Bowyer, S. L., Posey, D. J., Halberstadt, G. M. & McDougle, C. J. Increased prevalence of familial autoimmunity in probands with pervasive developmental disorders., **112**, e420–e420 (2003).
6. Goines, P. & Van de Water, J. The immune system's role in the biology of autism. *Current opinion in neurology*, **23**, 111 (2010).
7. Vargas, D. L., Nascimbene, C., Krishnan, C., Zimmerman, A. W. & Pardo, C. A. Neuroglial activation and neuroinflammation in the brain of patients with autism. *Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society*, **57**, 67–81 (2005).
8. Di Filippo, M., Sarchielli, P., Picconi, B. & Calabresi, P. Neuroinflammation and synaptic plasticity: theoretical basis for a novel, immune-centred, therapeutic approach to neurological disorders. *Trends in pharmacological sciences*, **29**, 402–412 (2008).
9. Pardo, C. A., Vargas, D. L. & Zimmerman, A. W. Immunity, neuroglia and neuroinflammation in autism. *International review of psychiatry*, **17**, 485–495 (2005).
10. Li, X. *et al.* Elevated immune response in the brain of autistic patients. *Journal of neuroimmunology*, **207**, 111–116 (2009).
11. Ashwood, P. *et al.* Associations of impaired behaviors with elevated plasma chemokines in autism spectrum disorders. *Journal of neuroimmunology*, **232**, 196–199 (2011).

12. Abdallah, M. *et al.* Neonatal levels of neurotrophic factors and risk of autism spectrum disorders. *Acta Psychiatry. Scand*, **128**, 61–69 (2013).
13. Han, Y. M. *et al.* Distinct cytokine and chemokine profiles in autism spectrum disorders. *Frontiers in immunology*, **8**, 11 (2017).
14. Ashwood, P. *et al.* Elevated plasma cytokines in autism spectrum disorders provide evidence of immune dysfunction and are associated with impaired behavioral outcome. *Brain, behavior, and immunity*, **25**, 40–45 (2011).
15. Grigorenko, E. L. *et al.* Macrophage migration inhibitory factor and autism spectrum disorders., **122**, e438–e445 (2008).
16. Ricci, S. *et al.* Altered cytokine and BDNF levels in autism spectrum disorder. *Neurotoxicity research*, **24**, 491–501 (2013).
17. Binder, D. K. & Scharfman, H. E. Brain-derived neurotrophic factor. *Growth factors (Chur, Switzerland)*, **22**, 123 (2004).
18. Huang, E. J. & Reichardt, L. F. Neurotrophins: roles in neuronal development and function. *Annual review of neuroscience*, **24**, 677–736 (2001).
19. Bekinschtein, P. *et al.* BDNF is essential to promote persistence of long-term memory storage. *Proceedings of the National Academy of Sciences* **105**, 2711–2716(2008)
20. Yamada, K. & Nabeshima, T. Brain-derived neurotrophic factor/TrkB signaling in memory processes. *Journal of pharmacological sciences*, **91**, 267–270 (2003).
21. Nickl-Jockschat, T. & Michel, T. The role of neurotrophic factors in autism. *Molecular psychiatry*, **16**, 478–490 (2011).
22. Angelucci, F., Gelfo, F., De Bartolo, P., Caltagirone, C. & Petrosini, L. BDNF concentrations are decreased in serum and parietal cortex in immunotoxin 192 IgG-Saporin rat model of cholinergic degeneration. *Neurochemistry international*, **59**, 1–4 (2011).
23. Morichi, S. *et al.* Brain-derived neurotrophic factor and interleukin-6 levels in the serum and cerebrospinal fluid of children with viral infection-induced encephalopathy. *Neurochemical research*, **39**, 2143–2149 (2014).
24. Green, M. J., Matheson, S., Shepherd, A., Weickert, C. & Carr, V. Brain-derived neurotrophic factor levels in schizophrenia: a systematic review with meta-analysis. *Molecular psychiatry*, **16**, 960–972 (2011).
25. Weickert, C. *et al.* Reduced brain-derived neurotrophic factor in prefrontal cortex of patients with schizophrenia. *Molecular psychiatry*, **8**, 592–610 (2003).
26. Peng, S., Wu, J., Mufson, E. J. & Fahnstock, M. Precursor form of brain-derived neurotrophic factor and mature brain-derived neurotrophic factor are decreased in the pre-clinical stages of Alzheimer's disease. *Journal of neurochemistry*, **93**, 1412–1421 (2005).
27. Michalski, B. & Fahnstock, M. Pro-brain-derived neurotrophic factor is decreased in parietal cortex in Alzheimer's disease. *Mol. Brain Res*, **111**, 148–154 (2003).
28. Bryn, V. *et al.* Brain derived neurotrophic factor (BDNF) and autism spectrum disorders (ASD) in childhood. *European Journal of Paediatric Neurology*, **19**, 411–414 (2015).

29. Just, M. A., Cherkassky, V. L., Keller, T. A., Kana, R. K. & Minshew, N. J. Functional and anatomical cortical underconnectivity in autism: evidence from an fMRI study of an executive function task and corpus callosum morphometry. *Cerebral cortex*, **17**, 951–961 (2007).
30. Kana, R. K., Libero, L. E. & Moore, M. S. Disrupted cortical connectivity theory as an explanatory model for autism spectrum disorders. *Physics of life reviews*, **8**, 410–437 (2011).
31. Correia, C. *et al.* Increased BDNF levels and NTRK2 gene association suggest a disruption of BDNF/TrkB signaling in autism. *Genes, Brain and behavior*, **9**, 841–848 (2010).
32. Zhang, Q., Jiang, L. & Lu, Y. J. Serum brain-derived neurotrophic factor levels in Chinese children with autism spectrum disorders: a pilot study. *International Journal of Developmental Neuroscience*, **37**, 65–68 (2014).
33. Nelson, P. G. *et al.* Selected neurotrophins, neuropeptides, and cytokines: developmental trajectory and concentrations in neonatal blood of children with autism or Down syndrome. *International journal of developmental neuroscience*, **24**, 73–80 (2006).
34. Garcia, K. L. *et al.* Altered balance of proteolytic isoforms of pro-brain-derived neurotrophic factor in autism. *Journal of neuropathology and experimental neurology*, **71**, 289–297 (2012).
35. Wechsler, D. Wechsler intelligence scale for children—Fourth edition, Hong Kong Version. *Zhuhai, Guangdong China: King-May Psychological Assessment Technology Development Limited Company* (2010)
36. American Psychiatric Association, D. & Association, A. P. *Diagnostic and statistical manual of mental disorders: DSM-5* (American Psychiatric Association, Washington, DC, 2013).
37. Lord, C., Rutter, M. & Le Couteur, A. Autism Diagnostic Interview-Revised: a revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of autism and developmental disorders*, **24**, 659–685 (1994).
38. Brickenkamp, R. & Zillmer, E. Test d2: Concentration-Endurance Test. *Gottingen Ger. CJ Hogrefe* (1998)
39. Llorente, A. M. *et al.* Children's Color Trails Test 1 & 2: test–retest reliability and factorial validity. *Clin. Neuropsychol*, **23**, 645–660 (2009).
40. Regard, M., Strauss, E. & Knapp, P. Children's production on verbal and non-verbal fluency tasks. *Perceptual and motor skills*, **55**, 839–844 (1982).
41. Van Eylen, L. *et al.* Cognitive flexibility in autism spectrum disorder: Explaining the inconsistencies? *Research in Autism Spectrum Disorders*, **5**, 1390–1401 (2011).
42. Bernstein, J. H. & Waber, D. P. *Developmental scoring system for the Rey-Osterrieth complex figure: DSS ROCF* (Psychological Assessment Resources, 1996).
43. Suzuki, K. *et al.* Plasma cytokine profiles in subjects with high-functioning autism spectrum disorders. *PloS one*, **6**, e20470 (2011).
44. Reaven, J. A., Hepburn, S. L. & Ross, R. G. Use of the ADOS and ADI-R in children with psychosis: importance of clinical judgment. *Clinical child psychology and psychiatry*, **13**, 81–94 (2008).
45. Cholemkey, H., Medda, J., Lempp, T. & Freitag, C. M. Classifying autism spectrum disorders by ADI-R: subtypes or severity gradient? *Journal of autism and developmental disorders*, **46**, 2327–2339 (2016).

46. Ahmad, S. F. *et al.* Resveratrol attenuates pro-inflammatory cytokines and activation of JAK1-STAT3 in BTBR T+ Itpr3tf/J autistic mice. *European Journal of Pharmacology*, **829**, 70–78 (2018).
47. Bakheet, S. A. *et al.* Resveratrol treatment attenuates chemokine receptor expression in the BTBR T+ tf/J mouse model of autism. *Molecular and Cellular Neuroscience*, **77**, 1–10 (2016).
48. Ahmad, S. F. *et al.* Immune alterations in CD8+ T cells are associated with neuronal CC and CXC chemokine receptor regulation through adenosine A2A receptor signaling in a BTBR T+ Itpr3 tf/J autistic mouse model. *Mol. Neurobiol*, **55**, 2603–2616 (2018).
49. Bitzer-Quintero, O. K. & González-Burgos, I. Immune system in the brain: a modulatory role on dendritic spine morphophysiology? *Neural plasticity* **2012**(2012)
50. Hutsler, J. J. & Zhang, H. Increased dendritic spine densities on cortical projection neurons in autism spectrum disorders. *Brain research*, **1309**, 83–94 (2010).
51. Weir, R., Bauman, M. D., Jacobs, B. & Schumann, C. Protracted dendritic growth in the typically developing human amygdala and increased spine density in young ASD brains. *Journal of Comparative Neurology*, **526**, 262–274 (2018).
52. McKinney, B. C., Grossman, A. W., Elisseou, N. M. & Greenough, W. T. Dendritic spine abnormalities in the occipital cortex of C57BL/6 Fmr1 knockout mice. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, **136**, 98–102 (2005).
53. Galvez, R., Gopal, A. R. & Greenough, W. T. Somatosensory cortical barrel dendritic abnormalities in a mouse model of the fragile X mental retardation syndrome. *Brain research*, **971**, 83–89 (2003).
54. Phillips, M. & Pozzo-Miller, L. Dendritic spine dysgenesis in autism related disorders. *Neuroscience letters*, **601**, 30–40 (2015).
55. Nakai, N., Takumi, T., Nakai, J. & Sato, M. Common defects of spine dynamics and circuit function in neurodevelopmental disorders: a systematic review of findings from in vivo optical imaging of mouse models. *Frontiers in neuroscience*, **12**, 412 (2018).
56. Giacobbo, B. L. *et al.* Brain-derived neurotrophic factor in brain disorders: focus on neuroinflammation. *Molecular neurobiology*, **56**, 3295–3312 (2019).
57. Hashimoto, K. *et al.* Reduced serum levels of brain-derived neurotrophic factor in adult male patients with autism. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, **30**, 1529–1531 (2006).
58. Abdallah, M. W. *et al.* Neonatal levels of neurotrophic factors and risk of autism spectrum disorders. *Acta Psychiatrica Scand*, **128**, 61–69 (2013).
59. Edelman, E., Lessmann, V. & Brigadski, T. Pre-and postsynaptic twists in BDNF secretion and action in synaptic plasticity., **76**, 610–627 (2014).
60. Panja, D. & Bramham, C. R. BDNF mechanisms in late LTP formation: a synthesis and breakdown., **76**, 664–676 (2014).
61. Horch, H. W. & Katz, L. C. BDNF release from single cells elicits local dendritic growth in nearby neurons. *Nature neuroscience*, **5**, 1177–1184 (2002).
62. Mamounas, L. A. *et al.* BDNF promotes the regenerative sprouting, but not survival, of injured serotonergic axons in the adult rat brain. *Journal of Neuroscience*, **20**, 771–782 (2000).

63. Lu, B. BDNF and activity-dependent synaptic modulation. *Learning & memory*, **10**, 86–98 (2003).
64. Martínez-Cerdeño, V. Dendrite and spine modifications in autism and related neurodevelopmental disorders in patients and animal models. *Developmental neurobiology*, **77**, 393–404 (2017).
65. Shukla, D. K., Keehn, B. & Müller, R. A. Tract-specific analyses of diffusion tensor imaging show widespread white matter compromise in autism spectrum disorder. *Journal of Child Psychology and Psychiatry*, **52**, 286–295 (2011).
66. Noriuchi, M. *et al.* Altered white matter fractional anisotropy and social impairment in children with autism spectrum disorder. *Brain research*, **1362**, 141–149 (2010).
67. Saghaazadeh, A. & Rezaei, N. Brain-derived neurotrophic factor levels in autism: a systematic review and meta-analysis. *Journal of autism and developmental disorders*, **47**, 1018–1029 (2017).
68. Zheng, Z. *et al.* Peripheral brain-derived neurotrophic factor in autism spectrum disorder: a systematic review and meta-analysis. *Scientific reports*, **6**, 1–8 (2016).
69. Geurts, H. M., de Vries, M. & van den Bergh, S. F. Executive functioning theory and autism. in *Handbook of executive functioning* 121-141 (Springer, 2014)
70. Diamond, A. Executive functions. *Annual review of psychology*, **64**, 135–168 (2013).
71. Bock, A. M., Gallaway, K. C. & Hund, A. M. Specifying links between executive functioning and theory of mind during middle childhood: Cognitive flexibility predicts social understanding. *Journal of Cognition and Development*, **16**, 509–521 (2015).
72. Arterberry, M. E., Bornstein, M. H., Midgett, C., Putnick, D. L. & Bornstein, M. H. Early attention and literacy experiences predict adaptive communication. *First Language*, **27**, 175–189 (2007).
73. Ozonoff, S., South, M. & Provencal, S. Executive functions. (2005)
74. Demetriou, E. *et al.* Autism spectrum disorders: a meta-analysis of executive function. *Molecular psychiatry*, **23**, 1198–1204 (2018).
75. Doesburg, S. M., Vidal, J. & Taylor, M. J. Reduced theta connectivity during set-shifting in children with autism. *Frontiers in human neuroscience*, **7**, 785 (2013).
76. Vissers, M. E., Cohen, M. X. & Geurts, H. M. Brain connectivity and high functioning autism: a promising path of research that needs refined models, methodological convergence, and stronger behavioral links. *Neuroscience & Biobehavioral Reviews*, **36**, 604–625 (2012).
77. Jeste, S. S. & Geschwind, D. H. Disentangling the heterogeneity of autism spectrum disorder through genetic findings. *Nature Reviews Neurology*, **10**, 74 (2014).