

Trends in Burnout and Psychological Distress in Hospital Staff Over 12 Months of The COVID-19 Pandemic: A Prospective Longitudinal Survey

Robert G. Maunder (✉ Robert.maunder@sinaihealth.ca)

Sinai Health, Canada & Department of Psychiatry, University of Toronto

Natalie D. Heeney

Sinai Health

Jonathan J. Hunter

Sinai Health, Canada & Department of Psychiatry, University of Toronto

Gillian Strudwick

Centre for Addiction and Mental Health, Canada & Institute of Health Policy, Management and Evaluation, University of Toronto

Lianne P. Jeffs

Lunenfeld-Tanenbaum Research Institute, Sinai Health

Leanne Ginty

Sinai Health

Jennie Johnstone

Sinai Health, Canada & Department of Laboratory Medicine and Pathobiology, University of Toronto

Alex Kiss

Department of Research Design and Biostatistics, Sunnybrook Research Institute, Canada & Dalla Lana School of Public Health, University of Toronto

Carla A. Loftus

Sinai Health

Lesley A. Wiesenfeld

Sinai Health, Canada & Department of Psychiatry, University of Toronto

Research Article

Keywords: COVID-19, pandemic, healthcare workers, nurses, burnout, resilience, psychological distress

Posted Date: November 12th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1065169/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background. The mental health effects of healthcare work during the COVID-19 pandemic have been substantial, but it is not known how long they will persist. This study aimed to determine if hospital workers' burnout and psychological distress increased monotonically over one year, during which waves of case numbers and hospitalizations waxed and waned, or followed some other pattern.

Methods. A prospective longitudinal survey was conducted at four time-points over one year in a cohort of 538 hospital workers and learners, which included validated measures of burnout (emotional exhaustion scale of Maslach Burnout Inventory) and psychological distress (K6). Repeated measures ANOVA tested changes over time and differences between subjects by occupational role and age. The direction and magnitude of changes over time were investigated by plotting rates of high scores (using cut-offs) at each time-point compared to case rates of COVID-19 in the city in which the study took place.

Results. There were significant changes in emotional exhaustion over time ($F = 4.6, p = .004$) and significant effects of occupational role ($F = 11.4, p < .001$) and age ($F = 12.3, p < .001$). The rate of high burnout was highest in nurses, followed by other healthcare professionals, other clinical staff, and lowest in non-clinical staff. Peak rates of high burnout occurred at the second or third measurement point for each occupational group, with lower rates at the fourth measurement point. Similarly to the results for emotional exhaustion, rates of high psychological distress peaked at the winter 2021 or spring 2021 measurement point in each occupational group and were highest in nurses.

Conclusions. Neither emotional exhaustion nor psychological distress was rising monotonically. Burnout and psychological distress were consistently related to occupational role and were highest in nurses. Although emotional exhaustion improved as the case rate of COVID-19 decreased, rates of high emotional exhaustion in nurses and other healthcare professionals remained higher than was typically measured in hospital-based healthcare workers prior to the pandemic. Ongoing monitoring of healthcare workers' mental health is warranted. Organizational and individual interventions to support healthcare workers continue to be important.

Background

A global pandemic of COVID-19 was declared by the World Health Organization on March 11, 2020, (1) and persists as of this writing, 20 months later. New cases have occurred in waves lasting months, which vary by geographic region (2). The mental health effects of working in healthcare during the pandemic are widely appreciated to be substantial, resulting from uncertainty, risk of infection, high volumes of acutely ill patients, and long hours, among other factors. Reports from cross-sectional studies have consistently indicated, for example, high rates of depressive and anxiety symptoms, sleep disturbance and burnout in healthcare workers (3–6).

Professional burnout is a particularly important outcome of occupational stress in healthcare because, in addition to its impact on individual professionals, burnout has adverse consequences for the healthcare system and patients, including absenteeism, higher workforce turnover, reduced productivity, increased medical errors, and reduced quality of care (7–10). As described by Maslach and colleagues (11), burnout is comprised of three components: emotional exhaustion, depersonalization (becoming indifferent or emotionally distant), and a diminished sense of personal achievement. Burnout was recognized prior to the COVID-19 pandemic to be a major occupational risk in health care (12,13). Workplace factors that were known to contribute to burnout include high workload, lack of scheduling flexibility, the burden of administrative tasks, and concern about workplace safety (14–20). In addition to workplace stressors that were present prior to the COVID-19 pandemic, additional potential sources of burnout have emerged, including constraints on care that cause moral distress (21,22) and redeployment to unfamiliar types of work (23).

While burnout is not considered to be a mental illness, rates of diagnoses of mood or anxiety disorders as defined in the *Diagnostic and Statistical Manual of Mental Disorders* (24), or measures that predict common mental illnesses, such as high levels of psychological distress, have also been measured to determine the impact of the COVID-19 pandemic on healthcare workers' mental health. A systematic review containing 65 studies that include 97,333 healthcare workers from 21 countries found pooled prevalence's of 21.7%, 22.1%, and 21.5% for moderate depression, anxiety, and post-traumatic stress disorder respectively during COVID-19 (25). Similarly, in a systematic review and meta-analysis including 29 studies and 22,380 hospital staff caring for COVID-

19 patients from countries in Europe and Asia, the prevalence of depression, anxiety, and stress was 24.3%, 25.8%, and 45%, respectively (6).

The long-term impact of the COVID-19 pandemic on healthcare workers' mental health is not known. Some factors suggest that the mental impact of the pandemic will be enduring. In particular, after the 2003 outbreak of severe acute respiratory syndrome (SARS) had resolved, healthcare workers in affected hospitals reported a range of stress effects that continued to be significantly elevated compared to workers in non-affected hospitals for as long as they were followed, which was 18-24 months (26). Of concern, the COVID-19 pandemic is far more severe, widespread and persistent than the SARS outbreak and has severely affected many aspects of life in the community as well as in the hospital, which suggests its mental impact will be greater, and may resolve more slowly. Furthermore, with respect to burnout, pre-pandemic rates of burnout in many settings were high, in the range of 20-40% (27–29) when defined by a score ≥ 27 on the emotional exhaustion scale of the Maslach Burnout Inventory, which is a commonly used operational definition (27), indicating that even a return to “normal” operations post-pandemic would not necessarily be conducive to quick recovery. On the other hand, there is much evidence that in general, the most common long-term response to exposure to highly aversive events is resilience (30).

As evidence emerges, prediction of future negative mental health consequences of the COVID-19 pandemic for healthcare workers will benefit from determining longitudinal trends in burnout and psychological distress among hospital workers. In particular, it would be useful to determine if negative mental health effects are rising monotonically over time, as one would expect to result from accumulating stress effects, or following a different pattern, possibly rising when COVID-19 cases are higher and improving as COVID-19 case rates decrease. Going forward, interventions to support healthcare workers' well-being could differ in these two scenarios.

Longitudinal data published thus far in the COVID-19 pandemic gives little insight into these long-term patterns. We note the following longitudinal studies, citing temporal trends rather than scores or case rates because these differ depending on operational definitions of burnout. Measures of burnout before and during the pandemic in intensive care settings have found increased burnout during the pandemic (21,31). An increase in burnout from April/May 2020 to July/August 2020 was found in oncology professionals (32), and from the fall of 2020 to the winter of 2021 in hospital personnel of all occupational types (33). Two studies have reported on multiple serial measurements over a period of approximately one month. Four measurements of burnout at weekly intervals in April 2020, revealed stable levels of burnout in emergency medicine providers (34). In contrast, burnout measured in physicians at five time-points over 25-31 days (following joining the COVID-19 treatment team) in Italy revealed variation over time (35). Studies of longer-term patterns of burnout and psychological distress during the pandemic have not yet been published.

We report on two mental health measures, namely the emotional exhaustion component of burnout and psychological distress, collected in the same cohort of hospital personnel at four time-points (approximately quarterly) from the fall of 2020 to the summer of 2021. Our primary question is whether indicators of these mental health problems are (i) rising monotonically, or (ii) following some other pattern, such as rising and falling in synchrony with the local epidemiological waves of COVID-19 cases. Our secondary questions are whether this pattern differs between personnel with different occupational roles, what proportion of hospital personnel participating in this survey are reporting levels of emotional exhaustion and psychological distress that are considered high at each time point, and how the depersonalization and personal achievement dimensions of burnout change over this period.

Methods

Study design and participants

A survey of the psychological well-being of a cohort of hospital staff, learners (nursing students, medical students, residents), and volunteers during the pandemic was conducted at two sites of Sinai Health (an urban acute care teaching hospital and a rehabilitation hospital, with > 6,000 employed staff) in Toronto, Canada at four time-points. The survey methods have been described previously (33). Briefly, the first survey (T1, “fall 2020”) was conducted from Sept 21-Nov 15, 2020. The first survey was open to all hospital employees, physicians, learners, volunteers, retail employees, and contractors. Potential participants were informed of an online survey link via emails from the hospital or from their chiefs and directors, as well as information posters in high-traffic areas of the hospital. All surveys were completed online using software that is compliant with privacy standards (the

Personal Health Information Protection Act, in this jurisdiction) (Alchemer, Louiseville, CO). Of 884 respondents who provided consent in a pre-survey recruitment phase, 538 (61.0%) completed a T1 survey to form the cohort for further follow-up. Subsequent surveys of this cohort were conducted in these time periods in 2021: Jan 25-Feb 15 (T2, “winter 2021”), Apr 26-May 16 (T3, “spring 2021”), and Jul 26-Aug 15 (T4, “summer 2021”). All members of the T1 cohort were invited to participate in all subsequent surveys, even if they were on leave or were not working at the hospital at the time of a survey. Each survey included measures of emotional exhaustion and psychological distress, among other measures. Participants were randomized to a shorter (Express) or longer (Enriched) version of the survey. Surveys varied in length over time due to instruments that were included at only some time-points. The total length of the Express survey varied from 45 to 83 items and the total length of the Enriched survey varied from 77 to 137 items. The participation rate at each time point (the numerator calculated as the number of surveys returned that included a valid measure of emotional exhaustion, psychological distress, or both) was: T2 N = 485 (90% of T1 cohort), T3 N = 424 (79%), T4 N = 409 (76%). A gift card (about US\$15 value) was provided for each completed survey. The study was approved by the Sinai Health Research Ethics Board.

Measures

Burnout was measured with the Maslach Burnout Inventory (MBI-HSS), which measures emotional exhaustion (9 items scored from 0-6, yielding a score from 0 to 54), depersonalization (5 items scored from 0-6, yielding a score from 0 to 30), and diminished personal achievement (8 items scored from 0-6, yielding a score from 0 to 48, with lower scores being worse) (36). Various case definitions of burnout based on MBI-HSS scores have been used previously, which differ in the choice of cut-off scores, and the choice of using the emotional exhaustion scale alone or various combinations of the three scales (27). In order to allow comparison to a broad range of prior studies, we used the emotional exhaustion scale as our primary continuous measure of burnout, and defined burnout using a cut-off of ≥ 27 , which is commonly used to identify high emotional exhaustion (27–29). In this cohort Cronbach’s alpha at the four time-points varied from .94 to .95. As secondary measures, to describe trends in other aspects of burnout we measured depersonalization and personal achievement in those participants randomly selected for the Enriched survey, using the most common cut-offs of depersonalization ≥ 10 and personal achievement ≤ 33 when reporting case numbers (27).

Psychological distress is comprised of depressive and anxiety symptoms measured as a screening test or as an index of severity of common mental disorders. Psychological distress was measured with the Kessler K6, which has 6 items scored from 0-4, yielding a range of 0-24. The K6 strongly discriminates between community cases and non-cases of psychiatric disorders diagnosed by structured interview and has acceptable sensitivity and specificity (37,38). A cut-off of ≥ 13 indicates likely serious mental illness (39). In this cohort Cronbach’s alpha at the four time-points varied from .85 to .87.

Analysis

At T1, participants were sorted into four categories of occupational role based on professional qualifications (i.e. classified as “professionals” if their job is regulated by a professional college, or equivalent) and whether they reported close patient contact (if they were within two metres of a patient for more than 15 minutes in the previous month). The four occupational role categories were nurses, other healthcare professionals (as listed in Table 1), other clinical staff (non-professionals who reported close patient contact), and non-clinical roles (non-professionals without close patient contact). Nurses were analyzed separately based on prior evidence that nurses experienced a greater burden of stress than other professionals during an outbreak of extraordinary infectious disease (26).

Participant characteristics were summarized using descriptive statistics. Continuous measures were summarized using means and standard deviations (SD). Categorical measures were summarized using counts and percentages. In order to provide context for changes in case rates of burnout and high psychological distress over time, daily rates of new cases of COVID-19 diagnosed in Toronto during the period of this study were downloaded from Public Health Ontario (40).

Changes over time in emotional exhaustion and psychological distress, as continuous variables, by occupational role were analyzed using repeated measures ANOVA, excluding subjects with missing data at any time point. Age was included as a covariate because lower age/less experience have been identified as correlates of higher burnout (41,42). Gender was not included because it is confounded with occupational role in hospital settings. Setting within the hospital (e.g. emergency department,

intensive care unit) was not included in the analysis because (i) many staff members work in multiple settings, and (ii) there were frequent dynamic changes in work settings for many staff members related to having multiple roles or having reassigned duties (e.g. medical nurses re-assigned to intensive care for a period of time and then returning to the medical unit). Within-subjects effects are reported using a Greenhouse-Geisser correction when Mauchly's test of sphericity indicated a significant deviation from the assumption of sphericity. All analyses were done with IBS SPSS Statistics 28 (Armonk, New York).

Results

The characteristics of personnel in the cohort at T1 are described in Table 1.

Table 1. Characteristics of participants

		N (%)
Role type*	Nursing	134 (24.9)
	Other clinical professionals	156 (29.0)
	Other clinical personnel	90 (16.7)
	Non-clinical personnel	158 (29.4)
Gender (Missing 1)	Female	422 (78.6)
	Male	85 (15.8)
	Other/Prefer not to say	30 (5.6)
Highest education	High school	13 (2.4)
	College diploma	79 (14.7)
	Undergraduate degree	176 (32.7)
	Professional/Graduate degree	270 (50.2)
Ethnic group (Missing 1)	African/Black	30 (5.6)
	Asian	148 (27.6)
	South Asian	35 (6.5)
	European/White	278 (51.8)
	Hispanic	15 (2.8)
Marital status	Other/Mixed/Multiple	31 (5.8)
	Single	211 (39.2)
	Married/Common-law	306 (56.9)
	Divorced/Separated/Widowed	21 (3.9)

**Specific job types, in descending order of number of participants. Groups with two or fewer members not listed. Some roles appear in both clinical and non-clinical lists as determined by patient contact as described by participant.*

Other clinical professionals: Physician, resident, dietician, occupational therapist, social worker, physiotherapist, manager of clinical area, speech language pathologist, pharmacist, respiratory therapist, spiritual care practitioner.

Other clinical positions: Administrative assistant, medical imaging technologist, assistant to physician/occupational therapist/physiotherapist, retail employee, porter, clinical research staff, volunteer.

Non-clinical positions: Research scientist, research staff, laboratory technician, corporate and administrative staff, administrative assistant, volunteer, manager of non-clinical area, building services staff, clerk, laboratory technologist, housekeeper.

Emotional exhaustion

Mean levels of emotional exhaustion for each occupational group at each time point are reported in Table 2. Mean levels peak at the winter 2021 or spring 2021 measurement point in each group and are lower by the summer 2021 measurement. Repeated measures ANOVA of emotional exhaustion for participants for whom there were no missing data over four time-points (82 nurses, 102 other healthcare professionals, 60 other clinical staff, and 118 non-clinical staff) demonstrated significant within-subjects changes in emotional exhaustion over time ($F = 4.6, p = .004$) and significant between-subject effects of occupational role ($F = 11.4, p < .001$) and age ($F = 12.3, p < .001$). The interaction of occupational role X time was not significant ($F = 1.4, p = .24$), indicating a similar pattern of change over time in all groups.

Table 2. Emotional exhaustion and psychological distress at four time-points in occupational groups.

Occupational Role	Time of survey	Emotional exhaustion		Psychological distress	
		Mean	Standard deviation	Mean	Standard deviation
Nurse	Fall 2020	28.8	12.4	7.4	4.2
	Winter 2021	32.4	12.6	8.5	4.2
	Spring 2021	31.3	13.7	8.7	4.5
	Summer 2021	30.6	13.4	7.1	4.7
Other healthcare professional	Fall 2020	24.4	12	5.3	3.6
	Winter 2021	28.3	11.9	6	4
	Spring 2021	29.8	11.5	6.7	3.9
	Summer 2021	25.3	11.9	5.3	3.8
Non-professional with patient contact	Fall 2020	24.9	13.2	6.8	5.3
	Winter 2021	25	12.7	6.8	5.1
	Spring 2021	27.3	13.1	8	6
	Summer 2021	22.2	12.5	6.2	5
Non-professional without patient contact	Fall 2020	22.4	11.6	6.7	4.3
	Winter 2021	24	12.2	7	4.3
	Spring 2021	24.4	12.6	6.9	4.5
	Summer 2021	20.9	12.2	5.9	4.4

Post hoc testing revealed that the relationship between age and emotional exhaustion was inverse (higher emotional exhaustion with younger age) and grew in strength over time (T1 $R = -.06, p = .14$, T2 $R = -.09, p = .05$, T3 $R = -.19, p < .001$, T4 $R = -.21, p < .001$). Regarding the direction and magnitude of differences in emotional exhaustion between staff with different occupational roles, the proportion of participants with high emotional exhaustion at each time point, by occupational role, is illustrated in Figure 1 (with the epidemiological curve of case rates in Toronto included for context). The rate of high burnout was highest in nurses, followed by other healthcare professionals, other clinical staff, and lowest in non-clinical staff.

Other burnout dimensions

Mean levels of depersonalization and personal achievement at each time point are provided in Table 3, as well as the proportion of participants whose depersonalization scores were high or personal achievement scores were low. Participants with high

depersonalization scores were rising over time, while the numbers with low personal achievement scores were falling. The number of subjects randomly selected to complete these extra measures (the 50% of subjects assigned to the Enriched survey), who completed measures of depersonalization and personal achievement at all four time-points included too few participants in one particular occupation category (non-professional staff without patient contact, N=8) to compare all occupational role groups. However, given the differences found for emotional exhaustion, we compared nurses (N= 41) to other healthcare professionals (N=55). With respect to depersonalization there were significant differences between these groups, with nurses reporting greater depersonalization ($F=7.4, p = .008$), with no significant effect of age. With respect to personal achievement there were also significant differences between nurses and other healthcare professionals ($F=9.6, p = .003$), with nurses reporting lower personal achievement, with no significant effect of age.

Table 3. Depersonalization and Personal Achievement dimensions of burnout at four time-points.

	Depersonalization				Personal Achievement			
	N	mean	SD	High (%)	N	mean	SD	High (%)
Fall 2020	277	4.80	5.59	18.1	270	34.42	10.99	37.0
Winter 2021	245	5.32	6.28	18.4	238	34.47	10.2	36.6
Spring 2021	154	6.42	6.86	23.4	148	37.85	7.05	20.3
Summer 2021	153	6.67	7.12	26.8	147	37.06	8.90	29.9

Psychological distress

Mean levels of psychological distress for each occupational group at each time point are reported in Table 2. Similarly to the results for emotional exhaustion, mean levels peak at the winter 2021 or spring 2021 measurement point in each group and are lower by the summer 2021 measurement. Repeated measures ANOVA of psychological distress for participants for whom there were no missing data over four time-points (77 nurses, 103 other healthcare professionals, 60 other clinical staff, and 110 non-clinical staff) demonstrated significant within-subjects changes in psychological distress over time ($F = 3.2, p = .024$) and significant between-subject effects of occupational role ($F = 3.6, p = .01$) and age ($F = 32.3, p < .001$). The interaction of occupation role X time was not significant ($F = 0.6, p = .83$), indicating a similar pattern of change over time in all groups.

Post hoc testing revealed that the relationship between age and psychological distress was inverse (higher psychological distress with younger age) and significant at each time point (T1 $R = -.24$, T2 $R = -.29$, T3 $R = -.24$, T4 $R = -.29, p < .001$ at each time point). The proportion of participants with high psychological distress at each time point, by occupational role, is illustrated in Figure 2. The severity of psychological distress was highest in nurses.

Discussion

In this longitudinal study of a single cohort of hospital-based health care workers over approximately one year during the COVID-19 pandemic, negative mental health indicators measured at three-month intervals changed significantly over time and varied between workers with different occupational roles. The severity of emotional exhaustion was greatest in nurses, followed by healthcare professionals of other disciplines, then non-professional staff whose work involves close patient contact, with the lowest levels in non-professional staff whose work does not involve close patient contact. Workplace factors that contributed to burnout in nurses prior to the pandemic, including high workload, lack of scheduling flexibility, and concern about workplace safety, have increased during the pandemic (43), which may contribute to nurses experiencing the highest levels of emotional exhaustion. In addition, the pandemic has been associated with the emergence of other stresses, including constraints on care that cause moral distress (21,22) and redeployment to unfamiliar types of work (23), which may have disproportionately affected nurses. Although not surprising, rates of high emotional exhaustion in nurses are nonetheless alarming. At the four measurement-points they were 54%, 62%, 63%, and 58% respectively, which is substantially greater than the pre-COVID-19 benchmark of 20-40% found using the same operational case definition (27). As nurses represent the largest single professional group in hospital care, this degree of mental health burden threatens the function of the healthcare system.

The rates of burnout in this study can be compared to those reported elsewhere, although most reports currently available relate to studies completed before 2021. In August – October 2020 a survey of Australian nurses using an abbreviated version of the MBI found high emotional exhaustion scores in 44% (44). A US study conducted over the same time span using a single item burnout measure found high burnout in 66% of nurses (45). A study of nurses in Uganda in May-June 2021 using the ProQOL instrument found high burnout in 40% (Kabunga & Okalo, 2021). The use of different measurement instruments precludes direct comparison, but these studies confirm multiple reports of high rates of burnout in nurses. A survey of internists at two hospitals in Vancouver, Canada from August – October 2020, found high emotional exhaustion in 63% using the same operational definition used in this study (47), which is consistent with our results. Burnout has not often been measured previously in hospital employees with roles other than regulated professionals, and so there is little historical context from which to interpret the degree of burnout found in that group.

The temporal pattern of change in emotional exhaustion did not differ significantly between occupational groups. Inspection of Figure 1 indicates that in three of the four groups this pattern consisted of a monotonic increase from fall 2020 to spring 2021, with a decrease from spring to summer 2021. The fourth group, non-professional staff with no close patient contact, appeared to differ only by beginning the descending trend sooner, peaking in winter 2021, an insignificant difference. Since the decrease in rates of high emotional exhaustion in summer 2021 corresponds to a period of low levels of new COVID-19 cases in the community, and corresponding low rates of COVID-19 related hospitalization, the trend may indicate that emotional exhaustion in hospital-based healthcare workers is able to recover somewhat as COVID-19 stresses subside. It is noteworthy that COVID-19 vaccination rates in the general public and in healthcare workers increased markedly in Ontario between the spring and summer 2021 surveys, which may have also contributed to reduced burnout. Importantly, the improvement in high emotional exhaustion during the summer of 2021 was an improvement to levels that remain much higher than the pre-COVID-19 benchmark.

Rates of high depersonalization and low personal achievement observed in this study are also concerning. Rates of depersonalization increased steadily over this time, which may indicate that depersonalization is a cumulative effect of pandemic related stress that is less responsive to decreasing case rates than emotional exhaustion. Our participants' sense of personal achievement may have been quicker to recover. Alternatively, the improvement in mean levels of personal achievement from winter 2021 to spring 2021 may have been the result of participants with low personal achievement scores in the winter 2021 dropping out. Post hoc testing reveals that at the winter 2021 measurement, the mean personal achievement scores of those who completed surveys in spring 2021 was 6.0 ± 7.0 , whereas the mean score in those who did not complete surveys in spring 2021 was 4.3 ± 4.9 ($p = .03$).

Psychological distress also changed over time in these hospital-based workers. As with emotional exhaustion, psychological distress was highest in nurses at each time point. Rates of psychological distress above the cut-off used in this study are clinically meaningful, because they indicate likely serious mental illness (39). Thus, it is concerning that the case rate of high psychological distress was 10.2% (13/128), 17.2% (20/116), 21.3% (20/94), and 13.1% (13/99) in nurses at the four measurement points. Rates of high psychological distress were lower in the other three occupational groups and were similar among these groups (the exception being spring 2021, when there was more variation). Although temporal patterns of psychological distress appeared more variable than the patterns of burnout, inspection of Figure 2 indicates that psychological distress was not monotonically rising in any occupational group.

The levels and patterns of emotional exhaustion and psychological distress measured in this hospital-based cohort raise concern for both the well-being of hospital-based healthcare workers and for the resilience of the healthcare system. Broad surveys of healthcare professionals in the same region during the COVID-19 pandemic indicate not only high rates of stress and burnout, but also that 43% of nurses surveyed were considering leaving healthcare work, and that this consideration was linked to feelings of high stress (48,49). While in other contexts the intention to leave one's profession does not necessarily translate into action, multiple media reports of emergency department closures, cancelled surgeries due to understaffing, and a "signing bonus" for new emergency department nursing hires, indicate that workforce shortages are emerging at this stage of the pandemic (50–54). This may be due to a vicious cycle of workplace conditions in which understaffing and increased workload are both a cause and consequence of high levels of emotional exhaustion, depersonalization, and psychological distress, as workers choose other alternatives (43).

Recognition of the mental health costs of healthcare work during the pandemic, and the related threat to the healthcare workforce, have led to recommendations for interventions to support healthcare workers that can be implemented at the level of healthcare systems and organizations, supplemented by individual support (43). Recommended interventions have included limiting shift lengths, maximizing scheduling flexibility, ensuring adequate training for unfamiliar tasks, providing support for moral distress, and promoting effective authentic leadership (43).

Our study's finding that lower age and, by implication, less experience in healthcare, is associated with greater emotional exhaustion and psychological distress is consistent with numerous previous studies (41,55–57). It is also relevant that younger adults may be more likely to have children at home, which is known to have added stress during the pandemic (33). Special efforts to support trainees and new graduates may include transition to practice programs that contain formal teaching and mentorship over several months (58). This consideration is especially relevant because widespread healthcare workforce shortages may lead to increases in training positions for healthcare professionals in order to increase the size of the pool of workers, leading to a younger healthcare workforce and a relative loss of senior mentors.

The strengths of this study include its prospective measurement of psychological distress and burnout with validated instruments over a one-year period in a single cohort of healthcare workers and its relatively high retention rate considering the extraordinary circumstances of hospital work during this time. It is also a strength that the survey extended to all workers and learners in the hospital, not just those in regulated professions. Its limitations include not using a sampling strategy that would ensure a representative sample, and evidence that psychological indices (i.e. burnout) differ in participants who completed all four surveys and those who did not. Additional considerations that may limit generalizability are the setting (an urban teaching hospital) and high rates of vaccination after the second survey. Despite enrolling participants based on their willingness to participate, the distribution of occupations and demographic measures suggest that most occupational groups are represented among the survey's participants.

Conclusion

In this cohort, neither emotional exhaustion nor psychological distress was rising monotonically, which provides hope that the negative mental health consequences of working in healthcare during the COVID-19 pandemic may improve as occupational stressors diminish. The depersonalization dimension of burnout, however, was found to rise consistently from one time point to the next. Burnout and psychological distress were consistently related to occupational role and were highest in nurses. Although emotional exhaustion improved as the case rate of COVID-19 decreased, rates of high emotional exhaustion remained much higher than was typically measured in hospital-based healthcare workers prior to the pandemic.

These results indicate that ongoing monitoring of healthcare workers' mental health is warranted to determine the rate of recovery of burnout and distress as the COVID-19 pandemic recedes. Attention to organizational and individual interventions to support healthcare workers to maintain the resilience of the health care system continues to be important.

Abbreviations

ANOVA Analysis of variance

COVID-19 Coronavirus disease

Declarations

Ethics approval and consent to participate

All participants provided consent to participate and the study was approved by the Sinai Health Research Ethics Board.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by a CIHR Canadian 2019 Novel Coronavirus (COVID-19) Rapid Research Grant (FRN 170642)

Authors' contributions

RGM, JJH, LPJ, LG, JJ, AK, CAL, LAW designed the study. RGM and NDH analyzed the data. All authors contributed to the interpretation of the data, provided critical feedback, and read and approved the final manuscript.

Acknowledgements

Not applicable

References

1. World Health Organization. Novel Coronavirus 2019 Interactive Timeline [Internet]. Vol. 2019. 2021. Available from: https://www.who.int/emergencies/diseases/novel-coronavirus-2019/interactive-timeline?gclid=CjwKCAjw7fuJBhBdEiwA2ILMYQPJ7BV5wKcj41OMFNsxfSQVg6hOMIJTAU8nuP9eKBoe7RXjov82fBoCJwsQAvD_BwE#
2. Worldometers. Worldometers Coronavirus [Internet]. Vol. 2021. 2021. Available from: <https://www.worldometers.info/coronavirus/worldwide-graphs/#daily-cases>
3. Lasalvia A, Amaddeo F, Porru S, Carta A, Tardivo S, Bovo C, et al. Levels of burn-out among healthcare workers during the COVID-19 pandemic and their associated factors: a cross-sectional study in a tertiary hospital of a highly burdened area of north-east Italy. *BMJ Open* [Internet]. 2021/01/19. 2021;11(1):e045127. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33455940>
4. Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsis E, Katsaounou P. Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain Behav Immun* [Internet]. 2020/05/22. 2020;88:901–7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32437915>
5. Shaukat N, Ali DM, Razzak J. Physical and mental health impacts of COVID-19 on healthcare workers: a scoping review. *Int J Emerg Med* [Internet]. 2020/07/22. 2020;13(1):40. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32689925>
6. Salari N, Khazaie H, Hosseini-Far A, Ghasemi H, Mohammadi M, Shohaimi S, et al. The prevalence of sleep disturbances among physicians and nurses facing the COVID-19 patients: a systematic review and meta-analysis. *Global Health* [Internet]. 2020/10/01. 2020;16(1):92. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32993696>
7. Dall'Ora C, Griffiths P, Ball J, Simon M, Aiken LH. Association of 12 h shifts and nurses' job satisfaction, burnout and intention to leave: findings from a cross-sectional study of 12 European countries. *BMJ Open* [Internet]. 2015/09/12. 2015;5(9):e008331. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26359284>
8. Hall LH, Johnson J, Watt I, Tsipa A, O'Connor DB. Healthcare Staff Wellbeing, Burnout, and Patient Safety: A Systematic Review. *PLoS One* [Internet]. 2016/07/09. 2016;11(7):e0159015. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27391946>
9. Dewa CS, Loong D, Bonato S, Thanh NX, Jacobs P. How does burnout affect physician productivity? A systematic literature review. *BMC Health Serv Res* [Internet]. 2014/07/30. 2014;14:325. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25066375>
10. Grossman Z, Chodick G, Kushnir T, Cohen HA, Chapnick G, Ashkenazi S. Burnout and intentions to quit the practice among community pediatricians: associations with specific professional activities. *Isr J Health Policy Res* [Internet]. 2019/01/06. 2019;8(1):2. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30609943>

11. Maslach C, Schaufeli WB, Leiter MP. Job burnout. *Annu Rev Psychol* [Internet]. 2001/01/10. 2001;52:397–422. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/11148311>
12. West CP, Dyrbye LN, Erwin PJ, Shanafelt TD. Interventions to prevent and reduce physician burnout: a systematic review and meta-analysis. *Lancet* [Internet]. 2016/10/04. 2016;388(10057):2272–81. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27692469>
13. Jun J, Ojemeni MM, Kalamani R, Tong J, Crecelius ML. Relationship between nurse burnout, patient and organizational outcomes: Systematic review. *Int J Nurs Stud* [Internet]. 2021/04/27. 2021;119:103933. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33901940>
14. Azam K, Khan A, Alam MT, Toqeer M. INTRODUCTION Causes and Adverse Impact of Physician Burnout: A Systematic Review. *Journal of the College of Physicians and Surgeons Pakistan*. 2017;27(8):495–501.
15. MacPhee M, Dahinten VS, Havaei F. The Impact of Heavy Perceived Nurse Workloads on Patient and Nurse Outcomes. *Administrative Sciences* 2017, Vol 7, Page 7 [Internet]. 2017 Mar 5 [cited 2021 Oct 31];7(1):7. Available from: <https://www.mdpi.com/2076-3387/7/1/7/htm>
16. Phillips C. Relationships between workload perception, burnout, and intent to leave among medical-surgical nurses. *International journal of evidence-based healthcare* [Internet]. 2020 Jun 1 [cited 2021 Oct 31];18(2):265–73. Available from: https://journals.lww.com/ijebh/Fulltext/2020/06000/Relationships_between_workload_perception,.11.aspx
17. Melnick ER, Dyrbye LN, Sinsky CA, Trockel M, West CP, Nedelec L, et al. The Association Between Perceived Electronic Health Record Usability and Professional Burnout Among US Physicians. *Mayo Clinic Proceedings* [Internet]. 2020 Mar 1 [cited 2021 Oct 31];95(3):476–87. Available from: <http://www.mayoclinicproceedings.org/article/S0025619619308365/fulltext>
18. Liu W, Zhao S, Shi L, Zhang Z, Liu X, Li L, et al. Workplace violence, job satisfaction, burnout, perceived organisational support and their effects on turnover intention among Chinese nurses in tertiary hospitals: a cross-sectional study. *BMJ Open* [Internet]. 2018 Jun 1 [cited 2021 Oct 31];8(6):e019525. Available from: <https://bmjopen.bmj.com/content/8/6/e019525>
19. Havaei F, Astivia OLO, MacPhee M. The impact of workplace violence on medical-surgical nurses' health outcome: A moderated mediation model of work environment conditions and burnout using secondary data. *International Journal of Nursing Studies*. 2020 Sep 1;109:103666.
20. Yoon HS, Sok SR. Experiences of violence, burnout and job satisfaction in Korean nurses in the emergency medical centre setting. *International Journal of Nursing Practice* [Internet]. 2016 Dec 1 [cited 2021 Oct 31];22(6):596–604. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/ijn.12479>
21. Kok N, van Gorp J, Teerenstra S, van der Hoeven H, Fuchs M, Hoedemaekers C, et al. Coronavirus Disease 2019 Immediately Increases Burnout Symptoms in ICU Professionals: A Longitudinal Cohort Study. *Crit Care Med* [Internet]. 2021/02/09. 2021;49(3):419–27. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33555778>
22. Sriharan A, West K, Almost J, Hamza A. COVID-19-Related Occupational Burnout and Moral Distress among Nurses: A Rapid Scoping Review. *Nursing Leadership (Toronto, Ont)* [Internet]. 2021 Mar 1 [cited 2021 Oct 24];34(1):7–19. Available from: <https://europepmc.org/article/med/33837685>
23. Morgantini LA, Naha U, Wang H, Francavilla S, Acar Ö, Flores JM, et al. Factors contributing to healthcare professional burnout during the COVID-19 pandemic: A rapid turnaround global survey. *PLOS ONE* [Internet]. 2020 Sep 1 [cited 2021 Oct 24];15(9):e0238217. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0238217>
24. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5)*. Fifth. Washington DC: American Psychiatric Association; 2013.
25. Li Y, Scherer N, Felix L, Kuper H. Prevalence of depression, anxiety and post-traumatic stress disorder in health care workers during the COVID-19 pandemic: A systematic review and meta-analysis. *PLOS ONE* [Internet]. 2021 Mar 1 [cited 2021 Oct 11];16(3):e0246454. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0246454>
26. Maunder RG, Lancee WJ, Balderson KE, Bennett JP, Borgundvaag B, Evans S, et al. Long-term psychological and occupational effects of providing hospital healthcare during SARS outbreak. *Emerging Infectious Diseases*. 2006;12(12).
27. Rotenstein LS, Torre M, Ramos MA, Rosales RC, Guille C, Sen S, et al. Prevalence of Burnout Among Physicians: A Systematic Review. *Jama* [Internet]. 2018/10/17. 2018;320(11):1131–50. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30326495>

28. Canadas-De la Fuente GA, Gomez-Urquiza JL, Ortega-Campos EM, Canadas GR, Albendin-Garcia L, de la Fuente-Solana EI. Prevalence of burnout syndrome in oncology nursing: A meta-analytic study. *Psychooncology* [Internet]. 2018/01/10. 2018;27(5):1426–33. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29314432>
29. Gomez-Urquiza JL, de la Fuente-Solana EI, Albendin-Garcia L, Vargas-Pecino C, Ortega-Campos EM, Canadas-De la Fuente GA. Prevalence of Burnout Syndrome in Emergency Nurses: A Meta-Analysis. *Crit Care Nurse* [Internet]. 2017/10/03. 2017;37(5):e1–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28966203>
30. Chen S, Bonanno GA. Psychological adjustment during the global outbreak of COVID-19: A resilience perspective. *Psychol Trauma* [Internet]. 2020/06/17. 2020;12(S1):S51–4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32538658>
31. Gomez S, Anderson BJ, Yu H, Gutsche J, Jablonski J, Martin N, et al. Benchmarking Critical Care Well-Being: Before and After the Coronavirus Disease 2019 Pandemic. *Crit Care Explor* [Internet]. 2020/11/03. 2020;2(10):e0233. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33134934>
32. Lim KHJ, Murali K, Kamposioras K, Punie K, Oing C, O'Connor M, et al. The concerns of oncology professionals during the COVID-19 pandemic: results from the ESMO Resilience Task Force survey II. *ESMO Open* [Internet]. 2021/07/04. 2021;6(4):100199. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34217129>
33. Maunder RG, Heeney ND, Kiss A, Hunter JJ, Jeffs LP, Ginty L, et al. Psychological impact of the COVID-19 pandemic on hospital workers over time: Relationship to occupational role, living with children and elders, and modifiable factors. *Gen Hosp Psychiatry* [Internet]. 2021/05/11. 2021;71:88–94. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33971518>
34. Kelker H, Yoder K, Musey Jr. P, Harris M, Johnson O, Sarmiento E, et al. Prospective study of emergency medicine provider wellness across ten academic and community hospitals during the initial surge of the COVID-19 pandemic. *BMC Emerg Med* [Internet]. 2021/03/26. 2021;21(1):36. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33761876>
35. Dionisi T, Sestito L, Tarli C, Antonelli M, Tosoni A, D'Addio S, et al. Risk of burnout and stress in physicians working in a COVID team: A longitudinal survey. *Int J Clin Pract* [Internet]. 2021/08/28. 2021:e14755. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34449957>
36. Maslach C, Jackson S, Leiter M. Maslach Burnout Inventory Manual [Internet]. Consulting Psychologists Press. 1996 [cited 2021 Oct 20]. Available from: <https://www.worldcat.org/title/maslach-burnout-inventory-manual/oclc/807246527>
37. Kessler RC, Andrews G, Colpe LJ, Mroczek DK. Short screening scales to monitor population prevalences and trends in non-specific psychological distress. 2002 [cited 2021 Nov 4]; Available from: <https://www.researchgate.net/publication/11174017>
38. Staples LG, Dear BF, Gandy M, Fogliati V, Fogliati R, Karin E, et al. Psychometric properties and clinical utility of brief measures of depression, anxiety, and general distress: The PHQ-2, GAD-2, and K-6. *Gen Hosp Psychiatry* [Internet]. 2018/12/07. 2019;56:13–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30508772>
39. Kessler RC, Barker PR, Colpe LJ, Epstein JF, Gfroerer JC, Hiripi E, et al. Screening for serious mental illness in the general population. *Arch Gen Psychiatry* [Internet]. 2003/02/13. 2003;60(2):184–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12578436>
40. Public Health Ontario. Ontario COVID-19 Data Tool | Public Health Ontario [Internet]. 2021 [cited 2021 Oct 24]. Available from: <https://www.publichealthontario.ca/en/data-and-analysis/infectious-disease/covid-19-data-surveillance/covid-19-data-tool?tab=trends>
41. Pulcrano M, Evans SRT, Sosin M. Quality of Life and Burnout Rates Across Surgical Specialties: A Systematic Review. *JAMA Surgery* [Internet]. 2016 Oct 1 [cited 2021 Oct 24];151(10):970–8. Available from: <https://jamanetwork.com/journals/jamasurgery/fullarticle/2533102>
42. Garrosa E, Moreno-Jiménez B, Liang Y, González JL. The relationship between socio-demographic variables, job stressors, burnout, and hardy personality in nurses: An exploratory study. *International Journal of Nursing Studies*. 2008 Mar 1;45(3):418–27.
43. Maunder RG, Heeney ND, Strudwick G, Shin HD, O'Neill B, Young N, et al. Burnout in Hospital-Based Healthcare Workers during COVID-19. 2021 Oct 7 [cited 2021 Oct 24]; Available from: <https://covid19-sciencetable.ca/sciencebrief/burnout-in-hospital-based-healthcare-workers-during-covid-19>
44. Smallwood N, Karimi L, Bismark M, Putland M, Johnson D, Dharmage S, et al. High levels of psychosocial distress among Australian frontline healthcare workers during the COVID-19 pandemic: a cross-sectional survey. *General psychiatry* [Internet].

- 2021 Sep 6 [cited 2021 Oct 24];34(5). Available from: <https://pubmed.ncbi.nlm.nih.gov/34514332/>
45. Melnyk B, Hsieh A, Tan A, Teall A, Weberg D, Jun J, et al. Associations Among Nurses' Mental/Physical Health, Lifestyle Behaviors, Shift Length, and Workplace Wellness Support During COVID-19: Important Implications for Health Care Systems. *Nursing Administration Quarterly*. 2021;Epub ahead of print.
 46. Kabunga A, Okalo P. Prevalence and predictors of burnout among nurses during COVID-19: a cross-sectional study in hospitals in central Uganda. *BMJ Open* [Internet]. 2021 [cited 2021 Oct 26];11:54284. Available from: <http://bmjopen.bmj.com/>
 47. Khan N, Palepu A, Dodek P, Salmon A, Leitch H, Ruzycki S, et al. Cross-sectional survey on physician burnout during the COVID-19 pandemic in Vancouver, Canada: the role of gender, ethnicity and sexual orientation. *BMJ open* [Internet]. 2021 May 10 [cited 2021 Oct 26];11(5). Available from: <https://pubmed.ncbi.nlm.nih.gov/33972345/>
 48. Ontario Medical Association. Healing the Healers: System-Level Solutions to Physician Burnout Recommendations of the Ontario Medical Association Burnout Task Force [Internet]. 2021 [cited 2021 Oct 31]. Available from: <https://www.oma.org/uploadedfiles/oma/media/pagetree/advocacy/health-policy-recommendations/burnout-paper.pdf>
 49. Registered Nurses Association of Ontario. Work and Wellbeing Survey Results [Internet]. 2021 [cited 2021 Oct 31]. Available from: https://rnao.ca/sites/rnao-ca/files/Nurses_Wellbeing_Survey_Results_-_March_31.pdf
 50. Signing bonuses are a sign of the times as desperate hospitals seek nurses | Ottawa Citizen [Internet]. [cited 2021 Oct 31]. Available from: <https://ottawacitizen.com/news/local-news/signing-bonuses-are-a-sign-of-the-times-as-desperate-hospitals-seek-nurses>
 51. Staff shortages at 2 N.B. hospitals lead to closures of labour and obstetrics units | CBC News [Internet]. [cited 2021 Oct 31]. Available from: <https://www.cbc.ca/news/canada/new-brunswick/hospital-labour-obstetrics-new-brunswick-1.6146501>
 52. Another Quebec ER had to shut down because of a lack of nurses, who are exhausted | CBC News [Internet]. [cited 2021 Oct 31]. Available from: <https://www.cbc.ca/news/canada/montreal/suroit-hospital-er-closes-overnight-1.6178786>
 53. Elective surgeries cancelled in Calgary, Fort Macleod emergency room closed as COVID-19 surge continues | CTV News [Internet]. [cited 2021 Oct 31]. Available from: <https://calgary.ctvnews.ca/elective-surgeries-cancelled-in-calgary-fort-macleod-emergency-room-closed-as-covid-19-surge-continues-1.5578056>
 54. Nova Scotia Health to close some surgery beds, redeploy nurses to areas with critical needs | CBC News [Internet]. [cited 2021 Oct 31]. Available from: <https://www.cbc.ca/news/canada/nova-scotia/health-care-nurses-emergency-department-staffing-1.6151994>
 55. Jalili M, Niroomand M, Hadavand F, Zeinali K, Fotouhi A. Burnout among healthcare professionals during COVID-19 pandemic: a cross-sectional study. *International Archives of Occupational and Environmental Health* 2021 94:6 [Internet]. 2021 Apr 17 [cited 2021 Oct 31];94(6):1345–52. Available from: <https://link.springer.com/article/10.1007/s00420-021-01695-x>
 56. Galanis P, Vraika I, Fragkou D, Bilali A, Kaitelidou D. Nurses' burnout and associated risk factors during the COVID-19 pandemic: A systematic review and meta-analysis. *Journal of Advanced Nursing* [Internet]. 2021 Aug 1 [cited 2021 Oct 31];77(8):3286–302. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/jan.14839>
 57. Abdelhafiz AS, Ali A, Ziady HH, Maaly AM, Alorabi M, Sultan EA. Prevalence, Associated Factors, and Consequences of Burnout Among Egyptian Physicians During COVID-19 Pandemic. *Frontiers in Public Health*. 2020 Dec 3;0:864.
 58. Brook J, Aitken L, Webb R, MacLaren J, Salmon D. Characteristics of successful interventions to reduce turnover and increase retention of early career nurses: A systematic review. *International Journal of Nursing Studies*. 2019 Mar 1;91:47–59.

Figures




Figure 1

Burnout by occupational role over time compared to daily COVID-19 case rates




Figure 2

Psychological distress by occupational role over time compared to daily COVID-19 case rates.