Program on high value cost-conscious education in Intensive Care: Educational program on prediction of outcome and cost awareness on Intensive Care admission

Linda van Wageningen (✉ l.vanwagenberg@umcutrecht.nl)

Universitair Medisch Centrum Utrecht <https://orcid.org/0000-0003-0702-6265>

Charlotte J. P. Beurskens

Amsterdam Universitair Medische Centra

Inge Stegeman

UMC Utrecht Brain Center Rudolf Magnus

Marcella C. A. Müller

Amsterdam Universitair Medische Centra

Research article

Keywords: High-value cost-conscious care, post-graduate education, curriculum development, evaluation of education

Posted Date: January 24th, 2020

DOI: <https://doi.org/10.21203/rs.2.20045/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Medical Education on June 8th, 2020. See the published version at <https://doi.org/10.1186/s12909-020-02100-w>.

Abstract

Introduction: Intensive Care (ICU) involves extended and long lasting support of vital functions and organs. However, current training programs of ICU residents mainly focus on extended support of vital functions and barely involve training on cost-awareness and outcome. We incorporated an educational program on high-value cost-conscious care for residents and fellows on our ICU and measured the effect of education. **Methods:** A cohort study with factorial survey design, in which residents and fellows ICU were asked to evaluate clinical vignettes, was performed on the mixed surgical-medical ICU of the Amsterdam University Medical Centre. Residents were offered an educational program focusing on outcome and costs of ICU care. Before and after the program they filled out a questionnaire, which consisted of 23 vignettes, in which known predictors of outcome of community acquired pneumonia (CAP), pancreatitis, acute respiratory distress syndrome (ARDS) and cardiac arrest were presented, together with varying patient factors (age, body mass index (BMI), acute kidney failure (AKI) and haemato-oncological malignancy). Participants were asked to either admit the patient or estimate mortality. **Results:** BMI, haemato-oncological malignancy and severity of pancreatitis were discriminative for admission to ICU in clinical vignettes on pancreatitis and CAP. After education, only severity of pancreatitis was judged as discriminative. Before the intervention only location of cardiac arrest (in- vs out of hospital) was distinctive for mortality, afterwards this changed to presence of haemato-oncological malignancy. **Conclusion:** We incorporated an educational program on high-value cost-conscious care in the training of ICU physicians. Based on our vignette study, we conclude that the improvement of knowledge of costs and prognosis after this program was limited.

Introduction

A growing elderly population, healthcare innovations and new treatment possibilities lead to a yearly rise in health care costs.¹ Intensive Care Units (ICU) offer opportunities for extended and long-lasting support of vital functions and organs. However, this type of care is expensive and does not automatically result in clinically beneficial outcomes.²

Careful decisions on the usage of critical care can only be made when physicians are fully acquainted with the potential clinical outcomes as well as the costs of ICU care. However, the level of awareness regarding the costs of common prescription drugs (especially high cost medication), and commonly used materials in the ICU is often lacking, as described by Hernu et al.³ Moreover, ICU physicians generally overestimate the long-term survival and quality of life of ICU survivors.⁴ This lack of knowledge may be attributed to the lack of training about cost awareness and outcome of ICU treatment in the current postgraduate medical training program.⁵

The Netherlands has one of the best healthcare systems in the world,⁶ however, maintaining this high-quality position is expensive, requiring about 10% of the Dutch Gross National Product in 2017.⁷ Wammes et al. showed in 2017 that 9,1% of the total cost for hospital health care is spend on ICU care.⁸

This makes ICU care very expensive, as only a small number of all Dutch patients were treated in the ICU. It seems only fair to educate future ICU physicians on the outcome of care and its related costs.

The Awareness project was launched in the Netherlands in 2017-2018 by the Federation of Medical Specialists and Maastricht University, to incorporate expedience and cost awareness in addition to the CANMED competencies in all post graduate medical training programs. Its goals were to preserve high quality care, while at the same time creating more awareness regarding the appropriateness of care and raise cost-consciousness.⁹ This is also called High Value Cost-Conscious care. All regional medical education programs were invited to apply for projects on these items. Hence, an educational program for ICU residents and fellows enhancing knowledge on outcome and cost awareness of ICU treatment was created at the Intensive Care Unit of the Amsterdam University Medical Centre.

The present study aims to assess the effect of this educational program on the knowledge of treatment outcomes and associated costs of ICU support.

Methods

Design

In this cohort study we used a factorial survey design in which participants were asked to evaluate clinical vignettes. Hereby importance of factors influencing decision making could be assessed. A vignette is a brief, written case history of a fictitious patient that is based on a realistic clinical situation. In the vignettes the factors of interest (in our study possible predictors of ICU outcome) were varied between the different vignettes, in which each combination was unique. We made combinations between categories of predictors, in which only one factor changed between each vignette. These vignettes were presented to the participant, residents and fellows in our ICU, in an online questionnaire using the Survey Monkey website.¹⁰ Formal approval of the institutional ethics committee was not requested, participants were informed about the anonymized use of the test results and participation was voluntary. Due to the nature of the project no sample size calculations were made.

The online questionnaire consisted of 23 clinical vignettes, in which four common ICU admission diagnoses were presented; community acquired pneumonia (CAP), pancreatitis, acute respiratory distress syndrome (ARDS) and cardiac arrest. The literature was studied for patient factors known to have impact on patient outcome in ICU for these conditions. With this information an expert panel of ICU physicians decided to add the following factors to the vignettes: age, body mass index (BMI), acute kidney injury (AKI) and presence of haematological malignancy for all diseases, and presence of chronic obstructive pulmonary disease (COPD) and severity of pneumonia measured by CURB-65 score for pneumonia, presence of chronic liver disease and severity of pancreatitis measured by Ranson score for pancreatitis, severity of ARDS and type of ARDS (distinguishing primary and secondary ARDS) for ARDS and for the cardiac arrest cases first rhythm (shockable versus non-shockable), location of arrest (in hospital versus out of hospital) and delay until start CPR.¹¹⁻¹⁸ All community acquired pneumonia cases were classified

as severe, according to the CURB-65 score¹¹. In cases for ARDS and cardiac arrest all patient had a BMI of 20-25 and an age of 60-80 years old. A complete overview of all factors involved is shown in table 1.

With the complete set of factors, 324 vignettes for community acquired pneumonia (2x2x3x3x3x3), 432 vignettes for pancreatitis (2x2x3x3x4x3), 288 vignettes for ARDS (2x2x3x3x2x4) and 288 vignettes for cardiac arrest (2x2x3x3x2x2x2) were created. Completing all 1332 vignettes would be too time-consuming for participants. All unrealistic vignettes were removed and a total of 23 clinically vignettes (6 pneumonia, 7 pancreatitis, 4 ARDS and 6 cardiac arrest) were randomly selected by a team of experts/ICU clinicians based on their realistic scenarios. In each vignette only one factor changed to the next vignette. The vignettes were written by one of the investigators (LW) and discussed by the other investigators (MM and CB) on clinical accuracy and realism. The complete questionnaire can be found in Supplement 1.

For the vignettes on community acquired pneumonia and pancreatitis, participants were asked to choose between admission or no admission to the ICU, taken into account chances on survival and outcome. For all vignettes with ARDS or cardiac arrest participants were asked to estimate mortality during first 30 days of ICU stay. The options were < 40% mortality, 40-80% mortality or > 80% mortality.

Also, five questions were added on the cost of five products, which are frequently used in the ICU. Each question was multiple choice and respondents were asked to pick the right price for the product.

Demographics of the participants were collected at the start of the questionnaire. Recorded variables included working experience in ICU care, previous experience in residency or fellowship and primary postgraduate medical education programs. The first questionnaire was carried out in January 2018, the second, after the educational intervention, in April 2018.

The questionnaire before and after the educational intervention was similar.

Setting

This study was executed at the Intensive care department of the Amsterdam University Medical Center, location Academic Medical Center, at the University of Amsterdam. The department is a 34-bed mixed medical-surgical ICU, where residents of various postgraduate medical education programs are trained (e.g. internal medicine, anaesthesiology, surgery, neurosurgery, cardiology, emergency medicine). In addition, yearly 7 fellows are trained to become an intensivist as a subspecialty of their training in anaesthesiology, internal medicine, cardiology or neurology.

Participants

The study population consisted of ICU residents with medical training in anaesthesiology, internal medicine, emergency medicine, neurosurgery, cardiothoracic surgery or general surgery, and fellows in Intensive Care Medicine, with a medical specialization in internal medicine, anaesthesiology or neurology. All physicians were employed on the ICU of the Amsterdam Medical Centre, a tertiary clinic, during the study period.

Intervention

Between January and April 2018 an educational program on outcome and cost of ICU treatment was implemented in the regular educational program. This program consisted of flipped classroom sessions on outcomes of patients admitted to the ICU with a certain illness, comorbidity or patient characteristic; lectures on costs of ICU care and cost reduction and organization of ICU and a weekly quiz.¹⁹ The flipped classroom sessions focused on COPD, haematological malignancies, pancreatitis, community acquired pneumonia, cardiac arrest, age and BMI. In total eight different flipped classroom sessions were organized. The classical lectures focused on organization of ICU care, assessing the outcomes of ICU care, cost reduction on laboratory measurements and financing ICU care. Since all of our physicians work on irregular shifts all lectures were held twice and handouts of the lectures and acquired information of the flipped classroom sessions were shared between all physicians.

The weekly quiz with a fictional but realistic case and a question on the actual costs of a certain investigation, medication or material was sent to all participants every Monday. After the first week a ranking scale was made with all participating competitors and updated every week.

Potential effect modifiers, confounders and bias

Potential modifiers of the measured effect of the intervention are previous experience in ICU care, outcome of ICU care, cost and already gathered knowledge on high-value cost-conscious care during general medical training or residency. For this reason, we included all residents and fellows employed at our ICU during the intervention and asked about their level of experience in ICU care. There was risk of inclusion bias, since participation was voluntary and this could select only participants interested in the subject of the study. To prevent selection bias we briefed all residents and fellows about the study and potential of high-value cost-conscious care education to gain interest in the study. We tried to minimize loss of follow up by asking to complete the final questionnaire on several occasions.

Statistical analysis

Descriptive statistics were used to describe the characteristics of the participating residents. We used logistic regression to determine the first outcome, relative importance of the factors for ICU admission, in which ICU admission was set as determinant and the factors as independent variables. Data are shown as odds ratios (OR) with 95% confidence interval.

For the second outcome, the importance of the factors for the estimate of mortality, we used univariate multi-nominal regressing, with percentage of survival as determinant and the factors as independent variables. Data are shown as odds ratios (OR) with 95% confidence interval.

The third outcome, the estimation of costs of products used regularly in ICU, was noted a percentage different from the true cost. Results are shown as means with standard error of the mean (SEM). A students t-test was performed to compare answers before and after education.

IBM Spss version 25 was used for the statistical analyses and a p value of < 0.05 was considered statistically significant.

Results

Participants

All residents and fellows employed at our ICU during the intervention were included in the study. The pre-education online questionnaire was completed by 32 (100%) participants and the post-education questionnaire by 27 (84%) participants. Loss to follow up of 5 participants was due to termination of the internship at ICU, after which they changed hospitals. There was no missing data. All the participants answered all the questions in the questionnaire. The majority of respondents had a background in anaesthesiology (56% and respectively 56%) and Internal medicine (25% and respectively 26%). Nine respondents had over one year of experience in Intensive Care, while four had no experience at all (table 2).

Predictive factors for ICU admission and mortality

Patients with a haematological malignancy (OR 2.084 (1.257-3.454), $p=0.04$), BMI > 25 (OR 2.931 (1.008-8.256), $p=0.04$) were more likely to be admitted to the ICU prior to the educational program and those with a pancreatitis with a Ranson score of 5-6 were less likely to be admitted to the ICU (OR 0.054 (0.019-0.156), $p< 0.01$), compared to pancreatitis with Ranson score 3-4. After the educational program the effects of factors haematological malignancy and BMI > 25 were attenuated. The effect of the Ranson score 5-6 remained after education (OR 0.129 (0.050-0.034), $p<0.01$) (table 3).

Outcomes of the multinomial regression for factors potentially influencing the estimate of mortality are shown in table 4. Prior to the educational program location of cardiac arrest (in- vs. out of hospital cardiac arrest) was perceived to be associated with mortality (estimated mortality $> 80\%$: OR 9.274 (1.2-

71.697), $p=0.033$). After the educational program only the presence of a haematological malignancy was considered to be an important factor in the estimation of mortality (estimated mortality 40-80%: OR 4.320 (2.089-8.933), $p<0.01$ and estimated mortality >80%: OR 3.124 (1.497-6.519), $p=0.02$).

Cost of ICU care

All participants filled out all the questions on costs of ICU products. After the educational program the questions on costs showed reduced deviation of the true costs (Figure 1). Answers on CT-brain (pre-education mean percentage $113.5\% \pm 10.8$, post-education $106.5\% \pm 8.4$, $p=0.049$) and immune-modulating enteral food (Impact) (pre-education $42.75\% \pm 3.7$, post-education $61.52\% \pm 5.7$, $p=0.0063$) both significantly showed reduced deviation in regard to the true product cost after education.

Discussion

We implemented an educational program for residents and fellows on the outcome of ICU care in a tertiary academic ICU in the Netherlands. The program was designed to enhance knowledge on the outcome and mortality of the most common ICU diagnoses and comorbidities. The program consisted of flipped class room sessions on outcome of care, classical lectures on benchmarking ICUs, organization of ICU and costs of ICU care, and a weekly quiz on costs of products often used in our ICU.

We used a clinical vignette study to evaluate the learning effect of our educational program. A vignette study can show which factors are pivotal in the decision whether to admit a certain patient to the ICU.

Interpretation

Participating physicians showed to make different decisions pre- versus post-education. A BMI over 25 and presence of a haematological malignancy were considered an important factor for ICU admission pre-education, however this effect was not measured after the educational program. In the program residents were taught that a BMI > 25 can be protective in certain situations.¹⁶⁻¹⁸ In addition, they also learned that the presence of any haemato-oncological disease results in high morbidity and mortality, and therefore reduces a positive effect of ICU care. The net effect of this increased awareness resulted in a lower odds ratio when considering a haematological-oncological disease as a factor for admittance after the educational program (table 3).²⁰⁻²² However, increased awareness on factors that influence the outcome of ICU admission was limited, and did not improve after the educational program. The latter was shown by the pancreatitis vignette where patients with more severe disease (expressed as Ranson's criteria) were less likely to be admitted to the ICU. The educational program did lead to a shift in knowledge on predictive factors in case of mortality in ARDS or cardiac arrest. While pre-education, the

location of the cardiac arrest (in hospital versus out of hospital) was marked as the most important factor, post education the most important factor shifted to the presence of a haematological malignancy.

The current project was carried out as a part of the Dutch Awareness project, which is a successor to the Choosing Wisely campaign, launched by the American Boards of Internal Medicine (ABIM) Foundation in 2012.²³ The influence of the latter was evaluated by Rosenberg in 2015, who studied the volume of change of seven low-value services mentioned in the Choosing Wisely campaign.²⁴ Low-value services are considered avoidable treatments and tests that are unnecessary or harmful. In only two recommendations a small decrease could be found (imaging for headache and cardiac imaging without history), showing the difficulty of implementing high-value cost-conscious care. Despite the short duration of our study and its small sample size, our study shows the same difficulty in implementing high-value cost-conscious care. In order to achieve a health system in which high-value cost-conscious care becomes the default mindset, a need for cultural change and involvement of the medical societies may be required as described by Kerr in 2017.²⁵ To deliver high-value cost-consciousness care, the theme should be implemented in (post-graduate) medical training programs and incorporated in daily work regimes in order to reach maximum effect.. A systematic review on how to educate physicians to deliver high-value, cost-conscious care demonstrated that effective transmission of knowledge, reflective practice, and a supportive environment with role-models contribute to the learning potential of physicians.²⁶

Limitations

There are several limitations to our study. For feasibility reasons the number of clinical vignettes was set to 23. In creating the vignettes only one of the factors could change between each case, this limited the number of factors used in our vignettes. Also, the range in answers was smaller than expected, this resulted in insufficient data to perform statistical analysis for some factors or categories. Future studies should therefore include a larger number of participants, so more vignettes can be developed and divided between participants.

The residents completing the questionnaire probably also experienced a learning effect from their ICU internship next to the supplied additional educational program. We cannot differentiate between the acquired knowledge from the educational program and the effect of their day to day job (being on call, treating patients etc.). Also, while this program was scheduled during normal working hours, some residents could not attend all classes. To this extent, all lectures were available online to the whole group, also those who did not attend the class in person. However, we do not have any data on which lectures were reviewed by those who did not attend. Hence, we were not able to relate the level of participation to the eventual learning effect of the program.

Decisions on whether a patient is admitted to the ICU are based on the expected benefit of admission. Moreover overuse and inappropriate use of ICU care should be limited as much as possible. The decision

to admit a patient is often based on clinical judgement, combined with knowledge on predictors of outcome. Valley et al. studied the consensus between physicians in the United States on ICU admission using clinical vignettes.²⁷ The consensus on whether to admit a patient was poor in this study. At most, only 69% of physicians agreed about the extent to which a patient would benefit from ICU care. The lack of overall agreement between ICU physicians is worrisome and may also lead to disproportionate health care costs. It also indicates that the question whether to admit a patient to ICU, aiming benefit for the patient, is not an easy one to answer.

Conclusion

The basic knowledge on ICU mortality and contributing factors is limited in residents and fellows at our ICU and there is a need to improve this knowledge to improve cost-conscious decision making in an era of rising health care costs. Our study shows the possible implications of an educational program focusing on knowledge about ICU admission. Based on our vignette study, we conclude that the improvement of knowledge of costs and prognosis after this program was limited. With this information we would like to improve our educational program and assess the outcomes within a larger sample size.

List Of Abbreviations

ABIM	American Board of Internal Medicine
ARDS	Acute Respiratory Distress Syndrome
AKI	Acute Kidney Injury
BMI	Body Mass Index
CAP	Community Acquired Pneumonia
COPD	Chronic Obstructive Pulmonary Disease
CPR	Cardiopulmonary Resuscitation
CT	Computed Tomography
ICU	Intensive Care Unit
OR	Odds Ratio
SEM	Standard Error of the Mean

Declarations

Ethics approval and consent to participate

Formal approval of the institutional ethics committee was not requested. Participants were informed about the anonymized use of the test results and participation was voluntary. All our residents and fellows were informed about this educational program and adjacent study design prior to the start of the study. We gave them the opportunity to object objected against the use of their test results or data, whenever they wanted.

The intervention was part of the educational sessions all residents and fellows attend during their employment at the ICU. We considered the online questionnaire as part of this, therefore we did not consult the ethics committee.

Consent for publication

Not applicable

Availability of data and materials

All data is anonymised. The data used and/ or analyzed can be available by the corresponding author on request.

Competing interests

The authors declare that they have no competing interests

Funding

No funding was used for this research

Authors' contributions

LW was involved in the design of the study and wrote the vignettes used for the study. LW was partially involved in teaching during the intervention and analyzed the data. LW was involved in making the draft of the manuscript.

CB was involved in the design of the study and improved the vignettes. CB was involved in the statistical analysis and drafted the manuscript.

IS was involved in the design of the study and statistical analysis of the results. IS was involved in drafting the manuscript.

MM was involved in the design of the study and improved the vignettes. MM was involved in teaching during the intervention and analyzed the data. MM was involved in drafting the manuscript.

Acknowledgements

We would like to thank all of the residents and fellows for their kind participation. We would like to thank K.E.M. Benders, MD PhD, for her support in English medical writing.

References

1. Lindemark F., Haaland ØA., Kvåle R., Flaatten H., Norheim OF., Johansson KA. Costs and expected gain in lifetime health from intensive care versus general ward care of 30,712 individual patients: a distribution weighted cost-effectiveness analysis. *Critical Care* 2017; 21:220-32
2. Talmo D., Shapiro N., Greenberg D., Stone PW., Neumann PJ. When is critical care medicine cost-effective? A systematic review of the cost-effectiveness literature. *Crit. Care* 2006 ;34:2738-47
3. Hernu R., Cour M., de la Salle S., Robert D., Argaud L. Cost awareness of physicians in intensive care units: a multicentric national study. *Intensive Care Med.* 2015;41:1402-10
4. Soliman IW., Cremer OL., de Lange DW. et al. The ability of intensive care unit physicians to estimate long-term prognosis in survivors of critical illness. *Journal of Critical Care* 2018;43:148-155.
5. CoBaTrICE, retrieved March 15th 2019 www.cobatrice.org/en/index/asp
6. Björnberg A. EuroHealth Consumer Index 2017. Health Consumer Powerhouse 2018
7. OECD Health Statistics 2018, retrieved January 13th 2019: <http://www.oecd.org/els/health-systems/health-data.htm>
8. Wammes JG., Tanke M., Jonkers W., Westert GP., van der Wees PJ. and Jeurissen PPT. Kenmerken en zorgverbruik van de patiënten met de hoogste zorgkosten. *Ned. Tijdschr Geneesk.* 2017;161:D2220
9. Bewustzijnsproject (Dutch Awareness Project), retrieved January 13th 2019 bewustzijnsproject.nl
10. Survey Monkey, retrieved January 13th 2019 <https://nl.surveymonkey.com>
11. Lim WS., van der Eerden MM., Laing R. et al. Defining community acquired pneumonia severity on presentation to hospital: an international derivation and validation study. *Thorax* 2003; 58: 377-382.
12. Ranson JH., Rifkind KM., Roses DF., Fink SD., Eng K., Spender FC. Prognostic signs and the role of operative management in acute pancreatitis. *Surg Gynecol Obstet.* 1974;139:69-81.
13. ARDS Definition Task Force: Ranieri VM, Rubenfeld GD, Thompson BT, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012;307:2526-33.
14. Martens PR., Mullie A., Calle P. van Hooijweghen R. Influence on outcome after cardiac arrest of time elapsed between call for help and start of bystander basic CPR. *Resuscitation* 1193;25:227-234
15. Hirlekar, Karlsson T., Aune S. et al. Survival and neurological outcome in the elderly after in-hospital cardiac arrest. *Resuscitation* 2017;118 101-106.
16. Pickkers P., de Keizer N., Dusseljee J., Weerheijm D., van der Hoeven JG., Peek N. Body Mass index is associated with hospital mortality in critically ill patients; an observational cohort study. *Care Med.* 2013;41: 1878-1883.
17. Ni YN., Luo J., Yu H. et al. Can body mass index predict clinical outcome for patients with acute lung injury/ acute respiratory distress syndrome? A meta-analysis. *Critical Care* 2017; 21: 36-44.

18. Tremblay A. and Bandi V. Impact of Body Mass index on outcomes following critical care. *Chest* 2003;123:1202-1207
19. Morgan H., McLean K., Chapman C., Fitzgerald J., Yousuf A. Hammoud M. The flipped classroom for medical students. *Clin Teacher* 2015;12: 155-160.
20. Hampshire PA., Welch CA., McCrossan LA., Francis K., Harrison DA. Admission factors associated with mortality in patients with haematological malignancy admitted to UK adult, general critical care units: a secondary analysis of the ICNARC Case Mix programme database. *Critical care* 2009;13,R137.
21. Azoulay E., Mokart D., Pene F. et al. Outcomes of critically ill patients with hematologic malignancies: prospective multicenter data from France and Belgium. *Journal of Clinical Oncology* 2013;31:2810-2818
22. Guideline Dutch Association of Intensive Care: Intensive Care opname, behandeling en ontslag van de kritisch zieke hemato-oncologische patient. *NVIC* 2016.
23. Morden NE., Colla CH., Sequist TD., Rosenthal MB. Choosing Wisely – The Politics and Economics of Labelling Low-Value Services. *NEJM* 2014;370:589-92.
24. Rosenberg A., Agiro A., Gottlieb M. et al. Early trends among seven recommendations from the Choosing Wisely campaign. *JAMA internal Medicine* 2015;175:1913-1920.
25. Kerr EA., Kullgren JT., Saini SD. Choosing Wisely: How to fulfil the promise in the next 5 years. *Health affairs* 2017;11:2012-2018.
26. Stammen LA., Stalmeijer RE., Paternotte E. et al. Training Physicians to Provide High-Value, Cost-Conscious Care. *JAMA*. 2015;314:2384-2400.
27. Valley TS., Admon AJ., Zahuranec DB., Garland A., Fagerlin., Washyna TJ. Estimating ICU benefit: A randomized study of physicians. *Crit. Care Med.*2019; 47:62-68.

Tables

Table 1: Variables in the clinical vignettes		
All cases	Presence of haematological malignancy	Yes/no
	Acute kidney failure	Yes/no
	Age	< 60 years/ 60-80 years/ >80 years
	BMI	< 20 / 20-25 / >25
Community acquired pneumonia	Presence of COPD	Absent / mild / severe
	Severity of Pneumonia (measured CURB -65 score)	CURB-65 score 0-1 / 2 / 3-5
Acute Pancreatitis	Presence of chronic liver disease	Absent/ Liver cirrhosis Child Pugh A/ B/ C
	Severity of pancreatitis, measured by Ranson score	Ranson Score 3-4/ 5-6/ 7-8
ARDS	Type of ARDS	Primary/ Secondary
	Severity of ARDS	None/ Mild/ Moderate/ Severe
Cardiac arrest	First rhythm	Shockable (VF/ VT)/ Non-shockable (PEA/ Asystole)
	Location of arrest	Out of hospital cardiac arrest/ In hospital cardiac arrest
	Delay until start CPR	< 3 minutes/ >3 minutes
BMI= body mass index, COPD = chronic obstructive pulmonary disease, CURB-65 score = confusion, urea, respiratory rate and blood pressure- score, ARDS = acute respiratory distress syndrome, VF= ventricular fibrillation, VT= ventricular tachycardia, PEA= pulseless electric activity, CPR= cardiopulmonary resuscitation		

Table 2: Demographics of participants of the online questionnaire			
Background		pre-education n=32 (100%)	post-education n=27 (84%)
	Anaesthesiology	18 (56%)	15 (56%)
	Internal Medicine	8 (25%)	7 (26%)
	Emergency Medicine	1 (3%)	1 (4%)
	Cardiology	2 (6%)	2 (7%)
	Neurology	1 (3%)	1 (4%)
	Cardiothoracic surgery	1 (3%)	1 (4%)
	Neurosurgery	1 (3%)	0 (0%)
Year of residency			
	1 st year of residency	0 (0%)	1 (4%)
	2 nd year of residency	10 (31%)	6 (22%)
	3 rd year of residency	14 (44%)	13 (48%)
	Fellowship ICU	8 (25%)	7 (26%)
Experience in ICU care			
	None	4 (13%)	3 (11%)
	0-3 months	6 (19%)	2 (7%)
	3-6 months	8 (25%)	8 (30%)
	6-12 months	5 (16%)	5 (19%)
	More than 1 year	9 (28%)	9 (33%)
ICU= Intensive Care Unit			

Table 3: Odds ratios for the admission of the patient to the ICU.					
		Pre-education		Post-education	
Factor		OR (95% CI)	p-value	OR (95% CI)	p-value
Presence of haematological malignancy		2.084 (1.257-3.454)	0.004	1.021 (0.582-1.789)	0.943
Acute kidney failure		0.694 (0.427-1.127)	0.140	0.803(0.465-1.388)	0.433
Age	<60 years	reference		reference	
	60-80 years	0.576 (0.231-1.438)	0.237	0.584 (0.214-1.590)	0.292
	>80 years	0.444 (0.142-1.394)	0.164	0.543 (0.153-1.931)	0.346
BMI > 25		2.931 (1.008-8.526)	0.048	3.228 (0.950-10.969)	0.060
Presence of COPD		0.890 (0.560-1.414)	0.622	0.866 (0.538-1.393)	0.553
Severity of Pancreatitis, measured by Ranson score ¹²					
	3-4	Reference			
	5-6	0.054 (0.019-0.156)	0.000	0.129 (0.050-0.334)	0.000
	7-8	0.664 (0.353-1.247)	0.203	0.626 (0.289-1.354)	0.234
Previous history of liver disease		0.633 (0.782-1.161)	0.633	1.000 (0.503-1.987)	1.000
OR= odds ratio, CI= confidence interval, BMI= body mass index, COPD= chronic obstructive pulmonary disease. The OR represents the average odd that the patient is likely to be not admitted to the ICU					

Table 4: Factors of importance in the estimation of mortality in ARDS and cardiac arrest								
	Pre- education				Post- education			
	Mortality 40-80%		Mortality > 80%		Mortality 40-80%		Mortality > 80%	
Factor	OR (CI)	p	OR (CI)	p	OR (CI)	p	OR (CI)	p
Presence of haematological malignancy	0.188 (0.042-0.847)	0.029	0.161 (0.036-0.720)	0.017	4.320 (2.089-8.933)	0.000	3.124 (1.497-6.519)	0.002
Acute kidney failure	0.056 (0.007-0.426)	0.005	0.023 (0.003-0.169)	0.000	0.79 (0.010-0.604)	0.014	0.024 (0.003-0.177)	0.000
Location of cardiac arrest	6.740 (0.846-53.708)	0.072	9.274 (1.2-71.697)	0.033	NA	NA	0.406 (0.168-0.981)	0.045
Delay until start CPR	0.045 (0.010-0.199)	0.00	0.021 (0.005-0.94)	0.000	0.038 (0.005-0.294)	0.02	0.013 (0.002-0.100)	0.000
First heart rhythm in cardiac arrest	0.610 (0.283-1.314)	0.207	0.631 (0.297-1.343)	0.232	0.541 (0.269-1.555)	0.330	0.541 (0.225-1.297)	0.168
Severity of ARDS	0.065 (0.018-0.236)	0.000	0.012 (0.003-0.055)	0.000	0.056 (0.007-0.443)	0.006	NA	NA
OR= odds ratio, CI= confidence interval 95%, CPR= cardiopulmonary resuscitation, ARDS= acute respiratory distress syndrome, OHCA = out of hospital cardiac arrest								

Figures

Figure 1

Percentage deviation of true cost of five different ICU products. Dots are pre-education, squares are post-education. Results are shown in means $\pm$ SEM. X-axis at 100% is the true cost of the product. All values close to the X-axis are approximating true product cost. Abbreviations: ABG= arterial blood gas analysis, CT= computed tomography, Tx= transfusion.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplement1.pdf](#)
- [STROBEchecklistcohortWagenbergetal.pdf](#)