Pseudomonas Species Prevalence, Protein Analysis, and Antibiotic Resistance: An Evolving Public Health Challenge

Ayman Elbehiry (✉ aymanella2007@yahoo.com)

University of Sadat City <https://orcid.org/0000-0001-7726-9532>

Eman Marzouk

Qassim University

Ihab Moussa

King Saud University College of Science

Adil Abalkhail

Qassim University

Mai Ibrahim

Cairo University Faculty of Veterinary Medicine

Mohamed Hamada

University of Sadat City

Mohammed Rawway

Al-Azhar University Faculty of Science

Original article

Keywords: Pseudomonas spp., chilled meat, identification, antimicrobial resistance

Posted Date: November 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1067810/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Psychrotrophic *Pseudomonas* is one of the significant microbes that lead to putrefaction in chilled meat. One of the biggest problems in the detection of *Pseudomonas* is that several species are seemingly identical. Antibiotic resistance is an alternative considerable challenge worldwide. Therefore, this study was designed to apply an accurate technique for the discrepancy of *Pseudomonas* species and to study their resistance against various antimicrobials. A total of 320 chicken meat specimens were cultivated, and phenotypically recognized the isolated bacteria. Protein analysis was carried out for cultured isolates via Microflex LT. The resistance of *Pseudomonas* isolates was recorded through Vitek® 2 AST-GN83 cards. Overall, 69 samples were identified as *Pseudomonas* spp. and included 18 *Pseudomonas lundensis* (*P. lundensis*), 16 *Pseudomonas fragi* (*P. fragi*), 13 *Pseudomonas oryzae* (*P. oryzae*), 10 *Pseudomonas stutzeri* (*P. stutzeri*), 5 *Pseudomonas fluorescens* (*P. fluorescens*), 4 *Pseudomonas putida* (*P. putida*), and 3 *Pseudomonas aeruginosa* (*P. aeruginosa*) isolates. Microflex LT identified all *Pseudomonas* isolates (100%) correctly with a score value ≥ 2.00 . PCA positively discriminated the identified isolates into various groups. The antimicrobial resistance levels against *Pseudomonas* isolates were 81.16% for nitrofurantoin, 71% for ampicillin and ampicillin/sulbactam, 65.22% for cefuroxime and ceftriaxone, 55% for aztreonam, and 49.28% for ciprofloxacin. The susceptibilities were 100% for cefotaxime, 98.55% for ceftazidime, 94.20% for each piperacillin/tazobactam and cefepime, 91.3% for cefazolin. In conclusion, chicken meat was found to be contaminated with different *Pseudomonas* spp., with high incidence rates of *P. lundensis*. Microflex LT is a potent tool for distinguishing *Pseudomonads* at the species level.

1. Introduction

There is no doubt that meat is one of the most important sources of nutrients such as proteins, essential fatty acids, vitamins, and minerals of various kinds and is characterized by its water activity, which plays a significant role in its rapid spoilage with various types of microorganisms (Doulgeraki et al. 2012; Wickramasinghe et al. 2019). Meat can incur microbial contamination during and after the slaughter process (Komba et al. 2012; Bantawa et al. 2018). Despite the strict procedures in place as means of quality in the meat industry, contamination with many microbes that threaten human health has become an important issue during the transport and storage process (Kennedy et al. 2005; Nychas et al. 2008; Rouger et al. 2017). Many microbial hazards can arise from meat and its products, leading to an increase in the prevalence of diseases and deaths, especially in developing countries (Mead et al. 1999). Spoilage bacteria habitually found in raw meat commonly exist before the slaughter process, or they can invade the meat throughout the handling, carriage, or packing process (Doulgeraki et al. 2012).

Among the important microbes that lead to the contamination of meat and meat products are *Pseudomonas*, *Brochothrix*, *Acinetobacter*, and *Shewanella*, which have been recovered from frozen meat (Wickramasinghe et al. 2019). Of the genus *Pseudomonas*, *P. fluorescens*, *P. fragi*, *P. lundensis*, *P. migulae*, and *P. putida* are considered frequent species found in chilled meat (Doulgeraki et al. 2012). *Pseudomonas lundensis* (*P. lundensis*) represents one of the most significant psychrotrophic microorganisms that leads to deterioration in frozen meat (Liu et al. 2015). *P. lundensis* is a gram-negative psychrotrophic motile bacterium that can grow at temperatures fluctuating from 0 to 33°C (Cox et al. 1989; Ercolini et al. 2010). In addition, previous studies carried out by Caldera et al. (2016) and Raposo et al. (2017) demonstrated that the majority of *Pseudomonas* species have the capacity to release different thermotolerant proteolytic and lipolytic enzymes that can seriously decrease the quality and shelf life of meat and its products.

Since *P. lundensis* has the ability to produce fluorescent pigments, the process of identifying it with nonpigment species (e.g., *P. fragi*) may be confusing. It has been further recognized that *P. lundensis* is strictly associated with both groups of *P. fragi* and *P. fluorescens*. These bacteria play a significant role in the deterioration of meat by forming unpleasant odours and slime throughout the stowage process (Wickramasinghe et al. 2019). Nevertheless, *P. lundensis* has predominantly been associated with milk and meat putrefaction, and new studies have utilized culture techniques and established that it exists in patients' lungs, particularly in patients suffering from cystic fibrosis (Scales et al. 2018). Nonetheless, the role of *P. lundensis* in lung deterioration and its probable role in respiratory distress are still unidentified. It is worth noting that in contrast to other microbes that cannot grow at low temperatures, psychrotrophic pseudomonads have the ability to adapt and rapidly colonize ice-cold foods, leading to deterioration and biofilm formation (De Jonghe et al. 2011; Quintieri et al. 2019; Orellana-Saez et al. 2019). This later action increases their adaptability and scattering capacity, and as a result, measures to eliminate them can provoke and increase their resistance to various antimicrobial drugs. As a consequence of these features, the existence of *Pseudomonas* spp. such as *P. lundensis*, *P. fragi*, *P. fluorescens*, *P. gessardii*, and *P. taetrolens* in chilled fresh food products has been developing great attention (Baruzzi et al. 2012; Guidone et al. 2016; Brasca et al. 2018; Quintieri et al. 2019).

Nevertheless, culture-independent techniques supply a broader viewpoint on bacterial assortment, and culturing novel isolates remains significant and habitually accomplished in the majority of microbiological laboratories. This is predominantly correct for ecological bacteria, which are considered a huge source of new natural products for feed flavours, and in the field of developing medicines and other manufacturing products (Stafsnes et al. 2013; Timperio et al. 2017). To recognize and categorize new isolates at the genus and species levels, numerous techniques, such as phenotypic and genotypic methods, exist for this purpose. However, while these methods are considered the gold standard for the identification of different types of microorganisms, they cannot provide adequate and reliable data regarding the detection and discrepancy of various bacteria at the species level (Pesciaroli et al. 2015).

Therefore, it is necessary to use a fast and accurate technique such as mass spectrometry technology to recognize and distinguish different microbes isolated from food products. MALDI-TOF MS is an extraordinary throughput-dependent tool utilized for protein analysis. Previous studies have proven that the analysis of complete bacterial cells using the technique of protein analysis is one of the most important methods employed in identifying different microbes in the past ten years. (Elbehiry et al. 2019). However, the challenge in the application of this technology in bacterial identification and grouping is the accessibility of cost-effective devices delivered with powerful datasets and easy software (Emonet et al. 2010; Elbehiry et al. 2017).

Nevertheless, the initial utilization of MALDI-TOF MS for the detection of several microorganisms, including bacteria and fungi, was not commonly used 30 years ago due to the absence of satisfactory databases (Tshikhudo et al. 2013). Recently, MALDI-TOF MS has been able to classify and discriminate

psychrotrophic bacteria at both the genus and species levels by matching the spectral profiles of field isolates with stored spectral proteins from reference strains (Vithanage et al. 2014; Dong 2020). The workflow of MALDI-TOF MS is based mainly on dissolving a fresh bacterial colony in an appropriate matrix compound and then inoculating it onto a target plate for protein analysis using laser shots in a measuring chamber. As a final point, a mass spectrum is obtained and displayed using particular software (Elbehiry et al. 2019).

Recently, antimicrobial resistance of different microorganisms has become an urgent matter due to their direct impact on public health worldwide. It is known that antibiotic-resistant bacteria have a close relationship with infections in hospitals, and *P. aeruginosa* is one of the most important opportunistic bacteria that affect human health, particularly in patients with defective immune systems (Chatterjee et al. 2016; Quintieri et al. 2019).

Furthermore, current suggestions emphasize that nonvirulent *Pseudomonads* not only have the ability to cause infections in the bloodstream of humans but also exhibit numerous types of multidrug resistance against various classes of antibiotics, which is a source of great danger to human health as a result of their high adaptability (Chatterjee et al. 2016; Cole and Singh 2017). In general, most of these types of bacteria are able to grow at temperatures below 70°C (psychrotrophic) and are characterized by their capabilities to contaminate fresh food and the surrounding environment, especially in the absence of the necessary sterilization procedures (Stellato et al. 2017; Quintieri et al. 2019). Numerous *Pseudomonas* spp. of food origin are competent to resist various antimicrobial agents of various classes, especially β -lactams such as penicillins, cephalosporins, carbapenems, and monobactams (King et al. 2014). The objective of the present investigation was to identify *Pseudomonas* spp. recovered from chicken meat samples using a Microflex LT device and to examine the antimicrobial resistance of *Pseudomonas* against various antimicrobial agents using AST GN83 cards.

2. Materials And Methods

2.1. Sample collection

A total of 320 frozen chicken meat products (200 g of each sample) represented by breast, thigh, burger, and nuggets (80 of each) were randomly collected from various retail stores in the Al-Qassim region, Saudi Arabia, at various intervals from June to December 2020. Each sample was preserved independently in a Ziploc® brand freezer bag at 4 °C, and all samples were then transferred directly to the microbiology laboratory in a heat-insulated ice box under the appropriate hygienic conditions for bacteriological examination.

2.2. Sample processing and isolation of *Pseudomonas* spp.

According to the guidelines provided by the FENG et al (2002), 25 g of each sample was moved into a sterilized container with 225 ml of sterile peptone water (0.1%) under strict hygienic measures, and homogenization was performed using a Denville Ultra EZgrind™ Tissue Homogenizer (Thomas Scientific, USA) at 14,000 rpm for 3 successive minutes and was then incubated for 5 minutes at 25°C. From this mixture, 1 ml was moved into a sterilized test tube containing 9 ml of sterile peptone water, from which tenfold (1:10) serial dilutions were processed. The prepared samples were subjected to isolation and determination of *Pseudomonas* counts (ISO, 2003). In brief, 0.1 ml of each homogenized sample was independently inoculated into duplicate Petri dishes on Thermo Scientific™ Remel™ *Pseudomonas* Isolation Agar supplemented with glycerol and uniformly distributed with a sterile plastic spreader (Thomas Scientific, USA). The inoculated plates were incubated at 4°C and 25°C for a couple of days. Then, greenish yellow colonies were detected and enumerated. The average count was considered and recorded from serial dilutions of 10^{-3} to 10^{-6} . Subculturing of all suspected colonies was carried out by streaking on nutrient agar and incubating at 4°C for 3 to 5 days to obtain purified colonies. All purified strains were kept in the CRYOBANK™ Bacterial Culture Freezing System (COPAN Diagnostics Inc., Murrieta, USA) for further investigations.

2.3. Phenotypic identification of *Pseudomonas* species

2.3.1. Microscopy and Gram stain properties

All purified strains kept in Cryobank vials were subcultured again for microscopic identification of *Pseudomonas* species using the Gram staining technique. In brief, a fresh colony of each strain was fixed onto a clean, dried slide, and then the colony was flooded with different chemicals. Crystal violet dye was first dropped onto the glass slide containing bacterial cells, and then iodine was added to fix the dye. Ethanol was then added to remove the dye from unspotted cells, and safranin was added as a final point to stain the gram-negative bacteria. The stained slides were scanned by light microscopy using an oil immersion lens to observe the morphology of bacteria. The isolates were recognized as gram-negative if they appeared pink.

2.3.2. Enzymatic activities

The enzymatic activities of *Pseudomonas* species were evaluated using certain biochemical tests, including oxidase, citrate utilization and indole tests. The oxidase test was performed by smearing a fresh colony of each isolate onto a sterile oxidase filter paper disc (Sigma–Aldrich, USA) soaked in distilled water. The positive oxidase activity results are indicated by the appearance of a purple colour. The capacity of *Pseudomonas* spp. to utilize citrate as a source of energy was carried out by streaking a fresh purified colony onto a Simmons citrate agar (Sigma–Aldrich, USA) slant and incubated at 37°C for 5–7 successive days. Citrate utilization was indicated by the appearance of a sky-blue colour. The capability of *Pseudomonas* spp. to convert tryptophan into indole was tested by inoculation of the suspected organism into tryptophan broth (Sigma–Aldrich Chemie GmbH, Germany), which was then incubated at 37°C for a couple of days. Thereafter, using gentle agitation, 0.5 ml of Kovacs reagent was added until a red–violet colour appeared, indicating positive results, while a yellow colour indicates negative results.

2.3.3. Identification and determination of antimicrobial resistance via the Vitek 2 Compact System

All gram-negative isolates that showed positive results for both citrate utilization and oxidase tests and negative indole tests were further examined by the Vitek 2 Compact system (bioMérieux, France) for confirmation of species identification and antimicrobial resistance. Briefly, preparation of the isolate suspension was carried out and then adjusted using McFarland standards (0.5 to 0.63) as stated in the company's instructions. AST-GN83 gram-negative identification cards, which included 18 antimicrobial drugs, were utilized as gram-negative antibiotic susceptibility cards for *Pseudomonas* spp. (Table 1). According to the company's instructions, Vitek®2 cards were inoculated, and then the isolate IDs were submitted to the device to permit the selection of precise interpretive standards. Based on the authorizations of the Clinical and Laboratory Standards Institute (CLSI) (Schreckenberger and Binnicker 2011), the minimum inhibitory concentration (MIC) was interpreted as susceptible, intermediate, or resistant. *Pseudomonas* sp. (ATCC® 19151™) was utilized as a reference strain throughout the experiment. All strains that exhibited intermediate reactions against antimicrobial drugs were considered resistant strains.

Table 1
MIC ranges (µg/ml) of the antibiotics used by Vitek 2
Compact AST GN83 cards

Antimicrobial drug	MIC range (µg/ml)
Amikacin	2–64
Amoxicillin/Clavulanic Acid	2/1–32/16
Ampicillin	2–32
Ampicillin/Sulbactam	2/1–32/16
Aztreonam	1–64
Cefazolin	4–64
Cefepime	1–645
Cefotaxime	0.25–64
Cefoxitin	4–64
Ceftazidime	1–64
Ceftriaxone	0.25–64
Cefuroxime	1–64
Ciprofloxacin	0.25–4
Gentamicin	1–16
Meropenem	0.25–16
Nitrofurantoin	16–512
Piperacillin/Tazobactam	4/4–128/4
Trimethoprim/Sulfamethoxazole	20–320

2.4. Proteomic screening for identification of *Pseudomonas* spp.

Microflex LT (Bruker Daltonics, Bremen, Germany) was applied for the recognition and classification of *Pseudomonas* species from chicken meat samples. All isolates were analysed using both FlexControl and Compass software (Flex Series version 1.3). All isolates were prepared by culturing on nutrient agar (Sigma–Aldrich, USA) followed by incubation for a couple of days at 37°C. The ethanol/formic acid extraction procedure was utilized as stated by Bruker Daltonics. Briefly, 2 pure colonies were relocated onto a clean Eppendorf tube holding 300 µl of sanitized water and 900 µl of ethanol (99.9%). The contents were mixed carefully by centrifugation at 13,000 rpm for two minutes. The obtained pellet was dried in air for 5 minutes after the removal of the supernatant. Fifty microlitres of formic acid (70%) was added to the air-dried pellet, and acetonitrile (70%) was then added after proper mixing for 2 minutes at 13,000 rpm. One microlitre of the supernatant for each isolate was then placed onto a target plate of Microflex LT and left to dry at 25°C. Subsequently, 1 µl of matrix solution acyano-4 hydroxy-cinnamic acid was added. The target plate was then submitted to the Microflex LT machine for identification and data reading. A bacterial test standard (*Escherichia coli*) was used as a positive control throughout the experiment.

The score values of indeterminate spectra were compared with the reference spectra stored in the databank. Microflex LT has the capacity to accurately recognize and distinguish different microorganisms when the values range from 2 to 3. However, misidentification can result if this value ≤ 1.69 . The diverse spectra created by IVD Compass Software were analysed via a m/z range from 2000 Da to 20,000 Da. A dendrogram could be produced from the main spectral profile (MSP) database, which includes > 6.989 different species of bacteria and fungi.

3. Results

3.1. Frequency and counting of *Pseudomonas* spp.

Out of the 320 chicken meat samples involved in the current investigation, 69 (21.56%) were determined to be positive for *Pseudomonas* spp. using culturing techniques. Out of 69 positive isolates, 8 (11.59%), 11 (15.94%), 25 (36.23%), and 25 (36.23%) were isolated from breast, thigh, burger, and nugget meat,

respectively. After statistical analysis, the mean values of colony-forming units (CFU/g) were $10.1 \times 10^3 \pm 1.45 \times 10^3$, $7.4 \times 10^3 \pm 0.89 \times 10^3$, $6.5 \times 10^3 \pm 6.43 \times 10^3$ and $4.3 \times 10^3 \pm 7.56 \times 10^3$ for the isolates recovered from the breast, thigh, burger, and nugget meat, respectively (Table 2).

Table 2
The mean values of colony forming units (CFU/g) for 69 *Pseudomonas* spp. detected in chicken meat products

Chicken meat product	Minimum	Maximum	Mean value $\pm$ standard error
Breast (n=8)	2.0×10^2	10.1×10^3	$10.1 \times 10^3 \pm 1.45 \times 10^3$
Thigh (n=11)	3.4×10^2	7.4×10^3	$7.4 \times 10^3 \pm 0.89 \times 10^3$
Hamburger (n=25)	3.9×10^2	6.5×10^3	$6.5 \times 10^3 \pm 6.43 \times 10^3$
Nuggets (n= 25)	3.7×10^2	4.3×10^3	$4.3 \times 10^3 \pm 7.56 \times 10^3$

3.2. Biochemical analysis of *Pseudomonas* spp.

A total of 69 *Pseudomonas* isolates that showed positive results for oxidase and citrate utilization and negative results for indole were examined by the Vitek 2 Compact system. Consistent with the interpreted results, 18 (26.09%) strains were identified as *Pseudomonas lundensis* (*P. lundensis*), 16 (23.19%) *Pseudomonas fragi* (*P. fragi*), 13 (18.84%) *Pseudomonas oryzihabitans* (*P. oryzihabitans*), 10 (14.49%) *Pseudomonas stutzeri* (*P. stutzeri*), 5 (7.25%) *Pseudomonas fluorescens* (*P. fluorescens*), 4 (5.8%) *Pseudomonas putida* (*P. putida*), and 3 (4.35%) *Pseudomonas aeruginosa* (*P. aeruginosa*). From the previous results, it was indicated that *P. lundensis* was the most common *Pseudomonas* spp. recovered from chicken meat samples, followed by *Pseudomonas fragi*, *P. oryzihabitans*, and *P. stutzeri* (Table 3).

Table 3
Pseudomonas spp. recovered from chicken meat products

<i>Pseudomonas</i> spp.	Chicken meat products								Total	
	Breast		Thigh		Hamburger		Nuggets			
	No.	%	No.	%	No.	%	No.	%	No.	%
<i>P. lundensis</i>	3		2		7		6		18	26.09
<i>P. fragi</i>	1		3		7		5		16	23.19
<i>P. oryzihabitans</i>	3		2		5		3		13	18.84
<i>P. stutzeri</i>	0		3		3		4		10	14.49
<i>P. fluorescens</i>	1		0		2		2		5	7.25
<i>P. putida</i>	0		1		1		2		4	5.80
<i>P. aeruginosa</i>	0		0		0		3		3	4.35
Total	8		11		25		25			

3.3. Proteomic analysis of *Pseudomonas* spp.

In the existing study, the isolated strains were screened by Microflex LT, and the spectra of the field isolates were parallel to the reference spectra. According to the results obtained, Microflex LT was capable of verifying all *Pseudomonas* spp. by 100%. Analysing these results shows that approximately 20 prominent ion peaks were detected in the original bands from the region varying from 2000 to 10,200 Daltons (Da) (Figure 1A), which were confirmed from the gel view (Figure 1B), and robust peaks were revealed at 3800, 3860, 4440, and 4550 Da (Figure 2A), which were confirmed from the gel view (Figure 2A). All 69 strains of *Pseudomonas* spp. were correctly identified as follows: 18 *P. lundensis*, 16 *P. fragi*, 13 *P. oryzihabitans*, 10 *P. stutzeri*, 5 *P. fluorescens*, 4 *P. putida*, and 3 *P. aeruginosa*. All of these species were synchronized with the *P. Lundensis* DSM 6252T HAM (Figure 3), *P. fragi* DSM 3456T HAM, *P. oryzihabitans* DSM 6835T, *P. stutzeri* V319 MCRF, *P. fluorescens* DSM 1976, *P. putida* ATCC 49128 THL, and *P. aeruginosa* DSM 1117 reference strains stored in Compass IVD software. All *Pseudomonas* strains were distinguished by matching their spectra with the Bruker database, which includes 70 strains obtained from the American Type Culture Collection (ATCC) and the German Collection of Microorganisms and Cell Cultures GmbH (DSMZ).

As summarized in Table 4, we found that 10/18 (55.56%) *P. lundensis*, 7/16 (43.75%) *P. fragi*, 6/13 (46.15%) *P. oryzihabitans*, 4/10 (40%) *P. stutzeri*, 3/5 (60%) *P. fluorescens*, 1/4 (25%) *P. putida*, and 2/3 (66.67%) *P. aeruginosa* were properly recognized, with log values ranging from 2.3 to 3.0. Likewise, 8/18 (44.44%) *P. lundensis*, 9/16 (56.25%) *P. fragi*, 7/13 (53.85%) *P. oryzihabitans*, 5/10 (50%) *P. stutzeri*, 2/5 (40%) *P. fluorescens*, 3/4 (75%) *P. putida*, and 1/3 (33.33%) *P. aeruginosa* were properly identified, with log values ranging from 2.0 to 2.29. Nonetheless, one isolate of *P. stutzeri* was detected at the genus level with a log value ranging from 1.70 to 1.99. Zero strains were not detected. An additional calculation tool termed principal component analysis (PCA) was created in the present investigation in Microflex LT Compass IVD software to determine the similarities and differences between the spectral proteins of *Pseudomonas* spp. Several spectral proteins of various colours are presented in three-dimensional (3d) images of PCA (Figure 4). The 3 loading values originating from the calculation of PC1, PC2, and PC3 were used for the identification of each peak. In our examination, the wide-ranging peaks recorded in the Microflex LT Compass IVD software were evaluated by the PCA mathematical tool, which was able to separate the *Pseudomonas* strains by placing them in groups

through dots of different colours (Figure 4) as follows: *P. lundensis* (blue), *P. fragi* (yellow), *P. oryzihabitans* (blue black), *P. stutzeri* (aqua), *P. fluorescens* (maroon), *P. putida* (red), and *P. aeruginosa* (green).

Table 4
Score values for 69 *Pseudomonas* spp. detected in chicken meat by PMFT

Species	Score value of identification							
	2.3–3		2–2.29		1.7–1.99		0–1.69	
	No.	%	No.	%	No.	%	No.	%
<i>P. lundensis</i>	10/18	55.56	8/18	44.44	0	0	0	0
<i>P. fragi</i>	7/16	43.75	9/16	56.25	0	0	0	0
<i>P. oryzihabitans</i>	6/13	46.15	7/13	53.85	0	0	0	0
<i>P. stutzeri</i>	4/10	40	5/10	50	1/10	10	0	0
<i>P. fluorescens</i>	3/5	60	2/5	40	0	0	0	0
<i>P. putida</i>	1/4	25	3/4	75	0	0	0	0
<i>P. aeruginosa</i>	2/3	66.67	1/3	33.33	0	0	0	0
3.4 Direct screening of the antimicrobial susceptibilities of certain <i>Pseudomonas</i> spp. by the VITEK AST GN83 cards								

A total of 69 different species of *Pseudomonas* were tested against 18 antimicrobial drugs utilized in the current investigation. According to the results labelled in Table 4, the majority of *Pseudomonas* species exhibited higher degrees of resistance against different classes of antibiotics. The highest degree of resistance was detected against nitrofurantoin (nitrofurantoin, 81.16%), followed by beta-lactam (ampicillin, 71%) & aztreonam, 55%), beta-lactam/beta-lactamase inhibitor (ampicillin/sulbactam, 71%), 2nd -generation cephalosporins (cefuroxime, 65.22%), 3rd -generation cephalosporins (ceftriaxone, 65.22%), ciprofloxacin (49.28%) and carbapenem (meropenem, 43.48%). In contrast, the most susceptible antimicrobial drugs were cefotaxime (100%) and ceftazidime (98.45%).

As shown in Table 5, of 22 *A. hydrophila* isolates, 21 (95.46%) were resistant to ampicillin as revealed in Table 5, of 18 *P. lundensis* strains, 18 (100%), 17 (94.44%), 17 (94.44%), 16 (88.89%), 16 (88.89%), 15 (83.33%) and 12 (66.67%) were resistant to ampicillin, amoxicillin/clavulanic acid, ampicillin/sulbactam, ciprofloxacin, nitrofurantoin, ceftriaxone, and cefuroxime, respectively. In contrast, all strains were sensitive to both cefotaxime and ceftazidime. In addition, out of 16 *P. fragi* strains, 16 (100%), 15 (93.75%), 15 (93.75%), 14 (87.5%), 13 (81.25%), 13 (81.25%), 12 (75%), and 12 (75%) were resistant to aztreonam, ceftriaxone, nitrofurantoin, ciprofloxacin, amoxicillin/clavulanic acid, cefuroxime, ampicillin, and cefoxitin, respectively. However, all strains of *P. fragi* showed higher degrees of susceptibility to only ceftazidime, cefotaxime, and ceftazidime.

Table 5 indicates that the majority of *P. oryzihabitans* isolates showed higher degrees of susceptibility for almost all antibiotics under study except nitrofurantoin, which exhibited a high resistance (100%), followed by cefuroxime (38.46%) and ceftriaxone (23.08%). Moreover, *P. stutzeri* strains exhibited a high resistance (100%) to amoxicillin/clavulanic acid, ampicillin, ampicillin/sulbactam, cefoxitin, ceftriaxone, cefuroxime, and nitrofurantoin. *P. fluorescens* strains also revealed a certain degree of resistance (60%) against amoxicillin/clavulanic acid, ampicillin, and ampicillin/sulbactam and 40% against ceftazidime and cefuroxime. Fifty percent of *P. putida* strains were resistant ampicillin and ampicillin/sulbactam. Furthermore, 100% of *P. aeruginosa* strains were unaffected by amoxicillin/clavulanic acid, ampicillin, and ceftazidime, and 66.67% were unaffected by ampicillin/sulbactam, cefoxitin, ceftriaxone, cefuroxime, and nitrofurantoin.

4. Discussion

The contamination of chilled meat with various microbes is one of the major causes of cost-effective problems in meat production and has an impact on public health worldwide (Wickramasinghe et al. 2019). Despite the use of modern methods of meat preservation, microbial contamination remains a major threat. Psychrotrophic *Pseudomonas* spp. are considered the main bacteria that lead to putrefaction of ice-cold meat under aerobic conditions. The genus *Pseudomonas* is one of the most polluting microbes and is characterized by its great ability to withstand difficult environmental conditions, which leads to the prevention of the growth of other microorganisms. To diminish contamination of meat, proper detection and treatment of various *Pseudomonas* spp. are critical. Therefore, this study concentrated on an accurate method of identification and differentiation of *Pseudomonas* spp. and evaluated their degrees of resistance and susceptibility to various antibiotics commonly used for treatment.

Based on our results, the mean *Pseudomonas* counts (CFU/g) were $10.1 \times 10^3 \pm 1.45 \times 10^3$, $7.4 \times 10^3 \pm 0.89 \times 10^3$, $6.5 \times 10^3 \pm 6.43 \times 10^3$ and $4.3 \times 10^3 \pm 7.56 \times 10^3$ for the isolates recovered from the breast, thigh, burger, and nuggets, respectively. Comparable findings were recorded by Hassan et al. (2020), who found that the mean *Pseudomonas* counts recovered from various chicken meat products (chilled breast, thigh, nuggets and burger) varied from $3.51 \times 10^3 \pm 0.76 \times 10^3$ to $8.44 \times 10^3 \pm 1.85 \times 10^3$. Other parallel records detected by Morshdy et al. (2018) and Abd El-Aziz (2015) were 3.6×10^3 and 2.6×10^4 , respectively. Although the research performed by Hinton et al. (2007) indicated that psychrotrophic bacteria were not recovered from carcasses washed with chlorinated water, different species of *Pseudomonas* were the most predominant psychrotrophs recovered from all carcasses when stored in refrigerators for two weeks. Several investigations have revealed that the initial *Pseudomonas* count is directly associated with the period of storing meat in the refrigerator, and meat spoilage occurs when the number of *Pseudomonas* ranges from 10^7 to 10^8 (Hassan et al. 2020).

In the current investigation, we identified 69 *Pseudomonas* spp. using biochemical analysis confirmed by proteomics methods. The identified isolates were represented as *P. lundensis* (18), *P. fragi* (16), *P. oryzihabitans* (13), *P. stutzeri* (10), *P. fluorescens* (5), *P. putida* (4), and *P. aeruginosa* (3). Chicken meat burgers (25/69) and nuggets (25/69) were the most contaminated chicken meat products of *Pseudomonas* species, which might be a result of mismanagement, extreme usage, and unsuccessful hygienic practices throughout processing and packing. Parallel findings were obtained by Hassan et al. (2020), who identified 166 isolates of *Pseudomonas* species, with a high prevalence of *P. fluorescens* followed by *P. alcaligenes*, *P. stutzeri*, *P. proteolytica*, and *P. fragi*, while low incidence rates were recorded for *P. aeruginosa*, *P. stutzeri*, and *P. acidovorans*. In another study, Arnaut-Rollier et al. (1999) detected 3 species of *Pseudomonas* (*P. fragi*, *P. lundensis*, *P. fluorescens biovars*) from both fresh and refrigerated chicken skin.

In addition, 11 strains of *Pseudomonas* species recovered from cooked chicken burgers were identified by Franzetti and Scarpellini (2007) as 8 strains of *P. fragi*, followed by 2 *P. chitorii* and 1 *P. fluorescens*. In contrast, *P. aeruginosa* was not detected in 100 chicken meat specimens (Iroha et al. 2011). In another study conducted by Caldera et al. (2016), the deterioration of food was commonly associated with *P. aeruginosa*, *P. fragi*, *P. lundensis* and *P. fluorescens* (Caldera et al. 2016). Moreover, Bellés et al. (2017) and Wang et al. (2017) clarified that the capability of these microorganisms to live at low temperatures may lead to trouble throughout the storage of foodstuffs. The existence of *Pseudomonas* spp. in various food samples is of high importance because this type of bacteria has a bad impact on human health and is considered a sign of food quality (Yagoub 2009).

Because phenotypic-based detection of various foodborne pathogens is difficult and takes a long time to be carried out, Microflex LT was meaningfully applied in our study for the initial detection and classification of numerous bacteria from chicken meat samples, as it is an easy, quick, specific, and inexpensive detection technique compared to other approaches (Singhal et al. 2015; van Belkum et al. 2017; Elbehiry et al. 2019). In recent times, Microflex LT has been discovered to be an imperative tool for the powerful recognition of microbial intimidations that may pollute both water and foodstuffs (Singhal et al. 2015; Elbehiry et al. 2019).

In the present investigation, the percentage of mass spectral identification of *Pseudomonas* strains was 100% for all 7 species of *Pseudomonas*. The interpreted results confirmed that all spectral profiles produced by Microflex LT IVD Compass Software were suitable to distinguish between *Pseudomonads* at the species level. The accurate identification observed in our study may be a result of the restructured database (Elbehiry et al. 2019). Comparable findings were noted by Böhme et al. (2011), who applied Microflex LT effectively in the exact identification of gram-negative bacteria (e.g., *Pseudomonas* and *Enterobacter*) of different species recovered from seafood. Consequently, Microflex LT has been demonstrated to be an authoritative instrument for microbial identification. Höll et al. (2016) also identified several microorganisms isolated from packaged poultry meat using Microflex LT, and they found that *Pseudomonas* spp. is one of the most common bacteria found after 7 days of storage at 4°C and 10°C.

In addition, principal component analysis (PCA) generated by the Microflex LT device magnificently divided *P. lundensis*, *P. fragi*, *P. oryzihabitans*, *P. stutzeri*, *P. fluorescens*, *P. putida*, and *P. aeruginosa* strains into different groups. Han (2010) and Elbehiry et al. (2019) indicated that PCA is usually applied as a mathematical tool to extract and demonstrate the modification in the spectral profiles within the database.

The spread of antimicrobial resistance amongst the genus *Pseudomonas* was also examined in the current study. Of late, antimicrobial resistance represents one of the common public health problems, as multidrug-resistant bacteria related to animals may be virulent and transferred simply to human beings through food chains and comprehensively dispersed through animal wastes to the environment (Manyi-Loh et al. 2018). Antimicrobial resistance is problematic and multifaceted and occurs as a consequence of the unreasonable use of antibiotics under poor hygienic measures (Osman et al. 2019).

In the present investigation, the AST GN83 card was applied to display the resistance and susceptibility of 69 *Pseudomonas* species recovered from various chicken meat samples against several antibiotics frequently utilized for the treatment of gram-negative pathogens. Based on our results, the majority of *Pseudomonas* isolates exhibited higher degrees of susceptibility to cefotaxime (100%), ceftazidime (98.55%), cefepime (94.2%), gentamycin (86.96%), and amikacin (82.16%). These findings were similar to those obtained by CLSI (2015). It was also observed that the majority of the *Pseudomonas* isolates were highly sensitive to meropenem (70%). Parallel results were achieved in previous studies carried out in Turkey by Shenoy et al. (2002) and Deniz Yilmaz et al. (2016) and in Kenya by Mwinyikombo (2018), who revealed that *Pseudomonas* isolates demonstrated higher degrees of susceptibility to both meropenem and imipenem. Nonetheless, other studies performed in India by Sivanmaliappan & Sevanan (2011) illustrated a higher degree of resistance to imipenem (66.6%), a finding that could be explained by the misuse of broad-spectrum antibiotics such as carbapenems.

The highest degree of resistance was detected against various classes of antibiotics, such as nitrofurantoin (81.16%), followed by beta-lactam [ampicillin (71%) and aztreonam (55%)], beta-lactam/beta-lactamase inhibitor [ampicillin/sulbactam (71%)], second-generation cephalosporins [cefuroxime (65.22%)], third-generation cephalosporins [ceftriaxone (65.22%)], ciprofloxacin (49.28%) and carbapenem [meropenem (43.48%)]. Similar findings regarding nitrofurantoin were obtained by Cunha et al. (2011) and Osei Sekyere et al. (2018), who indicated that various bacteria (e.g., *Pseudomonas*, *Salmonella*, *Proteus*) isolated from patients suffering from urinary tract infections are highly resistant to nitrofurantoin. It can be seen from our study that the majority of *Pseudomonas* isolates were found to directly develop resistance against various types of antibiotics. Among the *Pseudomonas* spp. detected in the present study, *P. lundensis*, *P. fragi*, *P. oryzihabitans*, *P. stutzeri* and *P. aeruginosa* can be proposed as reservoirs of antibiotic resistance.

The present study demonstrates the high incidence rates of *P. lundensis*, *P. fragi* and *P. oryzihabitans* among different species of *Pseudomonas* isolated from chicken meat samples. Throughout this investigation, we established that Microflex LT applied for the detection of *Pseudomonas* species is an authoritative, inexpensive, and truthful technique and was able to discriminate different species of *Pseudomonas* using PCA generated by the Microflex LT device. Future investigations will be required to determine the useful application of this technique for the proper recognition and discrepancies of *Pseudomonas* spp. in food samples. Furthermore, our results demonstrated that the various species of *Pseudomonas* had established multidrug resistance. Accordingly, the evolution of resistance might be foreseeable; hence, the number of effective antimicrobial drugs is decreasing. Since *Pseudomonas* can threaten public health, the broadcast of resistance may perhaps have negative influences on individuals.

Abbreviations

PCA: Principal Component Analysis; IVD: in vitro diagnostic device.

Declarations

Acknowledgement

The researchers would like to thank the Deanship of Scientific Research, Qassim University, for their support of this project.

Authors' contributions

AE and AA data curation, writing-original draft preparation. EM, AA, and IM conceptualization, methodology, and validation. MI and MH visualization, investigation. MR and AE supervision. AE, MH, and EM writing reviewing and editing. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

The data that support the findings of this study are available on request from the corresponding author.

Ethics approval and consent to participate

This article does not contain any studies with human participants or animals performed by any of the authors.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Abd El-Aziz DM (2015) Detection of *Pseudomonas* spp. in chicken and fish sold in markets of Assiut City, Egypt. *J Food Qual Hazards Control* 2(3): 86-89. Corpus ID: 56252088
2. Arnaut-Rollier I, Vauterin L, DeVos P, Massart DL, Devriese LA, De-Zutter L, Van-Hoof J (1999) A numerical taxonomic study of the *Pseudomonas* flora isolated from poultry meat. *J Appl Microbiol* 87(1): 15-28. doi: 10.1046/j.1365-2672.1999.00785.x
3. Bantawa K, Rai K, Subba Limbu D, Khanal H (2018) Food-borne bacterial pathogens in marketed raw meat of Dharan, eastern Nepal. *BMC Res Notes* 11(1), 618. doi: 10.1186/s13104-018-3722-x
4. Baruzzi F, Lagonigro R, Quintieri L, Morea M, Caputo L (2012). Occurrence of non-lactic acid bacteria populations involved in protein hydrolysis of cold-stored high moisture Mozzarella cheese. *Food Microbiol* 30(1): 37–44. doi: 10.1016/j.fm.2011.10.009
5. Bellés M, Alonso V, Roncalés P, Beltrán JA (2017) A review of fresh lamb chilling and preservation. *Small Rumin Res* 146: 41-7. <https://doi.org/10.1016/j.smallrumres.2016.12.003>
6. Böhme K, Fernández-No IC, Barros-Velázquez J, Gallardo JM, Cañas B, Calo-Mata P (2011) Rapid species identification of seafood spoilage and pathogenic Gram-positive bacteria by MALDI-TOF mass fingerprinting. *Electrophoresis* 32(21): 2951–2965. doi: 10.1002/elps.201100217
7. Brasca M, Decimo M, Morandi S, Machado SG, Bagliniére F, Vanetti MCD (2018) Psychrotrophic bacteria. In *Microbiology in Dairy Processing: Challenges and Opportunities*; Poltronieri P Ed, Wiley-Blackwell: Hoboken, NJ, USA, Chapter 37: 37–61. ISBN: 978-1-119-11480-2
8. Caldera L, Franzetti L, Van-Coillie E, De Vos P, Stragier P, De Block J, Heyndrickx M (2016) Identification, enzymatic spoilage characterization and proteolytic activity quantification of *Pseudomonas* spp. isolated from different foods. *Food Microbiol* 54:142-53. <https://doi.org/10.1016/j.fm.2015.10.004>
9. Chatterjee M, Anju C, Biswas L, Kumar VA, Mohan CG, Biswas R (2016) Antibiotic resistance in *Pseudomonas aeruginosa* and alternative therapeutic options. *Int J Med Microbiol* 306(1): 48-58. doi: 10.1016/j.ijmm.2015.11.004
10. CLSI (2015) Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fifth Informational Supplement. CLSI document M100-S25. Wayne, PA: Clinical and Laboratory Standards Institute. ISBN 1-56238-990-4
11. Cole ML, Singh OV (2017) Foodborne Pathogens and Their Apparent Linkage with Antibiotic Resistance. In *Foodborne Pathogens and Antibiotic Resistance*; Wiley: Hoboken, NJ, USA, chapter 11: 247-274. <https://doi.org/10.1002/9781119139188.ch11>
12. Cox JM, Mac Rae IC (1989) Identification of psychrotrophic pseudomonads from goats' milk by computer-assisted analysis of carbon source assimilation data. *J Appl Bacteriol* 67(4): 377-393. doi: 10.1111/j.1365-2672.1989.tb02508.x

13. Cunha BA, Schoch PE, Hage JR (2011) Nitrofurantoin: preferred empiric therapy for community-acquired lower urinary tract infections. 86(12): 1243-1244. doi: 10.4065/mcp.2011.0411
14. De Jonghe V, Coorevits A, Van Hoorde K, Messens W, Van Landschoot A, De Vos P, Heyndrickx M (2011) Influence of storage conditions on the growth of *Pseudomonas* species in refrigerated raw milk. Appl Environ Microbiol 77(2): 460-470. doi: 10.1128/AEM.00521-10
15. Deniz Yilmaz M, Eyigori H, Osma U, Tarik Selçuk O, Renda L, Pirtik I, Didem Yalcin A (2016) Prevalence of allergy in patients with benign lesions of the vocal folds. Acta Medica Mediterr 32(5): 195-201. DOI: 10.19193/0393-6384_2016_1_30
16. Dong Z (2020) The effect of psychrotrophic bacteria on the quality of UHT milk: Doctoral thesis, The School of Food and Advanced Technology, Massey University, Palmerston North, New Zealand. <http://hdl.handle.net/10179/15827>
17. Doulgeraki AI, Ercolini D, Villani F, Nychas G-JE (2012) Spoilage microbiota associated with the storage of raw meat in different conditions. Int J Food Microbiol 157(2): 130-141. <https://doi.org/10.1016/j.ijfoodmicro.2012.05.020>
18. Elbehiry A, Marzouk E, Abdeen E, Al-Dubaib M, Alsayeqh A, Ibrahim M, Hamada M, Alenzi A, Moussa I, Hemeg HA (2019) Proteomic characterization and discrimination of *Aeromonas* species recovered from meat and water samples with a spotlight on the antimicrobial resistance of *Aeromonas hydrophila*. Microbiologyopen 8(11): e782. doi: 10.1002/mbo3.782
19. Elbehiry A, Marzouk E, Hamada M, Al-Dubaib M, Alyamani E, Moussa IM, AlRowaidhan A, Hemeg HA (2017) Application of MALDI-TOF MS fingerprinting as a quick tool for identification and clustering of foodborne pathogens isolated from food products. New Microbiol 40(4): 269-278. Epub 2017 Aug 21. PMID: 28825446.
20. Emonet S, Shah HN, Cherkaoui A, Schrenzel J (2010) Application and use of various mass spectrometry methods in clinical microbiology. Clin Microbiol Infect 16(11):1604-13. doi: 10.1111/j.1469-0691.2010.03368.x
21. Ercolini D, Casaburi A, Nasi A, Ferrocino I, Di Monaco R, Ferranti P, Villani F (2010) Different molecular types of *Pseudomonas fragi* have the same overall behaviour as meat spoilers. Int J Food Microbiol 142(1-2): 120-131. doi: 10.1016/j.ijfoodmicro.2010.06.012
22. Feng P, Weagant SD, Grant MA, Burkhardt W (2002) Chapter 4. Enumeration of Escherichia coli and the Coliform Bacteria," In: Food and Drug Administration (FDA), Bacteriological Analytical Manual Online, 8th Edition, Silver Spring, Berlin, 1998.
23. Franzetti L, Scarpellini M (2007) Characterization of Pseudomonas spp. isolated from foods. Ann Microbiol 57: 39-47. <https://doi.org/10.1007/BF03175048>
24. Guidone A, Zotta T, Matera A, Ricciardi A, De Filippis F, Ercolini D, Parente E (2016) The microbiota of high-moisture mozzarella cheese produced with different acidification methods. Int J Food Microbiol 4(2016): 9-17. doi: 10.1016/j.ijfoodmicro.2015.09.002
25. Han H (2010) Nonnegative principal component analysis for mass spectral serum profiles and biomarker discovery. BMC Bioinformatics 11: S1. <https://doi.org/10.1186/1471-2105-11-S1-S1>
26. Hassan MA, Ibrahim HM, Shawky NA, Sheir SH (2020) Incidence of Psychotropic bacteria in frozen chicken meat products with special reference to *Pseudomonas* species. Benha Veterinary Medical Journal 39(1): 165-168. DOI: 10.21608/bvmj.2020.37744.1238
27. Hinton AJR, Northcutt JK, Smith DP, Musgrove MT, Ingram KD (2007) Spoilage microflora of broiler carcasses washed with electrolyzed oxidizing or chlorinated water using an inside-outside bird washer. Poult Sci 86(1): 123-127. doi: 10.1093/ps/86.1.123
28. Höll L, Behr J, Vogel RF (2016) Identification and growth dynamics of meat spoilage microorganisms in modified atmosphere packaged poultry meat by MALDI-TOF MS. Food Microbiol 60: 84-91. doi: 10.1016/j.fm.2016.07.003
29. ISO (2003) Microbiology of food and animal feeding stuffs - Horizontal method for the enumeration of microorganisms - Colony-count technique at 30 degrees C. International Standards Organization, Geneva. ISO 4833:2003. ICS: 07.100.30 Food microbiology
30. Iroha IR, Ugbo EC, Ilang DC, Oji AE, Ayogu TE (2011) Bacterial contamination of raw meat sold in Abakaliki, Ebonyi State Nigeria. J Public Health Epidemiol 3(2): 49-53. Available online at <http://www.academicjournals.org/jphe>
31. Kennedy J, Jackson V, Blair IS, McDowell DA, Cowan C, Bolton DJ (2005) Food safety knowledge of consumers and the microbiological and temperature status of their refrigerators. J Food Prot 68(7): 1421-1430. doi: 10.4315/0362-028x-68.7.1421
32. King DT, Sobhanifar S, Strynadka NCJ (2014) The Mechanisms of Resistance to β -Lactam Antibiotics. In: Gotte M., Berghuis A., Matlashewski G., Wainberg M., Sheppard D. (eds) Handbook of Antimicrobial Resistance. Springer, New York, NY. https://doi.org/10.1007/978-1-4939-0667-3_10-1
33. Komba EV, Komba EV, Mkupasi EM, Mbyuzi AO, Mshamu S, Mzula A, Luwumba D, Busagwe Z, Mzula A (2012) Sanitary practices and occurrence of zoonotic conditions in cattle at slaughter in Morogoro Municipality, Tanzania: implications for public health. Tanzan J Health Res 14(2): 131-8. DOI: 10.4314/thrb.v14i2.6
34. Liu YJ, Xie J, Zhao LJ, Qian YF, Zhao Y, Liu X (2015) Biofilm Formation Characteristics of *Pseudomonas lundensis* Isolated from Meat. J Food Sci 80(12): M2904-10. DOI: 10.1111/1750-3841.13142
35. Manyi-Loh C, Mamphweli S, Meyer E, Okoh A (2018) Antibiotic use in agriculture and its consequential resistance in environmental sources: potential public health implications. Molecules 23(4): 795. doi: 10.3390/molecules23040795
36. Mead PS, Slutsker L, Dietz V, McCaig LF, Bresee JS, Shapiro C, Griffin PM, Tauxe RV (1999) Food-related illness and death in the United States. Emerg Infect Dis 5(5): 607-625. DOI: 10.3201/eid0505.990502
37. Morshdy AM, Hussein MA, El-Arabay AE (2018) Chemical and Microbial Profile of Some Chicken Products. 5th International Food Safety Conference Damanhour University.
38. Mwinyikombo IS (2018) Isolation, antibiotic susceptibility, and molecular characterization of resistance genes in *Pseudomonas* isolates from selected hospitals in mombasa county, kenya. Master thesis in Infectious Diseases in Medical Bacteriology of Kenyatta University. <http://ir-library.ku.ac.ke/handle/123456789/19080>

39. Nychas G-JE, Skandamis PN, Tassou CC, Koutsoumanis KP (2008) Meat spoilage during distribution. *Meat Sci* 78(1-2): 77-89. <https://doi.org/10.1016/j.meatsci.2007.06.020>
40. Orellana-Saez M, Pacheco N, Costa JI, Mendez KN, Miossec MJ, Meneses C, Castro-Nallar E, Marcoleta AE, Poblete-Castro I (2019) In-Depth Genomic and Phenotypic Characterization of the Antarctic Psychrotolerant Strain *Pseudomonas* sp. MPC6 Reveals Unique Metabolic Features, Plasticity, and Biotechnological Potential. *Front Microbiol* 10: 1154. DOI: 10.3389/fmicb.2019.01154
41. Sekyere JO (2018) Genomic insights into nitrofurantoin resistance mechanisms and epidemiology in clinical Enterobacteriaceae. *Future Sci OA* 4(5): FS0293. doi: 10.4155/fsoa-2017-0156
42. Osman K, Orabi A, Elbehiry A, Hanafy MH, Ali AM, (2019) *Pseudomonas* species isolated from camel meat: quorum sensing-dependent virulence, biofilm formation and antibiotic resistance. *Future Microbiol* 14: 609-622. DOI: 10.2217/fmb-2018-0293
43. Pesciaroli C, Barghini P, Cerfolli F, Bellisario B, Fenice M (2015) Relationship between phylogenetic and nutritional diversity in Arctic (Kandalaksha Bay) seawater planktonic bacteria. *Ann Microbiol* 65(4): 2405-2414. DOI <https://doi.org/10.1007/s13213-015-1083-4>
44. Quintieri L, Zühlke D, Fanelli F, Caputo L, Liuzzi VC, Logrieco AF, Hirschfeld C, Becher D, Riedel K, Laura Q (2019) Proteomic analysis of the food spoiler *Pseudomonas fluorescens* ITEM 17298 reveals the antibiofilm activity of the pepsin-digested bovine lactoferrin. *Food Microbiol* 82: 177-193. <https://doi.org/10.1016/j.fm.2019.02.003>
45. Raposo A, Pérez E, de Faria CT, Ferrús MA, Carrascosa C (2017) Food Spoilage by *Pseudomonas* spp.-An overview. In *Foodborne Pathogens and Antibiotic Resistance*. Singh, O.V., Ed., John Wiley & Sons, Inc., Hoboken, N.J., USA. chapter 3: 41–71. <https://doi.org/10.1002/9781119139188.ch3>
46. Rouger A, Tresse O, Zagorec M (2017) Bacterial Contaminants of Poultry Meat: Sources, Species, and Dynamics. *Microorganisms* 5(3): 50. doi: 10.3390/microorganisms5030050
47. Scales BS, Erb-Downward JR, Falkowski NR, LiPuma JJ, Huffnagle GB (2018) Genome Sequences of 12 *Pseudomonas lundensis* Strains Isolated from the Lungs of Humans. *Genome Announc* 6(7): e01461–17. DOI: <https://doi.org/10.1128/genomeA.01461-17>
48. Schreckenberger PC, Binnicker MJ (2011) Optimizing Antimicrobial Susceptibility Test Reporting. *J Clin Microbiol* 49(9): 15-19. doi: 10.1128/JCM.00712-11
49. Shenoy S, Baliga S, Saldanha DRM, Prashanth HV (2002) Antibiotic sensitivity patterns of *Pseudomonas aeruginosa* strains isolated from various clinical specimens. *Indian J Med Sci* 56(9): 427–430. PMID: 12710338
50. Singhal N, Kumar M, Kanaugia PK, Virdi JS (2015) MALDI-TOF mass spectrometry: An emerging technology for microbial identification and diagnosis. *Front Microbiol* 6: 791-806. DOI: 10.3389/fmicb.2015.00791
51. Sivanmaliappan TS, Sevanan M (2011) Antimicrobial susceptibility patterns of *Pseudomonas aeruginosa* from diabetes patients with foot ulcers. *Int J Microbiol* 2011: 605195. DOI: 10.1155/2011/605195
52. Stafnes MH, Dybwad M, Brunsvik A, Bruheim P (2013) Large scale MALDI-TOF MS based taxa identification to identify novel pigment producers in a marine bacterial culture collection. *Antonie Leeuwenhoek* 103(3): 603-615. DOI: 10.1007/s10482-012-9844-6
53. Stellato G, Utter DR, Voorhis A, De Angelis M, Eren AM, Ercolini DA (2017) A Few *Pseudomonas* Oligotypes Dominate in the Meat and Dairy Processing Environment. *Front Microbiol* 8: 264. DOI: 10.3389/fmicb.2017.00264
54. Timperio AM, Gorrasí S, Zolla L, Fenice M (2017) Evaluation of MALDI-TOF mass spectrometry and MALDI BioTyper in comparison to 16S rDNA sequencing for the identification of bacteria isolated from Arctic Sea water. *PLoS One*. 12(7): e0181860. DOI: 10.1371/journal.pone.0181860
55. Tshikhudo P, Nnzeru R, Ntushelo K, Mudau F (2013) Bacterial species identification getting easier. *Afr J Biotechnol* 12(41): 5975-5982. DOI: 10.5897/AJB2013.12057
56. van Belkum A, Welker M, Pincus D, Charrier J, Girard V (2017) Matrix-assisted laser desorption ionization time-of-flight mass spectrometry in clinical microbiology: What are the current issues? *Ann Lab Med* 37(6): 475-483. doi: 10.3343/alm.2017.37.6.475
57. Vithanage NR, Yeager TR, Jadhav SR, Palombo EA, Datta N (2014) Comparison of identification systems for psychrotrophic bacteria isolated from raw bovine milk. *Int J Food Microbiol* 189: 26-38. doi: 10.1016/j.ijfoodmicro.2014.07.023
58. Wang G, Wang H, Han Y, Xing T, Ye K-p, Xu X-l, Zhou G-h (2017) Evaluation of the spoilage potential of bacteria isolated from chilled chicken in vitro and in situ. *Food Microbiol* 63: 139-146. <https://doi.org/10.1016/j.fm.2016.11.015>
59. Wickramasinghe NN, Ravensdale J, Coorey R, Chandry SP, Dykes GA (2019) The Predominance of Psychrotrophic *Pseudomonads* on Aerobically Stored Chilled Red Meat. *Compr Rev Food Sci Food Saf* 18(5): 1622-1635. <https://doi.org/10.1111/1541-4337.12483>
60. Yagoub SO (2009) Isolation of Enterobacteriaceae and *Pseudomonas* spp. from raw fish sold in fish market in Khartoum state. *J Bacteriol Res* 1(7): 85-88. Available online at <http://www.academicjournals.org/jbr>

Tables

Table 1. MIC ranges (µg/ml) of the antibiotics used by Vitek 2 Compact AST GN83 cards

Antimicrobial drug	MIC range (µg/ml)
Amikacin	2–64
Amoxicillin/Clavulanic Acid	2/1–32/16
Ampicillin	2–32
Ampicillin/Sulbactam	2/1–32/16
Aztreonam	1–64
Cefazolin	4–64
Cefepime	1–645
Cefotaxime	0.25–64
Cefoxitin	4–64
Ceftazidime	1–64
Ceftriaxone	0.25–64
Cefuroxime	1–64
Ciprofloxacin	0.25–4
Gentamicin	1–16
Meropenem	0.25–16
Nitrofurantoin	16–512
Piperacillin/Tazobactam	4/4–128/4
Trimethoprim/Sulfamethoxazole	20–320

Table 2. The mean values of colony forming units (CFU/g) for 69 *Pseudomonas* spp. detected in chicken meat products

Chicken meat product	Minimum	Maximum	Mean value ± standard error
Breast (n=8)	2.0 x 10 ²	10.1 x 10 ³	10.1 x 10 ³ ± 1.45 x10 ³
Thigh (n=11)	3.4 x 10 ²	7.4 x 10 ³	7.4 x 10 ³ ± 0.89 x 10 ³
Hamburger (n=25)	3.9 x 10 ²	6.5 x 10 ³	6.5 x 10 ³ ± 6.43 x10 ³
Nuggets (n= 25)	3.7 x 10 ²	4.3 x 10 ³	4.3 x 10 ³ ± 7.56 x10 ³

Table 3. *Pseudomonas* spp. recovered from chicken meat products

<i>Pseudomonas</i> spp.	Chicken meat products								Total	
	Breast		Thigh		Hamburger		Nuggets			
	No.	%	No.	%	No.	%	No.	%	No.	%
<i>P. lundensis</i>	3		2		7		6		18	26.09
<i>P. fragi</i>	1		3		7		5		16	23.19
<i>P. oryzihabitans</i>	3		2		5		3		13	18.84
<i>P. stutzeri</i>	0		3		3		4		10	14.49
<i>P. fluorescens</i>	1		0		2		2		5	7.25
<i>P. putida</i>	0		1		1		2		4	5.80
<i>P. aeruginosa</i>	0		0		0		3		3	4.35
<i>Total</i>	8		11		25		25			

Table 4. Score values for 69 *Pseudomonas* spp. detected in chicken meat by PMFT

Species	Score value of identification							
	2.3–3		2–2.29		1.7–1.99		0–1.69	
	No.	%	No.	%	No.	%	No.	%
<i>P. lundensis</i>	10/18	55.56	8/18	44.44	0	0	0	0
<i>P. fragi</i>	7/16	43.75	9/16	56.25	0	0	0	0
<i>P. oryzihabitans</i>	6/13	46.15	7/13	53.85	0	0	0	0
<i>P. stutzeri</i>	4/10	40	5/10	50	1/10	10	0	0
<i>P. fluorescens</i>	3/5	60	2/5	40	0	0	0	0
<i>P. putida</i>	1/4	25	3/4	75	0	0	0	0
<i>P. aeruginosa</i>	2/3	66.67	1/3	33.33	0	0	0	0

Table 5. Direct screening of the antimicrobial susceptibilities of certain *Pseudomonas* spp. by the VITEK AST GN83 cards

Antimicrobial agent	Percentage of antimicrobial resistance of <i>Pseudomonas</i> spp.															Total (n=6)
	<i>P. lundensis</i>		<i>P. fragi</i>		<i>P. oryzihabitans</i>		<i>P. stutzeri</i>		<i>P. fluorescens</i>		<i>P. putida</i>		<i>P. aeruginosa</i>			
	(n=18)		(n=16)		(n=13)		(n=10)		(n=5)		(n=4)		(n=3)			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	
Amikacin	5	27.78	5	31.25	0	0.00	2	20.00	0	0.00	0	0.00	1	33.33	13	
Amoxicillin/Clavulanic Acid	17	94.44	13	81.25	2	15.38	10	100.00	3	60.00	1	25.00	3	100.00	49	
Ampicillin	18	100.0	12	75.00	1	7.69	10	100.00	3	60.00	2	50.00	3	100.00	49	
Ampicillin/Sulbactam	17	94.44	3	18.75	1	7.69	10	100.00	3	60.00	2	50.00	2	66.67	38	
Aztreonam	2	11.11	16	100	0	0.00	1	10.00	0	0.00	0	0.00	1	33.33	20	
Cefazolin	1	5.56	0	0.00	0	0.00	0	0.00	2	40.00	0	0.00	3	100.00	6	
Cefepime	3	16.67	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	1	33.33	4	
Cefotaxime	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	
Cefoxitin	6	33.33	12	75.00	0	0.00	10	100.00	0	0.00	0	0.00	2	66.67	20	
Ceftazidime	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	1	33.33	1	
Ceftriaxone	15	83.33	15	93.75	3	23.08	10	100.00	0	0.00	0	0.00	2	66.67	45	
Cefuroxime	12	66.67	13	81.25	5	38.46	10	100.00	2	40.00	1	25.00	2	66.67	45	
Ciprofloxacin	16	88.89	14	87.50	0	0.00	3	30.00	0	0.00	0	0.00	1	33.33	34	
Gentamicin	2	11.11	4	25.00	0	0.00	2	20.00	0	0.00	0	0.00	1	33.33	9	
Meropenem	7	38.89	5	31.25	2	15.38	5	50.00	0	0.00	0	0.00	1	33.33	30	
Nitrofurantoin	16	88.89	15	93.75	13	100.00	10	100.00	0	0.00	0	0.00	2	66.67	56	
Piperacillin/Tazobactam	1	5.56	2	12.5	0	0.00	0	0.00	0	0.00	0	0.00	1	33.33	4	
Trimethoprim/Sulfamethoxazole	2	11.11	4	25.00	0	0.00	0	0.00	0	0.00	0	0.00	1	33.33	7	

Figures

Figure 1

Scattering of spectral proteins of 69 *Pseudomonas* spp.; (a) Spreading of peak intensities with a mass charge ratio from 2000 to 10,200 Da; (b) A gel view, in which the wide-ranging colour of dots represents the assembly of different spectra.

Figure 2
Scattering of spectral proteins of 69 *Pseudomonas* spp. (a) Higher peak intensities were distributed inside the line spectra with mass charge ratios from 3000 to 4600 Da; (b) A gel outline of spectral proteins scattered with the same pattern.

Figure 3

Mass spectral protein profiles of different *Pseudomonas* species recovered from chicken meat samples compared with 6 reference strains deposited in the IVD Compass software of Microflex LT. The stored spectral proteins are represented by blue colour in the lower part, while the green spectra in the upper part indicate matched peaks, red spectra indicate incompatible peaks, and yellow spectra indicate intermediate peaks.

Figure 4

A PCA dimensional image showing the grouping of 69 *Pseudomonas* spp.; (a) the alliance of *P. lundensis* (blue), *P. fragi* (yellow), *P. oryzihabitans* (blue black), *P. stutzeri* (aqua), *P. fluorescens* (maroon), *P. putida* (red), and *P. aeruginosa* (green) in the 1st three principal component models.