Short-Term Effects of Two Types of Goggles On Intraocular Pressure and Anterior Eye Segment Biometrics

Xu Zhang

Army Eye Center, Xinjiang Military General Hospital, Urumqi 830000

Huixian Wang

Army Eye Center, Xinjiang Military General Hospital, Urumqi 830000

Yuan Nie

Army Eye Center, Xinjiang Military General Hospital, Urumqi 830000

Wenjing Li (✉ 445263168@qq.com)

Army Eye Center, Xinjiang Military General Hospital, Urumqi 830000

Research Article

Keywords: Intraocular pressure, Central anterior chamber depth, Anterior chamber volume, Anterior chamber angle

Posted Date: November 17th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1069603/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Ophthalmology on February 12th, 2022. See the published version at <https://doi.org/10.1186/s12886-022-02308-y>.

Abstract

Background: To evaluate and compare the changes in intraocular pressure and anterior eye segment biometrics, during and after wearing two types of commonly used swimming goggles.

Methods: In a cross-sectional study, a total of 40 healthy adults aged between 18 and 60 years old were selected to wear two kinds of common swimming goggles (ocular socket and orbital goggles). Intraocular pressure and anterior segment biometry were evaluated before wearing, at 2 and 5 minutes of wearing, and at 5 minutes after removing the goggles. Intraocular pressure (IOP), corneal front keratometry values (K1, K2, Km), central corneal thickness (CCT), central anterior chamber depth (ACD), anterior chamber volume (ACV), and anterior chamber angle (ACA) were measured.

Results: The IOP at 2 minutes (21.0 ± 2.2 mmHg) and 5 minutes (21.2 ± 2.3 mmHg) was significantly higher than before wearing goggles (17.7 ± 2.1 mmHg). The IOP after the goggles were removed and at 5 minutes after the goggles were removed was 18.4 ± 2.3 mmHg and 17.7 ± 2.1 mmHg, respectively. ACV, ACD, and ACA values all decreased while the goggles were worn. After the goggles were removed, these changes gradually returned to baseline values, with no significant difference in the values before and after.

Conclusion: This study proves that wearing orbital goggles can lead to an acute increase in IOP and a slight decrease in ACV, ACD, and ACA. However, once the goggles are removed, these indicators return to baseline levels, showing that wearing orbital goggles has no significant lasting effect on IOP and anterior segment parameters.

Introduction

Swimming is a sport for all ages, its unique charm makes it one of the most popular fitness activities. Swimmers usually wear swimming goggles (SG) to prevent water from entering their eyes and to improve underwater visibility. The SG currently found on the market mainly come in two kinds: sport eye-socket goggles with a large frame and recreational eye-socket goggles with a small frame. Recent studies have shown that when goggles are worn, there is a brief increase in intraocular pressure (IOP) which returns to within the normal range once the goggles are removed.^[1-3]

However, until now, few studies have evaluated the changes in the biological parameters of the anterior segment while wearing SG, and there are no studies on the effects of different types of SG. As the increase of IOP may be related to the type of SG and the contact area between the goggles and the orbital tissues around the eye^{[3][4]}, this study aims to investigate whether differences exist in the short-term effects of wearing two types of SG with regard to the measurement of the anterior segment and IOP. Whether these changes are reversible after the goggles are removed is also investigated.

Objective And Methods

1.1 Objective

In this cross-sectional study, a total of 40 healthy Han Chinese men and women (22 females, 18 males) aged 18 to 60 years (mean age $\pm$ SD, 28.7 $\pm$ 7.6 years) were selected. All participants met the following inclusion criteria: (1) no history of refractive surgery or orthokeratology lens wear; (2) no general or ocular disease, and; (3) no current medications. The following exclusion criteria were used: refractive error (equivalent spherical lens) greater than ± 6 d, and; physical and cognitive understanding ability not sufficient to complete the test. This study was approved by the Ethics Committee of the General Hospital of Xinjiang Military Region and carried out in accordance with the Declaration of Helsinki. All participants signed an informed consent form prior to the study.

1.2 Methods

1.2.1 Interventions

All subjects underwent an initial ophthalmic examination that included best corrected visual acuity, slit lamp, optometry, IOP, Pentacam, and fundus ophthalmoscopy. Based on the examination results, subjects who met the inclusion criteria were selected. Informed consent was obtained to ensure the compliance of subjects.

Head measurement: A tape measure was used to measure the circumference from eye level to above the ears, and this measurement used to calculate the elastic force of the elastic band of the SG. The SG elastic bands were adjusted for each participant, with a fixed perimeter of 10cm smaller than the head circumference. The elastic compression force of the SG was calculated using Hooke's Law: $F_s = K \times 2X$, where: F_s =spring force; K is the elastic constant (rubber elastic band); X is the linear elongation of the material (multiplied by a factor of 2 as the SG have a double rubber belt), calculated as $X = (\text{head circumference} - \text{inside circumference of SG})$.

The experiment was divided into two stages, in which the two types of eye socket goggles (SG1 and SG2) were worn respectively. Both goggle types were made by the same manufacturer (Li Ning, China), consisting of two separate rigid plastic eye cups, each sealed around the edge by rubber buffering, with an adjustable elastic band on either side. The center part of the plastic mask for the right eye was removed in both SGs for inspection and measurement, and the elastic bands of both SGs were adjusted to a fixed circumference 10 cm smaller than the circumference of the subject's head. The SG1 cup measures 60 mm x 35 mm, and the SG2 cup 65 mm x 45 mm. Right eye data was collected from all subjects.

1.2.2 Observation indicators

IOP and anterior segment parameters were assessed in relation to wearing either eye-socket or wide vision swimming goggles SG (SG1 and SG2, respectively). Subjects randomly drew lots to determine which of the two SGs were tested first, with a 10-minute break between the two SGs tests. Each patient underwent Pentacam examination followed by non-contact tonometer examination. For data collection,

participants were seated in front of the Pentacam and a contactless tonometer, while two experienced technicians evaluated the anterior segment parameters and IOP, performing rapid data collection.

The SG1 test was performed on the subjects by taking Pentacam and IOP readings at 5 time points (2 minutes before wearing, at 2 minutes and 5 minutes during wearing, immediately after removal, and 5 minutes after removal). After an interval of 10 minutes, the procedure was repeated with SG2. A total of 10 measurements were taken for each subject, 5 for each SG. The choice of measurement intervals was based on a study by Morgan *et al.*^[1]

1.2.2.1 Intraocular pressure examination

The IOP examination was performed by taking the average of 3 consecutive measurements (in mmHg). Contactless tonometers have been shown to be affected by central corneal thickness (CCT)^[5]. In order to control for this influencing factor, all IOP data was corrected by the Pentacam using the Dresdner formula.^[6]

1.2.2.1 Pentacam check

The anterior segment parameters were obtained by the Pentacam high resolution rotating Scheimpflug camera (Oculus, Germany). This provided 3D anterior segment data and accurately measured the entire anterior segment from the anterior surface of the cornea to the posterior surface of the lens^[7]. In this study, we performed a 25-picture scan and recorded the following criteria: flat axial corneal curvature (K1); steep axial corneal curvature (K2); average corneal curvature (Km); CCT; anterior chamber depth (ACD); anterior chamber angle (ACA), and; anterior chamber volume (ACV).

1.2.3 Statistical analysis

The normal distribution of the data (Shapiro-Wilk test) and the homogeneity of variances (Levene's test) were first determined ($p > 0.05$). All values are presented as a mean $\pm$ standard deviation. Anova of repeated measurements was used for data involving repeated measurements (including the comparison of data at each time point before and after wearing SG), and independent sample T test was used for comparison between groups. For all analyses, an α value of .05 was adopted to define statistical significance and the Holm-Bonferroni correction was adopted for simultaneous comparisons.

Results

2.1 Subject demographics

A total of 22 females and 18 males were selected for this study, with the average age of the subjects being 28.7 ± 7.6 years. A total of 40 eyes (all right eyes) were analyzed, all subjects were Han nationality.

2.2 Comparison of indicators before and after wearing goggles

2.2.1 Comparison of IOP

When wearing SG1, the IOP at both 2 minutes (21.0 ± 2.2 mmHg) and 5 minutes (21.2 ± 2.3 mmHg) was significantly higher than before wearing SG1 (17.7 ± 2.1 mmHg). Immediately after (18.4 ± 2.3 mmHg) and at 5 minutes after SG1 removal (17.7 ± 2.1 mmHg), the IOP decreased significantly ($P < 0.001$), but was still higher than before wearing the goggles ($P = 0.009$). The intraocular pressure recovered to baseline levels by 5 minutes after removal but was statistically significant ($P < 0.001$).

There was no significant change in IOP during or after wearing SG2 ($P = 0.931$).

Table 1
Comparison of intraocular pressure between the two groups

indicators	group	Baseline	2 min	5 min	After removing	Recovery
IOP (mmHg)	SG1	17.7 ± 2.1	21.0 ± 2.2	21.2 ± 2.3	18.4 ± 2.3	17.7 ± 2.1
	SG2	17.6 ± 2.1	17.8 ± 2.0	17.7 ± 2.0	17.6 ± 2.0	17.6 ± 1.9
	t	0.392	6.940	7.382	1.519	0.145
	P	0.696	< 0.001	< 0.001	0.133	0.885

2.2 Comparison of anterior segment parameters

The ACV, ACD, and ACA values were smaller when SG1 was worn compared to before and after wearing and 5 minutes after SG1 was removed. There was no statistically significant change in K1, K2, and Km when SG1 was worn compared with before, after, and 5 minutes after SG1 was removed.

Wearing SG2 did not cause any significant changes in the K1, K2, Km, CCT, ACV, ACD, and ICA values of the subjects.

Table 2
Comparison of indicators before and after wearing two kinds of goggles (SG1 and SG2)

indicators	mirror	Baseline	2 min	5 min	After removing	Recovery
IOP (mmHg)	SG1	17.7 ±2.1	21.0 ±2.2	21.2 ±2.3	18.4 ±2.3	17.7 ±2.1
	SG2	17.6 ±2.1	17.8 ±2.0	17.7 ±2.0	17.6 ±2.0	17.6 ±1.9
K1 (D)	SG1	42.30 ±1.20	42.35 ±1.17	42.32 ±1.18	42.32 ±1.22	42.30 ±1.19
	SG2	42.31 ±1.20	42.30 ±1.21	42.30 ±1.19	42.31 ±1.21	42.31 ±1.19
K2 (D)	SG1	43.29 ±1.20	43.32 ±1.19	43.34 ±1.18	43.29 ±1.16	43.27 ±1.16
	SG2	43.29 ±1.19	43.31 ±1.17	43.30 ±1.21	43.29 ±1.18	43.26 ±1.17
Km (D)	SG1	43.29 ±1.16	43.32 ±1.13	43.34 ±1.11	43.29 ±1.13	43.27 ±1.14
	SG2	43.28 ±1.16	43.3 ±1.17	43.30 ±1.16	43.29 ±1.22	43.26 ±1.18
CCTS (mm)	SG1	536.6 ±28.2	534.2 ±29.4	533.9 ±29.1	536.4 ±27.6	536.2 ±28.3
	SG2	536.3 ±28.4	536.9 ±27.7	536.5 ±28.3	536.8 ±28.4	537.0 ±28.5
ACV (mm)	SG1	200.7 ±33.6	193.0 ±34.2	191.0 ±32.7	198.4 ±31.5	200.3 ±33.5
	SG2	198.9 ±33.1	199.2 ±33.7	199.7 ±34.1	198.4 ±33.4	199.1 ±33.7
ACD (mm)	SG1	3.14 ±0.30	3.08 ±0.29	3.07 ±0.29	3.12 ±0.29	3.15 ±0.29
	SG2	3.14 ±0.30	3.14 ±0.30	3.13 ±0.33	3.14 ±0.30	3.15 ±0.30
ACA (°)	SG1	40.84 ±5.15	39.90 ±5.39	39.37 ±5.30	40.43 ±5.67	41.11 ±5.11
	SG2	40.90 ±5.14	40.77 ±5.25	40.85 ±5.16	40.84 ±5.20	40.98 ±5.15

2.3 Comparison of indicators when wearing different goggles for 2 min and 5 min

When wearing SG1 for 2 min and 5 min, the IOP of subjects was significantly higher than when wearing SG2, and the difference was statistically significant (2 min: $t=6.940$, $P<0.001$; 5 min: $t=7.382$, $P<0.001$). There was no significant difference in ACV, ACD, and ACA values between the two goggles, as shown in Table 3 and Table 4.

Table 3
Comparison of each index between the two types of SG after 2 minutes of wearing

grouping	IOPe (mmHg)	ACD(mm)	ACV (mm)	ACA (°)
SG1	21.0 ±2.2	3.08 ±0.29	193.0 ±34.2	39.90 ±5.39
SG2	17.8 ±2.0	3.14 ±0.30	199.2 ±33.7	40.77 ±5.25
t	6.940	0.904	0.819	0.732
P	< 0.001	0.369	0.415	0.466

Table 4
Comparison of each index between the two types of SG after 5 minutes of wearing

grouping	IOP(mmHg)	ACD(mm)	ACV(mm)	ACA (°)
SG1	21.2 ±2.3	3.07 ±0.29	191.0 ±32.7	39.37 ±5.30
SG2	17.7 ±2.0	3.13 ±0.33	199.7 ±34.1	40.85 ±5.16
t	7.382	0.866	1.165	1.264
P	< 0.001	0.389	0.248	0.210

Discussion

Wearing goggles while swimming or diving can protect eyes from water and prevent the onset of acute conjunctivitis^{[8] [9]}, but the prevention of other diseases has not been reported. On the contrary, adverse reactions to wearing SG have been identified as follows: headache^[10]; nasal deformity^[11]; allergic dermatitis^{[12][13]}; different types of eye lesions, and; changes in intraocular pressure^{[14][15]}. A few studies have reported on the significant correlation between wearing goggles and intraocular pressure, with JesusVera *et al.* also finding that wearing goggles can reduce tear film rupture time and affect tear film stability. However, currently there are only a handful of reports on the influence of wearing goggles on the anterior segment^[16]. Raimundo *et al.* showed that corneal thinning (54.8±41.1 µm), atrial angle narrowing (2.6±2.6°) and an increase in intraocular pressure (4.0±1.9 mmHg) when wearing goggles had no effect on anterior chamber volume^[3]. Also, Dondu Melek Ulusoy *et al.* showed no significant change in anterior segment measurements, with the exception of ACV, while wearing goggles in their study of keratoconus patients, and concluded that the short-term use of goggles does not increase the risk of corneal parameter deterioration in this population^[17].

The results of this study showed that IOP increased significantly in the short term, while ACV, ACD, and ACA decreased, with all returning to the baseline level immediately after the goggles were removed. Corneal curvature, CCT, and other indicators showed no significant change. These findings may have implications for the prevention of a variety of eye diseases, as the increase and fluctuation of IOP is a

potential risk factor for the occurrence and development of glaucoma, as well as an influential factor in early refractive regression after corneal refractive surgery.

Our results are consistent with those of Raimundo *et al.*^[3], which also show an increase in intraocular pressure after wearing SG. However, contrary to our results, Raimundo *et al.* showed that CCT decreased by 55 µm while SG was worn. Both values returned to the pre-wear level after SG removal. The researchers concluded that SGs exert mechanical pressure on the orbital tissue and compress the eyeball, leading to significant thinning of the cornea. This discrepancy with our results may be related to the difference in the orbital margin area caused by ethnic differences.

In addition to verifying our hypothesis that wearing eye socket type goggles results in higher intraocular pressure, this research is novel showing that SGs with different size and design can produce different degrees of compression force on the eye and orbital tissue, and can have different influence on the biological characteristics of the anterior eye.

Drawing on the experimental design of previous studies, we removed the front part of the eyecup of each SG to assess both IOP and anterior segment morphology while they were being worn. Therefore, our experimental design did not accurately replicate realistic conditions, resulting in relatively conservative results. It can be expected that unmodified SG, combined with the effect of external water pressure, may contribute to a larger increase in IOP. In addition, all measurements were taken in the absence of physical activity, and it is important to consider that IOP and anterior segment parameters may vary based on exercise conditions and may also depend on a number of factors, such as exercise intensity, the participant's fitness level, and the time of measurement^{[18][19]}. A potential solution is to use Triggerfish (Sensimed), a new eye pressure sensing contact lens, to assess changes in IOP during swimming^[20]. Secondly, our study was carried out with healthy subjects, so our results cannot be directly applied to patients, such as those with glaucoma, keratoconus, or who have undergone corneal refractive surgery. Finally, the current study is limited to the short-term effects of wearing SG. While a recent study showed that adult swimmers who used SG regularly did not have a higher incidence of glaucoma compared to non-swimmers, further research is needed on the possible long-term effects of wearing SG on eye health^[21]. Despite the above limitations of this study, we believe that, depending on individual differences, some subjects are more sensitive than others to such factors affecting IOP, and the current findings may provide some guidance for the long-term care of some ophthalmic diseases and postoperative patients, especially those who are accustomed to swimming. However, according to our results, the use of eye socket type SG in swimming or water sports may produce negative effects on eye health, especially in patients with eye disease. Thus, we suggest that water sports enthusiasts, especially corneal refractive surgery patients and patients with glaucoma, consider wearing the larger framed orbital SG as opposed to ocular socket SG.

Abbreviations

IOP Intraocular pressure

SG swimming goggles

K1 flat axial corneal curvature

K2 steep axial corneal curvature

Km average corneal curvature;

CCT central corneal thickness

ACD anterior chamber depth

ACA anterior chamber angle

ACV anterior chamber volume

Declarations

Ethics approval and consent to participate

This study was approved by the Ethics Committee of the General Hospital of Xinjiang Military Region and carried out in accordance with the Declaration of Helsinki. All participants signed an informed consent form prior to the study.

- Consent for publication

All the patients included in this study gave written consent for their personal or clinical details to be published in this study.

- Availability of data and material

All data generated or analysed during this study are included in this published article.

- Competing interests

All authors have no competing interests.

- Funding

This study was one of Xinjiang Military General Hospital Beijing Road Medical District key supported scientific research projects. Associated grant numbers 2020jzbj1001

- Authors' contributions

Xu Zhang and Wenjing Li. wrote the main manuscript text. Huixian Wang prepared table 1-4 and Yuan Nie. prepared figures 1. All authors reviewed the manuscript.

- Acknowledgements

Not applicable

References

1. Morgan WH, Cunneen TS, Balaratnasingam C et al. Wearing swimming goggles can elevate intraocular pressure.[J]. Br J Ophthalmol, 2008, 92: 1218–21.
2. Paula Ana PB, Paula Jayter S, Silva Marcelo JL et al. Effects of Swimming Goggles Wearing on Intraocular Pressure, Ocular Perfusion Pressure, and Ocular Pulse Amplitude.[J]. J Glaucoma, 2016, 25: 860-864.
3. Raimundo J, Molina R, Antonio JR et al. Wearing Swimming Goggles Reduces Central Corneal Thickness and Anterior Chamber Angle, and Increases Intraocular Pressure.[J]. Curr Eye Res, 2020, 45: 535–541.
4. Ulusoy DM, Duru Z. Does wearing swimming goggles affect corneal morphology in keratoconic eyes? [J]. Arq Bras Oftalmol, 2020, 83: 225–228.
5. Kohli Poonam G, Randhawa BK, Singh KD et al. Relation between central corneal thickness and intraocular pressure in Punjabi population.[J]. J Med Eng Technol, 2010, 34: 1–6.
6. Janknecht P. Is it Possible to Derive the Dresdner Correction Formula Using a Finite Element Program? [J]. Klin Monbl Augenheilkd, 2021, 09: 56–59.
7. McAlinden C, Khadka J, Pesudovs K. A comprehensive evaluation of the precision (repeatability and reproducibility) of the Oculus Pentacam HR. Invest Ophthalmol Vis Sci. 2011; 52(10): 7731–7.
8. Bonadonna L, La RG. A Review and Update on Waterborne Viral Diseases Associated with Swimming Pools.[J]. Int J Environ Res Public Health, 2019, 16: 166.
9. Li J, Lu X, Sun Y et al. A swimming pool-associated outbreak of pharyngoconjunctival fever caused by human adenovirus type 4 in Beijing, China.[J]. Int J Infect Dis, 2018, 75: 89–91.
10. Ong Jonathan JY, Chan Amanda CY, Bharatendu Chandra et al. Headache Related to PPE Use during the COVID-19 Pandemic.[J]. Curr Pain Headache Rep, 2021, 25: 53.
11. Bodor RM, Breithaupt AD, Buncke GM, et al. Swimmer's nose deformity.[J]. Ann Plast Surg. 2008; 60: 658–660.
12. Tonelli SE, Goldminz AM. Allergic Contact Dermatitis of the Eyelids: Swimming Past Goggle Dermatitis.[J]. Dermatitis, 2020, 31: e12-e13.
13. Sakuragi Y, Sawada Y, Nakamura M. Leukoderma following allergic contact dermatitis caused by the silicone component silprene-30A/B in swimming goggles.[J]. Contact Dermatitis, 2017, 77: 418–419.
14. Jordan DR, Gilberg S, Khouri L. Eyelid masses associated with competitive swimming goggles.[J]. Can J Ophthalmol. 2001; 36: 339–340.
15. Peng KL, Chen KH, Hsu WM, et al. Corneal injury by antimisting agent in swim goggles: a case report. Cornea. 2006; 25: 228–231.

16. Vera J,Redondo B,Molina R et al.Effects of wearing swimming goggles on non-invasive tear break-up time in a laboratory setting.[J].J Optom,2021,388:1–6.
17. Ulusoy DM,Duru Z.Does wearing swimming goggles affect corneal morphology in keratoconic eyes? [J].Arq Bras Oftalmol,2020,83:225–228.
18. Vera J,Garcia-Ramos A,Redondo B,Cárdenas D,De Moraes CG, Jiménez R. Effect of a short-term cycle ergometer sprint training against heavy and light resistances on intraocular pressure responses.[J].J Glaucoma.2018;27(4):315–321.
19. Vera J,Redondo B, Molina R, Garcia-Ramos A, Jiménez R.Influence of holding weights of different magnitudes on intraocular pressure and anterior eye biometrics. [J].Graefe's Arch Clin Exp Ophthalmol.2019,257(10):1–6.
20. Gillmann K,Weinreb RN,Mansouri K.The effect of daily life activities on intraocular pressure related variations in open-angle glaucoma.[J].Sci Rep, 2021,11:6598.
21. McMonnies CW,The importance of and potential for continuous monitoring of intraocular pressure. [J].Clin Exp Optom,2017,100:203–207.

Figures

Figure 1

Eye-socket goggles SG1 (top) and wide vision swimming goggles SG2 (bottom), with the center part of the plastic eye mask for the right eye removed.

Figure 2

Comparison of intraocular pressure changes for SG2 and SG1