

Defining Kawasaki Disease and Pediatric Multi-Inflammatory Syndrome During SARS-CoV-2 Epidemic in Italy: Results From A National, Multicenter Survey.

Marco Cattalini

University of Brescia Department of Medical and Surgical Sciences: Università degli Studi di Brescia
Dipartimento Specialità Medico-Chirurgiche Scienze Radiologiche e Sanità Pubblica

Sara Della Paolera

University of Trieste Department of Surgical Medical and Health Sciences: Università degli Studi di
Trieste Dipartimento Universitario Clinico di Scienze Mediche e Chirurgiche e della Salute

Fiammetta Zunica

ASST Spedali Civili di Brescia: Azienda Socio Sanitaria Territoriale degli Spedali Civili di Brescia

Claudia Bracaglia

Ospedale Pediatrico Bambino Gesù: Ospedale Pediatrico Bambino Gesù

Manuela Giangreco

Institute for Maternal and Child Health: IRCCS materno infantile Burlo Garofolo

Lucio Verdoni

Papa Giovanni XXIII Hospital: Aziende Socio Sanitarie Territoriale Papa Giovanni XXIII

Antonella Meini

City Hospitals of Brescia Health Local Authority: Azienda Socio Sanitaria Territoriale degli Spedali Civili
di Brescia

Rita Sottile

Santobono Pausilipon Azienda Ospedaliera Pediatrica: Azienda Ospedaliera di Rilievo Nazionale
Santobono Pausilipon

Roberta Caorsi

Ospedale Pediatrico Gaslini: Istituto Giannina Gaslini

Gianvincenzo Zuccotti

Vittore Buzzi Children's Hospital: Ospedale dei Bambini Vittore Buzzi

Marianna Fabi

University Hospital of Bologna Sant'Orsola-Malpighi Polyclinic: Azienda Ospedaliero-Universitaria di
Bologna Policlinico Sant'Orsola-Malpighi

Davide Montin

University of Turin Department of Medical Sciences: Università degli Studi di Torino Dipartimento di
Scienze Mediche

Alessandra Meneghel

University of Padova Department of Medicine: Università degli Studi di Padova Dipartimento di Medicina

Alessandro Consolaro

Ospedale Pediatrico Gaslini: Istituto Giannina Gaslini

Rosa Maria Dellepiane

Policlinico of Milan: Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico

Maria Cristina Maggio

University of Palermo: Università degli Studi di Palermo

Francesco La Torre

Paediatric Hospital Giovanni XXIII: Ospedale Pediatrico Giovanni XXIII

Alessandra Marchesi

Bambino Gesù Pediatric Hospital: Ospedale Pediatrico Bambino Gesù

Gabriele Simonini

Children Hospital Meyer: Azienda Ospedaliero Universitaria Meyer

Alberto Villani

Bambino Gesù Pediatric Hospital: Ospedale Pediatrico Bambino Gesù

Rolando Cimaz

University of Milano–Bicocca: Università degli Studi di Milano-Bicocca

Angelo Ravelli

Ospedale Pediatrico Gaslini: Istituto Giannina Gaslini

Andrea Taddio (✉ andrea.taddio@burlo.trieste.it)

Institute of Child and Maternal Health "IRCCS Burlo Garofolo" and University of Trieste

<https://orcid.org/0000-0001-6817-4953>

Research article

Keywords: SARS-CoV-2, Kawasaki Disease, Pediatric Multi-Inflammatory Syndrome temporally associated to COVID-19, Myocarditis, Hypotension, Multisystem Inflammatory Syndrome associated with Coronavirus Disease, Coronary Artery Abnormalities

Posted Date: November 18th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-106975/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on March 16th, 2021. See the published version at <https://doi.org/10.1186/s12969-021-00511-7>.

Abstract

Background: There is mounting evidence on the existence of a Pediatric Multi-inflammatory Syndrome related to SARS-CoV-2 (PIMS-TS), sharing similarities with Kawasaki Disease (KD). The main outcome of the study were to better characterize the clinical features and the treatment response of PIMS-TS and to explore its relationship with KD determining whether KD and PIMS are two distinct entities.

Methods: The Rheumatology Study Group of the Italian Pediatric Society launched a survey to enroll patients diagnosed with KD (Kawasaki Disease Group – KDG) or KD-like (Kawacovid Group - KCG) disease. Demographic, clinical, laboratory data, treatment information, and patients' outcome were collected in an online anonymized database (RedCAP[®]). Relationship between clinical presentation and SARS-CoV-2 infection was also taken into account. Chi square test or exact Fisher test and non-parametric Wilcoxon Mann-Whitney test were used to study differences between two groups

Results: One-hundred-forty-nine cases were enrolled, (96 KDG and 53 KCG). KCG children were significantly older and presented more frequently from gastrointestinal and respiratory involvement. Cardiac involvement was more common in KCG, with 60,4% of patients with myocarditis and 37,8% with hypotension/non-cardiogenic shock. Coronary artery abnormalities (CAA) were more common in the KDG. The risk of ICU admission were higher in KCG. Lymphopenia, higher CRP levels, elevated Ferritin and Troponin-T characterized KCG. KDG received more frequently immunoglobulins (IVIG) and acetylsalicylic acid (ASA) (81,3% vs 66%; p=0.04 and 71,9% vs 43,4%; p=0.001 respectively) as KCG more often received glucocorticoids (56,6% vs 14,6%; p<0.0001). SARS-CoV-2 assay more often resulted positive in KCG than in KDG (75,5% vs 20%; p<0.0001). Short-term follow data showed minor complications. Comparing KDG with a KD-Historical Italian cohort (598 patients), no statistical difference was found in terms of clinical manifestations and laboratory data.

Conclusion: Our study suggests that SARS-CoV-2 infection might determine two distinct inflammatory diseases in children: KD and PIMS-TS. Older age at onset and clinical peculiarities like the occurrence of myocarditis characterize this multi-inflammatory syndrome. Our patients had an optimal response to treatments and a good outcome, with few complications and no deaths.

Background

Italy was the first Western Country to be hit by the SARS-CoV-2 epidemic. To date, more than 236000 cases have been diagnosed, with more than 32000 deaths. Children accounted for almost 2% of infections, with an estimated mortality rate of 0,2%¹. These data confirm previous reports on lower rates of SARS-CoV-2 infection and milder forms of the disease in children, compared to adults²⁻⁴. Nonetheless, few weeks after the epidemic peak, an abnormally high number of severely ill children were seen in those areas of the country with higher SARS-CoV-2 incidence⁵⁻⁷; these observations were then confirmed in other European countries⁸⁻¹⁰. There is now mounting evidence on the existence of a childhood multi-inflammatory syndrome related to SARS-CoV-2 infection sharing some similarities with Kawasaki

Disease (KD) and Toxic Shock Syndrome (TSS)⁹. This condition has been named such as Pediatric Multi-inflammatory Syndrome temporally associated with COVID-19 (PIMS-TS) or Multisystem Inflammatory Syndrome associated with Coronavirus Disease 2019 (MIS-C)^{11,12}.

However, the real extent of the clinical spectrum of disease, and the exact role of SARS-CoV-2 infection, are still poorly understood. Moreover, it is still not clear if SARS-CoV-2 might also be considered a trigger for KD development or if KD, during the SARS-CoV-2 epidemic, presented peculiar and unusual clinical manifestations.

Our study aimed to build a national survey for patients with KD or KD-like multisystemic disease during SARS-CoV-2 epidemic evaluating clinical manifestations, laboratory data, treatment, outcome, and relationship with virus outbreak.

Methods

Study Design and patient selection

This is an observational, retrospective, multicenter study. Institutional Review Board approval was achieved (IRCCS Burlo Garofolo-03/2020). The Rheumatology Study Group of the Italian Pediatric Society launched a national, online, survey on April 24th, 2020 to enroll those patients diagnosed with KD or KD-like multisystemic disease during SARS-CoV-2 epidemic. The children hospitalized between February 1st 2020, and May 31st 2020 with the clinical diagnosis of classical or incomplete-KD (iKD) as well as KD-like multi-inflammatory syndrome were enrolled.

The clinical classification was: 1) KD and iKD diagnosis, named as Kawasaki Disease Group (KDG), based on the fulfilment of the American Heart Association criteria¹³; 2) KD-like multi-inflammatory syndrome diagnosis, named as KawaCOVID Group (KCG), based on the presence of i) persistent fever, lymphopaenia and evidence of single or multi-organ dysfunction with other additional clinical, laboratory or imaging; ii) exclusion of any other microbial cause¹¹.

In consideration of the retrospective nature of the study, an expert panel of pediatric rheumatologists (AR, GS, CB, MC, RC), blinded for patients' recruiting center, was asked to review every patient included in the database to check the correct patient classification or eventually reclassify them properly. Those patients who did not fulfill any of the above-mentioned criteria were excluded from the study.

Data collection

Demographic, clinical, and laboratory data, treatment information, and patient outcome were collected in an online anonymized database which was built for the study purpose (RedCAP→). Data about complications and last follow-up after discharge were collected where available. Results from all tests for SARS-CoV-2 infection with an RT-PCR assay and/or with a serologic assay were also reported. Clinical data collected were part of the normal standard of care.

Statistical analysis

Categorical variables are described as absolute frequency and percentage while continuous variables as median and interquartile range. Chi square test or exact Fisher test were applied to evaluate the association between two categorical variables while non parametric Wilcoxon Mann-Whitney test was used to study differences between two groups of a categorical variable on a continuous variable. A p-value < 0.05 was considered as statistically significant. Statistical analysis was conducted using SAS software, Version 9.4 (SAS Institute Inc., Cary, NC, USA).

Results

Emails to more than 10000 members of the Italian Pediatric Society were sent. Data from 194 patients were entered into the database. After reviewing all the data for duplicates and files with missing information 159 cases were sent to the case definition committee. 10 patients were excluded by the committee since they didn't fulfill the inclusion criteria. 149 patients with a final diagnosis of KDG or KCG were included in the study. 69 patients satisfied KD criteria, 37 iKD criteria, and 53 KawaCOVID criteria. Among the 53 patients classified as KawaCOVID, 10 satisfied also KD or iKD criteria (7 and 3, respectively). The population consisted of 84 males and 65 females, the median age at the time of diagnosis was 3 years (IQR: 1–6 years). Sixty-four out of 149 patients reported at least one symptom in the three months before hospital admittance suggestive for SARS-CoV-2 infection; only one patient (with diarrhea and close contact to a confirmed SARS-CoV-2 patient) had a positive PCR nasal swab for SARS-CoV-2 one month before admission, while 29 patients were tested by nasal swab before disease onset for close contact with confirmed cases, but resulted negative.

More than half of the patients (76/149) were from highly endemic regions (≥ 30000 SARS-CoV-2 confirmed cases in the general population), while the others (73/149) came from regions with moderate-low numbers of SARS-CoV-2 confirmed cases (< 30000 cases in the general population). Distribution of KDG and KCG cases per endemic areas is represented in Fig. 1.

KDG vs KCG at time of diagnosis

KCG children were significantly older (7 years; IQR: 4,5–11 years vs 2 years; IQR: 1–4 years; $p < 0,0001$). Skin and mucosal involvement were more frequent among KDG, reaching statistical significance. Moreover, the presence of conjunctivitis was more frequent among KDG as well as the evidence of irritability. On the other hand, gastrointestinal and respiratory involvement were more often associated to KCG. Clinical manifestations are reported in details in Table 1 and presence of clinical manifestations among the two study groups in Fig. 2. ICU admission was more common in KCG (23,1% vs 1,1%; $p < 0,0001$) as well as the occurrence of secondary Hemophagocytic Lymphohistiocytosis (sHLH) (18,4% vs 1,2%; $p = 0,001$) defined as the presence of HLH-2004 diagnostic criteria¹⁴. Median length of hospital stay was 11 days (IQR: 8–15 days); KCG presented a longer hospitalization (12 days; IQR: 9–17 days vs 10 days; IQR: 7–14 days; $p = 0,02$). No death was reported.

Table 1
Clinical manifestations at disease onset of KawaCOVID Group (KCG) of patients and Kawasaki Disease Group (KDG) of patients.

	KCG (n = 53) N (%)	KDG (n = 96) N (%)	p value
Skin involvement			
Maculo-papular rash	21 (40)	59 (62)	0.001
Erythema multiforme	4 (8)	7 (7)	1
Scarlatiniform rash	0 (0)	8 (8)	0.05
Hands and feet erythema/oedema	15 (28)	38 (40)	0.17
Hands and feet desquamation	1 (2)	6 (6)	0.42
Perineal Erythema/Desquamation	0 (0)	8 (8)	0.05
Mucosal involvement			
Cheilitis	14 (26)	46 (48)	0.01
Pharyngeal and/or oral erythema	24 (45)	49 (51)	0.50
Strawberry tongue	4 (8)	14 (15)	0.29
Ocular involvement			
Non Secretive Conjunctivitis	27 (51)	67 (70)	0.02
Neurological involvement			
Headache	5 (9)	3 (3)	0.13
Consciousness abnormalities	4 (8)	1 (1)	0.05
Aseptic meningitis	3 (6)	1 (1)	0.13
Irritability	8 (15)	34 (35)	0.01
Gastrointestinal involvement			
Diarrhea	28(53)	11 (12)	< 0.0001
Vomiting	14 (26)	8 (8)	0.003
Abdominal pain	17 (32)	10 (10)	0.001
Gallbladder hydrops	2 (4)	3 (3)	1
Respiratory involvement			
Dyspnea	8 (15)	1 (1)	0.001
Tachypnea	12 (23)	4 (4)	0.001

	KCG (n = 53) N (%)	KDG (n = 96) N (%)	p value
Cough	3 (6)	10 (10)	0·38
Rhinitis	0 (0)	5 (5)	0·16
Lobar pneumonia	3 (6)	2 (2)	0·35
Interstitial pneumonia	20 (37)	17 (18)	0·01
O2 supplementation	14 (28)	2 (2)	< 0·0001
Ventilation	6 (12)	0 (0)	0·003
Osteoarticular involvement			
Arthritis and/or arthralgia	6 (11)	13 (14)	0·7

For what concerns the laboratory data, KCG presented lower count of leucocytes, lymphocytes, monocytes and platelets, while CRP levels were higher in KCG as well as Troponin, Troponin-T, Ferritin and D-dimer (Table 2).

Table 2
Laboratory data at disease onset of KawaCOVID Group (KCG) and Kawasaki Disease Group (KDG).

	KCG (n = 53)	KDG(n = 96)	p value
	Median (IQR)	Median (IQR)	
Leucocytes (n/mmc)	11500 (7310–15830)	15730 (11180–19810)	0,003
Neutrophils (n/mmc)	9156 (6480–14230)	10617 (6000–14780)	0,5
Lymphocytes (n/mmc)	940 (630–1570)	2790 (1680–4340)	< .0001
Monocytes (n°/mmc)	320 (160–530)	960 (540–1390)	< .0001
Hemoglobin (g/dL)	11,0 (10,1–11,8)	10,8 (10,0–11,7)	0,31
Platelets (n/mmc)	186000 (132000–292000)	402500 (242000–523000)	< .0001
ALT (U/L)	27 (18–63)	29 (18–81)	1
AST (U/L)	36 (20–65)	30 (21–50)	0,56
GGT (U/L)	31 (16–67)	28 (13–59)	0,59
Creatinine (mg/dL)	0,5 (0,4 - 0,7)	0,3 (0,2 - 0,4)	< .0001
Urea (mg/dL)	25,5 (20,5–47,5)	15,5 (10,5–21,5)	< .0001
Sodium (mEq/L)	134 (130,5-136)	135 (133–137)	0,04
Chlorine (mEq/L)	100 (95–102)	100 (98–102)	0,38
Potassium (mEq/L)	3,9 (3,4–4,4)	4,2 (3,8 - 4,5)	0,01
CRP (mg/L)	242 (138,5-300,5)	96,6 (40,6-195)	< .0001
IgA (mg/dL)	134,5 (98–150)	83 (48,5-160,5)	0,17
IgG (mg/dL)	857 (655–857)	821 (600-1.106)	0,89
IgM (mg/dL)	90 (75–116)	98,5 (90–152,5)	0,31
Total protein (g/dL)	6 (5,2–6,7)	6,6 (5,9 - 7,1)	0,03
Albumin (g/dL)	2,9 (2,6 - 3,5)	3,4 (2,8 - 3,7)	0,05
CPK (U/L)	57 (32–136)	52 (33–117)	0,55
Troponin-T (ng/L)	82,5 (20–126)	3,5 (3–4)	0,002
Cholesterol (mg/dL)	110,5 (87,5-128)	128,5 (93–167)	0,18
Fibrinogen (mg/dL)	643 (515–740)	570,5 (468–714)	0,14
INR (ratio)	1,2 (1,2 - 1,4)	1,2 (1,1–1,3)	0,01

	KCG (n = 53) Median (IQR)	KDG(n = 96) Median (IQR)	p value
aPTT (ratio)	1,2 (1,0–1,3)	1,1 (1,0–1,2)	0,12
C3 (mg/dL)	139,5 (138–141)	138 (121–196)	0,89
C4 (mg/dL)	27,5 (22–33)	36 (24–46)	0,89
Ferritin (ng/mL)	563,8 (250-1.068)	227 (147–449)	0,0004
D-dimer (ng/mL)	2514 (1.380–3.890)	1740 (730 – 2.530)	0,03
NT-BNP (pg/mL)	927 (701-1.734)	347 (81–350)	0,08
ESR (mm/hr)	59 (36–81)	66,5 (49,5–98)	0,18
aPTT (sec)	31,7 (30,9–32,4)	26,9 (23–30)	0,16
PT (sec)	14,3 (13,2–15,2)	75,5 (55–88)	0,08
Troponin (ng/L)	168 (80–672)	4 (3–24)	0,001
Triglycerides (mg/dL)	217,5 (137–261)	144,5 (122–196)	0,04

Cardiac involvement

Seventy-seven patients showed heart involvement during the course of the disease: 35 had myocarditis (60,4% KCG vs 3,1% KDG; $p < 0,0001$), 21 pericarditis (26,4% KCG vs 7,3 KDG; $p = 0,0013$), 19 valvular insufficiency (24,5% KCG vs 6,3% KDG; $p = 0,0036$), 34 coronary artery abnormalities (CAA) (13,2% KCG vs 28,1% KDG; $p = 0,0427$), 20 heart failure (35,8% KCG vs 1% KDG; $p < 0,00001$) and 21 hypotension/non-cardiogenic shock (37,8% KCG vs 1% KDG; $p < 0,0001$).

Data on heart ultrasonography at follow-up was available for 138 patients out of 149. Median time at follow/up was 39,9 days (SD \pm 20,79) after discharge. Normal echocardiograms were showed by 77,2% of patients; 15,4% showed persistent heart ultrasonography abnormalities (15,7% KCG vs 17,2% KDG; $p = 1$), in details 6 out of 8 KCG still had minimal signs of heart failure and 2 valvular insufficiency whether, among KDG, 10 out of 15 children had minimal CAA, 1 patient showed mild pericarditis, 1 patient heart failure and 3 had valvular insufficiency.

Treatment

One-hundred-twenty-eight patients received intravenous immunoglobulins (IVIG), 69 glucocorticoids (67 out of them received intravenous glucocorticoids), 123 acetylsalicylic acid (ASA), 4 hydroxychloroquine, 9 patients anakinra and 1 tocilizumab, 6 anti-viral agents, 93 antibiotics, 16 vasoactive agents and 27 heparins. At time of diagnosis IVIG and ASA were used more frequently in KDG (81,3% vs 66%; $p = 0,04$ and 71,9% vs 43,4%; $p = 0,001$ respectively) while glucocorticoids were more often used in patients with KCG (56,6% vs 14,6%; $p < 0,0001$). Three patients with KCG received anti-IL1 treatment (anakinra) and 1

patient hydroxychloroquine. Antiviral treatment was used only in three patients (2 in the KCG and 1 in KDG), while antibiotics, vasoactive drugs and heparin were used more frequently in KCG (73,6% vs 35,4%, $p < 0,0001$; 20,8% vs 0,0%; $p < 0,0001$ and 28,3% vs 1,0%; $p < 0,0001$ respectively).

Among KCG the use of IVIG alone, glucocorticoids alone or combined with IVIG at the time of diagnosis was not associated with a higher incidence of clinical worsening and/or presence of cardiac involvement at follow-up (data not shown).

Relationship between SARS-CoV-2 infection and KawaCOVID

One-hundred-thirty-six patients out of 149 were tested for SARS-CoV-2 with RT-PCR assay and/or with serologic assay. Fifty-four patients resulted positive (14 only for RT-PCR, 31 only for serologic assay and 9 for both). The positive assays for SARS-CoV-2 were higher in KCG (75,5% vs 20%; $p < 0,0001$).

Since SARS-CoV-2 spread with different prevalence through Italy, with Piedmont and Lombardy being the most heavily hit regions, we compared the data of patients from these two regions (76 patients) (Group A) with the data of patients from the other regions (73 patients) (Group B) irrespectively from diagnostic classification. Group A patients presented more often with diarrhea (36,8% vs 15,1%; $p = 0,003$), myocarditis (29,0% vs 9,6%; $p = 0,003$), pericarditis (15,8% vs 2,7%; $p = 0,01$) and hypotension/non-cardiogenic shock (17,1% vs 2,7%; $p = 0,01$). Again, Group A patients were confirmed to be significantly older (4 years; IQR: 2–8 years vs 2,5 years; IQR: 1–4 years; $p = 0,004$). Laboratory test showed lower WBC (11700/mm³; IQR: 7960–16900/mm³ vs 16250/mm³; IQR: 11180–19810/mm³; $p = 0,03$), lymphocytes (1390/mm³; IQR: 790–2570/mm³ vs 2562,5/mm³; IQR: 1515–4255/mm³; $p = 0,0003$) and platelets (225000/mm³; IQR: 148000–389000/mm³ vs 361000/mm³; IQR: 244000–471000/mm³; $p = 0,01$) in group A. Patients from group A also had higher levels of CRP (195 mg/L; IQR: 82,2-266,7 mg/L vs 96,6 mg/L; IQR: 40,6-197,9 mg/L; $p = 0,003$), and lower levels of ferritin (449 ng/mL; IQR: 227–745 ng/mL vs 218 ng/mL; IQR: 145–458 ng/mL; $p = 0,02$).

While the KDG cases were spread during the study period, the KCG cases were mainly concentrated towards the end of the observation period with about 1 month of delay compared to the peak of the SARS-CoV-2 epidemic (Fig. 3).

Kawasaki Disease during SARS-CoV-2 epidemic

Clinical characteristics of KDG during SARS-CoV-2 epidemic (KDG-CoV2) were compared to 598 Kawasaki Disease patients (KDG-Historical) seen in three different Italian tertiary pediatric hospitals (Institute for Maternal and Child Health, IRCCS “Burlo Garofolo”, Trieste; AOU Meyer, Florence; IRCCS Istituto Giannina Gaslini, Genoa) from January 1st 2000 to December 31st 2019.

No difference was found in age at time of diagnosis, Female/Male ratio, classical/incomplete Kawasaki Disease and clinical manifestations (conjunctivitis, lymphadenopathy, cheilitis, skin rash, extremities changes, sterile pyuria, gastrointestinal involvement, respiratory symptoms, musculoskeletal symptoms,

neurological symptoms, prevalence of CAA, myocarditis, pericarditis, and valvular insufficiency) between KDG and KD-Historical group respectively. No difference was also found in laboratory data available (data not shown). KDG patients from highly endemic areas (Piedmont and Lombardy) were compared to those of medium-low endemic areas. No statistical difference was found in clinical manifestations and laboratory data between the two groups (data not shown).

Discussion

The occurrence of a severe pediatric inflammatory disease, with some characteristics of KD, has been described since late march, in areas with high SARS-CoV-2 incidence^{5,7,9,15-18}. Whether this is a particularly aggressive form of KD triggered by SARS-CoV-2 or a completely different entity is still a matter of debate.

Although some case series have already been reported¹⁵⁻¹⁸, this is a national survey aimed to collect all patients affected by KD and those with KD-like inflammatory symptoms during SARS-CoV-2 epidemic.

Notably, after the first cases of SARS-CoV-2 in the northern Italy regions, the lock-down policy reduced the spread of the infection through other regions.

We confirmed some clinical evidence already reported in the literature; KCG patients are older at disease onset and present more frequently gastrointestinal and respiratory symptoms³, while the classic mucocutaneous symptoms of KD were less common. Children in the KCG also showed higher markers of inflammation, with lower WBC, and platelets^{7,8}. Indeed, lymphopenia is very common in patients with SARS-CoV-2. Accordingly, patients from the KCG were at higher risk of developing sHLH which is indeed rarely described in children with KD²⁰.

Other authors compared patients with PIMS-TS or analogs with a historic cohort of KD patients obtaining similar results^{7,9,16}; the importance of our observation relies on the fact that our KDG was recruited during the same period of KCG with a controlled patients selection.

Patients from the KCG also had significantly longer hospitalization and higher probability to need ICU admittance, for the occurrence of shock, need of vasoactive agents, and invasive ventilation. In the UK, it has been recently reported a higher incidence of ICU admittance of patients with unexplained inflammatory conditions suggestive of PIMS-TS; it is important to underline that in this cohort 2 out of 78 children died²¹. No death was reported in our cohort. Although we did not find any significant correlation between initial treatment and outcome at the end of follow-up, it is important to underline that the use of glucocorticoids was more frequent in KCG and that all KCG patients received glucocorticoids, IVIG or both within 24 hours since admission. This is to confirm that maybe a more aggressive treatment at the time of admission might prevent clinical complications or even death. We would suggest, in absence of stronger clinical evidence, that early treatment with both IVIG (2 gr/kg/day) and glucocorticoids (methylprednisone 2 mg/kg/day) is indicated in KawaCOVID patients; in our experience treatment with

parenteral anakinra (from 2 mg/kg/day to 10 mg/kg/day) is indicated in those patients with rapid clinical worsening or if clinical improvement is not reached within 48 hours since starting IVIG and corticosteroids. Our hypothesis seems to be confirmed from Navallo-Millan *et al.*, suggesting that anakinra could be beneficial in COVID-19 adult patients with evidence of cytokine storm syndrome when initiated early after the onset of Acute Hypoxic Respiratory Failure²².

Heart involvement is peculiar to both KDG and KCG^{6,23}. The major complication in KDG patients was the occurrence of CAA while KCG patients were at higher risk to develop myocarditis with heart insufficiency and related manifestations, such as valvular insufficiency. One of the main unanswered questions on PIMS-TS/MIS-C is if patients with this disease are at risk of developing CAA in the follow-up, as described for KD. For the first time we were able to provide short-term follow-up data on our KCG patients, showing that a minority of patients had minor complications, sequelae of the acute myocarditis, while the 3 KCG patients that developed CAA during hospital stay (aneurysms in 2 patients, dilation in 1), had resolution of the coronary involvement at follow-up. This was in contrast with data from the KDG, where 10/15 patients still presented at follow-up with CAA.

Although PIMS-TS and analogs are considered a complication of SARS-CoV-2 infection, the results from nasal swabs PCR and serologies are usually variable^{14,24}. Data on SARS-CoV-2 exposure were present in 136/149 patients from our cohort. Evidence of SARS-CoV-2 infection was more frequent in patients from the KCG. As already mentioned, SARS-CoV-2 spread with different incidence through Italy. When comparing patients from high endemic regions vs patients from low endemic ones, results were very similar to the analysis of KCG vs KDG, confirming the hypothesis that KawaCOVID is strictly related to SARS-CoV-2 infection.

Similar to other reports, there was a 4-weeks delay in our population between the number of patients with COVID-19 in the general population and the peak of KawaCOVID⁹. Very interestingly, while the number of KawaCOVID cases in the observation period showed the pattern already described, the number of KD was constant throughout the same period. KD incidence is estimated as 5,7 per 100000 children 0–14 y/o in Italy²⁵. Considering only KD diagnosis, the estimated incidence in our study was 3,99 per 100000 children < 14y/o, 5,48/100000 children < 14 y/o considering only patients from highly endemic regions (Piedmont and Lombardy). Notably, since participation in our registry was voluntary, cases may have been under-reported.

Since some of our patients with KD had evidence of SARS-CoV-2 infection, we compared patients from KDG with a historic cohort of Italian KD patients to better define if SARS-CoV-2 could be a trigger of KD and determine a specific pattern of disease. There was no difference in clinical or laboratory characteristics and outcome between the two populations, suggesting that, if SARS-CoV-2 could trigger KD, it does not determine specific disease features.

There is still a debate whether KawaCOVID may be considered a separate clinical entity or a more severe form of KD²⁶, specifically triggered by SARS-CoV-2. The stronger evidence to suggest a disease

continuum between KD and KSS is the remarkable overlapping of clinical features, mainly the mucocutaneous manifestations and the possible formation of coronary artery aneurysms. This overlapping is even more obvious when considering a rare subtype of KD, described by Kanegaye *et al*²⁷ and called Kawasaki Shock Syndrome (KSS), whose main feature is the occurrence of shock. Another important point towards this hypothesis is the good treatment response to systemic glucocorticoids and IVIG²⁸.

On the other hand, previous data suggested that patients with PIMS-TS have some peculiar characteristics when compared to classic KD patients or even KSS^{7,9}. We suggest that SARS-CoV-2 might determine two types of inflammatory diseases in children: the first manifestation is the classic KD, that could be triggered by the coronavirus, as already suggested²⁹. According to our results, the occurrence of this SARS-CoV-2-triggered KD is rare and does not impact significantly on the annual incidence of the disease. The second manifestation related to SARS-CoV-2 exposure during childhood is the multisystem inflammatory syndrome, which affects older children and presents mainly with myocarditis, gastrointestinal symptoms, and the occurrence of shock. Since it appears this latter disease is distinct from KD we recognize the term KawaCOVID could be misleading and both PIMS-TS or MIS-C seem more appropriate. Age at SARS-CoV-2 infection, previous coronaviruses exposure²³, host immunity, and predisposing genetics might have a role in determining which patients would develop a SARS-CoV-2 related disease.

Conclusions

Even though PIMS-TS/MIS-C is a severe disease, the majority of patients respond to treatments, with few complications and a good outcome. As prospective data on larger series are not available, our experience suggests a prompt treatment with glucocorticoids and IVIG. The use of anakinra is also to take into account if clinical worsening is present or in case of a lack of clinical response. Ongoing studies on the treatment of inflammatory conditions related to SARS-CoV-2 might help to define the best options for PIMS-TS patients³⁰.

List Of Abbreviations

ASA: acetylsalicylic acid

CAA: Coronary artery abnormalities

CRP: C-Reactive Protein

iKD: incomplete Kawasaki Disease

IVIG: intravenous immunoglobulins

MIS-C: Multisystem Inflammatory Syndrome associated with Coronavirus Disease

KCG: KawaCovid Group

KDG: Kawasaki Disease Group

PIMS-TS: Pediatric Multi-Inflammatory Syndrome related to SARS-CoV-2

RT-PCR: Real Time-Polimerase Chain Reaction

sHLH: Secondary Hemophagocytic Lymphohistiocytosis

TSS: Toxic Shock Syndrome

Declarations

Ethics approval and consent to participate

Institutional Review Board approval and consent to participate was achieved (IRCCS Burlo Garofolo-03/2020).

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

AR has received grant support and/or speaking or consultant fees from AbbVie, Angelini, Bristol-Myers Squibb, Novartis, Pfizer, Reckitt-Benkiser, Roche and Johnson & Johnson; MC, SDP, FZ, CB, LV, AM, RS, RC, GZ, MF, DM, AM, AC, RMD, MCM, FLT, AM, GS, AV, RC and AT do not have any conflict of interest to declare.

Funding

There no funding to declare

Authors' Contributions

MC, SDP, FZ, CB LV, AM, RS, RC GZ and MF conceptualized, designed the study and drafted the initial manuscript. MG, AT, DM, AM, AC, RD, CM, FL and AM designed the data collection instruments, collected data and performed the statistical analysis. AT, MC, GS, AV, RC and AR conceived of the study, and participated in its design and coordination and helped to draft the manuscript. All authors read and approved the final manuscript

Acknowledgment

Collaborating Authors' names of the Rheumatology Study Group of the Italian Pediatric Society

Paolo Adamoli: Dipartimento Materno Infantile , Ospedale Moriggia Pelascini – Gravedona, Como.

Maria Concetta Alberelli: UOC Pediatria, Marche-Nord; Clotilde Alizzi: Department of Health Promotion Sciences Maternal and Infantile Care, Internal Medicine and Medical Specialities "G. D'Alessandro", University of Palermo, Palermo, Italy; Patrizia Barone: Unità Operativa Complessa di Broncopneumologia Pediatrica AOU "Policlinico - Vittorio Emanuele Via Santa Sofia 78 Catania; Lucia Augusta Baselli: Pediatric Intermediate Care Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; Veronica Bennato: U.O. Pediatria, Ospedale A, Manzoni, Lecco; Francesca Biscaro: UOC Pediatria, Ospedale Ca' Foncello, Treviso; Grazia Bossi: UOC Pediatria, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy; Andrea Campana: Bambino Gesù Children's Hospital, Rome, Italy; Maurizio Carone: UO Malattie Infettive; Ospedale Pediatrico 'Giovanni XXIII', Bari, Italy; Adele Civino: U.O.C. Pediatria P.O. Vito Fazzi, Lecce; Giovanni Conti: Nefrologia e Reumatologia Pediatrica con Dialisi, Azienda Ospedaliero-Universitario "G. Martino", Messina; Eleonora Dei Rossi: University of Trieste, Trieste, Italy; Emanuela Del Giudice: Department of Maternal Infantile and Urological Sciences, Sapeinza University of Rome, Polo Pontini, Rome, Italy; Alice Dell'Anna: U.O.C. Pediatria P.O. Vito Fazzi Lecce; Maia De Luca: Bambino Gesù Children's Hospital, Rome, Italy, Piazza S. Onofrio n. 4, 00165, Rome, Italy; Enrico Felici: Pediatric and Pediatric Emergency Unit, The Children Hospital, AO SS Antonio e Biago e C. Arrigo, Alessandria, Italy; Giovanni Filocamo: Fondazione IRCCS Cà Granda, Ospedale Maggiore Policlinico, Milano; Ilenia Floretta: Pediatria, Ospedale Santa Chiara, Trento, Italy; Maria Loreta Foschini: SC Pediatria, PO SAN MICHELE AOBrotzu, Cagliari, Italy; Marcello Lanari: Department of Pediatrics, University of Bologna, IRCCS S. Orsola-Malpighi Hospital, Bologna, Italy; Bianca Lattanzi: SOD Pediatria, Ospedali Riuniti, Ancona, Italy; Alessandra Lazzerotti: Clinica Pediatrica; Università Milano Bicocca; Fondazione MBBM - onlus c/o Ospedale San Gerardo, Monza, Italy; Francesco Licciardi: Department of Pediatrics and Public Health, University of Turin, Turin, Italy; Alessandra Manerba: Child Cardiology, ASST Spedali Civili di Brescia and University of Brescia, Brescia, Italy; Savina Mannarino: Division of Cardiology, Children's Hospital V Buzzi, ASST FBF Sacco, Milan, Italy; Achille Marino: Department of Pediatrics, Desio Hospital, ASST Monza, Desio, Italy; Agostina Marolda: Pediatrics and Neonatology Department, ASST Ovest Milanese, "G. Fornaroli" Hospital, Magenta, Milan; Laura Martelli: Paediatric Department, Hospital Papa Giovanni XXIII, Bergamo, Italy; Giorgia Martini; Department of Woman's and Child's Health, University of Padova, Padua, Italy; Angela Mauro: Department of Paediatrics, Emergency Department, Santobono-Pausilipon Children's Hospital, Naples, Italy.

Maria Vincenza Mastrolia: Pediatric Rheumatology Unit, AOU Meyer, University of Florence, Florence, Italy.

Angelo Mazza: Paediatric Department, Hospital Papa Giovanni XXIII, Bergamo, Italy; Angela Miniaci: Clinica Pediatrica, Reumatologia, Azienda Ospedaliero-Universitaria di Bologna, IRCCS S. Orsola-Malpighi Hospital, Bologna, Italy; Francesca Minoia: Fondazione IRCCS Cà Granda, Ospedale Maggiore Policlinico, Milano; Alma Olivieri: Dipartimento della donna, del bambino e di chirurgia generale e specialistica,

Università della Campania, “L Vanvitelli, Napoli; Guido Pennoni: Dipartimento Materno-Infantile; Gubbio-Gualdo Tadino, Italy; Rossana Pignataro: UOC Pediatria e Neonatologia, ASST Lodi, Lodi; Francesca Ricci, Clinica Pediatrica, ASST Spedali Civili di Brescia e Università degli Studi di Brescia, Brescia, Italy; Donato Rigante: Department of Pediatrics, Università Cattolica Sacro Cuore, Rome, Italy; Matilde Rossi: UOC di Pediatrai e Neonatologia, Ospedale di Macerata, Macerata; Claudia Santagati: Dipartimento di Pediatria, Ospedale di Rovigo, Rovigo; Martina Soliani: Pediatria ASST Cremona, Italy; Silvia Sonego: University of Trieste, Trieste, Italy; Domenico Sperli: UOC di Pediatria, S.O. “Annunziata” - A.O. di Cosenza; Sara Stucchi: Maternal and Child Health, Division of Paediatrics, ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy; Barbara Teruzzi: Maternal and Child Health, Division of Paediatrics, ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy; Elpidio Tierno: UOC di Pediatria, Dipartimento della Salute della Donna e del Bambin, AORN “Sant’Anna e San Sebastiano”- Caserta; Tatiana Utyatnikova: Dipartimento Materno-Infantile, Pediatria, ASST Bergamo-EST, Seriate, Bergamo; Piero Valentini; Department of Pediatrics, Università Cattolica Sacro Cuore, Rome, Italy; Gianluca Vergine; UOC Pediatria Rimini, Ospedale Infermi, ASL Romagna, Rimini, Italy.

References

1. https://www-epicentro-iss-it.proxy.unibs.it/coronavirus/bollettino/Infografica_10giugno_ITA.pdf (accessed June 12, 2020).
2. Garazzino S, Montagnani C, Donà D, Meini A, Felici E, Vergine G et al. Multicentre Italian study of SARS-CoV-2 infection in children and adolescents, preliminary data as at 10 April 2020. *Euro Surveill.* 2020;25:2000600.
3. Parri N, Lenge M, Buonsenso D. Coronavirus Infection in Pediatric Emergency Departments (CONFIDENCE) Research Group. Children with Covid-19 in Pediatric Emergency Departments in Italy. *N Engl J Med.* 2020;383:187-190.
4. Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z, et al. Epidemiological Characteristics of 2143 Pediatric Patients With 2019 Coronavirus Disease in China. *Pediatrics* 2020;145:e20200702.
5. Licciardi F, Pruccoli G, Denina M, Parodi E, Taglietto M, Rosati S, et al. SARS-CoV-2-Induced Kawasaki-Like Hyperinflammatory Syndrome: A Novel COVID Phenotype in Children. *Pediatrics.* 2020;146:e20201711.
6. Wolfler A, Mannarino S, Giacomet V, Camporesi A, Zuccotti G. Acute myocardial injury: a novel clinical pattern in children with COVID-19. *Lancet Child Adolesc Health.* 2020;4:e26-e27.
7. Verdoni L, Mazza A, Gervasoni A, Martelli L, Ruggeri M, Ciuffreda M, et al. An outbreak of severe Kawasaki-like disease at the Italian epicentre of the SARS-CoV-2 epidemic: an observational cohort study. *Lancet.* 2020;395:1771-8.
8. Riphagen S, Gomez X, Gonzalez-Martinez C, Wilkinson N, Theocharis P. Hyperinflammatory shock in children during COVID-19 pandemic. *Lancet.* 2020;395:1607-1608.
9. Whittaker E, Bamford A, Kenny J, Kaforu M, Jones CE, Shah P, et al. Clinical Characteristics of 58 Children With a Pediatric Inflammatory Multisystem Syndrome Temporally Associated With SARS-

- CoV-2. JAMA. 2020;324:259-269.
10. Grimaud M, Starck J, Levy M, Marais C, Chareyre C, Khraiche D, et al. Acute myocarditis and multisystem inflammatory emerging disease following SARS-CoV-2 infection in critically ill children. *Ann Intensive Care*. 2020;10:69.
 11. Royal College of Pediatrics and Child Health. Guidance: Paediatric multisystem inflammatory syndrome temporally associated with COVID-19. <https://www.rcpch.ac.uk/sites/default/files/2020-05/COVID-19-Paediatric-multisystem-inflammatory-syndrome-20200501.pdf> (accessed May 8, 2020).
 12. Centre for Disease Control and Prevention. Emergency Preparedness and Response Multisystem Inflammatory Syndrome in Children (MIS-C) Associated with Coronavirus Disease 2019. 2020 DOI:10.1016/S0140-6736(20)31094.
 13. McCrindle BW, Rowley AH, Newburger JW, Burns JC, Bolger AF, Gewitz M, et al. Diagnosis, Treatment, and Long-Term Management of Kawasaki Disease: A Scientific Statement for Health Professionals From the American Heart Association [published correction appears in *Circulation*. 2019 Jul 30; 140: e181-e184]. *Circulation*. 2017; 135:e927-e999.
 14. Henter JI, Horne A, Aricó M, Egeler RM, Filipovich AH, Imashuku S, et al. HLH-2004: Diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer*. 2007;48:124-131.
 15. Belot A, Antona D, Renolleau S, Javouhey E, Hentgen V, Angoulvant F, et al. SARS-CoV-2-related paediatric inflammatory multisystem syndrome, an epidemiological study, France, 1 March to 17 May 2020. *Euro Surveill*. 2020;25:2001010.
 16. Pouletty M, Borocco C, Ouldali N, Caseris M, Basmaci R, Lachaume N, et al. Paediatric multisystem inflammatory syndrome temporally associated with SARS-CoV-2 mimicking Kawasaki disease (Kawa-COVID-19): a multicentre cohort. *Ann Rheum Dis*. 2020;79:999-1006.
 17. Dufort EM, Koumans EH, Chow EJ, Rosenthal EM, Muse A, Rowlands J, et al. Multisystem Inflammatory Syndrome in Children in New York State. *N Engl J Med*. 2020;383:347-358.
 18. Feldstein LR, Rose EB, Horwitz SM, Collins JP, Newhams MM, Son MBF et al. Multisystem Inflammatory Syndrome in U.S. Children and Adolescents. *N Engl J Med*. 2020;383:334-346.
 19. European Centre for Disease Control and Prevention. Paediatric inflammatory multisystem syndrome and SARS-CoV-2 infection in children. 2020 <https://www.ecdc.europa.eu/sites/default/files/documents/covid-19-risk-assessment-paediatric-inflammatory-multisystem-syndrome-15-May-2020.pdf>.
 20. García-Pavón S, Yamazaki-Nakashimada MA, Báez M, Borjas-Aguilar KL, Murata C. Kawasaki Disease Complicated With Macrophage Activation Syndrome: A Systematic Review. *J Pediatr Hematol Oncol*. 2017;39:445-451.
 21. Davies P, Evans C, Kanthimathinathan HK, Lillie J, Brierly J, Waters G, et al. Intensive care admissions of children with paediatric inflammatory multisystem syndrome temporally associated with SARS-

- CoV-2 (PIMS-TS) in the UK: a multicentre observational study *Lancet Child Adolesc Health*. 2020; S2352-4642(20)30215-7. DOI:10.1016/S2352-4642(20)30215-7
22. Navarro-Millán I, Sattui SE, Lakhanpal A, Zisa D, Siegel CH, Crow MK. Use of Anakinra to Prevent Mechanical Ventilation in Severe COVID-19: A Case Series. *Arthritis Rheumatol*. 2020;10.1002/art.41422. DOI:10.1002/art.41422
 23. Belhadjer Z, Méot M, Bajolle F, Khraiche D, Legendre A, Abakka S, et al. Acute heart failure in multisystem inflammatory syndrome in children (MIS-C) in the context of global SARS-CoV-2 pandemic. *Circulation*. 2020;DOI:10.1161/CIRCULATIONAHA.120.048360.
 24. Toubiana J, Poirault C, Corsia A, Bajolle F, Fourgeaud J, Angoulvant F, et al. Kawasaki-like multisystem inflammatory syndrome in children during the covid-19 pandemic in Paris, France: prospective observational study. *BMJ*. 2020; 369: m2094. Published 2020 Jun 3.
 25. Cimaz R, Fanti E, Mauro A, Voller F, Rusconi F. Epidemiology of Kawasaki disease in Italy: surveillance from national hospitalization records. *Eur J Pediatr*. 2017;176:1061-5.
 26. Ravelli A, Martini A. Kawasaki disease or Kawasaki syndrome?. *Ann Rheum Dis*. 2020; 79: 993-5.
 27. Kanegaye JT, Wilder MS, Molkara D, Frazer JR, et al. Recognition of a Kawasaki disease shock syndrome. *Pediatrics*. 2009;123:e783-e789.
 28. Taddio A, Rossi ED, Monasta L, Pastore S, Tommasini A, Lepore L, et al. Describing Kawasaki shock syndrome: results from a retrospective study and literature review. *Clin Rheumatol*. 2017;36:223-8.
 29. Jones VG, Mills M, Suarez D, Hogan CA, Yeh D, Segal JB, et al. COVID-19 and Kawasaki Disease: novel virus and novel case. *Hosp Pediatr*. 2020;10:537-40.
 30. [No author] Best available treatment study for inflammatory conditions associated with COVID-19. <https://doi.org/10.1186/ISRCTN69546370>.

Figures




Figure 1

SARS-CoV-2 infection and KD/KawaCOVID patients in Italy. Distribution of KDG patients (blue numbers) and KCG (green numbers) in different areas per SARS-CoV-2 incidence (right squares indicate SARS-CoV-2 number of patients/general population). KDG = Kawasaki Disease Group; KCG = KawaCOVID Group. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.




Figure 2

Trend of clinical manifestations among the KawaCOVID Group (KCG) and the Kawasaki Disease Group (KDG)




Figure 3

Relationship between SARS-CoV-2 infection trend and new cases of Kawasaki Disease and KawaCOVID during SARS-CoV-2 infection in Italy from February 1st 2020 to May 31st 2020.