Periodic Sea-Level Oscillation in Tokyo Bay Detected with the Tokyo-Bay Seafloor Hyper-Kilometric Submarine Deep Detector (TS-HKMSDD)

Hiroyuki K. M. Tanaka (✉ ht@eri.u-tokyo.ac.jp)

University of Tokyo

Masaatsu Aichi

University of Tokyo

Szabolcs József Balogh

Wigner Research Centre for Physics

Cristiano Bozza

University of Salerno

Rosa Coniglione

Istituto Nazionale di Fisica Nucleare-Laboratori Nazionali del Sud, Catania, Italy.

Jon Gluyas

Durham University

Naoto Hayashi

University of Tokyo

Marko Holma

Muon Solutions Oy Ltd

Jari Joutsenvaara

Muon Solutions Oy Ltd

Osamu Kamoshida

NEC (Japan)

Yasuhiro Kato

University of Tokyo

Tadahiro Kin

Kyushu University

Pasi Kuusiniemi

Muon Solutions Oy Ltd

Giovanni Leone

The University of Atacama

Domenico Lo Presti

University of Catania

Jun Matsushima

University of Tokyo

Hideaki Miyamoto

University of Tokyo

Hirohisa Mori

University of Tokyo

Yukihiro Nomura

University of Tokyo

Naoya Okamoto

Kyushu University

László Oláh

University of Tokyo

Sara Steigerwald

International Virtual Muography Institute (VMI), Global

Kenji Shimazoe

University of Tokyo

Kenji Sumiya

International Virtual Muography Institute (VMI), Global

Hiroyuki Takahashi

University of Tokyo

Lee F. Thompson

University of Sheffield

Tomochika Tokunaga

University of Tokyo

Yusuke Yokota

University of Tokyo

Sean Paling

Boulby Underground Laboratory

Dezső Varga

Wigner Research Centre for Physics

Research Article

Keywords: Periodic, oscillation, Hyper-Kilometric, TS-HKMSDD

Posted Date: December 3rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1069909/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Meteorological-tsunami-like (or meteotsunami-like) periodic oscillation was muographically detected with the Tokyo-Bay Seafloor Hyper-Kilometric Submarine Deep Detector (TS-HKMSDD) deployed in the underwater highway called the Trans-Tokyo Bay Expressway or Tokyo Bay Aqua-Line (TBAL). It was detected right after the arrival of the 2021 Typhoon-16 that passed through the region 400 km south of the bay. The measured oscillation period and decay time were respectively 3 hours and 10 hours. These measurements were found to be consistent with previous tide gauge measurements. Meteotsunamis are known to take place in bays and lakes, and the temporal and spatial characteristics of meteotsunamis are similar to seismic tsunamis. However, their generation and propagation mechanisms are not well understood. The current result indicates that a combination of muography and trans-bay or trans-lake underwater tunnels will offer an additional tool to measure meteotsunamis at locations where tide gauges are unavailable.

Introduction

Meteotsunamis or meteorological tsunamis are tsunami-like sea-level oscillations that take place in closed or semi-closed water bodies in bays or lakes with periods ranging from minutes to several hours¹. As the generation of these tsunamis is related to one of the natural oscillation modes of the bay or the lake, their period and amplitude are a function of the size, depth and the configuration of the coastline². The temporal and spatial characteristics of meteotsunamis and seismic tsunamis are similar, and shifting atmospheric disturbances, which are usually caused by sudden atmospheric pressure and/or wind changes, are the significant factors that will induce oscillations in water bodies. Atmospheric energy is transferred to a body of water more efficiently and in a more concentrated manner when the propagation speed of the atmospheric disturbance is approximately equal to the local free wave speed. This process is even more efficient if the water depth is within the optimal range³. Stronger atmospheric disturbances usually generate larger scale oscillations. Meteotsunamis are associated with frontal passages⁴, cyclones^{5,6}, atmospheric gravity waves⁷, and mesoscale convective systems^{8,9}, including derechos¹⁰, and have been reported worldwide^{1,11}. Importantly, their impacts on human communities and infrastructure are often severe¹¹ due to their high wave runup and strong associated currents^{12,13,14,15}. Even meteotsunamis with moderate heights (~0.3 m) generate hazardous currents^{16,17}. Owing to the ubiquitous nature of atmospheric disturbances, associated meteotsunamis can add to the risk posed by seismic tsunamis¹⁸ or can increase the risk to regions not traditionally recognized as seismic-tsunami-prone¹⁹. However, our quantitative understanding of the associated risks and frequency, generation, and propagation mechanisms related to meteotsunamis is limited^{20,21}.

A typhoon is defined as a tropical cyclone (TC) when it develops in the Northwestern Pacific Basin, which has been recognized as one of the most active tropical cyclone areas in the world²². In Japan, typhoon measurements have been recorded since 1951²³. In Tokyo Bay, meteotsunamis induced by typhoons and the sea surface currents induced by meteotsunamis have been respectively measured with a tide gauge²⁴

and High Frequency Radar (HFR)²⁵. Results from these measurements are consistent with each other, and have differentiated two oscillation modes (OM) in Tokyo Bay, more specifically, OM1 (usually 2-3 hours duration) and OM2 (usually 5-6 hours duration). It has been interpreted that OM1 occurs when the tsunami is confined in the northern part of Tokyo Bay, whereas OM2 is associated with the tsunamis that occur throughout the entire longitudinal length of Tokyo Bay²⁵.

Muography is similar to x-ray imagery, but it utilizes the strong penetration capability of high-energy muons (> a few tens of GeV) and their relativistic effect. Since the number of muons that pass through gigantic bodies reflects the interior spatial distribution of density, this distribution can be mapped by identifying where these muons passed through the object and subsequently creating a plot of the number of penetrating muons on a 2-dimensional plane. The origin of these high-energy muons is galactic cosmic rays (GCR) which are accelerated by high-energy events such as supernovas in our galaxy. The GCRs mainly consist of protons and alpha particles. These charged particles are generally accelerated to close to the speed of light. Although the galactic magnetic field is mainly aligned with the spiral galaxy, there is also a random component. The direction that cosmic rays travel is strongly affected by this random component of the galactic magnetic field. As a result, cosmic rays travel for millions of years (depending on their energy) before arriving at the Solar system. Consequently, by the time they arrive here, their initial direction of origin is completely lost as they have obtained an isotropic distribution of arrival directions. These cosmic rays interact with the Earth's atmosphere, and muons are generated. Muography takes advantage of the characteristics of muons, particularly their penetrative nature and universality, for a wide variety of applications, including visualizing the internal structure of volcanoes, tunnels, natural caves, and cultural heritage. So far, applications have focused in Africa^{26,27}, the Americas^{27,29,30}, Asia^{31,32,33,34,35,36,37,38,39,40,41}, and Europe^{42,43,44,45,46,47,48}.

The Tokyo-Bay Seafloor Hyper-Kilometric Submarine Deep Detector (TS-HKMSDD) consists of a linear array of several particle detectors located inside the underwater tunnel called the Tokyo Bay Aqua-Line (TBAL). The first and second segments of TS-HKMSDD were respectively installed in March 2021²³ and June 2021. The current total length of TS-HKMSDD is 200 m with a total active area of 3 m². This article reports the results found thus far, including a meteotsunami-like periodical oscillation in muon flux as observed with TS-HKMSDD right after the typhoon approach to Tokyo.

Results

The 2021-Typhoon-16 originated at 13.6° N and 143.3° E. at 21:00 September 23, 2021. Figures 1A, 1B, and 1C show the meteorological history of Typhoon-16. The Japan Meteorological Agency (JMA) reported that this typhoon tracked first westward but transiently shifted to a more northerly course on September 26²³. At 09:00 on September 26, Typhoon-16 was estimated to have attained Category 3 winds (178-208 km/h). At 15:00 JST on September 26, Typhoon-16 achieved its minimum barometric pressure at 920 hPa. This fall in barometric pressure indicated a 78 hPa pressure drop in the preceding 48 hours. The typhoon's winds continued to increase before peaking at 15:00 JST on September 26, with its

maximum winds reaching 198 km/h²³. Typhoon-16 only maintained peak intensity for 15 hours, but even after this, it remained a powerful tropical cyclone²³. With very little change in barometric pressure from 920 hPa to 945 hPa, the typhoon was tracked to be moving in a northeastward direction throughout October 1. Meanwhile, the storm area (>54km/h) had grown from E: 390 km W: 280 km to E: 750 km W: 560 km. At that moment, Typhoon-16 was located 400 km south of Tokyo. Following this approach of Typhoon-16 to Tokyo Bay, the barometric pressure observed in Yokohama, Japan, had dropped by ~20 hPa in 24 hours between 17:00 JST on September 30 and 17:00 JST on October 1 (Figure 1D). At 21:00 JST on October 1, the typhoon had weakened to tropical storm intensity and transformed into an extratropical cyclone at 09:00 JST on October 2.

The Trans-Tokyo Bay Expressway, also known as TBAL, is a combined bridge and tunnel structure spanning the entire width of Tokyo Bay, Japan (Figure 2). It consists of a 4.4-km long bridge and a 9.6-km long tunnel underneath the bay. The tunnel section is called the Aqua-Tunnel. The average sea depth is 20 m in most of the region where the Aqua-Tunnel was constructed. The tunnel was constructed at a depth of 20 m underneath the seafloor. In this work, 20 muographic sensor modules (MSMs) were deployed inside the Aqua-Tunnel to construct a linear array of MSMs called the Tokyo-Bay Seafloor Hyper-Kilometric Submarine Deep Detector (TS-HKMSDD). Since a detailed description of HKMSDD can be found elsewhere⁴⁹, it is only briefly introduced here. Each MSM consisted of two scintillation detectors, and only the events producing signals at both detectors in coincidence were considered muon events. A 2-cm-thick lead block was inserted between these plastic scintillators to reduce the number of random coincidences resulting from electronic noise or the gamma-rays emitted from the tunnel's concrete wall.

Figure 3A shows the muon count rate (MCR) as recorded every 5 minutes at TS-HKMSDD for 3 weeks, including the period when Typhoon-16 approached and occupied the region 400 km south of Tokyo Bay. At TS-HKMSDD, a large portion of the inverse barometric effect (IBE) is cancelled except for the small residual IBE coming from the muon's different energy loss rate between air and water⁵³. There had been a clear anti-correlation with the astronomical tide height (ATH) variations (Figure 3C) except for the period right after the severe atmospheric pressure drop observed at Tokyo Bay on October 1. Figure 3B focuses on the MCR data recorded for 12 days including the period when Typhoon-16 approached the region 400 km south of Tokyo Bay. As can be seen in the red box of Figure 3B, this disturbance was an oscillation with a period of ~3 hours. The starting point and the duration of the red box were respectively corresponding to the timing when the minimum pressure was observed in Yokohama (17:00 JST on October 1) and the oscillation decay time (10 hours) calculated based on the damping coefficient ($2.45 \times 10^{-5} \text{ s}^{-1}$) of Tokyo Bay that was modelled for the oscillation observed during the passage of Typhoon-15 on September 11, 2001²⁴.

Discussion

Oscillation decay time. Figure 4A focuses on the time region within the red box in Figure 3. The observed oscillation period was ~3 hours and is consistent with the period (155 minutes) measured right after the

passage of 2015-Typhoon-15²⁵, which was associated with the confined mode (OM1). Figures 4B and 4C focuses on two other periods that follow the time region shown in Figure 4A (03:00-13:00 on October 2 and 13:00-23:00 on October 2).

For validation of the current FFT computation, Figure 5 compares the results of the FFT of MCR and ATH⁴⁹ as they were recorded during the previous 2 months (August 1 - September 30). Four fundamental tidal constants are well reproduced, and the spectrum shapes are almost identical between MCR and ATH. The tidal constituents were those of lunar diurnal A (O_1), lunar diurnal B (K_1), principal lunar semidiurnal (M_2), and principal solar semidiurnal (S_2). Figure 5 essentially shows five major tidal constituents of tide levels at Tokyo Bay, Japan⁵⁰.

Figure 6 shows the time series of the abnormal tides measured muographically between 15:50 on October 1 and 07:20 on October 2, 2021. These time series were derived by subtracting ATH from the tide levels converted from MCR⁴⁹. The decay curves (Eq. 1) with a damping coefficient modeled for Tokyo Bay and Lake Geneva are overlaid.

$$H=H_0\exp(-\beta t), (1)$$

where the damping coefficient β is $2.43 \times 10^{-5} \text{ s}^{-1}$ for Tokyo Bay and $6.83 \times 10^{-6} \text{ s}^{-1}$ for Lake Geneva²⁵. Based on Figures 4-6, it can be concluded that the oscillation decay time is consistent with the value previously modeled for Tokyo Bay.

In conclusion, it has been shown that muography conducted inside an underwater tunnel has the potential to become a valuable tool for monitoring meteotsunamis in bays and lakes in regions where no other measurement tools are available. Understanding the regional tides of inner bays is not only vital for navigation safety, but also necessary for environmental hazard assessments. The concept of HKMSDD is deployable at any given underwater tunnel of the appropriate size and depth worldwide. For example, the Transbay Tube at San Francisco Bay, CA, where a well-known meteotsunami event was induced by a moving pressure pulse on November 21, 1910⁵¹, has similar characteristics to the TBAL site currently used for TS-HKMSDD. Other meteotsunami examples can be found in the English Channel^{9,52} and the Gulf of Finland⁵³. The Channel Tunnel connects the UK to France and could be used for similar measurements. Similarly, the underwater tunnel across the Gulf of Finland to connect Finland and Estonia could be a good candidate location if the tunnel project is realized. Various trans-bay and under-lake tunnels exist or are under construction globally. It is anticipated that the same muography HKMSDD configuration could be installed in several underwater tunnels worldwide to serve as a local and global sea level monitoring array.

Declarations

Acknowledgments

This project was supported by the East Nippon Expressway Company Limited, the Geospatial Information Authority of Japan, the Ministry of Land, Infrastructure, Transport and Tourism, Ministry of Education, Culture, Sports, Science and Technology, and the Japan Society for the Promotion of Science. Fruitful discussions with the Japan Coast Guard are also acknowledged. The authors presented this work on behalf of the MAGMA-HKMSDD Collaboration (see Supplementary Information 1).

Contributions

H.K.M.T., C.B., R.C., J.G., M.H., J.J., L.O. L.F.T. and S.S. wrote the text. H.K.M.T. prepared the figures. All authors reviewed the manuscript.

Corresponding author

Correspondence to Hiroyuki K. M. Tanaka.

Competing interests

The authors declare no competing interests.

References

1. Monserrat, S., Vilibić, I. & Rabinovich, A. B. Meteotsunamis: atmospherically induced destructive ocean waves in the tsunami frequency band. *Nat. Hazards Earth Syst. Sci*, **6**, 1035–1051 <https://doi.org/10.5194/nhess-6-1035-200> (2006).
2. Rueda, F. J. & Schladow, S. G. Surface seiches in lakes of complex geometry. *Limnol. Oceanogr*, **47**, 906–910 <https://doi.org/10.4319/lo.2002.47.3.0906> (2002).
3. Proudman, J. Atmospheric pressure and tides. *Geophys. suppl. mon. not. R. Astron. Soc*, **2**, 197–209 (1929).
4. Tanaka, K. Atmospheric pressure-wave bands around a cold front resulted in a meteotsunami in the East China Sea in February 2009. *Nat. Hazards Earth Syst. Sci*, **10**, 2599–2610 <https://doi.org/doi:10.5194/nhess-10-2599-2010> (2010).
5. Mercer, D., Sheng, J. Y., Greatbatch, R. J. & Bobanovic, J. Barotropic waves generated by storms moving rapidly over shallow water. *J. Geophys. Res*, **107**, 3152 <https://doi.org/doi:10.1029/2001jc001140> (2002).
6. Ozsoy, O., Haigh, I. D., Wadey, M. P., Nicholls, R. J. & Wells, N. C. High-frequency sea level variations and implications for coastal flooding: A case study of the Solent, UK. *Cont. Shelf Res*, **122**, 1–13 <https://doi.org/doi:10.1016/j.csr.2016.03.021> (2016).
7. Monserrat, S., Ibbetson, A. & Thorpe, A. J. Atmospheric gravity-waves and the rissaga phenomenon. *Q. J. R. Meteorol. Soc*, **117**, 553–570 <https://doi.org/doi:10.1002/qj.49711749907> (1991).
8. Wertman, C. A. *et al.* Mesoscale convective system surface pressure anomalies responsible for meteotsunamis along the US East Coast on June 13th, 2013. *Sci. Rep*, **4**, <https://doi.org/doi:>

- 10.1038/srep07143 (2014).
9. Tappin, D. R. *et al.* The English Channel 'tsunami' of 27 June 2011 - a probable meteorological source. *Weather*, **68**, 144–152 <https://doi.org/doi: 10.1002/wea.2061> (2013).
 10. Sepic, J. & Rabinovich, A. B. Meteotsunami in the Great Lakes and on the Atlantic coast of the United States generated by the “derecho” of June 29-30, 2012. *Nat. Hazards*, **74**, 75–107 <https://doi.org/doi: 10.1007/s11069-014-1310-5> (2014).
 11. Pattiaratchi, C. B. & Wijeratne, E. M. S. Are meteotsunamis an underrated hazard? *Philos. Trans. R. Soc. A Math., Phys. Eng. Sci*, **373**, 2053 (2015).
 12. Ewing, M., Press, F. & Donn, W. J. An explanation of the Lake Michigan wave of 26 June 1954., **120**, 684–686, doi: <https://doi.org/10.1126/science.120.3122.684> (1954).
 13. Orlic, M., Belusic, D., Janekovic, I. & Pasaric, M. Fresh evidence relating the great Adriatic surge of 21 June 1978 to mesoscale atmospheric forcing. *J. Geophys. Res*, **115**, C06011 <https://doi.org/doi: 10.1029/2009jc005777> (2010).
 14. Cho, K. H. *et al.* A synoptic study on tsunami-like sea level oscillations along the west coast of Korea using an unstructured-grid ocean model. *J. Coast. Res*, **65**, 678–683 <https://doi.org/doi: 10.2112/si65-115.1> (2013).
 15. Haslett, S. K., Mellor, H. E. & Bryant, E. A. Meteo-tsunami hazard associated with summer thunderstorms in the United Kingdom. *Phys. Chem. Earth*, **34**, 1016–1022 <https://doi.org/doi: 10.1016/j.pce.2009.10.005> (2009).
 16. Wilson, R. I. *et al.* Observations and Impacts from the 2010 Chilean and 2011 Japanese Tsunamis in California (USA). *Pure Appl. Geophys*, **170**, 1127–1147 <https://doi.org/doi: 10.1007/s00024-012-0527-z> (2013).
 17. Wilson, R. I., Dengler, L. A., Legg, M. R., Long, K. & Miller, K. M. The Chilean tsunami on the California coastline. *Seismol. Res. Lett*, **81**, 545–546 (2010).
 18. Pattiaratchi, C. B. & Wijeratne, E. M. S. Are meteotsunamis an underrated hazard? *Philos. Trans. R. Soc. A Math., Phys. Eng. Sci*, **373**, 20140377 <https://doi.org/doi: 10.1098/rsta.2014.0377> (2015).
 19. ten Brink, U. S., Chaytor, J. D., Geist, E. L., Brothers, D. S. & Andrews, B. D. Assessment of tsunami hazard to the US Atlantic margin. *Mar. Geol*, **353**, 31–54 <https://doi.org/doi: 10.1016/j.margeo.2014.02.011> (2014).
 20. Geist, E. L., ten Brink, U. S. & Gove, M. A framework for the probabilistic analysis of meteotsunamis. *Nat. Hazards*, **74**, 123–142 <https://doi.org/doi: 10.1007/s11069-014-1294-1> (2014).
 21. Bechle, A. J., Kristovich, D. A. R. & Wu, C. H. Meteotsunami occurrences and causes in Lake Michigan. *J. Geophys. Res*, **120**, 8422–8438 <https://doi.org/doi: 10.1002/2015jc011317> (2015).
 22. Peduzzi, P. *et al.* Global trends in tropical cyclone risk. *Nat. Clim. Chang*, **2**, 289–294 (2012).
 23. Japan Meteorological Agency. Weather, Climate & Earthquake Information, <http://www.jma.go.jp/jma/indexe.html> (2021)

24. Kanari, S. Damping Coefficient and Lunar Frictional Coefficient of Tokyo Bay during the Passage of Typhoon No. 15 on 11 Sept. 2001. <http://ris-geo.jp/pdf/publication/05-043.pdf> (2003)
25. Suzuki, K. & Tsubokawa, Y. Detection of seiche caused by typhoon using current velocity data. https://www.jstage.jst.go.jp/article/kaigan/71/2/71_I_127/_article/-char/ja/ (2015).ã€€
26. Alvarez, L. W. *et al.* Search for hidden chambers in the pyramid., **167**, 832–739 (1970).
27. Morishima, K. *et al.* Discovery of a big void in Khufu's Pyramid by observation of cosmic-ray muons., **552**, 386–390 (2017).
28. Lesparre, N. *et al.* Density muon radiography of La Soufrière of Guadeloupe volcano: Comparison with geological, electrical resistivity and gravity data. *Geophys. J. Int*, **190**, 1008–1019 (2012).
29. Jourde, K. *et al.* Muon dynamic radiography of density changes induced by hydrothermal activity at the La Soufrière of Guadeloupe volcano. *Sci. Rep*, **6**, 33406 (2016).
30. Rosas-Carbajal, M. *et al.* Three-dimensional density structure of La Soufrière de Guadeloupe lava dome from simultaneous muon radiographies and gravity data. *Geophys. Res. Lett*, **44**, 6743–6751 (2017).
31. Tanaka, H. K. M. *et al.* High resolution imaging in the inhomogeneous crust with cosmic-ray muon radiography: The density structure below the volcanic crater floor of Mt. Asama, Japan. *Earth Planet. Sci. Lett*, **263**, 104–113 (2007).
32. Tanaka, H. K. M. *et al.* Imaging the conduit size of the dome with cosmic-ray muons: The structure beneath Showa-Shinzan Lava Dome, Japan. *Geophys. Res. Lett*, **34**, 053007 (2007).
33. Tanaka, H. K. M., Uchida, T., Tanaka, M., Shinohara, H. & Taira, H. Cosmic-ray muon imaging of magma in a conduit: Degassing process of Satsuma-Iwojima Volcano, Japan. *Geophys. Res. Lett*, **36**, L01304 (2009).
34. Tanaka, H. K. M., Kusagaya, T. & Shinohara, H. Radiographic visualization of magma dynamics in an erupting volcano. *Nat. Commun*, **5**, 3381 (2014).
35. Tanaka, H. K. M. Muographic mapping of the subsurface density structures in Miura, Boso and Izu peninsulas, Japan. *Sci. Rep*, **5**, 8305 (2015).
36. Tanaka, H. K. M. Instant snapshot of the internal structure of Unzen lava dome, Japan with airborne muography. *Sci. Rep*, **6**, 39741 (2016).
37. Olah, L., Tanaka, H. K. M., Ohminato, T. & Varga, D. High-definition and low-noise muography of the Sakurajima volcano with gaseous tracking detectors. *Sci. Rep*, **8**, 3207 (2018).
38. Tanaka, H. K. M., Sumiya, K. & Oláh, L. Muography as a new tool to study the historic earthquakes recorded in ancient burial mounds. *Geosci. Instrum. Method Data Syst*, **9**, 357–364 <https://doi.org/10.5194/gi-9-357-2020> (2020).
39. Tanaka, H. K. M. Development of the muographic tephra deposit monitoring system. *Sci. Rep*, **10**, 14820 <https://doi.org/10.1038/s41598-020-71902-1> (2020).
40. Oláh, L., Tanaka, H. K. M. & Hamar, G. Muographic monitoring of hydrogeomorphic changes induced by post-eruptive lahars and erosion of Sakurajima volcano. *Sci. Rep*, **11**, 17729

- <https://doi.org/10.1038/s41598-021-96947-8> (2021).
41. Oláh, L., Hamar, G., Miyamoto, S., Tanaka, H. K. M. & Varga, D. The first prototype of an MWPC-based borehole-detector and its application for muography of an underground pillar. *Geophys. Explor*, **71**, 161–168 <https://doi.org/10.3124/segj.71.161> (2018).
 42. Oláh, L. *et al.* CCC-based muon telescope for examination of natural caves. *Geosci. Instrum. Method Data Syst*, **1**, 229–234 (2012).
 43. Tioukov, V. *et al.* First muography of Stromboli volcano. *Sci. Rep*, **9**, 6695 (2019).
 44. Cimmino, L. *et al.* 3D muography for the search of hidden cavities. *Sci. Rep*, **9**, 2974 <https://doi.org/10.1038/s41598-019-39682-5> (2019).
 45. Lo Presti, D. *et al.* Muographic monitoring of the volcano-tectonic evolution of Mount Etna. *Sci. Rep*, **10**, 11351 <https://doi.org/10.1038/s41598-020-68435-y> (2020).
 46. Tompson, L. F. *et al.* Muon tomography for railway tunnel imaging. *Phys. Rev. Res*, **2**, 023017 <https://doi.org/10.1103/PhysRevResearch.2.023017> (2020).
 47. Ambrosino, F. *et al.* Joint measurement of the atmospheric muon flux through the Puy de Dôme volcano with plastic scintillators and Resistive Plate Chambers detectors. *J. Geophys. Res*, **120**, 7290–7307 (2015).
 48. Saracino, G. *et al.* Imaging of underground cavities with cosmic-ray muons from observations at Mt. Echia (Naples). *Sci. Rep*, **7**, 1181 (2017).
 49. Tanaka, H. K. M. *et al.* First results of undersea muography with the Tokyo-Bay Seafloor Hyper-Kilometric Submarine Deep Detector. *Sci. Rep*, **11**, 19485 <https://doi.org/10.1038/s41598-021-98559-8> (2021).
 50. Japan Coast Guard. Chiba real time tide gauge data (2021). https://www1.kaiho.mlit.go.jp/TIDE/gauge/gauge_eng.php?s=0053
 51. Wilson, B. W. & Seiches *Adv. Hydrosci.* **8**, 1–94 (1972)
 52. Williams, D. A. *et al.* Examination of Generation Mechanisms for an English Channel Meteotsunami: Combining Observations and Modeling. *J. Phys. Oceanogr*, **49**, 103–120 (2019).
 53. Pellikka, H. *et al.* Meteotsunami occurrence in the Gulf of Finland over the past century. *Nat. Hazards Earth Syst. Sci*, **20**, 2535–2546 (2020).

Figures

Figure 1

Meteorological history of 2021-Typhoon-16. The trajectory of the Typhoon-16 (A), the central pressure (B), and the maximum wind velocity (C) are shown as a function of the time since the moment of its development on September 24. The severe pressure drop observed on October 1 in Tokyo Bay (Yokohama) is also shown (D). HKMT drew the map based on the data in Reference23 and the image with Microsoft PowerPoint software and holds the copyright.

Figure 2

Location of the Tokyo Bay Aqua-Line (TBAL). The inset shows the magnified view of the southern central part of Japan that includes Tokyo Bay (A). The red line indicates the location of TBAL. The cross-sectional view of TBAL is shown in the middle panel (B). The symbol "HKMSDD-SEG" indicates the location of the currently installed TS-HKMSDD that spans 200 m along the tunnel. The photograph of TS-

HKMSDD is also shown (C). HKMT drew the map and the image with Microsoft PowerPoint software and holds the copyright of the images and photographs.

Figure 3

Muon count rate (MCR) as recorded with TS-HKMSDD. The time series plots are shown for 3 weeks (A) and 12 days (B), including the period when Typhoon-16 approached and occupied the region 400 km south of Tokyo Bay. The astronomical tide height (ATH) variations (C) are also shown for the same period as in Panel A. The red boxes indicate the time region when the periodic oscillation was observed right after the pressure drop observed in Tokyo bay.

Figure 4

Time series showing the muon count rate (MCR) in different time windows (A-C) and their Fast Fourier Transformation (FFT) results (D-F). Vertical bars in (A-C) indicate the one standard deviation error bars. Figures 4D, 4E and 4F show the result of the Fast Fourier Transformation (FFT) of MCR recorded in the time regions shown in Figures 4A, 4B and 4C, respectively. An oscillation with a frequency of ~ 100 micro-Hz (a period of ~ 3 hours) can be seen only in Figure 4D. The first peak seen in all of the figures (Figures 4D, 4E and 4F) is associated with ATH (for more details, see Figure 5).

Figure 5

Power spectra of MCR (A) and ATH (B), recorded between August 1 and September 30. The power spectrum of MCR is superimposed to that of ATH (C). Symbols O_1 , K_1 , N_2 , M_2 , and S_2 respectively indicate the lunar diurnal A, lunar diurnal B, larger lunar elliptic semi diurnal, principal lunar semi diurnal, and principal solar semi diurnal constituents. In these plots, the amplitude of the muographic M_2 peak in (A) was normalized to the astronomical M_2 peak (B).

Figure 6

Time series of the abnormal tides muographically measured between 15:50 on October 1 and 07:20 on October 2, 2021. The decay curves calculated for Tokyo Bay (orange) and Lake Geneva (green) are overlaid. Vertical bars indicate the standard deviation error bars.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [41598202198559MOESM1ESM10272021.docx](#)