

Crowdsourcing to Develop Open-Access Learning Resources on Antimicrobial Resistance

Eneyi Kpokiri (✉ eneyi.kpokiri@lshtm.ac.uk)

London School of Hygiene & Tropical Medicine

Randall John

University of North Carolina at Chapel Hill

Dan Wu

London School of Hygiene & Tropical Medicine

Noah Fongwen

London School of Hygiene & Tropical Medicine

Jehan Budak

University of Washington

Christina Chang

Massachusetts General Hospital

Jason Ong

Monash University

Joseph Tucker

University of North Carolina at Chapel Hill

Research Article

Keywords: antimicrobial resistance, infectious diseases educational stewardship, medical education, curriculum development, crowdsourcing

Posted Date: February 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-107133/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Objectives

Antimicrobial resistance (AMR) is a significant threat to global public health. Many medical curricula have limited clinical cases and materials focused on AMR. Enhanced AMR education and training are needed to support antimicrobial stewardship programmes. We used crowdsourcing methods to develop open-access, learner-centred AMR resources. Crowdsourcing is the process of having a large group, including experts and non-experts, solve a problem and then share solutions with the public.

Methods

We organised a global crowdsourcing contest soliciting AMR-related multiple-choice questions, infographics, and images. First, we convened a diverse steering committee group to finalise a call for entries. Second, we launched the contest and disseminated the call for entries using social media, blog posts, email, and an in-person event. Partner institutions included two digital healthcare platforms: Fig. 1[®] and Ding Xiang Yuan. Both organizations serve as online communities for healthcare specialists and professionals to report and comment on clinical information. At the end of the call, solicited entries were screened for eligibility and judged on merit and relevance to AMR learning and education. Exceptional entries were recognised, awarded prizes, and further reviewed for sharing with the public via open-access platforms.

Results

We received 59 entries from nine countries. These included 54 multiple-choice questions, four infographics, and one image. Eligible entries (n = 56) were reviewed and assigned a score on a 1–10 scale. Eight entries received mean scores greater than 6.0 and were selected as finalists. The eight finalist entries consisted of three infographics and five multiple-choice questions. They were disseminated through open-access publications and online medical communities.

Conclusions

We demonstrate that crowdsourcing challenge contests can be used to identify infectious disease teaching materials. Medical educators and curriculum developers can adapt this method to solicit additional teaching content for medical students.

Background

Antimicrobial resistance (AMR) is a major global health problem. Antimicrobial stewardship programmes are increasingly designed to enhance and expand medical school infectious disease curricula.^{1–3} Educational interventions have been shown to improve antimicrobial use and practices.^{4,5} Some clinicians, however, still have limited understanding of antimicrobial stewardship and AMR. Studies from the United States and Europe suggest gaps in medical student exposure to appropriate antimicrobial

prescribing practices and AMR.^{6,7} In addition, separate surveys in the Congo and Ethiopia suggest poor levels of AMR understanding among healthcare providers and students.^{8,9} Other studies from the United Kingdom and Belgium also demonstrate that there are inconsistencies between antibiotic prescribing guidelines and clinician practices.^{10,11} In response, we sought to identify medical education resources on AMR using crowdsourcing methods. Crowdsourcing has a large group, including experts and non-experts, solve a problem and then share solutions with the public.¹² In the past, crowdsourcing has been used to expand existing medical curricula and develop flashcard study tools for preclinical education.^{13–15} Additionally, researchers have successfully crowdsourced challenging, high quality, and complex multiple-choice questions (MCQs) from medical students.¹⁶

MCQs are preferred by some students as study tools, and surveys among healthcare providers suggest a preference for communicating clinical information through infographics in comparison to conventional text reports.^{16,17} The purpose of this study was to examine a crowdsourcing contest soliciting AMR infographics, MCQs, and images for medical teaching. Medical educators and curriculum developers can adopt this method in the future to create AMR-focused learning materials for educational and antimicrobial stewardship efforts.

Methods

The crowdsourcing contest design was based on the framework provided by the UNICEF/UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases (TDR) Practical Guide on Crowdsourcing in Health and Health Research.¹² Crowdsourcing has six steps: selecting crowdsourcing as the methodology, convening a steering committee, engaging communities to participate, receiving and judging contributions, recognising finalists, and sharing solutions (Table 1).

The open challenge contest

The International Diagnostics Centre at the London School of Hygiene and Tropical Medicine and SESH (Social Entrepreneurship to Spur Health) organised this contest. The contest was officially launched in April 2019, and the call was open for two months. An open call for entries was provided on a contest website and promoted using the online medical learning platforms and other social media channels. All entries were screened for eligibility. Eligible entries were evaluated by three clinical experts who were identified by the steering committee. Eight entries that achieved a mean score of 6.0 or greater emerged finalists and were awarded a total of 1000 USD in prize money through Amazon gift cards and cash. The MCQs were arranged into a slide deck similar to an AMR learning module developed through a challenge contest.¹⁸ The finalist infographics were published as posters on F1000Research, a life sciences-focused digital publishing platform.^{19–22} We asked participants to specify the AMR learning objective(s) (Table 2) that their entry addressed. The contest was organised in line with terms and conditions as specified by the legal committee of the London School of Hygiene and Tropical Medicine (LSHTM). As part of the conditions of the contest, participants were made to obtain informed consent from subjects where any

personal data was included in the entry. Ethics approval was deemed unnecessary by the institutional review board at LSHTM.

Results

We received 59 entries with 56 eligible entries that came from nine countries: Cameroon (n = 30), the United States (n = 10), Nigeria (n = 5), the United Kingdom (n = 3), Australia (n = 3), Jordan (n = 2), Singapore (n = 1), India (n = 1), and China (n = 1). Of the 56 eligible entries, there were 51 MCQs, four infographics, and one image. Of the 56 eligible entries, 54 were solicited through the official contest website, and two were solicited through the online learning platforms.

The average score of all entries (n = 56) was 4.84. Score distributions showed that the majority of entry scores fell within a range of 3.00 to 5.99 (n = 46). Breakdown by entry type shows that infographics (n = 4) had an average score of 7.00, images (n = 1) had an average score of 5.00, and MCQs (n = 51) had an average score of 4.67.

Finalist entries centred on a wide range of topics in AMR education and research. MCQs selected as finalists focused primarily on AMR background information and prevention/treatment. Topics included effective infection control in health institutions, multi-drug resistant organisms, antibiotic usage in animal farming, antibiotic treatment options in response to persistent symptoms, and mechanisms of AMR. Submitters also identified veterinary practitioners and farmers as a target audience due to the increasing prevalence of antibiotics in agriculture practices and raising livestock.²³ The finalist infographics addressed AMR background information, treatment, and diagnosis. Finalist submissions are included in the **Supplementary materials**.

Discussion

We have demonstrated that crowdsourcing methods can be used to identify open-access medical education materials on antimicrobial resistance (AMR). Our findings support existing literature demonstrating that crowdsourcing is an efficient method to develop educational resources in the medical and public health fields.^{13,15,16} This challenge contest is a unique example of how to implement crowdsourcing methods to create medical education curricula specifically for the purposes of enhancing antimicrobial stewardship efforts.

This study draws on insights and examples from different settings. The contest was global in scope, as we received 56 eligible entries from nine different countries across five continents. We were able to solicit entries from both high-income and low-income countries, as well as entries from different practice areas, such as human medicine, veterinary medicine, and hospital- and community-based medicine. In addition, we were able to identify relevant online platforms to support contest implementation. The use of online platforms facilitated broad dissemination to an international audience, and spurred engagement surrounding AMR and antibiotic prescribing practices. The inclusion of several social media metrics in

our study also offers key insight into the use of digital platforms in crowdsourcing challenge contests and can guide future contest-organisers who wish to interact with online platforms for contest organisation and promotion.

This challenge contest received high-quality submissions, consistent with other crowdsourcing studies.²⁴ In our contest, eight entries (representing 14% of all submissions) achieved a mean score of 6.0 or greater, which was similar to the frequency of high-quality submissions in another global health innovation contest on Hepatitis B and C.²⁴ In terms of entry dissemination, three infographics were identified for online publication through F1000Research, nine MCQs were shared through the online learning platforms and 22 MCQs were selected for inclusion in a study slide deck on AMR. 25 out of 56 eligible entries (representing 43% of the submissions) were selected for dissemination, which was slightly higher than the dissemination frequency from other challenge contests.^{24,25}

Crowdsourcing has several advantages. First, through contest promotion, we were able to spur creativity and awareness surrounding AMR and acquired a diverse and global range of ideas. Soliciting MCQs, infographics, and images on AMR from medical students, physicians, and other healthcare professionals in multiple countries suggests that crowdsourcing is also feasible across different settings. Second, medical curricula development can be a time-consuming and challenging task for a small number of individuals.^{13,26} We demonstrate that a bottom-up crowdsourcing approach can be a cost-effective method to develop medical teaching materials rapidly, decreasing the potential burden on educators and curriculum developers. This suggests that similar to other studies, crowdsourced materials from both experts and non-experts can be used in medical education.^{14,15,27} However, consistent with existing literature, one disadvantage to our crowdsourcing approach was the need for solicited contributions to be significantly edited and refined.¹³

Although many people viewed the contest promotion announcements on online platforms, we received few submissions from them: only two entries were submitted through the online platforms. This data suggests that paid online platforms to promote participation in challenge contests may be less effective, indicating a need for additional crowdsourcing interventions studying the use of paid online platforms. Given that in-person promotion of challenge contests has been associated with a greater volume of entries, more attention to in-person activities may also be useful for promotion.^{28,29}

Our study has implications for research and policy surrounding medical education and curriculum development. While our study shows how crowdsourcing is an effective strategy to develop additional study resources in medical education, there is a need for more research to evaluate the impact and effectiveness of these educational resources. Robust programmes are essential in evaluating the extent to which the use of medical education stewardship approaches translate into improvements in clinical practice and understanding. There is also a need to review current curricula on AMR to identify content gaps and inform future projects.

Our contest had some limitations. First, we heavily targeted medical student groups. However, the timing of the call for entries overlapped with the examination calendar of many medical and public health schools, while others were already on break. Second, our participation may have been limited as we only promoted the contest at one in-person AMR event. Third, although our contest was global in scope, our entries were not entirely representative of all global settings, as there were no entries from the Latin American region. Last, entries accepted through the online learning platforms were limited to those in the English language and Chinese (Ding Xiang Yuan). This may have affected participation from non-English and non-Chinese speaking countries.

Conclusions

This study enhances our understanding of crowdsourcing in the context of medical education. Our contest demonstrates that crowdsourcing can be used to increase study materials available for medical students and physicians. Clinical educators should consider adopting crowdsourcing approaches to enhance medical education and mitigate traditional barriers associated with curriculum development. There is a need for additional research testing the impact and efficacy of crowdsourced clinical training resources for students and practitioners.

Declarations

Ethics approval and consent to participate

Not applicable. The institutional review board at the London School of Hygiene and Tropical Medicine deemed this work a non-human subject research and ethics was not necessary. All participants in the open call completed an informed consent statement at the point of submitting the entries.

Consent for publication

Not applicable. Where patients' medical records were used, informed consent was obtained by participants prior to submission and all identifying information has been removed.

Availability of data and materials

Additional data collected from entries are available in the supplementary files

Conflicts of Interest

The authors declare that they have no competing interests

Funding

This project was supported by the Academy of Medical Sciences and the Newton Fund (Grant number NIF\R1\181020).

Author Contributions

JDT conceived the study design with input from JWB, CCC, and EEK. RJ, DW and NF facilitated promotion of the call. JWB, CCC, and JJO reviewed finalist entries. EEK and RJ prepared the results and drafted the manuscript, and all authors reviewed and approved the final version.

Acknowledgements

The authors wish to thank SESH (Social Entrepreneurship to Spur Health) for creating and hosting the contest website, and the online platforms (Figure 1[®] and Ding Xiang Yuan) for providing support in promoting the open call for entries. We also thank members of the steering committee for providing guidance at all stages of the contest and all the judges who volunteered to review submissions.

References

1. Hsu JL. Building an Antibiotic Stewardship Program: An Interactive Teaching Module for Medical Students. *MedEdPORTAL: the Journal of Teaching and Learning Resources*.14.
2. Silverberg SL, Zannella VE, Countryman D, et al. A review of antimicrobial stewardship training in medical education. *International journal of medical education*. 2017;8:353–374.
3. Laks M, Guerra CM, Miraglia JL, Medeiros EA. Distance learning in antimicrobial stewardship: innovation in medical education. *BMC medical education*. 2019;19(1):191.
4. Kandeel A, Palms DL, Afifi S, et al. An educational intervention to promote appropriate antibiotic use for acute respiratory infections in a district in Egypt- pilot study. *BMC Public Health*. 2019;19(3):498.
5. Roque F, Herdeiro MT, Soares S, Teixeira Rodrigues A, Breitenfeld L, Figueiras A. Educational interventions to improve prescription and dispensing of antibiotics: a systematic review. *BMC Public Health*. 2014;14(1):1276.
6. Abbo LM, Cosgrove SE, Pottinger PS, et al. Medical students' perceptions and knowledge about antimicrobial stewardship: how are we educating our future prescribers? *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America*. 2013;57(5):631–638.
7. Dyar OJ, Pulcini C, Howard P, Nathwani D. European medical students: a first multicentre study of knowledge, attitudes and perceptions of antibiotic prescribing and antibiotic resistance. *The Journal of antimicrobial chemotherapy*. 2014;69(3):842–846.
8. Thriemer K, Katuala Y, Batoko B, et al. Antibiotic prescribing in DR Congo: a knowledge, attitude and practice survey among medical doctors and students. *PloS one*. 2013;8(2):e55495.
9. Abera B, Kibret M, Mulu W. Knowledge and beliefs on antimicrobial resistance among physicians and nurses in hospitals in Amhara Region, Ethiopia. *BMC pharmacology & toxicology*. 2014;15:26.
10. Palin V, Mölter A, Belmonte M, et al. Antibiotic prescribing for common infections in UK general practice: variability and drivers. *Journal of Antimicrobial Chemotherapy*. 2019;74(8):2440–2450.

11. Mainjot A, D'Hoore W, Vanheusden A, Van Nieuwenhuysen J-P. Antibiotic prescribing in dental practice in Belgium. 2009;42(12):1112–1117.
12. Han L, Chen A, Wei S, Ong JJ, Iwelunmor J, Tucker JD. *Crowdsourcing in Health and Health Research: A Practical Guide*. Geneva: World Health Organization;2018.
13. Blackwell KA, Travis MJ, Arbuckle MR, Ross DA. Crowdsourcing medical education. *Medical education*. 2016;50(5):576.
14. Tackett S, Raymond M, Desai R, et al. Crowdsourcing for assessment items to support adaptive learning. *Medical teacher*. 2018;40(8):838–841.
15. Bow HC, Dattilo JR, Jonas AM, Lehmann CU. A crowdsourcing model for creating preclinical medical education study tools. *Academic medicine: journal of the Association of American Medical Colleges*. 2013;88(6):766–770.
16. Grainger R, Dai W, Osborne E, Kenwright D. Medical students create multiple-choice questions for learning in pathology education: a pilot study. *BMC medical education*. 2018;18(1):201.
17. Turck CJ, Silva MA, Tremblay SR, Sachse SL. A Preliminary Study of Health Care Professionals' Preferences for Infographics Versus Conventional Abstracts for Communicating the Results of Clinical Research. 2014;34(S1):S36-S38.
18. SESH Global. The clinical diagnostics and AMR learning module. <https://www.seshglobal.org/AMRLearningModule>. Accessed.
19. Khan S, John R, Wu D, Kpokiri E. Multidrug resistant fungal infections: challenges arising with treatment options [version 1; not peer reviewed] (poster). 2020;9. doi:<https://doi.org/10.7490/f1000research.1117825.1>.
20. Khan S, John R, Wu D, Kpokiri E. Carbapenem resistant gram negative CNS infections: threats to commonly used antimicrobials in treatment of MDR gram negatives [version 1; not peer reviewed] (poster). 2020;9. doi:<https://doi.org/10.7490/f1000research.1117823.1>.
21. Khan S, John R, Wu D, Kpokiri E. Ceftazidime-Avibactam-Aztreonam synergy testing: The use of combination ceftazadime-avibactam and aztreonam in the treatment of metallobeta-lactamase producing carbapenem resistant gram negative infections [version 1; not peer reviewed] (poster). 2020;9. doi:<https://doi.org/10.7490/f1000research.1117824.1>.
22. F1000. <https://f1000research.com/>. Accessed.
23. Stop using antibiotics in healthy animals to prevent the spread of antibiotic resistance. World Health Organization. <https://www.who.int/news-room/detail/07-11-2017-stop-using-antibiotics-in-healthy-animals-to-prevent-the-spread-of-antibiotic-resistance>. Published 2017. Accessed.
24. Tucker JD, Meyers K, Best J, et al. The HepTestContest: a global innovation contest to identify approaches to hepatitis B and C testing. *BMC infectious diseases*. 2017;17(Suppl 1):701.
25. Tang W, Wei C, Cao B, et al. Crowdsourcing to expand HIV testing among men who have sex with men in China: A closed cohort stepped wedge cluster randomized controlled trial. *PLoS medicine*. 2018;15(8):e1002645.

26. Shappell E, Chan TM, Thoma B, et al. Crowdsourced Curriculum Development for Online Medical Education. *Cureus*. 2017;9(12):e1925.
27. Dai JC, Lendvay TS, Sorensen MD. Crowdsourcing in Surgical Skills Acquisition: A Developing Technology in Surgical Education. 2017;9(6):697–705.
28. Zhang Y, Kim JA, Liu F, et al. Creative Contributory Contests to Spur Innovation in Sexual Health: 2 Cases and a Guide for Implementation. *Sexually transmitted diseases*. 2015;42(11):625–628.
29. Zhang Y, Tang S, Li K, et al. Quantitative evaluation of an innovation contest to enhance a sexual health campaign in China. *BMC infectious diseases*. 2019;19(1):112.

Tables

Table 1 Stages employed in the contest

Organising a steering committee

The purpose of the steering committee was to provide guidance and outline an overall framework for contest execution. We convened a steering committee composed of ten individuals from five countries that met periodically through 60-minute teleconference meetings over the duration of the contest to discuss design, organisation, and implementation. Members included experts in medicine, public health, infectious diseases, and medical education, as well as representatives from partner organisations who helped with contest promotion and dissemination.

Engaging the community to contribute

The contest was officially launched on April 15, 2019. The website provided detailed information on the contest, including the purpose, categories of participation, rules, steering committee members, and partner organisations. In order to foster creativity and avoid cognitive fixation, we did not provide examples of entries. Promotional information was disseminated through social media (Instagram and Twitter), blog posts, email, and personal contacts. Emails were sent to relevant entities and individuals, including medical student interest groups at multiple institutions (such as interest groups in internal medicine, infectious diseases, and global public health), researchers, and medical specialists across multiple countries. We promoted the contest using Figure 1[®] and Ding Xiang Yuan (DXY). Both are digital platforms that bring together medical students, healthcare providers, and other healthcare-oriented professionals to share, distribute, and comment on medical cases.¹⁸ DXY is China's largest online healthcare community, with more than four million registered users. On DXY's main platform website, we paid for four banner advertisements and one push notification to registered users, which reached 16,323 individuals and had 99 unique opens. We also created an official Figure 1[®] profile to facilitate contest promotion and developed a promotional infographic that was distributed via the Figure 1[®] app and made using Canva, an online graphic-design tool

In order to encourage participation on Figure 1[®], we posted two MCQs on their platform. Each question focused on correct antibiotic treatment options based on presenting symptoms. As of November 9, 2019, the first promotional MCQ received 31 user-comments and 13,762 views and the second MCQ received 54 user-comments and 17,429 views. Promotion was also conducted through the official Figure 1[®] app email account.

Finally, we promoted the contest through one in-person event. Contest flyers were distributed at an AMR-focused conference in Belfast, United Kingdom where over 150 healthcare professionals working in the field of antimicrobial resistance and infection prevention and control convened.

We analysed metrics from both Figure 1[®] and the contest website. Figure 1[®] metrics showed that there were 85 comments, 107 saves, and 31,191 views from both promotional MCQs. Email analytics from the Figure 1[®] official email account suggest that details regarding the contest were disseminated to a high number of individuals. This first email had 764 opens, 491 unique opens, and 50 clicks. A digest email that consisted of both AMR contest and Figure 1[®] app content and was then sent to primary care physicians, nurses, and medical students. This email had 13,485 email opens, 8,979 unique clicks, and 74 clicks on AMR contest content. A third email was sent on May 22, 2019 specifically asking Figure 1[®] users to contribute AMR related MCQs. This email had 19,298 opens, 12,243 unique opens, and 6781 clicks.

Using online analytical tools, official contest website metrics were obtained between April 30 and June 9, for a total of 41 days. During this period, the website received 578 total clicks, for an average of 14.1 clicks per day, or 98.7 clicks per week. By reviewing IP addresses, it was determined that website clicks originated from 36 different countries.

Receiving and evaluating contributions

Participants were given the option to submit their entries through a digital form made using Qualtrics[®] Survey Software (Qualtrics, Provo, UT) or upload their entry through either the Figure 1[®]

app or DXY, provided that they previously had a platform account. Individuals could submit multiple entries. In addition to entries, we collected the following sociodemographic details regarding participants: name, institution, and country. We asked participants to specify a target audience for their entry (general and primary care physicians, medical students, pharmacists, infectious disease specialists, nurses, etc.) and identify any AMR learning objective(s) that were addressed, whether it be in regards to background, prevention, diagnosis, or treatment of AMR (**Table 2**). Consent was sought from participants regarding the modification and use of solicited entries for learning purposes, along with records of proper citation of external sources, and documentation of patient confidentiality.

Eligibility was determined based on pre-specified criteria: that the entry focused on AMR, was in English, and was in the correct format (as either a MCQ accompanied by answers, an infographic accompanied by brief explanatory captions, or an image). After determining eligibility, entries were transferred to evaluators for phase 1 judging. Evaluation was conducted by three clinical experts who were identified by the steering committee. Their participation as a judge was voluntary.

During phase 1, each judge awarded every entry an individual score between 1 and 10 based on predetermined criteria. Predetermined evaluation criteria included adherence to the required format, contribution to existing learning resources, relevance and effectiveness in enhancing awareness and understanding of AMR, and focus on one or more of four AMR learning objectives (**Table 2**). The three individual scores were then averaged to determine one single score. Comments and revisions to further improve and develop finalist entries were sent to participants.

Recognising finalists

The eight finalists with mean scores of 6.0 or greater were awarded a total of 1000 USD in prize money through Amazon gift cards and cash. Entries were awarded differing cash prize amounts based on the strength of the submission. All submitters were awarded a commendation certificate in recognition of their participation. The judges also received a thank you letter.

Sharing solutions

After further review by expert judges, entries were shared with the public. The MCQs were arranged into a slide deck and included as an additional study material to an AMR learning module that was developed through a similar challenge contest in 2018.¹⁸ Nine MCQs were selected by Figure 1[®] to be shared on their platform via posts, reaching a total of 68 comments, 126 saves, and 81,928 views. Three finalist infographics were published as posters on F1000Research, a life sciences-focused digital publishing platform.

We asked participants to identify a target audience for their entry. Target audiences identified by submitters included medical students, general practitioners and physicians, internal medicine specialists, infectious disease specialists, microbiologists, prescribers in low- and middle-income countries, junior doctors, and nurses. Submitters also identified veterinary practitioners and farmers as a target audience due to the increasing prevalence of antibiotics in agriculture practices and raising livestock.

Table 2. AMR Learning Objectives.

Overall objectives	Individual objectives
Background Information	1. Interpret local epidemiologic data or antibiograms to determine local rates of AMR infections 2. List key risk factors for drug-resistant infections
Prevention	3. Describe factors that may lead to unnecessary antibiotic prescribing by healthcare providers 4. Describe the types of precautions needed/infection control measures for AMR organisms
Diagnosis	5. Interpret susceptibility of testing results to select the most appropriate antibiotic regimen 6. Utilise the local (and regional, if available) microbiology lab to help interpret patient test results
Treatment	7. Identify infections that do not require antibiotic therapy 8. Recognise that treatment of infections may require both antibiotic therapy and source control 9. Recognise the concept of using the narrowest spectrum antibiotic for the shortest period of time 10. Utilise a multidisciplinary healthcare approach when managing AMR organism 11. List resources that can be useful in the treatment of patients with AMR infections 12. Describe the incidence and spectrum of adverse antibiotic effects

Figures

Image not available with this version

Figure 1

Figure 1

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AMRPaperSupplementaryMaterial1.docx](#)