

BMP1 is Not Required for Lung Fibrosis in Mice

Hsiao-Yen Ma

Roche (United States)

Elsa-Noah N'Diaye

Roche (United States)

Patrick Caplazi

Roche (United States)

Zhiyu Huang

Roche (United States)

Alexander Arlantino

Roche (United States)

Surinder Jeet

Roche (United States)

Aaron Wong

Roche (United States)

Hans Brightbill

Roche (United States)

Qingling Li

Roche (United States)

Weng Ruh Wong

Roche (United States)

Wendy Sandoval

Roche (United States)

Lucinda Tam

Roche (United States)

Robert Newman

Roche (United States)

Merone Roose-Girma

Roche (United States)

Ning Ding (✉ ding.ning@gene.com)

Roche (United States)

Research Article

Keywords: BMP1 , lung fibrosis , mice

Posted Date: November 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1072394/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on March 31st, 2022. See the published version at <https://doi.org/10.1038/s41598-022-09557-3>.

Abstract

Bone morphogenetic protein 1 (BMP1) belongs to the astacin/BMP1/tolloid-like family of zinc metalloproteinases, which play a fundamental role in the development and formation of extracellular matrix (ECM). BMP1 mediates the cleavage of carboxyl terminal (C-term) propeptides from procollagens, a crucial step in fibrillar collagen fiber formation. Blocking BMP1 by small molecule or antibody inhibitors has been linked to anti-fibrotic activity in the preclinical models of skin, kidney and liver fibrosis. Therefore, we reason that BMP1 may be important for the pathogenesis of lung fibrosis and BMP1 could be a potential therapeutic target for progressive fibrotic disease such as idiopathic pulmonary fibrosis (IPF). Here, we observed the increased expression of BMP1 in both human IPF lungs and mouse fibrotic lungs induced by bleomycin. Furthermore, we developed an inducible *Bmp1* conditional knockout (cKO) mouse strain. We found that *Bmp1* deletion does not protect mice from lung fibrosis triggered by bleomycin. Moreover, we found no significant impact of BMP1 deficiency upon C-term propeptide of type I procollagen (C1CP) production in the fibrotic mouse lungs. Based on these results, we propose that BMP1 is not required for lung fibrosis in mice and BMP1 may not be considered a candidate therapeutic target for IPF.

Introduction

Characterized by excessive deposition of excessive collagen matrix proteins, idiopathic pulmonary fibrosis (IPF) is a progressive fibrotic disease that is associated with high morbidity and mortality¹. IPF is the pathological result of repetitive microinjuries to lung airway and alveolar epithelium, which provoke the accumulation of myofibroblasts and the deposition of collagen fibrils in the interstitial regions^{2,3}. As the common and final step for IPF as well as other fibrotic diseases, excessive collagen fibril formation and deposition lead to loss of lung function and, ultimately, death^{4,5}. Hence, the mediators of this critical biological process may represent appealing therapeutic targets for IPF^{4,5}.

Collagen fibril formation is a complex process involving multiple posttranslational modifications^{6,7}. One of the central steps in this process is the proteolytic processing of procollagens^{6,7}. Once synthesized within cells, monomeric collagen chains are packaged and secreted into triple-helix procollagen molecules with amino (N-term) and carboxyl terminal (C-term) propeptides, which need to be cleaved to enable collagen maturation and the subsequent formation of supramolecular collagen fibrils in the extracellular matrix (ECM). At this step, C-term propeptide removal seems to be more important than N-term processing in their ability to trigger collagen fibrillogenesis⁸⁻¹¹.

Multiple proteinases have been discovered to exhibit enzymatic activity to cleave C-term propeptides from procollagens within the past few decades¹²⁻¹⁶. Among them, the bone morphogenetic protein 1 (BMP1) appears to be the main enzyme responsible for the fibrillar procollagen maturation, particularly type I procollagen^{14,15,17,18}. Human genetics studies identified several loss-of-function *BMP1* mutations associated with Osteogenesis Imperfecta (OI), an inheritable brittle bone disease largely due to

autosomal dominant mutations in *COL1* genes^{19–28}, supporting a critical role of BMP1 in type I procollagen maturation and processing during human bone development. It has also been shown that inhibition of BMP1 by small molecules or antibody reduces skin^{29,30}, kidney^{31,32} and liver fibrosis³³ in the animal models. In line with these reports, we have recently shown that genetic deletion or pharmacological inhibition of BMP1 abrogates type I C-terminal procollagen propeptide (CICP) production and collagen deposition from primary lung fibroblasts³⁴.

Considering the important role of BMP1 in procollagen processing, maturation and deposition, we hypothesized that BMP1 may be a critical determinant of lung fibrosis and could be a potential therapeutic target for progressive fibrotic diseases such as IPF. Upon analyses of fibrotic lung samples from human and mouse, we found BMP1 expression is increased in fibrotic lesions. Next, we developed a mouse model where efficient deletion of both *Bmp1* alleles can induced by tamoxifen treatment. However, when these *Bmp1* conditional knockout (cKO) mice were tested in the bleomycin model of lung fibrosis, we observed that BMP1 deficiency does not prevent lung fibrosis in mice. Further examination of CICP production also revealed that BMP1 is not necessary for type I procollagen C-term processing and maturation during lung fibrosis. Taken together, we provided the first genetic evidence that BMP1 is not required for the development of lung fibrosis in mice, thus arguing against the drug target candidacy of BMP1 for IPF.

Results

BMP1 expression is increased in IPF lungs

We first assessed the protein level of BMP1 by immunohistochemical staining in lung samples collected from normal control and IPF patients. We only overserved a minimal signal of BMP1 protein in normal lung. On the contrary, the immunoreactivity of BMP1 was strongly detected in IPF lungs. Specifically, BMP1 protein appears to be enriched in the fibroblastic foci (FF), a histological feature of IPF, and the epithelium lining (Fig. 1).

BMP1 expression is increased in the bleomycin model of lung fibrosis

To complement our observation in human samples, we sought to determine whether BMP1 expression is increased in the mouse model of lung fibrosis. Intratracheal (IT) instillation of bleomycin model has been widely used as an animal model to induce experimental lung fibrosis. Mechanistically, bleomycin damages lung epithelium, which results in an immune response and fibroblasts activation, differentiation and ECM deposition³⁵. Through analysis of a previously published transcriptomics data in the bleomycin model³⁶, we found that *Bmp1* expression was trending slightly lower, though not significant, in the bleomycin-challenged moue lungs at early time points (day 2 and 7) but was significantly increased at the advanced fibrotic stage of the model (day 14-28) (Fig. 2A). We also examined the expression of other

procollagen C-term proteinases including tolloid-like 1 (*Tll1*), tolloid-like 2 (*Tll2*), meprin A (*Mep1a*) and meprin β (*Mep1b*) in the same dataset. The expression of these genes was not significantly increased in the bleomycin injured mouse lungs throughout the experiment except that *Mep1a* expression was significantly increased only at day 14 (Fig. 2A). *Mep1b* expression was below the detection limit (data not shown). Consistent to the results from transcriptomic analyses, we found that BMP1 protein level was dramatically elevated in mouse lungs 24 days after the challenge of the bleomycin (Fig. 2B). These results thus suggest that the bleomycin model may be suitable to investigate the *in vivo* role of BMP1 in lung fibrosis.

Inducible and efficient deletion of *Bmp1* alleles in *Bmp1* cKO mice

Bmp1 constitutional knockout mice die soon after birth from the failure of ventral body wall closure due to abnormal collagen fibrillogenesis³¹. To bypass this postnatal lethality, we generated mice homozygous for the floxed (cKO) or wild-type (WT) *Bmp1* alleles and hemizygous for a Cre-ERT2 fusion transgene driven by the human Rosa26 promoter. Thus, both *Bmp1* alleles can be deleted when these cKO mice are treated with tamoxifen. In order to confirm the efficient deletion of *Bmp1* alleles, we administered tamoxifen to *Bmp1* cKO and WT mice through intraperitoneal (IP) injection a week prior to IT injection of bleomycin or saline (Fig. 3A). Lung tissues were then harvested for gene expression analyses on day 24 after the initial bleomycin challenge (Fig. 3A). The data showed that tamoxifen induced more than 90% loss of *Bmp1* transcript in lungs in the presence or absence of bleomycin when compared to WT controls (Fig. 3B). In addition, we observed that *Tll1*, *Tll2* and *Mep1a* expression was not significantly affected by *Bmp1* deletion, suggesting that the loss of BMP1 is unlikely compensated by the increased expression of other C-proteinases as examined.

Bmp1 cKO mice are not protected from bleomycin-induced lung fibrosis

Next, we investigated whether inducible deletion of *Bmp1* protects mice from the development of lung fibrosis in the bleomycin model. To this end, we first performed the histological analysis to assess the severity of fibrotic lesions in the lungs of *Bmp1* cKO and WT mice. The data indicate there is no difference of the severity of fibrotic lesions in the lungs between *Bmp1* WT and cKO mice (Fig. 4A and B). Furthermore, using hydroxyproline assay, we quantified the overall collagen deposition in lungs and found no significant impact of *Bmp1* deletion on total hydroxyproline level in lungs (Fig. 4C and D). We also measured the newly synthesized hydroxyproline and found that *Bmp1* deletion does not affect this readout either, suggesting that BMP1 is unlikely to contribute to active collagen synthesis and production. Consistent to the histology and hydroxyproline data, we observed that the representative pro-fibrotic gene (*Col1a1*, *Col3a1*, *Serpine1* and *Timp1*) expression was not significantly impacted by *Bmp1* deletion

(Fig. 5). Collectively, these results suggest that BMP1 does not contribute to the pathogenesis of lung fibrosis in the bleomycin model.

Bmp1 deletion does not reduce C1CP production during bleomycin-induced lung fibrosis

The lack of anti-fibrotic effects in *Bmp1* cKO mice prompted us to explore whether BMP1 modulates procollagen, particularly type I procollagen, C-term processing during the development of lung fibrosis in the bleomycin model. We performed western blot (WB) in lung extracts from a satellite group of *Bmp1* WT and cKO mice treated with bleomycin or saline to assess the level of C1CP production. While the BMP1 protein level appears to be efficiently depleted in the lungs of cKO mice and the C1CP level was substantially increased in bleomycin injured lungs, we found no significant reduction of C1CP level upon BMP1 depletion (Fig. 6A and B). Therefore, these results suggest that BMP1 is not necessary for type I procollagen C-term processing and maturation during lung fibrosis in mice, which may explain the redundant role of BMP1 in lung fibrosis.

Discussion

As IPF has the worst prognosis among all fibrotic diseases and most IPF patients survive less than 3 years from diagnosis, this fatal and progressive fibrotic lung disease placed a huge burden on the global public health system¹⁻³. However, very few experimental therapies exhibited clinical benefits in human patients and the current standard of care medications, pirfenidone and nintedanib, only slow down the decline of lung function without significantly improving patient survival^{37,38}. New treatment options are therefore urgently needed to prevent or even reverse the progression of IPF toward disease mortality^{4,5}.

Since its discovery as the first C-term proteinase in early 90s^{14,15}, BMP1 has long been proposed to be an appealing therapeutic target for progressive fibrotic diseases³⁹. Within the past few decades, several potent hydroxamic acid-containing small molecule BMP1 inhibitors have been reported, but no advancement to clinical trial has been documented^{29,40-42} possibly due to poor selectivity and/or poor metabolic stability⁴³. More recently, BMP1 antibody inhibitors have been reported^{31,34} and one of them exhibited anti-fibrotic efficacy in preclinical models of kidney and liver fibrosis^{31,33}. However, to the best of our knowledge, there are no genetics studies that have ever validated the role of BMP1 in the pathogenesis of tissue fibrosis.

In this end, we developed a *Bmp1* cKO mouse model where both *Bmp1* alleles can be deleted in the adulthood of the animals. This unique mouse model allows us to fully employ a genetic approach to determine the contribution of BMP1 to lung fibrosis in animals. To our surprise, we found that BMP1 is clearly not required for lung fibrosis in the bleomycin model and BMP1 does not contribute to type I procollagen C-term processing. These findings not only suggest that BMP1 may not be a suitable drug target for progressive lung fibrotic diseases but also represent the first instance that BMP1 is not the

major C-term proteinase for procollagen. These results are somewhat contradictory, but not mutually exclusive, with previous reports of BMP1's dominant role in collagen deposition or procollagen C-term processing using BMP1 inhibitors and in vitro cell culture system^{17,18, 29–31,33,34}. Herein, we suggest that there could be three explanations to reconcile these discrepancies.

First of all, a few non-BMP1 proteinases are also able to cleave C-term propeptide of procollagen. For instance, due to the similar domain structures, TLL1 and TLL2 were identified as BMP1-like proteinases and all these BMP1/tolloid-like proteinases (BTPs) can process C-term procollagen efficiently^{12,13}. MEP1A and MEP1B have also been identified as procollagen C-term proteinases¹⁶. Hence, the functional redundancy between these C-proteinases may well compensate the depletion of BMP1 *in vivo*. In support of this notion, genetic deletion of *Bmp1* or *Bmp1/Tll1* in mouse embryonic fibroblasts (MEFs) suggested that both BMP1 and TLL1 contribute to C-term procollagen processing¹⁷. On the other hand, as tissue fibrosis ensues from a pathological microenvironment, multiple cell types can be involved in this process. Thus, it is possible that proteinase activities from non-fibroblastic cell types may contribute to C-term processing of procollagens as well. Therefore, future studies using genetic models or unbiased biochemical analyses are needed to identify which proteinases are truly responsible for procollagen C-term processing and maturation during lung fibrosis *in vivo*.

In addition to the functional redundancy, we cannot ignore the possibility that BMP1 may play a tissue specific role in the pathogenesis of fibrosis. It is noteworthy that, while previous studies tested BMP1 inhibitors in the animal models of skin^{29,30}, kidney^{31,32} and liver fibrosis³³, the anti-fibrotic efficacy of BMP1 inhibitors in lung fibrosis models has not been documented. Thus, it is plausible that BMP1 may play a much more important role in skin, kidney and liver fibrosis than in lung fibrosis. This possibility can be further supported by human genetics data that the collagen abnormality caused by BMP1 loss-of-function mutations appears to be largely restricted to bone tissue^{19–28}. Nonetheless, additional studies on *Bmp1* cKO mice in the fibrosis models of other tissues are mandatory to fully test this hypothesis.

Thirdly, as the selectivity for both small molecule and antibody BMP1 inhibitors has not been fully characterized, it may be reasonable to speculate that these molecules may hit targets beyond BMP1 *in vivo*, if the anti-fibrotic effects of these BMP1 inhibitors can be independently reproduced. Given the high sequence and structure similarity between BTP family members⁴⁴, it is highly likely that BMP1 inhibitors may target the entire BTP family to achieve their anti-fibrotic effects. However, as the CICP level was not determined in the previous pharmacology studies in animal models, we cannot exclude another possibility that the anti-fibrotic effects of these inhibitors may, at least partially, come from non-BTP targets, which may be functionally relevant to tissue fibrosis.

Overall, our study provided the first genetic evidence that BMP1 is not required for lung fibrosis and BMP1 is not the major proteinase to cleave C-term of type I procollagen in the bleomycin model. Based on these findings, we thus propose that BMP1 should not be considered as a candidate therapeutic target for fibrotic lung diseases like IPF. More importantly, our study exemplifies the necessity of leveraging genetic

models to validate drug targets when possible and suggests that the role of BMP1 in the pathological tissue fibrosis may need to be re-evaluated using the similar genetic approaches.

Materials And Methods

Human samples.

Explanted lung tissues were obtained from patients with a pathological diagnosis of usual interstitial pneumonia and a consensus clinical diagnosis of IPF assigned by multidisciplinary discussion and review of clinical materials. Written informed consent was obtained from all subjects and the study was approved by the University of California, San Francisco (UCSF) institutional review board. Human lungs not used by the Northern California Transplant Donor Network were used as controls; studies indicate that these lungs are physiologically and pathologically normal⁴⁵. All procedures were carried out according to the Helsinki Declaration and its later amendments.

Immunohistochemistry staining.

4µm sections of formalin-fixed and paraffin-embedded specimens were deparaffinated followed by antigen retrieval using Target Retrieval (Dako #S1700) and quenching endogenous peroxidase activity. The sections were subsequently blocked with Vector Avidin Biotin Blocking Kit, stained in PBS plus 10% rabbit serum and 3% BSA with anti-BMP1 (0.5 µg/ml, R&D, #AF1297) for lung sections and secondary biotinylated antibodies, incubated with Vectastain ABC Elite reagent and Pierce Metal Enhanced DAB solution and counterstained with Mayer's hematoxylin. Human sections were imaged with a 20x Plan Apo DIC M objective (NA: 0.75, Nikon) on a Nikon Ti-Eclipse inverted microscope equipped with an Andor Neo scMOS camera (Andor, Oxford Instruments), a linear-encoded automated stage (Applied Scientific Instrumentation), and a SOLA LED light engine (Lumencor) all run by NIS Elements software (Nikon).

Induced excision of Bmp1 in adult mice

All animal studies were conducted in accordance with the Guide for the Care and Use of Laboratory Animals, published by the National Institutes of Health (NIH) (NIH Publication 8523, revised 1985). The Institutional Animal Care and Use Committee (IACUC) at Genentech reviewed and approved all animal protocols. To generate *Bmp1* conditional KO mice, *Bmp1*-flox/flox mice⁴⁶ were bred to the *Rosa26*-CreERT2 mouse strain⁴⁷ to produce *Bmp1*-flox/flox;*Rosa26*-Cre-ERT2 mice. *Bmp1*-wt/wt;*Rosa26*-Cre-ERT2 mice were used as controls in this study. Mice from both groups were injected intraperitoneally daily for 5 consecutive days with tamoxifen (Cat# T5648; Sigma) in sunflower oil (S5007-1L; Sigma) at 80 mg/kg. Animals then rest for 2 days and received second dose of tamoxifen again for five consecutive days. Subsequent studies were initiated at least one week after the completion of tamoxifen treatment.

Bleomycin model of lung fibrosis

The study was carried out in compliance with the ARRIVE guidelines. Adult mice (>12 weeks) were randomized based on pre-study weights to minimize variance between experimental and control groups. For intratracheal (i.t.) dosing, all mice were lightly anesthetized with isoflurane in an induction chamber. Once anesthetized, the animals were removed from the chamber, manually restrained, the mouth of the animal was opened and the tongue set aside. A 1 ml syringe with 50 microliters of sterile injectable isotonic saline or bleomycin (0.72U/kg (DNC# 0703-3155-01; TEVA) in 50 µl sterile isotonic saline) was connected to a 24 gauge gavage needle. The gavage needle was inserted into the trachea and a dose of either vehicle or bleomycin was delivered intratracheally. After delivery, animals were monitored continuously until fully awake and ambulatory. Mouse whole lungs were harvested at day 24 after bleomycin administration, n=5 in the saline control group, n=30 in the bleomycin group in *Bmp1*-cKO and WT control littermates respectively.

Histology analysis.

Formalin-fixed samples of mouse lungs were embedded as a whole and processed to 1 slide per animal, stained with Masson's Trichrome. The extent of pulmonary fibrosis was scored according to the following criteria:

- (A) Interstitial fibrosis pattern – number of foci: 0, none detected; 1, ≤ 10 ; 2, ≤ 15 ; 3, > 15 / all sections, but distinct; 4, multifocally coalescent or locally extensive; 5, diffuse;
- (B) Interstitial fibrosis – size of foci: 0, none detected; 1, largest focus \leq area of ~ 2 alveolar spaces; 2, largest focus \leq area of ~ 4 alveolar spaces; 3, coalescent (> 4 patent alveolar spaces); 4, locally extensive (60-90% of an entire lobe); 5, diffuse ($> 90\%$ of an entire lobe);
- (C) Total scores: number of foci x size of foci.

Hydroxyproline and measurement.

Total hydroxyproline was analyzed as previously described⁴⁸ and mass spectrometry and analysis were performed by Metabolic Solutions. Deuterated water labeling was used to assess new collagen synthesis. In brief, mice were injected with deuterated water (DLM-4-99.8-1000; Cambridge Isotope Laboratories) intraperitoneally two weeks prior to the end of the study at 35 mL/kg in 2 divided doses 4 hours apart. Afterward, 8% deuterated water in drinking water was provided ad lib in water bottle until the end of the study. Mass spectrometry and analysis were performed by Metabolic Solutions.

RT-qPCR.

Total RNA was purified using RNeasy kit (Qiagen) and treated with DNaseI (LifeTechnologies). Complementary DNA synthesis was carried out with iScript RT Supermix (Bio-Rad). Quantitative PCR was performed in technical triplicates using SYBR Green reagent (Bio-Rad). The relative standard curve

method was used for quantitation and expression levels were calculated by normalization to *Hrpt*. The sequences of primers are listed in Extended Table 1.

Western blot.

Western blot was carried out in total protein extracts as previously described⁴⁹. Equal amounts of protein lysates were separated by SDS–PAGE, transferred to a nitrocellulose membrane, and subjected to immunoblotting analysis using the following primary antibodies: BMP1 and C1CP antibodies were reported previously³⁴, GAPDH (1:1000, CST, #5174), β -actin (1:3000, Sigma, a5441). The raw image data are provided in Supplemental Figures 1–2.

RNA-seq analysis and plotting

Expression was measured in normalized reads per kilobase per million total reads (nRPKM). RNAseq data on mouse lungs from bleomycin model was published in the NCBI GEO database under accession GSE168529³⁶.

Quantification and Statistical Analysis

GraphPad Prism was utilized for statistical analysis on Figures 3-6. Statistical details of experiments can be found in figure legends, including the statistical tests used and value and definition of n. Differences were considered to be statistically significant when $p < 0.05$. Western blot band densitometry was performed using ImageJ software.

Declarations

Acknowledgements

We thank the Genentech microinjection, animal production, genetic analysis, and histology laboratories for technical assistance; the Genentech Center for Advanced Light Microscopy for imaging. We also thank Rajita Pappu for critical review of this work; Linda Rangell and Debra Dunlap for immunohistochemistry support; and Mariela Del Rio and Bridget Hough for animal husbandry.

Author contributions

H.Y.M., E.N-D. designed the study, performed the experiments, analyzed the data, and wrote the manuscript; P.C. contributed to the histological analysis of fibrotic tissues; Z.H., A.A., S.J., A.W. H.B. performed the tamoxifen and bleomycin model experiments; Q.L., W.R.W., W.S. performed the hydroxyproline analyses; L.T., R.N., M.R-G. designed the targeting vector and supervised the generation of the *Bmp1* cKO mouse line; N.D. designed the study, analyzed the data, wrote the manuscript with H.Y.M. and supervised the study.

Competing interests

All authors are or were employees of Genentech.

References

1. Kim, D. S., Collard, H. R. & King, T. E. Jr. Classification and natural history of the idiopathic interstitial pneumonias. *Proc Am Thorac Soc*, **3**, 285–292 <https://doi.org/doi:10.1513/pats.200601-005TK> (2006).
2. Wynn, T. A. Integrating mechanisms of pulmonary fibrosis. *J Exp Med*, **208**, 1339–1350 <https://doi.org/doi:10.1084/jem.20110551> (2011).
3. Coward, W. R., Saini, G. & Jenkins, G. The pathogenesis of idiopathic pulmonary fibrosis. *Thor Adv Respir Dis*, **4**, 367–388 <https://doi.org/doi:10.1177/1753465810379801> (2010).
4. Henderson, N. C., Rieder, F. & Wynn, T. A. Fibrosis: from mechanisms to medicines., **587**, 555–566 <https://doi.org/doi:10.1038/s41586-020-2938-9> (2020).
5. Friedman, S. L., Sheppard, D., Duffield, J. S. & Violette, S. Therapy for fibrotic diseases: nearing the starting line. *Sci Transl Med*, **5**, 167161 <https://doi.org/doi:10.1126/scitranslmed.3004700> (2013).
6. Kadler, K. E. Fell Muir Lecture: Collagen fibril formation in vitro and in vivo. *Int J Exp Pathol*, **98**, 4–16 <https://doi.org/doi:10.1111/iep.12224> (2017).
7. Mouw, J. K., Ou, G. & Weaver, V. M. Extracellular matrix assembly: a multiscale deconstruction. *Nat Rev Mol Cell Biol*, **15**, 771–785 <https://doi.org/doi:10.1038/nrm3902> (2014).
8. Kadler, K. E., Hojima, Y. & Prockop, D. J. Assembly of collagen fibrils de novo by cleavage of the type I pC-collagen with procollagen C-proteinase. Assay of critical concentration demonstrates that collagen self-assembly is a classical example of an entropy-driven process. *J Biol Chem*, **262**, 15696–15701 (1987).
9. Canty, E. G. & Kadler, K. E. Procollagen trafficking, processing and fibrillogenesis. *J Cell Sci*, **118**, 1341–1353 <https://doi.org/doi:10.1242/jcs.01731> (2005).
10. Hulmes, D. J. Building collagen molecules, fibrils, and suprafibrillar structures. *J Struct Biol*, **137**, 2–10 <https://doi.org/doi:10.1006/jsbi.2002.4450> (2002).
11. Kadler, K. E., Holmes, D. F., Trotter, J. A. & Chapman, J. A. Collagen fibril formation. *Biochem J*, **316** (Pt 1), 1–11 <https://doi.org/doi:10.1042/bj3160001> (1996).
12. Scott, I. C. *et al.* Mammalian BMP-1/Tolloid-related metalloproteinases, including novel family member mammalian Tolloid-like 2, have differential enzymatic activities and distributions of expression relevant to patterning and skeletogenesis. *Dev Biol*, **213**, 283–300 <https://doi.org/doi:10.1006/dbio.1999.9383> (1999).
13. Takahara, K., Brevard, R., Hoffman, G. G., Suzuki, N. & Greenspan, D. S. Characterization of a novel gene product (mammalian tolloid-like) with high sequence similarity to mammalian tolloid/bone morphogenetic protein-1., **34**, 157–165 <https://doi.org/doi:10.1006/geno.1996.0260> (1996).

14. Li, S. W. *et al.* The C-proteinase that processes procollagens to fibrillar collagens is identical to the protein previously identified as bone morphogenic protein-1. *Proc Natl Acad Sci U S A*, **93**, 5127–5130 <https://doi.org/doi:10.1073/pnas.93.10.5127> (1996).
15. Kessler, E., Takahara, K., Biniaminov, L., Brusel, M. & Greenspan, D. S. Bone morphogenetic protein-1: the type I procollagen C-proteinase., **271**, 360–362 <https://doi.org/doi:10.1126/science.271.5247.360> (1996).
16. Broder, C. *et al.* Metalloproteases meprin alpha and meprin beta are C- and N-procollagen proteinases important for collagen assembly and tensile strength. *Proc Natl Acad Sci U S A*, **110**, 14219–14224 <https://doi.org/doi:10.1073/pnas.1305464110> (2013).
17. Pappano, W. N., Steiglit, B. M., Scott, I. C., Keene, D. R. & Greenspan, D. S. Use of Bmp1/Tll1 doubly homozygous null mice and proteomics to identify and validate in vivo substrates of bone morphogenetic protein 1/tolloid-like metalloproteinases. *Mol Cell Biol*, **23**, 4428–4438 <https://doi.org/doi:10.1128/mcb.23.13.4428-4438.2003> (2003).
18. Suzuki, N. *et al.* Failure of ventral body wall closure in mouse embryos lacking a procollagen C-proteinase encoded by Bmp1, a mammalian gene related to Drosophila tolloid. *Development*, **122**, 3587–3595 (1996).
19. Cho, S. Y. *et al.* Identification and in vivo functional characterization of novel compound heterozygous BMP1 variants in osteogenesis imperfecta. *Hum Mutat*, **36**, 191–195 <https://doi.org/doi:10.1002/humu.22731> (2015).
20. Hoyer-Kuhn, H. *et al.* Hyperosteoidosis and hypermineralization in the same bone: bone tissue analyses in a boy with a homozygous BMP1 mutation. *Calcif Tissue Int*, **93**, 565–570 <https://doi.org/doi:10.1007/s00223-013-9799-2> (2013).
21. Martinez-Glez, V. *et al.* Identification of a mutation causing deficient BMP1/mTLD proteolytic activity in autosomal recessive osteogenesis imperfecta. *Hum Mutat*, **33**, 343–350 <https://doi.org/doi:10.1002/humu.21647> (2012).
22. Pollitt, R. C. *et al.* Phenotypic variability in patients with osteogenesis imperfecta caused by BMP1 mutations. *Am J Med Genet A*, **170**, 3150–3156 <https://doi.org/doi:10.1002/ajmg.a.37958> (2016).
23. Sangsin, A. *et al.* Two novel compound heterozygous BMP1 mutations in a patient with osteogenesis imperfecta: a case report. *BMC Med Genet*, **18**, 25 <https://doi.org/doi:10.1186/s12881-017-0384-9> (2017).
24. Syx, D. *et al.* Defective Proteolytic Processing of Fibrillar Procollagens and Prodecorin Due to Biallelic BMP1 Mutations Results in a Severe, Progressive Form of Osteogenesis Imperfecta. *J Bone Miner Res*, **30**, 1445–1456 <https://doi.org/doi:10.1002/jbmr.2473> (2015).
25. Valencia, M. *et al.* Report of a newly identified patient with mutations in BMP1 and underlying pathogenetic aspects. *Am J Med Genet A*, **164A**, 1143–1150 <https://doi.org/doi:10.1002/ajmg.a.36427> (2014).
26. Xu, X. J. *et al.* Novel mutations in BMP1 induce a rare type of osteogenesis imperfecta. *Clin Chim Acta*, **489**, 21–28 <https://doi.org/doi:10.1016/j.cca.2018.11.004> (2019).

27. Asharani, P. V. *et al.* Attenuated BMP1 function compromises osteogenesis, leading to bone fragility in humans and zebrafish. *Am J Hum Genet*, **90**, 661–674 <https://doi.org/doi:10.1016/j.ajhg.2012.02.026> (2012).
28. Lindahl, K. *et al.* COL1 C-propeptide cleavage site mutations cause high bone mass osteogenesis imperfecta. *Hum Mutat*, **32**, 598–609 <https://doi.org/doi:10.1002/humu.21475> (2011).
29. Fish, P. V. *et al.* Potent and selective nonpeptidic inhibitors of procollagen C-proteinase. *J Med Chem*, **50**, 3442–3456 <https://doi.org/doi:10.1021/jm061010z> (2007).
30. Reid, R. R., Mogford, J. E., Butt, R., deGiorgio-Miller, A. & Mustoe, T. A. Inhibition of procollagen C-proteinase reduces scar hypertrophy in a rabbit model of cutaneous scarring. *Wound Repair Regen*, **14**, 138–141 <https://doi.org/doi:10.1111/j.1743-6109.2006.00103.x> (2006).
31. Grgurevic, L. *et al.* Circulating bone morphogenetic protein 1-3 isoform increases renal fibrosis. *J Am Soc Nephrol*, **22**, 681–692 <https://doi.org/doi:10.1681/ASN.2010070722> (2011).
32. Bai, M. *et al.* BMP1 inhibitor UK383,367 attenuates renal fibrosis and inflammation in CKD. *Am J Physiol Renal Physiol*, **317**, F1430–F1438 <https://doi.org/doi:10.1152/ajprenal.00230.2019> (2019).
33. Grgurevic, L. *et al.* Systemic inhibition of BMP1-3 decreases progression of CCl₄-induced liver fibrosis in rats., **35**, 201–215 <https://doi.org/doi:10.1080/08977194.2018.1428966> (2017).
34. N'Diaye, E. N. *et al.* Extracellular BMP1 is the major proteinase for COOH-terminal proteolysis of type I procollagen in lung fibroblasts. *Am J Physiol Cell Physiol*, **320**, C162–C174 <https://doi.org/doi:10.1152/ajpcell.00012.2020> (2021).
35. Tashiro, J. *et al.* Exploring Animal Models That Resemble Idiopathic Pulmonary Fibrosis. *Front Med (Lausanne)*, **4**, 118 <https://doi.org/doi:10.3389/fmed.2017.00118> (2017).
36. Sun, T. *et al.* TGFbeta2 and TGFbeta3 isoforms drive fibrotic disease pathogenesis. *Sci Transl Med*, **13**, <https://doi.org/doi:10.1126/scitranslmed.abe0407> (2021).
37. King, T. E. Jr. *et al.* A phase 3 trial of pirfenidone in patients with idiopathic pulmonary fibrosis. *N Engl J Med*, **370**, 2083–2092 <https://doi.org/doi:10.1056/NEJMoa1402582> (2014).
38. Richeldi, L. *et al.* Efficacy and safety of nintedanib in idiopathic pulmonary fibrosis. *N Engl J Med*, **370**, 2071–2082 <https://doi.org/doi:10.1056/NEJMoa1402584> (2014).
39. Turtle, E. D. & Ho, W. B. Inhibition of procollagen C-proteinase: fibrosis and beyond. *Expert Opinion on Therapeutic Patents*, **14**, 1185–1197 <https://doi.org/doi:10.1517/13543776.14.8.1185> (2005).
40. Kallander, L. S. *et al.* Reverse Hydroxamate Inhibitors of Bone Morphogenetic Protein 1. *ACS Med Chem Lett*, **9**, 736–740 <https://doi.org/doi:10.1021/acsmedchemlett.8b00173> (2018).
41. Dankwardt, S. M. *et al.* Amino acid derived sulfonamide hydroxamates as inhibitors of procollagen C-proteinase. Part 2: Solid-phase optimization of side chains. *Bioorg Med Chem Lett*, **12**, 1233–1235 [https://doi.org/doi:10.1016/s0960-894x\(02\)00117-8](https://doi.org/doi:10.1016/s0960-894x(02)00117-8) (2002).
42. Turtle, E. *et al.* Design and synthesis of procollagen C-proteinase inhibitors. *Bioorg Med Chem Lett*, **22**, 7397–7401 <https://doi.org/doi:10.1016/j.bmcl.2012.10.067> (2012).

43. Vandembroucke, R. E. & Libert, C. Is there new hope for therapeutic matrix metalloproteinase inhibition? *Nat Rev Drug Discov*, **13**, 904–927 <https://doi.org/doi:10.1038/nrd4390> (2014).
44. Mac Sweeney, A. *et al.* Structural basis for the substrate specificity of bone morphogenetic protein 1/tolloid-like metalloproteases. *J Mol Biol*, **384**, 228–239 <https://doi.org/doi:10.1016/j.jmb.2008.09.029> (2008).
45. Ware, L. B. *et al.* Assessment of lungs rejected for transplantation and implications for donor selection., **360**, 619–620 [https://doi.org/doi:10.1016/s0140-6736\(02\)09774-x](https://doi.org/doi:10.1016/s0140-6736(02)09774-x) (2002).
46. Muir, A. M. *et al.* Induced ablation of Bmp1 and Tll1 produces osteogenesis imperfecta in mice. *Hum Mol Genet*, **23**, 3085–3101 <https://doi.org/doi:10.1093/hmg/ddu013> (2014).
47. Seibler, J. *et al.* Rapid generation of inducible mouse mutants. *Nucleic Acids Res* **31**, e12, [doi:10.1093/nar/gng012](https://doi.org/doi:10.1093/nar/gng012) (2003)
48. Decaris, M. L. *et al.* Proteomic analysis of altered extracellular matrix turnover in bleomycin-induced pulmonary fibrosis. *Mol Cell Proteomics*, **13**, 1741–1752 <https://doi.org/doi:10.1074/mcp.M113.037267> (2014).
49. Ding, N. *et al.* Mediator links epigenetic silencing of neuronal gene expression with x-linked mental retardation. *Mol Cell*, **31**, 347–359 <https://doi.org/doi:10.1016/j.molcel.2008.05.023> (2008).

Table

Due to technical limitations, table 1 is only available as a download in the Supplemental Files section.

Figures




Figure 1. BMP1 expression is increased in human IPF lungs
Representative images of BMP1 immunohistochemistry staining in human IPF and normal lung specimens. EP: epithelium. FF: fibroblastic foci.

Figure 1

Please See image above for figure legend.





Figure 2. BMP1 expression is increased in the bleomycin model
 (A) mRNA levels of *Bmp1*, *Tll1*, *Tll2* and *Mep1a* in mouse lungs on indicated days (0, 2, 4, 7, 14, 21 and 28, n=5-7) after bleomycin challenge. * $p < 0.05$, ** $p < 0.01$. Data represents mean \pm S.E.M. p-value is calculated using one-way ANOVA. (B) Representative western blot of BMP1 in lung lysates after 24 days bleomycin challenge compare to saline group. β -actin is a loading control.

Figure 2

Please See image above for figure legend.




Figure 3. Inducible and efficient deletion of *Bmp1* in the bleomycin model

(A) Schematic regime of tamoxifen-induced *Bmp1* deletion following by intratracheal bleomycin challenge. Tissue harvesting on day 24 after bleomycin challenge. (B) mRNA expression of *Bmp1*, *Tll1*, *Tll2* and *Mep1a*. *Bmp1* WT: Saline n=5, bleomycin n=30; *Bmp1* cKO: saline n=5, bleomycin n=21. *** p < 0.001. Data represents mean \pm S.D. p-value is calculated using one-way ANOVA.

Figure 3

Please See image above for figure legend.




Figure 4. *Bmp1* cKO mice are not protective against bleomycin-induced lung fibrosis

Representative images of trichrome histology staining (A) in lung tissues with quantification of fibrosis lesion (B). Quantification of fibrosis by total hydroxyproline (C) and newly synthesized hydroxyproline (D). *Bmp1* WT: Saline n=5, bleomycin n=30; *Bmp1* cKO: saline n=5, bleomycin n=21. Data represents mean ± S.D.

Figure 4

Please See image above for figure legend.




Figure 5. *Bmp1* deletion has no effects on pro-fibrotic gene expression

mRNA expression of selective pro-fibrotic genes from lung tissue. *Bmp1* WT: Saline n=5, bleomycin n=30; *Bmp1* cKO: saline n=5, bleomycin n=21. Data represents mean ± S.D. p-value is calculated using one-way ANOVA.

Figure 5

Please See image above for figure legend.




Figure 6. *Bmp1* deletion does not reduce CICP production during bleomycin-induced lung fibrosis

Equal amounts of total proteins were subject to western blot for the detection of BMP1 and CICP in lung tissues. GAPDH was used as a loading control. (B) Quantification of relative protein expression of BMP1 and CICP in lung tissues by normalizing with GAPDH. Data represents mean \pm S.D. p-value is calculated using one-way ANOVA.

Figure 6

Please See image above for figure legend.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Bmp1supfigures.pdf](#)
- [Tab1.jpg](#)