Reporting Quality of Clinical Practice Guidelines Regarding Gout and Hyperuricemia According to the Right Checklist

Can Wang

The Affiliated Hospital of Qingdao University

Xufei Luo

School of Public Health of Lanzhou University

Maichao Li

The Affiliated Hospital of Qingdao University

Lingling Cui

The Affiliated Hospital of Qingdao University

Xinde Li

The Affiliated Hospital of Qingdao University

Lin Han

The Affiliated Hospital of Qingdao University

Xuefeng Wang

The Affiliated Hospital of Qingdao University

Wei Ren

The Affiliated Hospital of Qingdao University

Yuwei He

The Affiliated Hospital of Qingdao University

Wenyan Sun

The Affiliated Hospital of Qingdao University

Changgui Li

The Affiliated Hospital of Qingdao University

Yaolong Chen (✉ gssrmywl2016@163.com)

Institute of Health Data Science of Lanzhou University

Zhen Liu

The Affiliated Hospital of Qingdao University

Methodology

Keywords: clinical practice guideline, gout, hyperuricemia, RIGHT check list, reporting quality

Posted Date: November 18th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-107411/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Systematic Reviews on April 5th, 2021. See the published version at <https://doi.org/10.1186/s13643-021-01645-1>.

Abstract

Objectives The Reporting Items for Practice Guidelines in Healthcare (RIGHT) checklist was used to assess the reporting quality of 2009–2019 Clinical Practice Guidelines (CPGs) regarding gout and hyperuricemia, aimed to improve the reporting quality of future guidelines.

Methods We searched PubMed, the Chinese Biomedical Literature database, the Wan Fang Database, and the China National Knowledge Infrastructure from January 2009 to June 2019 for relevant guidelines. We also searched the websites of guideline development organizations (the Guidelines International Network, the National Institute for Health and Clinical Excellence, the American College of Rheumatology, and the European League Against Rheumatism) (EULAR). Furthermore, supplementary guidelines reported in included articles were systematically searched, as well as Medlive and Google Scholar.

Results Seventeen guidelines were included, of which one was in Chinese and 16 were in English. The mean reporting rate of the 35 items specified was 14.9 (42.5%), only five CPGs (29.4%) had a reporting rate >50%. Of the 35 items, three were very frequently reported. The reporting proportion of the seven domains (Basic information, Background, Evidence, Recommendations, Review and quality assurance, Funding and declaration and management of interests, and Other information) were 64.7%, 36.8%, 50.6% 50.6%, 42.9%, 8.82%, 33.8%, and 31.4%, respectively.

Conclusion The reporting quality of the present guidelines for gout and hyperuricemia is relatively poor. We suggest that the RIGHT reporting checklist should be used by CPG developers to ensure higher reporting quality of future guidelines.

Background

Gout is a group of heterogeneous diseases caused by long-term disturbance of purine metabolism, which results in a high serum uric acid concentration. The prevalence of gout is >1% in most developed countries, and with the recent improvement in living standards, the prevalence of gout is likely to increase[1]. This chronic disease is associated with substantial morbidity and mortality, making it a major global social and economic burden[2]. Furthermore, gout and hyperuricemia can induce and exacerbate metabolic diseases, such as hypertension, diabetes, and disorders of lipid metabolism, besides they are also independent risk factors for stroke and myocardial infarction[3, 4]. Thus, gout and hyperuricemia have become common conditions that seriously affect human health.

With recent advances in pharmaceutical therapies, auxiliary diagnostic methods, and novel treatment approaches, a number of clinical practice guidelines (CPGs) have been developed to standardize the diagnosis and treatment of gout[5-9]. On the basis of evidence provided by systematic reviews, these CPGs have aimed to provide patients with the optimal medical treatment strategy [10]. Decisions regarding diagnosis and therapy are made on the basis of the CPGs, which standardize the behavior of clinicians, with the aims of improving clinical success and reducing cost [11]; therefore, the quality of CPGs is crucially important. There are two quality evaluation methods suitable for CPGs: the Appraisal of Guidelines for Research and Evaluation tool (AGREE) and the Reporting Items for Practice Guidelines in Healthcare (RIGHT) tool [12]. A number of studies have been published that used AGREEII to assess the quality of the gout guidelines, which showed that the current guidelines are of poor quality [8, 13-15]. However, AGREEII does not separate the quality of the report contents from the quality of the methodology of CPGs, Yao et.al recommended RIGHT checklist providing detailed information that lacked in AGREEII [16], meaning that the reporting quality of the gout and hyperuricemia guidelines has not been fully assessed [12]. Therefore, in the present study, we evaluated the reporting quality of gout and hyperuricemia CPGs using the RIGHT tool, to compensate previous study using AGREEII, and permit the standardization reporting of future guidelines.

Materials And Methods

Data source and search strategy

We searched PubMed, the Chinese Biomedical Literature database (CBM), the Wan Fang Database, and the China National Knowledge Infrastructure (CNKI) from January 2009 to June 2019 for the relevant guidelines. We also searched the websites of the organizations responsible for guideline development: the Guidelines International Network (GIN), the National Institute for Health and Clinical Excellence (NICE), the American College of Rheumatology (ACR), and the European League Against Rheumatism (EULAR). Furthermore, supplementary guidelines reported in the included articles were systematically searched, and we also searched Medlive and Google Scholar for additional material. [8, 9, 13].

Study selection and data extraction

Inclusion criteria

CPGs regarding gout and hyperuricemia, written in English or Chinese.

Exclusion criteria

1) old versions of guidelines, if an updated version was available; 2) interpretations and translations of guidelines; 3) repeatedly published guidelines; and 4) guidelines for which full text access was not available.

Data extraction

Two researchers (Zhen Liu and Can Wang) searched the database, independently screened the guidelines, and determined whether the retrieved guidelines met the study criteria. Then, independent screening and cross-checking of the guidelines were carried out. Differences between reviewers were resolved through discussion or consultation with a third party (Yaolong Chen and his team).

Reporting quality assessment

The RIGHT tool (supplementary material or <http://www.right-statement.org>) was used to evaluate the eligibility guidelines included in this study. The tool consists of 22 key items and 35 sub-items, which are divided into the following seven areas: Basic information (items 1–4), Background (items 5–9), Evidence (items 10–12), Recommendations (items 13–15), Review and quality assurance (items 16–17), Funding and declaration and management of interests (items 18–19), and Other information (items 20–22). Each item was independently scored by two researchers, and most of the items were graded dichotomously, as “reported” (Y) or “unreported” (N). “Reported” meant that the relevant information was fully reported, whereas “unreported” meant the relevant information was unavailable. However, “partially reported” (P) was also used to indicate that the guideline contained only partial information, and when the guideline evaluation was not applicable to the item, “not applicable” (NA) was recorded. We reported the results for each item as absolute quantities and percentages. For each item, we also reported the number and percentage of projects reported by each guideline. If the reporting proportion of the guidelines was < 50%, the quality of the item was regarded as low. The data were extracted and analyzed using Excel 2016 (Microsoft Corp., Redmond, WA, USA).

Results

Results of the selection

A total of 765 records were identified through database searching. Four other records were identified *via* the guideline development organization websites and Google Scholar. Of these, 17 guidelines that met the criteria were analyzed (Fig. 1).

Characteristics of the included CPGs

Of the 17 CPGs, one was in Chinese and 16 were in English. Nine were developed by European countries (52.9%), of which four were developed by EULAR (23.5%); six originated from the United States (35.3%), of which three were developed by ACR (17.6%) and three (17.6%) by universities. The remaining two (11.8%) were developed by the Evidence, Expertise, and Exchange initiative. The features of each are summarized in Table 1.

Table 1
Characteristics of included CPGs

Serial number	Title	Published date	Developer	Nation	Published journal
1	2018 updated European League Against Rheumatism evidence-based recommendations for the diagnosis of gout[18]	2019	European League Against Rheumatism.	Europe	Annals of the Rheumatic Diseases
2	Management of gout and hyperuricemia: Multidisciplinary consensus in Taiwan[19]	2018	Astellas, Taiwan	Taiwan,China	SPECIAL EDITORIAL REVIEW
3	The British Society for Rheumatology Guideline for the Management of Gout[20]	2017	British Society for Rheumatology Standards, Audit and Guidelines Working Group.	UK	RHEUMATOLOGY
4	1.Diagnosis of Acute Gout: A Clinical Practice Guideline From the American College of Physicians[21] 2.Management of Acute and Recurrent Gout: A Clinical Practice Guideline From the American College of Physicians[22]	2017	American College of Physicians	USA	Annals of Internal Medicine
5	[2016 China Gout Clinical Practice Guideline][23]	2016	Chinese Rheumatology Association	China	Zhonghua Nei Ke Za Zhi
6	2016 updated EULAR evidence-based recommendations for the management of gout[24]	2016	European League Against Rheumatism	Europe	Clinical and epidemiological research
7	Treat-to-target (T2T) recommendations for gout[25]	2016	European League Against Rheumatism	Europe	Annals of the Rheumatic Diseases
8	Australian and New Zealand recommendations for the diagnosis and management of gout: integrating systematic literature review and expert opinion in the 3e Initiative[26]	2015	APLAR, Asia Pacific League of Associations for Rheumatology	Australian and New Zealand	International Journal of Rheumatic Diseases
9	2015 Gout classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative[27]	2015	ACR/EULAR	USA and Europe	Annals of the Rheumatic Diseases
10	Portuguese recommendations for the diagnosis and management of Gout[28]	2014	A panel of 78 international rheumatologists in 3e (Evidence, Expertise, Exchange) Initiative	Portugal	PRÁTICA CLÍNICA
11	Clinical Practice Guidelines for Management of Gout[29]	2013	Spanish Society of Rheumatology	Spain	guipclingot
12	Multinational evidence-based recommendations for the diagnosis and management of gout: integrating systematic literature review and expert opinion of a broad panel of rheumatologists in the 3e initiative[30]	2013	3e (Evidence, Expertise, Exchange) Initiative/A panel of international rheumatologists	International	Annals of the Rheumatic Diseases
13	Italian Society of Rheumatology recommendations for the management of gout[31]	2013	Italian Society of Rheumatology	Italy	Reumatismo
14	Management of chronic gout in adults[32]	2012	University of Texas	USA	National Guideline Clearinghouse
15	1.2012 American College of Rheumatology Guidelines for Management of Gout. Part 1: Systematic Nonpharmacologic and Pharmacologic Therapeutic Approaches to Hyperuricemia[33] 2.2012 American College of Rheumatology Guidelines for Management of Gout. Part 2: Therapy and Anti-inflammatory Prophylaxis of Acute Gouty Arthritis[34]	2012	American College of Rheumatology	USA	Arthritis Care & Research

Serial number	Title	Published date	Developer	Nation	Published journal
16	Japanese Guideline for the Management of Hyperuricemia and Gout: Second Edition[35]	2011	Tokyo Women's Medical University	Japan	Nucleosides, Nucleotides and Nucleic Acids
17	Management of initial gout in adults[36]	2009	University of Texas	USA	National Guideline Clearinghouse

Overall reporting quality

The mean reporting rate of the 35 items was 14.9 (42.5%) and it ranged from nine (25.7%) to 22 (62.9%) for the 17 CPGs (Fig. 2). Of the 17 CPGs, only five (29.4%) had a reporting rate > 50%. The best one reported 62.9% of the items, while the poorest two reported < 26% of the 35 items.

Of the 35 items, the most frequently reported were #11a (Indicate whether the guideline is based on new systematic reviews done specifically for this guideline or whether existing systematic reviews were used), #12 (Describe the approach used to assess the certainty of the body of evidence), and #13c (Indicate the strength of the recommendations and the certainty of the supporting evidence), which were reported in 16 (94.1%) guidelines. These were followed by #13a (Provide clear, precise, and actionable recommendations), which was reported in 15 (88.2%) CPGs. Items #8b (Describe the setting(s) for which the guideline is intended, such as primary care, low- and middle-income countries, or inpatient facilities) and #17 (Indicate whether the guideline was subjected to a quality assurance process. If yes, describe the process) were reported in none of the CPGs (Fig. 3) (Table 2).

Table 2
Quality of the guidelines, according to the reporting of each item on the RIGHT checklist

		17	16	15	14	13	12	11	10	9	8	7	6
		2009 FNPP	2011 TWMU	2012 ACR	2012 FNPP	2013 SIR	2013 3E	2013 SER	2014 PORTUGAL	2015ACR/EULAR	2015 APLAR	2016 T2T	2016 EULAR
basic information	1a	N	Y	Y	N	Y	Y	Y	Y	N	Y	Y	Y
	1b	N	N	Y	N	N	N	N	N	Y	N	N	Y
	1c	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	Y
	2	P	Y	Y	P	Y	Y	N	Y	Y	Y	Y	Y
	3	N	N	P	N	P	Y	Y	Y	N	N	Y	P
	4	N	Y	Y	N	Y	Y	N	Y	Y	Y	Y	Y
total		1	4	5	1	4	5	3	5	4	4	4	5
background	5	N	N	Y	N	Y	P	Y	N	Y	Y	N	P
	6	N	Y	Y	Y	N	N	Y	P	N	P	P	N
	7a	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	Y
	7b	N	Y	N	N	N	N	N	N	N	N	N	Y
	8a	Y	N	Y	Y	N	N	Y	N	N	N	N	P
	8b	N	N	N	N	N	N	N	N	N	N	N	N
	9a	N	N	Y	N	N	P	Y	P	N	Y	P	Y
	9b	N	N	P	P	P	P	Y	P	P	P	Y	Y
total		2	3	5	3	1	0	5	0	2	3	2	4
Evidence	10a	N	N	Y	N	Y	Y	P	Y	N	Y	N	P
	10b	N	N	N	N	Y	N	N	N	N	N	Y	N
	11a	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
	11b	P	N	N	N	N	Y	P	N	N	N	N	N
	12	Y	Y	Y	Y	Y	Y	Y	Y	N	Y	Y	Y
total		2	2	3	2	4	4	2	3	1	3	3	2
Recommendations	13a	P	Y	Y	P	Y	Y	Y	Y	Y	Y	Y	Y
	13b	N	Y	N	N	N	N	N	N	N	N	N	Y
	13c	Y	Y	Y	Y	Y	Y	Y	Y	N	Y	Y	Y
	14a	N	Y	N	N	N	N	N	N	N	Y	N	N
	14b	N	N	N	N	Y	N	N	N	N	N	N	N
	14c	N	N	N	N	Y	N	N	N	N	N	N	N
	15	N	Y	Y	N	N	Y	Y	Y	Y	Y	Y	Y
total		1	5	3	1	4	3	3	3	2	4	3	4
review and quality assurance	16	Y	Y	N	N	N	N	N	N	N	N	P	P
	17	N	N	N	N	N	N	N	N	N	N	N	N
total		1	1	0	0	0	0	0	0	0	0	0	0
Funding and declaration and management of interests	18a	Y	N	Y	Y	N	Y	Y	N	Y	N	Y	N
	18b	N	NA	N	N	NA	N	Y	NA	N	NA	N	NA
	19a	N	N	Y	N	N	Y	Y	N	Y	N	Y	Y
	19b	NA	NA	N	NA	NA	N	N	NA	N	N	N	P
total		1	0	2	1	0	2	3	0	2	0	2	1
other information	20	Y	Y	Y	Y	N	Y	N	N	Y	N	Y	Y
	21	N	N	Y	N	N	N	N	N	N	N	N	Y

	17	16	15	14	13	12	11	10	9	8	7	6
22	N	N	Y	N	N	Y	N	N	N	P	N	N
Total	1	1	3	1	0	2	0	0	1	0	1	2
	9	16	21	9	13	16	16	11	12	14	15	18

According to the RIGHT tool, the reporting rates of the seven domains (Basic information, Background, Evidence, Recommendations, Review and quality assurance, Statements and management of funds and interests, and Other information) were 64.7%, 36.8%, 50.6%, 50.6%, 42.9%, 8.82%, 33.8%, and 31.4%, respectively.

Subgroup analysis

The mean number of reporting items of the nine guidelines developed by European countries was 15.2, and only three guidelines had a compliance rate of > 50% (33.3%); the mean number of reporting items in the four EULAR-generated CPGs was 15.8. For the six guidelines developed in the United States, the mean number of reporting items was 14.8. Two guidelines were prepared by the ACR, with a mean number of reporting items of 16.5, and two guidelines, which were published on NGC (National Guideline Clearinghouse), reported only nine items. Twenty-two items were reported in the American College of Physicians (ACP) guideline, which was the guideline with the largest number of reporting items among the 17 included in the analysis.

Eight guidelines were published between 2009 and 2014, which had a mean of 13.9 reporting items, and only one (12.5%) guideline had a compliance rate of more than 50%. Between 2015 and 2019, a total of nine guidelines were published, with a mean of 15.8 items being reported, but only four (44.4%) had a compliance rate of > 50%.

Analysis of each section of the RIGHT checklist

Basic information

With regard to the basic information, most of the CPGs could be judged by their title (82.4%), and the contact information for at least one author could be found (82.4%). The classification of the guideline could be easily discerned in 76.5% of the CPGs and a summary was presented in 70.5%. However, only 29.4% of the CPGs reported a publication date. Forty-one-point-two percent of the CPGs defined new terms and gave corresponding acronyms.

Background

The 17 CPGs did not comprehensively describe the background. More than half reported the epidemiology of gout (52.9%). Six (35.3%) guidelines reported the overall objectives of the guidelines and the specific objectives they were designed to achieve. Most of the guidelines described the main target population (76.5%), but only 17.6% described the subgroups that should be considered. Fewer than half (47.1%) of the guidelines described the key users and other potential users of the guidelines, and none described the specific target environment. Six (35.3%) CPGs listed all the contributors and their roles, and six (35.3%) gave the titles and affiliations of all the participants in their development, but only four (23.5%) contained both sets of information.

Evidence

Seven (41.2%) of the guidelines described the key findings on which the recommendations were based, but only two described the selection and classification of outcomes, and only one reported both. Sixteen (94.1%) guidelines described whether the systematic reviews on which the guidelines were based had been recently completed, but only two (11.8%) presented references, described how they were retrieved and evaluated, and stated whether they had been updated. Only one CPG did not describe the method of evaluation and grading of the quality of evidence.

Recommendations

Most of the guidelines (15, 80.2%) provided clear, accurate, and enforceable recommendations. Sixteen (94.1%) CPGs described the strength of the recommendation and the quality of the evidence supporting it, but only four provided recommendations for subgroups. Four CPGs took into account the preferences and values of the target group, two (11.8%) took into account cost and resource utilization, two (11.8%) took into account fairness, feasibility and acceptability, when creating their recommendations. Most of the guidelines (15, 88.2%) described the decision-making process and the methods used by the working group that created the guideline.

Review and quality assurance

Only three (17.6%) CPGs described whether they had been sent for review or not, and none described the level of supervision involved.

Funding and declaration and management of interests

More than half of the CPGs (9, 52.9%) described the sources of funding at each stage of their development, but only one described the role of the sponsor in the different stages of guideline formulation, as well as in the dissemination and implementation of the recommendations. Similarly, most of the CPGs (11, 64.7%) described the types of conflict of interest associated, but only two (11.8%) described the evaluation and management of these conflicts of interest and how users of the guidelines could obtain this information.

Other information

Eleven (64.7%) CPGs described where the guidelines, corresponding attachments, and other related documents could be obtained. However, only three (17.6%) articles described the differences between current practice and that recommended by research evidence, and/or provided recommendations for future research. In addition, only two CPGs described all the limitations associated with the formulation of the guidelines and their possible impact on the effectiveness of the recommendations.

Discussion

The quality of CPGs about gout and hyperuricemia were assessed before with AGREE II[5], whereas AGREE II and RIGHT checklist had unique items by themselves [16]; the RIGHT checklist added new and detailed items that AGREE II lacked. We assessed reporting quality of CPGs in the field of gout and hyperuricemia, using the RIGHT checklist in our study, to help guideline developers better standardizing the reporting quality of future CPGs. Furthermore, it also could help healthcare professionals better understand and implement the contents of such CPGs [12]. We used the RIGHT checklist to evaluate 17 guidelines for gout and hyperuricemia of various types that had been prepared in different geographical regions over the past 10 years, and found that they varied in quality. Of the 17 CPGs, only five reported > 50% of the items in the RIGHT tool, suggesting that many were of low quality. This finding is similar to that made previously using the AGREE tool [13].

Among the seven sections of the RIGHT tool, the reporting rate of Basic information was the highest (64.7%), followed by Evidence (50.6%), but the reporting rate of the other sections was < 50%. The most poorly reported section was the Declaration and management of funds and interests, with only 8.8% of CPGs reporting these items. This suggests that the guideline creators are generally good at including basic information, but that review and quality assurance are easily ignored.

Most of the CPGs were written by European and American organizations, and the overall completeness of the guidelines was slightly better in Europe than that in the United States. However, the scores for the CPGs written by the ACR and ACP, which are influential guideline-writing organizations in the United States, were higher than those for EULAR-generated CPGs. This shows that there are differences in the quality of the CPGs written by different organizations in different geographical regions. With regard to the relationship between publication date and score, the compilation of guidelines has greatly improved in recent years. The guideline writers have aimed to correct the deficits of the previous CPGs with regard to the practical applications of the guidelines, but the emergence of guideline-based systematic reviews in recent years have also made them consider writing their guidelines in accordance with certain norms.

Analysis of the reporting quality of guidelines regarding gout and hyperuricemia

Basic information

Most of the CPGs had a high reporting rate for basic information, but there were still some in which the year of publication was not clear and there was ambiguity in the definitions of acronyms. These issues are very important for researchers and practitioners who wish to know how up-to-date guidelines are, and it is necessary to interpret vague terms and acronyms for more accurate understanding of the guidelines.

Background

In the background to the CPGs, the writers usually described the main target groups for the guidelines and discussed the epidemiology of the problems described. However, the other items were frequently not well reported. For example, the inclusion of 8a (the main users of the guide should be described) and 8b (the specific environment for which the guide should be described) demonstrate that the application of the guidelines will vary for different users and different environments. For example, for low- and middle-income people, it is necessary to consider the economic benefits [17]. Therefore, the background to the guidelines should be described in detail.

Evidence

About half of the items in the Evidence section were reported, and items 10b (the methodology for the selection and classification of outcomes should be described) and 11b (if guide makers use published systematic reviews, references should be given and describe how they were retrieved and evaluated) had low reporting rates. Items 10b and 11b can greatly help researchers and practitioners to understand the evidence and assess the accuracy of the guidelines. Furthermore, they are important for peer review, because they permit shortcomings in the CPGs to be found and remedied. Items 10b and 11b can also be supplemented as new evidence accumulates over time.

Recommendations

Most of the guidelines did not make recommendations for subgroups, suggesting that the writers did not regard subgroups as important. Most writers did not consider items 14a (whether to consider the preferences and values of the target population), 14b (whether to consider cost and resource utilization), or 14c (whether to consider fairness, feasibility, and acceptability), or they considered these issues but did not included them in the CPGs. This means that healthcare professionals cannot easily adapt the CPGs to differing clinical situations.

Review and quality assurance

Items in this section were described in only a small number of guidelines, such that the reporting rate was the lowest for all the sections. Some of the guidelines may have been independently reviewed and quality-controlled, but the absence of such information is likely to make guide users and peer reviewers doubt the quality of the guidelines.

Funding and declaration and management of interests

The RIGHT checklist showed that the quality of the published CPGs was low with regard to this aspect, suggesting that the writers did not pay enough attention to it. A lack of information in this domain may lead to the inference that the recommendations of the CPGs could have been influenced by multiple interested parties, which would lower the credibility of the guidelines, especially with regard to specific treatment recommendations.

Other information

Most of the CPGs failed to describe the differences between practice and research evidence, or to provide recommendations for future research (or both). They also failed to describe the limitations in the formulation of the CPGs and the possible impact of these limitations on the effectiveness of the recommendations. Such content could have provided a reference for guide users to appropriately use the recommendations, and provided guidance for future updates and other researchers.

Strengths And Limitation

Strengths

This is the first study to use the RIGHT tool to evaluate the quality of CPGs in the field of gout and hyperuricemia. In addition, the methods of systematic retrieval, screening, and evaluation were adopted, and the quality of the outcome measurements was strictly controlled.

Limitations

First, only guidelines written in Chinese and English were included. Second, only CPGs published in journals and online regarding gout and hyperuricemia were included; CPGs published in the form of books or government documents were not analyzed.

In conclusion, by using the RIGHT checklist to evaluate CPGs in the field of gout and hyperuricemia, we found that their standards substantially differed, unfortunately most of the current CPGs had relatively low scores. The RIGHT evaluation form is a comprehensive tool for the evaluation of the quality of such guidelines. We suggest organizations that participate in the reporting of guidelines regarding gout and hyperuricemia to use this tool as to improve standardization of reporting.

Abbreviations

RIGHT

Reporting Items for Practice Guidelines in Healthcare

CPGs

Clinical Practice Guidelines

EULAR

European League Against Rheumatism

AGREE

Appraisal of Guidelines for Research and Evaluation tool

CBM

Chinese Biomedical Literature database

CNKI

China National Knowledge Infrastructure

GIN

Guidelines International Network

NICE

National Institute for Health and Clinical Excellence

ACR

American College of Rheumatology

NGC

National Guideline Clearinghouse

ACP

American College of Physicians

Declarations

Funding: This study was supported by the National Natural Science Foundation of China (grant numbers 81520108007, 81770869, and 81801434), the Natural Science Foundation of Shandong Province (grant number ZR2018ZC1053), the Qingdao Applied Basic Research Project (grant number 19-6-2-53-cg), and the Shandong Province Key Research and Development Program (grant number 2018CXGC1207).

Conflicts of interest: The authors declare that they have no conflict of interest.

Ethics approval: This article does not contain any studies conducted on human participants or animals that required ethics approval.

Consent for publication All authors have approved the manuscript and agree with this submission .

Authors' contributions: Can Wang, Xufei Luo, and Maichao Li contributed equally to this work. Xufei Luo and Yaolong Chen designed the study. Can Wang, Zhen Liu, and Maichao Li conducted the evaluation, wrote the manuscript, and complement the methods of this study. Xufei Luo and Yaolong Chen supplied methodological assistance. Changgui Li supplied funding and helped revise the manuscript. All authors provided input regarding the direction of the study and the content of the paper. All authors approved the final version of the paper.

Acknowledgments

We thank colleagues from the Evidence-Based Medicine Center, School of Basic Medical Sciences, and Lanzhou University (Lanzhou, China), including Xufei Luo and Yaolong Chen, for their kind suggestions and assistance with the study.

References

1. Kuo CF, Grainge MJ, Zhang W, Doherty M. Global epidemiology of gout: prevalence, incidence and risk factors. *Nat Rev Rheumatol* 2015;11(11):649-62.
2. Rai SK, Burns LC, De Vera MA, Haji A, Giustini D, Choi HK. The economic burden of gout: A systematic review. *Semin Arthritis Rheum* 2015;45(1):75-80.
3. Shiozawa A, Szabo SM, Bolzani A, Cheung A, Choi HK. Serum Uric Acid and the Risk of Incident and Recurrent Gout: A Systematic Review. *J Rheumatol* 2017;44(3):388-96.
4. MacFarlane LA, Kim SC. Gout: a review of nonmodifiable and modifiable risk factors. *Rheum Dis Clin North Am* 2014;40(4):581-604.
5. Wang D, Yu Y, Chen Y, et al. Assessing the Quality of Global Clinical Practice Guidelines on Gout Using AGREE II Instrument. *J Clin Rheumatol* 2020;26(2):54-9.
6. Yu Y, Wang D, Zhou Q, et al. Recommendations in clinical practice guidelines on gout: systematic review and consistency analysis. *Clin Exp Rheumatol* 2020.
7. Ughi N, Prevete I, Ramonda R, et al. The Italian Society of Rheumatology clinical practice guidelines for the diagnosis and management of gout. *Reumatismo* 2019;71(S1):50-79.
8. Li Q, Li X, Wang J, et al. Diagnosis and treatment for hyperuricemia and gout: a systematic review of clinical practice guidelines and consensus statements. *BMJ Open* 2019;9(8):e026677.
9. Yang N, Yu Y, Zhang A, et al. Reporting, presentation and wording of recommendations in clinical practice guideline for gout: a systematic analysis. *BMJ Open* 2019;9(1):e024315.
10. Guidelines IoMCoSfDTCP, Graham R, Mancher M. *Clinical practice guidelines we can trust*. National Academies Press Washington, DC; 2011.
11. Grilli R, Magrini N, Penna A, Mura G, Liberati A. Practice guidelines developed by specialty societies: the need for a critical appraisal. *Lancet* 2000;355(9198):103-6.
12. Chen Y, Yang K, Marusic A, et al. A Reporting Tool for Practice Guidelines in Health Care: The RIGHT Statement. *Ann Intern Med* 2017;166(2):128-32.
13. Wang D, Yu Y, Chen Y, et al. Assessing the Quality of Global Clinical Practice Guidelines on Gout Using AGREE II Instrument. *J Clin Rheumatol* 2018.
14. Collaboration A. Development and validation of an international appraisal instrument for assessing the quality of clinical practice guidelines: the AGREE project. *Qual Saf Health Care* 2003;12(1):18-23.
15. Oxman AD, Schunemann HJ, Fretheim A. Improving the use of research evidence in guideline development: 16. Evaluation. *Health Res Policy Syst* 2006;4:28.
16. Yao X, Ma J, Wang Q, Kanters D, Ali MU, Florez ID. A Comparison of AGREE and RIGHT: which Clinical Practice Guideline Reporting Checklist Should Be Followed by Guideline Developers? *J Gen Intern Med* 2020;35(3):894-8.
17. Xiao Y, Jiang L, Tong Y, et al. Evaluation of the quality of guidelines for assisted reproductive technology using the RIGHT checklist: A cross-sectional study. *Eur J Obstet Gynecol Reprod Biol* 2019;241:42-8.
18. Richette P, Doherty M, Pascual E, et al. 2018 updated European League Against Rheumatism evidence-based recommendations for the diagnosis of gout. *Annals of the rheumatic diseases* 2019;annrheumdis-2019-215315.
19. Yu KH, Chen DY, Chen JH, et al. Management of gout and hyperuricemia: multidisciplinary consensus in Taiwan. *International journal of rheumatic diseases* 2018;21(4):772-87.
20. Hui M, Carr A, Cameron S, et al. The British Society for Rheumatology Guideline for the Management of Gout. *Rheumatology (Oxford)* 2017;56(7):e1-e20.
21. Qaseem A, McLean RM, Starkey M, Forciea MA, Clinical Guidelines Committee of the American College of P. Diagnosis of Acute Gout: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med* 2017;166(1):52-7.
22. Qaseem A, Harris RP, Forciea MA, Clinical Guidelines Committee of the American College of P. Management of Acute and Recurrent Gout: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med* 2017;166(1):58-68.
23. Chinese Rheumatology A. [2016 China gout clinical practice guideline]. *Zhonghua Nei Ke Za Zhi* 2016;55(11):892-9.
24. Richette P, Doherty M, Pascual E, et al. 2016 updated EULAR evidence-based recommendations for the management of gout. *Ann Rheum Dis* 2017;76(1):29-42.
25. Kiltz U, Smolen J, Bardin T, et al. Treat-to-target (T2T) recommendations for gout. *Ann Rheum Dis* 2017;76(4):632-8.
26. Graf SW, Whittle SL, Wechalekar MD, et al. Australian and New Zealand recommendations for the diagnosis and management of gout: integrating systematic literature review and expert opinion in the 3e Initiative. *International journal of rheumatic diseases* 2015;18(3):341-51.
27. Neogi T, Jansen TL, Dalbeth N, et al. 2015 Gout classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Ann Rheum Dis* 2015;74(10):1789-98.

28. Araújo F, Cordeiro I, Teixeira F, et al. Portuguese recommendations for the diagnosis and management of gout. *Acta reumatologica portuguesa* 2014;39(2):158-71.
29. Spanish Society of Rheumatology (SER). Clinical Practice Guidelines for Management of Gout. In. www.formacionsanitaria.com; 2013.
30. Sivera F, Andres M, Carmona L, et al. Multinational evidence-based recommendations for the diagnosis and management of gout: integrating systematic literature review and expert opinion of a broad panel of rheumatologists in the 3e initiative. *Ann Rheum Dis* 2014;73(2):328-35.
31. Manara M, Bortoluzzi A, Favero M, et al. Italian Society of Rheumatology recommendations for the management of gout. *Reumatismo* 2013;4-21.
32. The University of Texas at Austin SoN, Family Nurse Practitioner Program, . Management of chronic gout in adults. In. <https://www.ahrq.gov/gam/index.html>; 2012.
33. Khanna D, Fitzgerald JD, Khanna PP, et al. 2012 American College of Rheumatology guidelines for management of gout. Part 1: systematic nonpharmacologic and pharmacologic therapeutic approaches to hyperuricemia. *Arthritis Care Res (Hoboken)* 2012;64(10):1431-46.
34. Khanna D, Khanna PP, Fitzgerald JD, et al. 2012 American College of Rheumatology guidelines for management of gout. Part 2: therapy and antiinflammatory prophylaxis of acute gouty arthritis. *Arthritis Care Res (Hoboken)* 2012;64(10):1447-61.
35. Yamanaka H, Japanese Society of G, Nucleic Acid M. Japanese guideline for the management of hyperuricemia and gout: second edition. *Nucleosides Nucleotides Nucleic Acids* 2011;30(12):1018-29.
36. The University of Texas at Austin SoN, Family Nurse Practitioner Program, . Management of initial gout in adults. In. <https://www.ahrq.gov/gam/index.html>; 2009.

Figures

Figure 1

Literature selection flow chart

Figure 1
Literature selection flow chart

Figure 2
Number of reported items in each guideline

Figure 2

Number of reported items in each guideline

Figure 3

Number of guidelines reporting each item Notes: Items listed according to the reporting frequency of "Y".

Figure 3

Number of guidelines reporting each item Notes: Items listed according to the reporting frequency of “Y”.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupportingFile.pdf](#)
- [SupportingFile.pdf](#)