Single-Base Resolution Methylomes of Somatic Embryogenesis in *Theobroma Cacao* L. Reveal Epigenome Modifications Associated With Somatic Embryo Abnormalities

Claudia Garcia (✉ garcia.claudia2000@gmail.com)

Mars Center for Cocoa Science

Alex-Alan Furtado de Almeida

Department of Biological Sciences, State University of Santa Cruz

Marcio Costa

Department of Biological Sciences, State University of Santa Cruz

Dahyana Britto

Mars Center for Cocoa Science

Fabio Correa

Department of Statistics, Rhodes University

Pedro Mangabeira

Department of Biological Sciences, State University of Santa Cruz

Lidiane Silva

Mars Center for Cocoa Science

Jose Silva

Department of Biological Sciences, State University of Santa Cruz

Stefan Royaert

Mars Incorporate

Jean-Philippe Marelli

Mars Incorporate

Research Article

Keywords: Cacao, epigenetic effects, hypoxia, abnormalities, somatic embryogenesis, programmed cell death, oxidative stress

Posted Date: December 29th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1074593/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Propagation by somatic embryogenesis in *Theobroma cacao* has some issues to be solved, as many morphologically abnormal somatic embryos that do not germinate into plants are frequently observed, thus hampering plant production on a commercial scale. For the first time the methylome landscape of *T. cacao* somatic embryogenesis was examined, using whole-genome bisulfite sequencing technique, with the aim to understand the epigenetic basis of somatic embryo abnormalities. We identified 873 differentially methylated genes (DMGs) in the CpG context between zygotic embryos, normal and abnormal somatic embryos, with important roles in development, programmed cell death, oxidative stress, and hypoxia induction, which can help to explain the morphological abnormalities of somatic embryos. We also identified the role of ethylene and its precursor 1-aminocyclopropane-1-carboxylate in several biological processes, such as hypoxia induction, cell differentiation and cell polarity, that could be associated to the development of abnormal somatic embryos. The biological processes and the hypothesis of ethylene and its precursor involvement in the somatic embryo abnormalities in cacao are discussed.

Introduction

Theobroma cacao L. (2n= 20) is a preferentially allogamous plant, classified in the Malvaceae family. It has been grown in the lowlands of tropical regions (South and Central America, West Africa and South East Asia) with social and economic importance because it is the income of thousands of small-holder farmers around the world ¹. Cacao beans are used as raw material to produce cocoa powder (cacao byproduct), liquor and cocoa butter, which are the main ingredients of chocolate, and to produce additives for pharmaceuticals ^{2,3}. Cocoa chemical compounds have nutritional value mainly the flavanols because they are important substances for human health ⁴. Despite its importance, cacao propagation has been done by traditional methods with sometimes low efficiencies, such as grafting and rooted cuttings, or by the seedlings generating genetic depression in the parental lines that is a negative characteristic for breeding programs ⁵. Other propagation technologies, such as somatic embryogenesis, have been developed with the aim to obtain plants with genetic uniformity through the cloning process and rapid and efficient propagation of cacao plants ⁶. Despite the tremendous efforts to develop a viable somatic embryogenesis protocol for cacao plant propagation, there are still some issues to be solved, including the main one related to the high frequencies of abnormal embryos that cannot be converted into normal plantlets ⁷.

Embryogenesis in plants can take place sexually (zygotic embryogenesis) or asexually (apomixis). Zygotic embryogenesis starts with double fertilization of the male and female gametes ^{8,9}. Zygotic embryos (ZE) go through different stages of development, i.e., globular, heart, torpedo and cotyledonal in dicotyledonous plants versus, globular, scutellar and coleoptilar stages in monocotyledonous plants; and early and late embryogenesis in gymnosperm plants ¹⁰. Asexual embryogenesis, on the other hand is a complicated process that can be induced naturally (gametophytic or sporophytic) or artificially (somatic embryogenesis), where the stages of development are the same between ZE and somatic embryos (SE) ⁹. Although ZE and SE show similarities in the stages of differentiation, their morphologies can be quite different ¹¹. Besides, ZE are protected by endosperm and SE do not have this structure ¹⁰.

Somatic embryos in *T. cacao* have been developed in tissue culture from different explant sources, such as leaves, flowers, and nucellar tissue, with the highest success using flower explants (petals and staminodes) ⁶. Cacao somatic embryogenesis process is divided in six well defined steps as follows: induction, expression, multiplication, maturation, germination and plant conversion ¹². All these steps are mediated by a complex regulatory network where different genes are regulated at the transcriptional level by both genetic and epigenetic mechanisms ^{11,13}.

Chromatin structure is affected by epigenetic mechanisms such as histone modifications, small RNAs (sRNAs) and DNA methylation, ¹⁴. The lysine residues at the N-terminal tails of the histone proteins can be covalently modified by acetylation or methylation, thereby changing the chromatin configuration. Acetylation allows gene transcription, whereas methylation inhibits or enhances it depending where the methyl group is positioned in the gene region. Although methylation of the histone H3K9 is related to transcriptional repression, methylation of the histone H3K4 is linked to enhancement of gene expression¹⁵. sRNAs act independently of histone modification in targeting DNA methylation by RNA-directed DNA methylation mechanism (RdDM) or at the post-transcriptional level (leading sequence-specific transcript degradation and translational repression). Inherited epigenetic variation in plants is led by the RdDM by targeting the de novo DNA methylation.

DNA methylation has been widely described as a mechanism implicated in genomic imprinting, repression of gene expression and transposon silencing. DNA methylation in plants occurs when a methyl group is attached to the carbon five of the cytosine. This takes mainly place at cytosines in CpG islands (long repetition of C and G), CHG and CHH (H= A, T or C) sites ¹⁶. DNA methylation profiles may change under variation in environmental conditions that can be stressful for the plants ¹⁷. It has been known that CG methylation is higher in animals, while methylation in plants has been found in the three contexts (CG, CHG, and CHH) ¹⁴. The position of the methyl group in the gene fulfills important and different functions. If the methyl group is positioned at the regulatory sequences, gene expression is repressed, but methylation also plays an important function when it is placed in the gene body. It has been known that exons are more highly methylated than introns, where methylation in the first exon of the gene blocks gene transcription and methylation in the internal exons does not affect it seriously ¹⁸. On the other hand, methylation in introns is associated with gene transcription enhancement ¹⁹

In somatic embryogenesis, the tissue is subjected to different environmental stimuli, such as plant grow regulators (PGRs), where the explants and the somatic embryos produced in culture undergo stress. A key PGR used in somatic embryogenesis is 2,4-dichlorophenoxyacetic acid (2,4-D), a synthetic auxin that is known as an important cell stressor and has an important role in cell reprogramming through DNA methylation ²⁰. 2,4-D plays a role in cell polarity and asymmetric cell division that influence changes in the cell morphology, physiology, metabolism and gene expression ^{21–23}. 2,4-D is also known as a potent inducer of ethylene production in plant tissues ^{24,25}. Ethylene is an important phytohormone that works together with jasmonic acid (JA), salicylic acid (SA), abscisic acid (ABA), among others, to regulate several biological processes, including oxidative stress, hypoxia and programmed cell death (PCD) ²⁶. Furthermore, it is known to act as an important mediator in aerenchyma formation under hypoxic environments ²⁷.

In order to understand the epigenetic basis of abnormal somatic embryo production in *T. cacao*, whole-genome bisulfite sequencing (WGBS) was carried out with the aim to identify differentially methylated genes (DMGs), and their functions, among ZE, normal and abnormal SE. In addition, scanning and transmission electron microscopy analyses were carried out to identify morphological and structural differences between ZE and SE. Understanding the main factors that influence the abnormal SE formation could provide guidance towards somatic embryos development processes to improve the quality and efficiency of the plantlets produced. Additionally, our results provide new knowledge in the somatic embryogenesis regulatory network during ZE and SE differentiation in *T. cacao*, providing insights in the basic biology of the somatic and zygotic embryogenesis development.

Results

Morphological characterization of somatic embryos

We first compared the phenotypic characteristics among ZE, normal and abnormal SE at the late torpedo stage to identify the types of abnormalities present in primary SE and secondary SE from CCN 10 (Figures 1A-I). In a population of 118 primary SE and 530 secondary SE, we were able to identify seven different abnormal types (AT) of embryos. Depending on their morphologies, the AT were classified as AT1-7 (Figures 1A-G). AT1 has more than two fused embryo axes; AT2 has no root and cotyledon formation; AT3 is translucent with curved axis, single cotyledon and without apical shoot formation; AT4 is translucent with more than two cotyledons, thick axes without root and shoot meristem formation; AT5 contains long and slender axes, poor cotyledon development and without root and shoot meristem formation; AT6 is translucent with an amorphous axes and a single amorphous cotyledons; AT7 is long and slender cotyledons with curved and amorphous axes.

Concerning the primary SE, 18.64% of the embryos were normal type (Figure 1H) and the other 81.36% were abnormal, where the AT2 was predominant, with frequency of 20.34% from the total population. Of the total secondary SE, on the other hand, only 18.11% were normal and 81.89% were abnormal, with 27.92% of AT2 and 20.75% of AT3 as the main ones (Table 1). The seven anomalous phenotypes of secondary SE were subjected to germination and only 1% of AT2 embryos generated an apical shoot and 4% of AT3 embryos generated axillar shoots and adventitious roots with callus formation in the axis base, while the other SE abnormal types did not germinate (Supplementary Figures S1A and S1B).

Table 1
Number of normal and abnormal somatic embryos and percentage in primary and secondary somatic embryogenesis

Type of embryo	Primary somatic embryogenesis		Secondary somatic embryogenesis	
	Number of embryos	Percentage (%)	Number of embryos	Percentage (%)
AT1	24	20.34	49	9.25
AT2	42	35.59	148	27.92
AT3	3	2.54	110	20.75
AT4	12	10.17	22	4.15
AT5	0	0.00	20	3.77
AT6	13	11.02	60	11.32
AT7	2	1.69	25	4.72
Normal Embryos	22	18.64	96	18.11
Total	118	100	530	100

Microscopic analysis

AT4 embryos were analyzed using scanning electron microscopy (SEM) due to their similarities with normal SE. Transversal and longitudinal cuts revealed several cavity formations between intercellular spaces along the embryo body (Figures 2A-D). In contrast, transverse and longitudinal cuts from normal SE showed asymmetric cell divisions and no cavities were formed (Figures 2E-H). The analysis shows that in an area of 400 mm² there are no cavities in normal SE, but there are on average 3.0 cavities in the same area in AT4 SE, with an average of 217 µm in size.

Transmission electron microscopy (TEM) analysis of ZE and normal SE showed that their cells have nuclei with normal appearances, whereas some cells in abnormal SE showed a lot of vesicles and large lytic vacuoles without nucleus presence (Figures 3A-C).

Normal and abnormal SE exhibited PCD with some differences (Figures 4A-H). Red fluorescence color in abnormal SE was more intense than in normal SE in the fragmented DNA inside cells (Figures 4A-H). While abnormal SE showed cell death in apical and root meristems (Figures 4C-D and 4G-H respectively), normal SE showed cell death only in the epidermal layer in cotyledons (Figure 4B).

Normal SE showed an average of 97 cells with PCD in an area of 34.72 mm² in its cotyledons. On the other hand, abnormal SE showed in its cotyledons an average of 195 cells with PCD in the same area. Statistical analysis showed significant differences with p-values < 0.05 in fluorescence area and fluorescence intensity in cells of cotyledon, root and shoot regions between normal and abnormal SE. Cotyledons in abnormal SE showed the highest fluorescence area, with an average of 7.01 µm² (Figure 5), compared with cells of cotyledons in normal SE, which showed a fluorescence area of 2.49 µm² (Figure 5). Cells in root meristem region of abnormal SE showed a fluorescence area with an average of 5.78 µm² (Figure 5), whereas cells in the same region of normal SE showed a fluorescence area of 0.09 µm² (Figure 5). Cells in the shoot meristem region of normal SE were not stained in red fluorescence, showing no cellular death, but the same region in abnormal SE showed a fluorescence area of 0.36 µm² (Figure 5).

Whole-genome DNA methylation maps of zygotic and somatic embryos

Unbiased DNA methylation maps were generated from a set of mixed embryos samples from ZE, normal and abnormal SE by WGBS. We sequenced a total of 57.2 Gb from ZE, 74.1 Gb from normal SE and 72.5 Gb from abnormal SE. From the total base pairs, we mapped from each sample 86.14%, 67.08% and 68.30% of sites with methylation in ZE, normal and abnormal SE, respectively, with Phred quality score over 30, where the CpG content in percent (%) is 21.04%, 24.88% and 24.3% in each sample, respectively (Supplementary Table S1). After trimming process to eliminate adapter sequences and bases with low quality from each read, we generated a total of 53.8 Gb (ZE), 57.1 Gb (normal SE) and 54.6 Gb (abnormal SE), which were aligned to the *Cacao_Matina1-6* genome sequence used as reference genome. In the basic statistics of the alignment for ACGT content by chromosome, 86.58-104.03% were covered by at least one unambiguously mapped read with Phred quality score over 30 and standard deviation between 322.33 - 452.18, resulting in an overall average sequencing depth of 30-fold (Supplementary Table S2).

For methylation calling we used a total read of 203,071,419 from ZE, 263,500,094 from normal SE and 245,099,294 from abnormal SE (Supplementary Table S3). Methylation calling was different between the contexts. From the total sequencing coverage of CpG, CHG and CHH contexts across all the genome, only 50.14 - 53.95% (CpG), 28.23 - 31.58% (CHG) and 5.34 - 9.91% (CHH) were methylated. Normal SE showed the highest methylation percentages over the three contexts. By contrast, ZE showed the lowest and abnormal SE the intermediate values (Supplementary Table S4).

A total of 7,391,277 CpG, 10,706,841 CHG and 66,034,574 CHH methylation sites were identified in this study. For all the contexts (CpG, CHG, and CHH), the lower sequencing depth cut-off was 10. 7,361,253 CpG, 10,679,910 CHG and 65,894,628 CHH methylation sites to be analyzed. In the CpG context we found a binomial distribution in the level of methylation for most of the data (Supplementary Figure S2). ZE methylation landscape showed that 35.02% of the CpG were with DNA methylation levels between 0.0 - 0.1 (low methylation values) and 35.82% were with values between 0.9 - 1.0 (high methylation values). For normal SE, 36.33% of the CpG sites were low methylated and 46.11%

were high methylated. Finally, for abnormal SE 35.86% of CpG sites were low methylated and 43.43% were high methylated. These results show that normal and abnormal SE had a higher percentage of high levels of methylated CpG sites in comparison with ZE.

The CHG and CHH contexts were analyzed as bimodal and right-skewed distribution, respectively (Supplementary Figures S3 and S4). The CHG context in ZE, normal and abnormal SE samples showed the maximum data as low methylated, with values between 0.0-0.1, and with frequencies of 60.11%, 60.15% and 60.08% respectively. CHH context analysis showed methylation levels between 0.0-0.1 in ZE, normal and abnormal SE, with frequencies of 82.42%, 71.85% and 74.65%, respectively. The degree of reproducibility between samples, the normal distribution between samples and the density plot to compare distribution of methylation ratio before and after normalization were analyzed using Pearson's correlation (Range: $-1 \leq r \leq 1$), box plot and density plot, respectively (Supplementary Figures S5A-C, S6A-C and S7A-C).

DNA methylation dynamics in non-CpG contexts

Analyzing the CHG context ZE, normal SE and abnormal SE, the hierarchical clustering the Euclidian distances showed that normal and abnormal SE are grouped in the same cluster and ZE are grouped in a different cluster (Figure 6A) indicating that normal and abnormal SE have similar DNA methylation profiles while the DNA methylome of ZE is different. The frequency histograms represent the distribution of differentially methylated regions between samples, where the highest frequencies of the data are grouped to the left showing values with levels of methylation between 0.0-0.1 for ZE and normal SE (Figure 6B), ZE and abnormal SE (Figure 6C) and normal and abnormal SE (Figure 6D), indicating that the regions in the CHG context in the DNA methylome are mainly sparsely methylated. The analysis regarding the functional regions differentially methylated in the CHG context showed that introns and intergenic regions are the regions differentially methylated with the highest frequencies when compared to the DNA methylomes between ZE and normal SE (Figure 6E), between ZE and abnormal SE (Figure 6F) and between normal and abnormal SE (Figure 6G). Interestingly, the CHH context has the same profile as the CHG context in the DNA methylome of the ZE, normal and abnormal SE. In the CHH context the samples were grouped in the hierarchical clustering similarly as in the CHG context, normal and abnormal SE are together in the same cluster and ZE is grouped in a different cluster than SE. Differentially methylated regions have the lowest levels of methylation in this context, with values between 0.0-0.1, and the high frequencies of functional regions differentially methylated between samples are also the introns and intergenic regions (Supplementary Figures S8A-G).

DNA methylation dynamics in CpG context

The dynamics in the CpG context are different compared to the CHG and CHH contexts. The methylated CpG region coverage per base followed a normal distribution for ZE, normal and abnormal SE with high frequencies of values between 1-2 log 10 of read coverage per base and with percentages between 30.4 and 37.3 of coverage (Figure 7A). The percentage of methylation in all CpG regions presented a bimodal distribution for ZE, normal and abnormal SE with high frequencies of low methylated regions (0-1%) and highly methylated regions (90-100%) (Figure 7B). The DNA methylome comparison of ZE, normal and abnormal SE revealed high percentages of CpG regions differentially hypermethylated for all of the 10 chromosomes where the chromosomes 4, 6 and 9 showed the highest percentages of regions hypermethylated. In the whole CpG context the proportion of regions that are hypo and hyper methylated are quite similar comparing between samples (Figures 7C and E). Chromosome 6 shows the highest proportion of hypermethylation comparing ZE with abnormal and normal SE with values superior to 0.5. For the comparison between normal and abnormal SE, chromosomes 4 and 5 presented the highest proportions of hypermethylation, with

values superior to 0.6 (Figure 7E). In the hierarchical clustering CpG methylated regions of normal and abnormal SE are grouped in the same cluster and ZE is grouped in a different cluster (Figure 7D).

From a total of 7,361,253 CpG regions, 545,358 were differentially methylated between normal SE and ZE; 467,448 were differentially methylated between abnormal SE and ZE and 172,607 were differentially methylated between abnormal SE and normal SE. Comparison of the level of methylation between samples was carried out to estimate the level of methylation frequencies. In the histograms (Figures 8A-C) we can observe that there is a bimodal distribution of the levels of methylation for the comparison between ZE and normal SE, between ZE and abnormal SE, and between normal and abnormal SE, respectively. Analyzing the functional regions differentially methylated in the CpG context we found that intronic, intergenic and exonic regions are differentially methylated with the highest frequencies when compared to the DNA methylomes between ZE and normal SE (Figure 8D), between ZE and abnormal SE (Figure 8E) and between normal and abnormal SE (Figure 8F).

Functional annotation of differentially methylated sequences

Whole genomic sequences from ZE, normal and abnormal SE were analyzed and 873 out of 15,873 (between ZE and normal SE, and between normal and abnormal SE) and 23,957 (between ZE and abnormal SE) were significantly differentially methylated gene regions in CpG context found in this study. These sequences were functionally annotated as retrotransposons, transcriptional factors involved in autophagy, hypoxia cellular response, DNA repair, auxin response, auxin transport, stress response, disease resistance, cell polarity, PCD, cell wall biogenesis, cell transport, cell division, cellular homeostasis, cell signaling, cellular metabolism, and developmental processes such as embryo morphogenesis (Supplementary Table S5 - S8).

Methylation profiles in some regions of these genes were different for ZE and normal and abnormal SE when they are compared across the ten chromosomes (Figure 9A). Interestingly, ZE and abnormal SE have some regions with similar methylation level, whereas normal SE is generally different from ZE (Figure 9B). In the heatmap methylated regions are grouped in four hierarchical clusters (Figure 9B). The first cluster shows genes in ZE and abnormal SE that are hypomethylated, whereas in normal SE these genes are hypermethylated. The second cluster shows genes hypermethylated for ZE and hypomethylated for normal and abnormal SE. The third cluster contains genes in ZE that are hypomethylated and low methylated while in normal SE are hypermethylated and for abnormal SE the same genes are low methylated or not methylated. Finally, in the fourth cluster ZE have genes low and not methylated while normal SE has genes hypomethylated and abnormal SE has genes hypermethylated and low methylated (Figure 9B). It seems like the gene regulation by the methylation effect is complex and works modulating specific genes in different ways at the same time.

Furthermore, we focused on 22 out of 873 genes that are differentially methylated among ZE, normal and abnormal SE in CpG contexts because they are related to embryos development, PCD, defense and stress response mechanisms probably involved in inducing SE abnormalities as was noted in the microscopy and PCD analysis in this study (Table 2 and Figure 9C). On chromosome 10, the gene Thecc1EG045590 encodes the GroES-like zinc-binding alcohol dehydrogenase family protein with important function in the PCD process. This gene has a high level of methylation in intron and exon regions for ZE (Figure 9D) but for normal and abnormal SE the methylation level is low (Figures 9E and F respectively). Those types of embryos were analyzed for PCD, and they presented cells in the meristem zones with DNA damage indicating activity of this gene. Another important gene on the same chromosome is the Thecc1EG044386 that encodes a FASCICLIN-like arabinogalactan protein 16 precursor associated with shoot embryo development. Its intronic region is hypermethylated for ZE (Figure 9D) but it has low levels of methylation in normal and abnormal SE (Figures 9E and F respectively). Similarly, Thecc1EG031714 encodes a global transcription factor

group E4 on the chromosome 7 involved in embryo root development. ZE has a 0.75 methylation level in its intron and a 0.8 methylation level in his exon, while normal and abnormal SE have a 0.5 methylation level in their intron and 1.0 and 0.9 levels of methylation in their exon, respectively. Methylation in introns is associated with increased gene expression and methylation in exons is associated with gene repression. Intermediated values in the level of methylation of exons in ZE may explain that these embryos are in active development in torpedo stage inside the seed, but high levels of methylation in the same gene region in normal and abnormal SE may associated with development suppression of root meristems where abnormal SE is more repressed than normal SE. This was evaluated in the germination test performed in this work where different types of abnormal SE were submitted to germination and compared with normal SE the abnormal SE does not generate taproot.

Genes that cod for WRKY domain class transcription factor, jasmonate-zim-domain protein 3 and WRKY DNA-binding protein 70 (Thecc1EG010128, Thecc1EG012913 and Thecc1EG045556 respectively) important in defense response in plants are differentially methylated between the different type of embryos. The Thecc1EG010128 is methylated in the exon for ZE with 0.75 level of methylation, whereas the methylation levels for normal and abnormal SEs are 1.0 and 0.9, respectively. The exon and intron of the gene Thecc1EG012913 are methylated with level of 0.30 and 1.0, respectively for ZE, in normal and abnormal SE the intron is 0.60 and 0.1 respectively, but exon in normal SE is unmethylated, while for abnormal SE exon is 0.1 level of methylation. The intronic region of the gene Thecc1EG045556 is highly methylated for ZE (1.0 level of methylation) and for normal and abnormal SE is 0.30 and 0.70 level of methylation, respectively. Those methylation profiles might be associated with the low stress environment ZE are exposed inside the fruit, in difference normal and abnormal SE are more exposed to inappropriate environment in the *in vitro* culture conditions.

The different regions of genes that code for the Nucleic acid-binding proteins superfamily (Thecc1EG021358), Nudix hydrolase (Thecc1EG024914), Chromatin remodeling factor18 (Thecc1EG025136) and Peroxidase superfamily protein (Thecc1EG045425) between others are highly methylated for ZE but low methylated for normal and abnormal SE. These genes code for proteins important in stress response. The exonic region of Thecc1EG021358 and Thecc1EG024914 are more methylated in ZE than the same regions in normal and abnormal SE. In addition, the intronic regions of Thecc1EG024914, Thecc1EG025136 and Thecc1EG045425 are more methylated in ZE than the same regions in those genes for normal and abnormal SE. Methylation in exons is associated with induction or repression of gene expression depending on the position of the exon, but methylation in intronic region is associated with gene expression induction. The low level of methylation in the exon region of those genes in normal and abnormal SE might be associated with their high stress response to the *in vitro* culture conditions.

Thecc1EG039622, Thecc1EG020308 and Thecc1EG000039 are genes located on the chromosomes 1, 4 and 9 respectively, and play a role in the codification of Auxin efflux carrier family protein (*PIN* proteins). Methylation levels in ZE are 0.6 in the intronic region of the gene Thecc1EG039622 and 1.0 level of methylation for both normal and abnormal SE in the same region of the gene. ZE have 0.8 level of methylation in the UTR3 region for Thecc1EG020308 gene and in normal and abnormal SE it reached values of methylation levels 0.0 and 0.25, respectively, in the same regions of the gene. ZE have a methylation level of 0.75 in upstream of the transcriptional start site and 0.8 in the intronic region of the gene Thecc1EG000039. This is quite different from the methylation levels of 0.0 and 0.25 in the upstream region, and the levels of 0.0 and 0.1 in the intronic region in normal and abnormal SE. Auxin efflux carrier proteins fulfill important function in cellular transport where ZE seem to be active differently in this function compared to normal and abnormal SE. Another important gene that codes for an auxin transport protein is Thecc1EG019010 which is located on chromosome 4. In ZE, methylation levels are 0.7 in the exons and 0.75 in introns. Interestingly for normal and abnormal SE both regions have a methylation level of 1.0 which might indicate that probably the auxin trafficking in these cells is blocked (Figures 9D-F).

Regarding cellular signaling the genes Thecc1EG029410 (Auxin response factor 6) and Thecc1EG005126 (Auxin response factor 11) are differently regulated in ZE, normal and abnormal SE. In ZE the Auxin response factor 6 is methylated in its exonic region and upstream of the transcriptional start site with values of 0.8 and 0.9, respectively. In normal SE, the same regions have methylation levels of 0.1 and 0.2 (upstream and exonic regions). In abnormal SE, the methylation levels are 0.25 for both regions. In the Auxin response factor 11, methylation levels are 0.3 in upstream of the transcriptional start site and 0.7 in exon regions in ZE. In normal SE, methylation levels are less than 0.2 for both of regions. In abnormal SE, the methylation level is 0.2 for upstream transcriptional start site region and 0.4 for the exon region (Figure 9D-F). The high activity of those auxin response factors in normal and abnormal SE probably can occur because they are exposed to the auxin 2,4-D in the culture conditions.

Table 2

Differentially methylated genes in ZE, normal and abnormal SE with important functions in cellular process implicated in embryos development, stress response, defense response and PCD.

Cacao Gene ID	Chromosome	Position	Context	Protein	Function
Thecc1EG000039	1	168106..170565	CG	Auxin efflux carrier family protein	cellular transport
Thecc1EG005126	1	35923454..35927475	CG	Auxin response factor 11	cellular signaling
Thecc1EG011892	2	42804438..42817324	CG	Phosphoenolpyruvate carboxylase 4	photosynthesis
Thecc1EG010128	2	31421994..31432423	CG	WRKY domain class transcription factor	defense response
Thecc1EG012913	3	7678634..7683914	CG	Jasmonate-zim-domain protein 3	defense response
Thecc1EG016700	3	35110634..35118337	CG	Xyloglucanase 113	cell wall biogenesis
Thecc1EG015074	3	27103765..27106730	CG	GroES-like zinc-binding alcohol dehydrogenase family protein	programmed cell death
Thecc1EG020308	4	28823116..28825480	CG	Auxin efflux carrier family protein	cellular transport
Thecc1EG019010	4	20387381..20408383	CG	Auxin transport protein (BIG)	cellular transport
Thecc1EG021358	4	33815542..33819710	CG	Nucleic acid-binding proteins superfamily	stress response
Thecc1EG024914	5	30161616..30166159	CG	Nudix hydrolase	stress response
Thecc1EG025136	5	31557145..31566806	CG	Chromatin remodeling factor18	stress response
Thecc1EG029410	6	22823715..22832247	CG	Auxin response factor 6	cellular signaling
Thecc1EG029951	6	25524446..25527828	CG	Bromodomain 4	stress response
Thecc1EG031773	7	6985775..6991143	CG	Phosphoenolpyruvate carboxylase 3	photosynthesis
Thecc1EG031714	7	6606601..6611209	CG	Global transcription factor group E4	embryo root development
Thecc1EG035691	8	11465349..11468083	CG	Peroxidase	stress response
Thecc1EG039622	9	19300621..19303650	CG	Auxin efflux carrier family protein	cellular transport
Thecc1EG044386	10	16464327..16469069	CG	FASCIKLIN-like arabinogalactan protein 16 precursor	embryo shoot system development

Cacao Gene ID	Chromosome	Position	Context	Protein	Function
Thecc1EG045425	10	24923124..24924516	CG	Peroxidase superfamily protein	stress response
Thecc1EG045590	10	25845100..25847907	CG	GroES-like zinc-binding alcohol dehydrogenase family protein	programmed cell death
Thecc1EG045556	10	25728048..25729425	CG	WRKY DNA-binding protein 70	defense response

Non-CpG contexts in the embryo cacao epigenome had 15,942 (between ZE and normal SE and normal and abnormal SE) and 19,783 (between ZE and abnormal SE) differentially methylated gene regions in CHG context. In the CHH context, there were 24,658 (between ZE and normal SE), 23,820 (between ZE and abnormal SE) and 22,893 (between normal and abnormal SE) methylated gene regions. All these differentially methylated gene regions are part of genes that code for proteins with functions in autophagy, hypoxia cellular response, DNA repair, auxin response, auxin transport, stress response, disease resistance, cell polarity, PCD, cell wall biogenesis, cell transport, cell division, cellular homeostasis, cell signaling, cellular metabolism, and developmental processes such as embryo morphogenesis as is the case in the CpG context (Supplementary tables S9 - S14). The more interesting thing is that the non-CpG context were rich in regions of retrotransposon differentially methylated between ZE, normal and abnormal SE with values between 905 and 1,002 in the CHG context, where 122 are hypermethylated for ZE and 157 are hypomethylated for normal and abnormal SE. In the CHH context we found between 1,062 and 1,074 retrotransposons differentially methylated being 38 hypermethylated for ZE and 86 and hypomethylated for normal and abnormal SE. In contrast, the CpG context only had between 707 and 945 differentially methylated retrotransposons being 275 hypermethylated for ZE and 268 hypomethylated for normal and abnormal SE (Supplementary tables S6 – S14) where 14 were significantly differentially methylated when the methylome of ZE, normal and abnormal SE are compared (Supplementary table S5). Another important result in this work is the high number of methylated regions of unknown genes in the CpG context (between 3,933 and 5,902), CHG context (between 4,422 and 5,249) and CHH context (between 5,815 and 6,137) (Supplementary tables S6 - S14).

Discussion

Theobroma cacao somatic embryos produced in *in vitro* conditions were classified morphologically in normal and abnormal embryos, with seven different morphologies identified for abnormal SE in our work. Abnormal SE were subjected to germination and only AT2 embryos developed an apical meristem. AT3 embryos were able to develop axillar meristems and adventitious roots, but they were not able to develop into normal plants (Figure 1B-C). Abnormalities in somatic embryos of *T. cacao* are frequently found in other works, where the researchers identified somatic embryos with more than two cotyledons, fasciation and fusion of more than two cotyledons, but the behavior of these abnormal SE in germination is not known²⁸. The histological (Figure 4C and G) and scanning electron microscopy analysis of abnormal SE (Figure 2A-D) revealed abnormal cell division in the provascular tissue originating from the provascular initial cells²⁹ and some intercellular cavity formation in the embryo body similar to cavity formation through both schizogenous or lysigenous aerenchyma formation in plants when they are exposed to flooding conditions³⁰. Schizogenous aerenchyma is formed by cell division and enlargement of intercellular spaces of cortical cells where there is a separation between the adjacent cell files, whereas lysigenous aerenchyma is formed by the collapse or lysis of the cortical cell files via programmed cell death (PCD)³⁰.

PCD in plants is induced by biological processes essential for embryo development, formation and maturation of many tissues and stress response to environmental conditions for adaptation or reaction to biotic and abiotic stimuli³¹. PCD in abnormal SE was notoriously higher than in normal SE (Figure 4A-H). The TUNEL analysis showed that there was DNA fragmentation because the cell nuclei stained fluorescent red in the tissue analyzed, mainly in the embryo body in abnormal SE (Figure 4H)³². The TUNEL analysis is characterized for the label of free 3'OH ends in the fragmented genomic DNA with a fluorescein-dUTP or TMR-dUTP (Tetramethylrhodamine-dUTP) that bright red fluorescent at fluorescence microscope. Another indication of PCD in abnormal SE was the presence of lytic vacuoles in their cells (Figure 3C) and the absence of a nucleus. These signals are characteristic of vacuolar cell death in plants, which can be stimulated by recycling part of their cells during normal development or stress response³³. In this study we identified genes differentially methylated in ZE, normal and abnormal SE with important functions in PCD, specifically in cell response to hypoxia, calcium signaling and production of peroxidases which is one of the important reactive oxygen species (ROS) key modulator of plant growth and development, and stress adaptation (Supplementary Table S5). Those findings can explain the presence of cavities in the embryo body of abnormal SE, a probably product of lysigeny via PCD for lack of oxygen in the *in vitro* culture environment, inducing the death of the tissue mediated by ROS³⁰.

The methylome landscape of zygotic and normal and abnormal SE from *T. cacao* was developed for first time with the aim to identify gene regions differentially methylated and methylation profiles among them using WGBS. Levels of DNA methylation are higher in the CpG context than in the CHG and CHH contexts in many plants species³⁴. For cacao, we found a global distribution of methylation in the three contexts between 50.14 - 53.95% (CpG), 28.23 - 31.58% (CHG) and 5.34 - 9.91% (CHH). Non-CpG sequences in ZE, normal and abnormal SE were hypomethylated in high frequencies (Figure 6B-D and Supplementary Figure S8B-D), whereas CpG sequences displayed similar percentages of hypo and hypermethylation (Figure 7E and 8A-C).

In several studies was reported that methylation in CpG, CHG and CHH contexts is associated with activation or inactivation of transposon elements (TEs) and repetitive DNA sequences, as well as in the silencing or enhancing of genes depending where the methyl group is positioned in the gene body or in its surrounding^{35,36}. It has been known that TEs and repetitive DNA sequences are mainly located in heterochromatin where DNA is highly compacted avoiding the transcription of these regions³⁷. TEs are classified in two groups, depending on their transposition mechanism. The class I are retrotransposons, which are transposed by copy-and-paste mechanism via RNA; the class II are DNA transposons, which are transposed by copy-and-paste mechanism via DNA³⁸. Intergenic region of the DNA is heavily associated with closed heterochromatin state, in which its compaction is mainly regulated by the level of methylation in the CpG and CHG contexts, and less in the CHH context³⁹. In our study we found that the intergenic region is the second most differentially methylated region between ZE, normal and abnormal SE in the three contexts and we identified 14 retrotransposons significantly differentially methylated out of 945 spread over the chromosomes 3 to 10 in the CpG context, 1,002 in the CHG context spread over all 10 chromosomes and 1,074 in the CHH context also spread over all 10 chromosomes when methylome of ZE and normal and abnormal SE are compared (Supplementary Table S5 - S14). From all these differentially methylated retrotransposons we identified 275, 122 and 38 hypermethylated for ZE and hypomethylated for normal and abnormal SE in the CpG, CHG and CHH contexts, respectively. In many studies was identified that low levels of methylation in retrotransposons can be associate with their activation⁴⁰, as a result, retrotransposons act as mediator to the stress conditions in the *in vitro* culture of SE, but retrotransposon activation can also induce abnormal phenotypes SE, as it was identified in other works where the activation of TEs is associated with adaptative response to environmental stress and abnormal phenotypes in plants^{38,41,42}.

Methylation in DNA can inhibit or enhance gene expression depending on where the methyl group is placed. Methyl group can be placed in exon or introns in the gene bodies, or in the transcription start sites, which regulated the expression by enhancing or blocking the gene transcription depending where the methyl group position⁴³. Our results demonstrated that introns in ZE, normal and abnormal SE are differentially methylated in more than 40% of the genome comparing with the other regions of the gene, followed by intergenic and exonic regions with more than 20% and 15% respectively in all the three contexts (Supplementary figures 9S-11S). Methylation in exons has been linked to splicing where the mature mRNA is formed by removing introns from the transcribed mRNA precursor⁴⁴. Splicing occurs cotranscriptionally when the transcript is still attached to the DNA and a large fraction of introns are removed from the mRNA precursor by RNA polymerase II⁴⁵. The analysis of DNA methylome in humans showed twice density of methylation in exon when is compared with introns, but the differences were dissolved when the number of CpG sequences between exon and introns was calculated⁴⁶. The high CpG sequences quantity in exons influence in the amount of the total CpG methylation content in the methylome, being exons recognized by the RNA polymerase II and the spliceosome machinery from introns in the splicing⁴⁴. In difference of the studies reported above, in our work we identified high frequencies of differential methylation regions in introns in all of three contexts (CpG, CHG and CHH) when is compared ZE, normal and abnormal SE methylomes, being exons the third gene site more methylated, important finding because those differences have a strong association with the abnormalities in SE and probably influence the abnormal splicing in the abnormal SE phenotypes.

Considerable increase of gene expression is associated to intronic regions, such as the enhancement of ubiquitin C gene expression in humans⁴⁷ and other genes in yeast⁴⁸ where genes with methylation in the 5' splice sites (5'SSs) of intron regions revealed highest expression compared with genes without methylation in the same regions^{49,50}. In plants, introns play an important role in gene regulation as enhancer element^{51,52}. The mechanism of intron-mediated enhancement occurs when introns are located next to the promoter in the transcribed region, thereby increasing the levels of mature mRNA in the cytosol or when the 5'SSs (highly methylated) of introns are recognized by U1 Spliceosomal RNA (small nuclear RNA) in the alternative splicing, causing a 3-fold increase in the transcription products⁵⁰. In mammals, intronic methylation is an important factor in gene expression regulation of cancer, where hypomethylated motifs in the intron region are associated with upregulation of oncogene expression and hypermethylation is associated with downregulation of tumor suppressive expression⁵³. In our work, part of the differential methylation of introns among ZE, normal and abnormal SE (Supplementary tables S5-S14) was probably due to the different phenotypes presented among them. In addition, differences in the methylation profiles of important genes involved in embryo structure development, stress response to the environmental conditions, and the effects of cellular compounds production can be the consequence of abnormalities in SE (Figure 1A-1I). A similar observation was mentioned in a study that was made in non-small cell lung cancer, where methylation profiles were different in two types of cell lung cancer and can be divided in two phenotypically distinct subtypes of tumors using methylation profile as a biomarker platform⁵⁴.

The main differentially methylated genes in this work were those involved in oxidative stress, auxin transport, auxin responsive genes, disease resistance, senescence, hypoxia, and developmental processes. Oxidative stress can start when homeostasis in the cells is disrupted⁵⁵. The principal ROS that are crucial inducer of oxidative stress are mainly singlet oxygen ($^1\text{O}_2^*$), hydroxyl radical ($\cdot\text{OH}$), hydrogen peroxide (H_2O_2), superoxide radical ($\text{O}_2^{\cdot-}$) and nitric oxide (NO)⁵⁶. Hydrogen peroxide is one of the major ROS compounds produced in the cells during abiotic and biotic stress conditions⁵⁷. ROS production can increase and accumulate in the cells, creating an imbalance between ROS production and detoxification, thereby generating abnormal chemical signaling⁵⁶. Another important function of ROS is the direct effect in the biosynthesis, transport, metabolism and signaling of auxin in plants⁵⁸. Genome regions

coding for peroxidase superfamily proteins were identified in our work as differentially methylated, it may be associated with the oxidative stress presented in ZE and normal and abnormal SE in different ways due to abiotic stress induced by the environmental conditions. Hypermethylation in intergenic regions of genes that code for peroxidases in ZE and low levels of methylation in normal and abnormal SE (more in abnormal than normal SE) confirmed that ZE have different environmental conditions than normal and abnormal SEs. In theory, low methylation in intergenic regions is associated with TEs activation when plants are under stress conditions. At the same time, we can infer that abnormal SE presented a particular methylation profile regarding to genes involved in oxidative stress, auxin transport, and developmental processes. Microscopy analysis of the abnormal SE showed abnormal apical-radical axial cell divisions and PCD in the apical shoot and root meristem. Compared to normal SE, this indicates a relationship between ROS and their effect in the wrong auxin polar transport in cells of the abnormal SE.

Oxidative stress induces chemical signals such as abscisic acid, ethylene, brassinosteroids, auxin, gibberellic acid, methyl jasmonate, salicylic acid etc.⁵⁶. The genes that codified the proteins precursor of these compounds were identified in our work as differentially methylated between ZE, normal and abnormal SE. One of the most important is ethylene or its precursor 1-aminocyclopropane-1-carboxylate (ACC) since their production can be associated with the use of 2,4-dichlorophenoxyacetic acid (2,4-D) in the culture medium. Ethylene in high concentration can be accumulated, incrementing respiration rate in the tissue and inducing hypoxia in the culture environment^{59,60}. Ethylene is necessary to stimulate hypoxic induction and it can also regulate aerenchyma formation in root tips of maize plants exposed to hypoxic conditions by schizogeny or lysigeny mediated by PCD and dissolution of cells^{61,62}. ACC the direct precursor of ethylene has a negative effect in *Arabidopsis thaliana*, in both rosette development and hypocotyl growth, and inhibits primary root elongation independently of ethylene perception when it was applied exogenously⁶³. In our research we identified in the methylome of normal and abnormal SE the precursor of ethylene (ACC) differentially methylated downstream of transcription start site of the gene Thecc1EG037684 when is compared with the methylome of ZE, being 0.8 level of methylation for ZE and 0.2 for abnormal SE and 0.5 for normal SE (Supplementary table S6-S7). This leads us to speculate that SE are exposed to hypoxic environments in *in vitro* conditions due to ACC accumulation. Hypoxic environment in the tissue culture can affect some cells in the same tissue generating abnormal development of SE induced by ethylene or ACC.

As hypothesis, the provascular tissue abnormalities in abnormal SE might be associated with and interruption of the auxin transport from the shoot meristem throughout the embryo axis to the root meristem due to the significative differential of methylation profiles between ZE, normal and abnormal SE of genes that encode the *PIN* proteins in the chromosomes 1, 4 and 9. As mentioned above, abnormal SE presented a disorganized growing, without polarity between shoot and root meristems (4E-H), suggesting problems in auxin efflux transportation⁶⁴. The *PIN*-formed proteins are transporters acting in the efflux of signal molecules as auxin in plants⁶⁵. *PIN* proteins are asymmetrically localized within cells and their polarity determines the direction of intercellular auxin efflux⁶⁵. *PIN* proteins are located in the plasma membrane and they have influence in plant development by mediating auxin efflux through cells compartments⁶⁵. When the gene that encodes the *PIN7* protein is mutated in the *A. thaliana* genome, induced anomalies in zygotic embryos because the auxin efflux and polarity in early embryogenesis is disrupted⁶⁴. Ethylene is an important regulator in auxin influx and efflux by basipetal auxin transportation. Exogenous application of ethylene inhibits the auxin transportation by proteins as *PIN1*, *PIN2*, *PIN3* and *PIN4* in consequence root formation is disrupted in *A. thaliana*. Ethylene was also reported as important hormone that downregulates the genes that encodes the lateral auxin movement in *Zea mays* and it downregulates the genes that encodes root tips and epicotyls formation in *Pisum sativum* confirming that there is a crosstalk between ethylene and *auxin efflux carrier proteins* in cell polarity⁶⁶

In conclusion, for the first time the methylomes of torpedo ZE and SE in cacao are reported, revealing a unique profile in abnormal SE accompanying the morphologies presented in those embryos. Although abnormalities in SE prevent embryo germination, we identified seven abnormal phenotypes in a SE production of CCN10 genotype where some of them germinated partially but did not convert in normal plants. The methylome landscape of ZE and normal and abnormal SEs are different showing a specific methylation profile for each one. Different methylation levels were found in the three contexts (CpG, CHG and CHH) in ZE, normal and abnormal SE in cacao where CpG and CHG had similar bimodal distribution of hypo and hypermethylated regions with similar frequencies of hypo and hypermethylation in the CpG context and higher frequencies of hypomethylation than hypermethylation in the CHG context. In contrast, the CHH context had the highest frequencies of hypomethylation in a right-squeezed distribution.

The main differentially methylated regions in the three contexts were intronic and intergenic regions. Methylation in intronic regions the highest region methylated in this study is in theory associated with gene expression enhancing, and probably in the wrong identification of introns in splicing process to form the mRNA, important finding in our work that might explain the abnormal phenotypes in SE. Intergenic regions were the second highest differentially methylated regions when ZE are compared with normal and abnormal SE, being less methylated in the abnormal ones. Intergenic regions are heavily associated with heterochromatin remodeling where generally TEs placed. The TEs are implicated in the regulation of stress in the tissue culture environment. Stress environment in the tissue culture was revealed in this study because genes related to oxidative stress were found differentially methylated between ZE, normal and abnormal SE. Microscopy analysis of Abnormal SE revealed cavity formation in their axes. In addition, cotyledons, shoots, and roots in abnormal SE had the highest area of fluorescence values due to PCD. Cavity formation in abnormal SE can be associated with lysigeny via vacuolar cell death probably as a response of the cell to the lack of oxygen in tissue culture environment. Hypoxia induction in abnormal SE can be associated with genes that codes for ethylene and its precursor ACC, which were differentially methylated in this work when compared with ZE. Ethylene production in *in vitro* environment reduces oxygen concentration and helps stimulating hypoxia induction. Differentiation and cellular auxin transportation of abnormal SE could also be influenced by ethylene in the cell environment due to the *in vitro* culture condition. Ethylene production in tissue culture could be mediated by 2,4-D (the main synthetic auxin used in the SE culture in this work) that in high concentration is accumulated in somatic tissue affecting embryos development by ethylene effect. An in-depth study of the gene expression of a core selection of the 873 genes found differentially methylated in this work would be required to confirm our results.

Methods

Plant material

Zygotic embryos

Zygotic embryos at the torpedo stage from cacao pods were obtained by controlled hand pollination of CCN 10 (CCN-*Colección Castro Naranja*) with CCN 51 as pollen donor in the field at the Mars Center for Cocoa Science (MCCS) germplasm collection. Cacao pods were harvested 13 weeks after pollination (before maturation), which corresponds to the later torpedo stage when embryos maturation has not started yet as previously described¹⁰. A total of 10 g of ZE were extracted from each pod, frozen in liquid nitrogen and stored at - 80°C for DNA and RNA extraction.

Somatic embryos

Primary SE generated from flowers parts (2,000 petals) and secondary SE generated from primary SE cotyledons (60 primary SE) of CCN 10 were obtained using the Li et al. (1998) and Maximova et al. (2002) protocols respectively with some modifications previously described in⁷. The somatic embryos in late torpedo stage as formerly described¹¹

were classified in normal and abnormal SE depending on different morphologies presented. Ten grams of secondary normal and 10 grams of abnormal SE were frozen in liquid nitrogen and stored at - 80°C for DNA and RNA extraction.

Morphological characterization of somatic embryos

Morphological evaluation was done with a representative sample of 118 primary SE (produced from 1,000 petals as explants) and 530 secondary SE (produced from cotyledons of 20 primary SE as explants) of CCN 10 in late torpedo stage and using as a control zygotic embryos in the same stage of development and following the references ratings previously described^{10,67}. The analysis was performed in a digital microscope (model AM7013MZT, Dino-Lite premier brand). The percentage of normal and abnormal SE was calculated counting the number of embryos (normal and abnormal) into the total of embryos present in each sample and multiplying by hundred. In addition, the percentage of conversion of normal and abnormal secondary SE to plantlets was also calculated to assess the viability of embryos and the capacity to generate normal plantlets. Embryo maturation and conversion were performed using the protocol previously described⁷. Embryos were considered as converted when the shoot apex developed and formed radical extension.

Microscopic analysis

Structure analysis of intercellular spaces in somatic embryos by Scanning Electron Microscope (SEM)

Samples of secondary SE (normal and abnormal) in late torpedo stage (three biological replicates) of CCN 10 were collected and fixed in 2.5% glutaraldehyde, in a sodium cacodylate buffer at 0.1 M, pH 6.8, during 12 h. Subsequently, embryos were washed in the same buffer and were dehydrated in an increasing acetone series (50, 60, 70, 80, 90 and 100% each one for 10 min). Immediately after dehydration, the material was taken to a critical point dryer (model CPD 030, BAL-TEC brand), which allowed the full elimination of water in the plant tissue. Afterward, the samples were placed in a metal stub with double-sided tape on it for subsequent metallization in a sputter COATER, (model SCD 050, BAL-TEC brand), followed by subsequent observation in *SEM* (FEI Company, model Quanta 250, Eindhoven, Netherlands)⁶⁸. The pictures were captured with CCD and Nav-CamTM cameras and processed by the Scandium Image software (SIS).

Ultrastructural analysis of zygotic and somatic embryos by Transmission Electron Microscope (TEM)

Samples of ZE and secondary SE (normal and abnormal) in late torpedo stage (three biological replicates) of CCN 10 were collected and fixed in 3% glutaraldehyde, in a sodium cacodylate buffer at 0.1 M, pH 6.8, during 12 h. The samples were subjected to a series of washes (3 times for 15 min) in a sodium cacodylate buffer at 0.1 M, pH 7.2, and post-fixed in 1% osmium tetroxide, prepared in the same buffer, during 1 h at 4°C. Next, the samples were washed 3 times in the same sodium cacodylate buffer for 15 min each one and were then dehydrated in an increasing ethanol series (30, 50, 70, 80, and 90% each one for 15 min), followed by two washes in 100% ethanol (each one for 30 min). Shortly after, the samples were soaked in a mixture of 100% ethanol and LR White resin in the proportions of 3:1(2 h), 1:1 (2 h), 1:3 (overnight), followed by two changes of pure LR White resin every 4 h, always under slow agitation. Afterwards, the samples were placed in gelatin capsules and covered with pure LR White resin. Resin polymerization was completed after 24 h at 60°C. The ultrathin sections (70 nm) were made with a diamond knife, using a Leica ultramicrotome (model UC6, Nussloch, Germany). The cut sections were deposited on copper grids (300 mesh), contrasted with uranyl acetate in aqueous solution for 25 min, and then with lead citrate for 30 min. Subsequently, they were observed in *TEM* MORGAGNI and wide angle Olympus MegaView III camera and the iTEM software (FEI Company, model 268 D, Eindhoven, Netherlands)⁶⁹.

Cell death detection and histological analysis

Cell death was detected using the “*In situ*” cell death detection fluorescein kit provided by ROCHE. Samples of normal SE and abnormal SE in late torpedo stage of CCN 10 were collected and fixed in FAA solution (Combine 10 parts 37-40% formaldehyde, 70 parts 95% ethanol, 15 parts dH₂O, and 5 parts acetic acid) during 12 h. The samples were submerged in paraffin and sectioned in slices of 5µm using Leica ultramicrotome (model UC6, Nussloch, Germany). The tissue sections were pretreated with proteinase K diluted in Tris-HCl 10 mM (20 mg mL⁻¹) pH 7.4 during 20 min. After that we followed the manufacturer recommendations. Tissue sections were analyzed in LEICA fluorescence microscope model DM2500 identifying red fluorescence with the Leica Application Suit (LAS X) software (Cells-stained fluorescent red are positive for cell death). Pictures were captured with the QImaging Retiga 4000R monochrome camera, **using the Texas Red fluorescent filter** (Leica # TX2). Same tissue sections were treated with 1% toluidine in a solution of 0.1 M sodium phosphate and pH 6.8 for histological analysis. Area and intensity of fluorescence were calculated using the ImageJ program (<https://imagej.nih.gov/ij/download.html>), and the statistical analysis was performed with the software R 4.0.2.

DNA extraction

Genomic DNA from a population of 200 normal SE, 200 abnormal SE and 200 ZE at the late torpedo stage was extracted using the DNeasy extraction kit Plant Mini Kit (250) from QIAGEN for future methylation analysis. A tissue sample of 100 mg was disrupted in liquid N₂ to a fine powder using a mortar and pestle, transferred to a 2 µL Eppendorf microtube and DNA immediately extracted, following the kit protocol (DNeasy® Plant Handbook, QIAGEN). The extraction procedure was repeated two times to ensure the DNA quality and quantity. After the second extraction, DNA was precipitated with 1:1 volume of isopropanol and 2 µL of glycogen from a stock of 20 mg mL⁻¹ for 20 min at -20°C. The samples were centrifuged in a bench-top centrifuge (Mikro 220R-V1.20, Hettich Zentrifugen, Germany) at 8000 x g for 30 min at 20°C. The supernatant was discarded, the resulting pellet was left to dry for 30 min in environmental conditions. Afterward, the pellet was re-suspended in 50 µL of 100% ethanol. The DNA quality was checked on a 1% agarose gel in 1X TBE (2.5 g L⁻¹ boric acid, 10.8 g L⁻¹ Tris base, 4 mL of 0.5M EDTA, pH 8.0) using a molecular marker (Hi DNA mass Ladder, Invitrogen brand) as reference. DNA quantification was performed using a spectrophotometer Multiskan GO, Thermo Scientific brand by absorbance at a wavelength of 260/230 and 260/280 nm.

Global DNA methylation by whole-genome bisulfite sequencing

The Whole-Genome Bisulfite Sequencing was performed on the total DNA extracted from two replicates of a set of mixed embryos from each of the three representative samples (normal SE, abnormal SE and ZE), using the protocol to prepare samples for Illumina sequencing platform (WBS for methylation analysis preparing samples protocol (KK8201 & 59104)). For a few minutes, genomic DNA (5 µg) was normalized with unmethylated lambda (λ) DNA 5% (5ng λ DNA per microgram of genomic DNA, Promega). DNA was fragmented by sonication using a Covaris S220/E220 sonicator, and fragment selection (150–300 bp) was performed with AMPure XP beads (Agencourt Bioscience). Genomic DNA libraries were constructed using the Illumina TruSeq Sample Preparation kit, considering the Illumina's standard protocol. Repair was performed at the ends of the DNA; at the 3' end of each fragment an adenine was added, and the Illumina TruSeq adapters were also ligated to both ends. After ligation of the adapters, DNA was treated with sodium bisulfite using the EpiTect Bisulfite kit (QIAGEN). The bisulfite conversion was performed two times to ensure a conversion rate over 99%. Seven PCR cycles were carried out to ensure the enrichment of the adaptor-ligated DNA using PfuTurboCx Hot-Start DNA polymerase (Stratagene). Library quality and concentration were checked using the

Agilent 2100 Bioanalyzer. Paired-end DNA sequencing (2 × 100 bp) with 30X coverage was then performed using the Illumina HiSeq 2000 platform.

Read mapping and estimation of cytosine methylation levels in CpG, CHH and CHG contexts

Before mapping and estimation of cytosine methylation levels, the raw sequence reads were filtered to make the quality check using the FASTQC bioinformatics tool with Phred quality score over 20 at each cycle. Also the adapter sequences were trimmed off the raw sequence reads using Trimmomatic program⁷⁰. Read mapping was carried out using the BSMAP which is based on the SOAP (Short Oligo Alignment Program) against to the Cacao_Matina1-6 (Tcacao_genome_v1.1) genome as a reference. The only uniquely mapped reads were selected to sort; index and PCR duplicates were removed with SAMBAMBA (v0.5.9). The evaluation of the quality of the alignment data (a BAM file) and the basic statistics of the alignment (ACGT Content, mean and standard coverage by chromosome, insert distribution etc.) was performed with Qualimap 2.2. The methylation ratio of every single cytosine location was called from the mapping results using 'methylation.py' script in BSMAP. The results of the coverage profiles were calculated as the number # of C/effective CT counts for each cytosine in CpG, CHH and CHG contexts⁷¹.

Differential of methylation in CpG, CHH and CHG contexts

DNA methylation ratio values of single cytosine obtained from BSMAP program were used as the original raw data from the three samples (normal SE, abnormal SE and ZE). The methylation ratio values of all samples were normalized using median scaling normalization to compare the methylation level between samples. During data preprocessing, low coverage and high coverage bases were filtered. Lower read cut-off of 10 means that bases with coverage below 10X were discarded because a high enough read coverage will increase the power of the statistical tests. Higher read cut-off of 500 means that bases with more than 500X in each sample were also discarded.

Statistical analysis was performed using the difference of two groups (delta_mean) per comparison pair. The significant results were selected on conditions of $|\text{delta_mean}| \geq 0.2$ with p-value of 0.05. Level of methylation was displayed as hypermethylation when the delta mean ≥ 0.2 ; and hypomethylation when the delta mean ≤ -0.2 . For significant CpG list, hierarchical clustering analysis was performed to group the similar samples grouped in CpGs, CHGs and CHHs. A cluster analysis was performed considering the correlations between the samples and grouping with Ward's method⁷². Differential methylation was obtained from the regions significantly different using Logistic regression (P < 0.001). The analyzes were performed with the aid of packages methylKit, complexHeatmap, circlize and annotate using software R 4.0.2.

Genomic and functional annotations of CpG, CHH and CHG sites

Each cytosine locus in CpG, CHH and CHG contexts was annotated using the Table Browser tool of the University of California Santa Cruz (UCSC) Genome Browser. Annotation included functional location of each gene (promoter regions which are defined as -2kb upstream of the transcription start site, exons and introns, intergenic, and UTRs regions), transcripts identification (ID), gene ID, strand, and sequence context⁷³. The gene function was identified by making the alignments with expectation values (E-value) < 0.001 with the aim to infer gene homology comparing with others genomes such as *A. thaliana* using the cacao genome database in the Phytozome V13 (https://phytozome-next.jgi.doe.gov/info/Tcacao_v2_1) and the translated-DNA: protein alignments by BLASTX similarity search tool in the National Center for Biotechnology Information (NCBI) databases. The protein metabolic function was identified using the collection databases from Kyoto Encyclopedia of Genes and Genomes (KEGG).

This research was carried out in Brazil and the plant material used in it was donated from the germplasm collection of the Mars Center for Cacao Science (MCCS), which complies with the relevant national guidelines and the legislation of the Conselho de Gestão do Patrimônio Genético - CGEN permit for access to genetic resources number 02000.000070/2015-05.

Declarations

Acknowledgments

We thank the Microscopy Center of the State University of Santa Cruz due to the availability of facilities where the analysis was carried out.

Author contributions statement

A-A.F.A., D.B., M.C., J-P. M. and S.R. supported, oriented the project research, and advised all its execution. L.S. executed all the tissue cultures to obtain the SE and the extraction of ZE used as samples. J. S. executed the samples preparation for microscopy analysis. P.M. processed the ultramicrotome cuts and analyzed the samples in the TEM. F.C. processed and analyzed WGBS data. C.G. executed all the steps of the research with help of the authors mentioned above and wrote the manuscript.

Competing financial interests

The authors declare no competing financial interests.

References

1. Franzen, M. & Borgerhoff Mulder, M. Ecological, economic and social perspectives on cocoa production worldwide. *Biodivers. Conserv.* **16**, 3835–3849 (2007).
2. Almeida, A.-A. F. de & Valle, R. R. Ecophysiology of the cacao tree. *Braz. J. Plant Physiol.* **19**, 425–448 (2007).
3. Scholey, A. & Owen, L. Effects of chocolate on cognitive function and mood: a systematic review. *Nutr. Rev.* **71**, 665–681 (2013).
4. Pucciarelli, D. Cocoa and Heart Health: A Historical Review of the Science. *Nutrients* **5**, 3854–3870 (2013).
5. Marinato, C. S., Galeas Aguilar, M. A., Spagiari Souza, C. A. & Valim Olmo, F. J. Enxertia precoce em cacau (*Theobroma cacao* L.). *Enciclopédia Biosf. Cent. Científico Conhecer* **9**, 1285–1292 (2013).
6. Maximova, S. N. *et al.* Efficiency, genotypic variability, and cellular origin of primary and secondary somatic embryogenesis of *Theobroma cacao* L. *Vitro Cell. Dev. Biol. - Plant* **38**, 252–259 (2002).
7. Garcia, C. *et al.* Optimization of somatic embryogenesis procedure for commercial clones of *Theobroma cacao* L. *Afr. J. Biotechnol.* **15**, 1936–1951 (2016).
8. Fan, Y.-F., Jiang, L., Gong, H.-Q. & Liu, C.-M. Sexual Reproduction in Higher Plants I: Fertilization and the Initiation of Zygotic Program. *J. Integr. Plant Biol.* **50**, 860–867 (2008).
9. *In Vitro Embryogenesis in Higher Plants*. vol. 1359 (Springer New York, 2016).
10. Alemanno, L., Berthouly, M. & Michaux-Ferriere, N. A comparison between *Theobroma cacao* L. zygotic embryogenesis and somatic embryogenesis from floral explants. *Vitro Cell. Dev. Biol.-Plant* **33**, 163–172 (1997).
11. Maximova, S. N. *et al.* Genome-wide analysis reveals divergent patterns of gene expression during zygotic and somatic embryo maturation of *Theobroma cacao* L., the chocolate tree. *BMC Plant Biol.* **14**, 185 (2014).

12. Zavattieri, M. A., Frederico, A. M., Lima, M., Sabino, R. & Arnholdt-Schmitt, B. Induction of somatic embryogenesis as an example of stress-related plant reactions. *Electron. J. Biotechnol.* **13**, (2010).
13. Pila Quinga, L. A., Pacheco de Freitas Fraga, H., do Nascimento Vieira, L. & Guerra, M. P. Epigenetics of long-term somatic embryogenesis in *Theobroma cacao* L.: DNA methylation and recovery of embryogenic potential. *Plant Cell Tissue Organ Cult. PCTOC* (2017) doi:10.1007/s11240-017-1284-6.
14. Miguel, C. & Marum, L. An epigenetic view of plant cells cultured in vitro: somaclonal variation and beyond. *J. Exp. Bot.* **62**, 3713–3725 (2011).
15. Zhang, H., Lang, Z. & Zhu, J.-K. Dynamics and function of DNA methylation in plants. *Nat. Rev. Mol. Cell Biol.* **19**, 489–506 (2018).
16. Santos, D. & Fevereiro, P. Loss of DNA methylation affects somatic embryogenesis in *Medicago truncatula*. *Plant Cell Tissue Organ Cult.* **70**, 155–161 (2002).
17. Peng, H. & Zhang, J. Plant genomic DNA methylation in response to stresses: Potential applications and challenges in plant breeding. *Prog. Nat. Sci.* **19**, 1037–1045 (2009).
18. Brenet, F. *et al.* DNA Methylation of the First Exon Is Tightly Linked to Transcriptional Silencing. *PLOS ONE* **6**, e14524 (2011).
19. Unoki, M. & Nakamura, Y. Methylation at CpG islands in intron 1 of EGR2 confers enhancer-like activity. *FEBS Lett.* **554**, 67–72 (2003).
20. Gaj, M. Factors influencing somatic embryogenesis induction and plant regeneration with particular reference to *Arabidopsis thaliana* (L.) Heynh. *Plant Growth Regul.* **43**, 27–47 (2004).
21. Ardebili, S. H. *et al.* Effect of 2, 4-D as a Novel Inducer of Embryogenesis in Microspores of *Brassica napus* L. *Czech J Genet Plant Breed* **47**, 114–122 (2011).
22. Fehér, A., Pasternak, T. & Dudits, D. Transition of somatic plant cells to an embryogenic state. *Plant Cell Tissue Organ Cult.* **74**, 201–228 (2003).
23. Garcia, C. *et al.* Abnormalities in somatic embryogenesis caused by 2,4-D: an overview. *Plant Cell Tissue Organ Cult. PCTOC* 1–20 (2019) doi:10.1007/s11240-019-01569-8.
24. Gamborg, O. L. & Larue, T. A. G. Ethylene Production by Plant Cell Cultures: The Effect of Auxins, Absciscic Acid, and Kinetin on Ethylene Production in Suspension Cultures of Rose and Ruta Cells. *PLANT Physiol.* **48**, 399–401 (1971).
25. Garcia, F. & Einset, J. W. Ethylene and ethane production in 2, 4-D treated and salt treated tobacco tissue cultures. *Ann. Bot.* **51**, 287–295 (1983).
26. Wang, K. L.-C., Li, H. & Ecker, J. R. Ethylene Biosynthesis and Signaling Networks. *Plant Cell* **14**, S131–S151 (2002).
27. Yang, C.-Y., Hsu, F.-C., Li, J.-P., Wang, N.-N. & Shih, M.-C. The AP2/ERF Transcription Factor AtERF73/HRE1 Modulates Ethylene Responses during Hypoxia in *Arabidopsis*. *PLANT Physiol.* **156**, 202–212 (2011).
28. Vaca, C. I. C. Inducción de la embriogénesis somática en clones superiores de cacao (*Theobroma cacao* L.), con resistencia a enfermedades fungosas. (2004).
29. De Rybel, B., Mähönen, A. P., Helariutta, Y. & Weijers, D. Plant vascular development: from early specification to differentiation. *Nat. Rev. Mol. Cell Biol.* **17**, 30–40 (2016).
30. Takahashi, H., Yamauchi, T., Colmer, T. D. & Nakazono, M. Aerenchyma Formation in Plants. in *Low-Oxygen Stress in Plants* (eds. van Dongen, J. T. & Licausi, F.) vol. 21 247–265 (Springer Vienna, 2014).
31. Gadjev, I., Stone, J. M. & Gechev, T. S. Chapter 3: Programmed Cell Death in Plants. in *International Review of Cell and Molecular Biology* vol. 270 87–144 (Elsevier, 2008).

32. M-Hamvas, M. Cylindrospermopsin induces biochemical changes leading to programmed cell death in plants. *11*.
33. Van Doorn, W. G. *et al.* Morphological classification of plant cell deaths. *Cell Death Differ.* **18**, 1241–1246 (2011).
34. Bartels, A. *et al.* Dynamic DNA Methylation in Plant Growth and Development. *Int. J. Mol. Sci.* **19**, 2144 (2018).
35. Bewick, A. J. & Schmitz, R. J. Gene body DNA methylation in plants. *Curr. Opin. Plant Biol.* **36**, 103–110 (2017).
36. Yan, X. *et al.* DNA methylation signature of intergenic region involves in nucleosome remodeler DDM1-mediated repression of aberrant gene transcriptional read-through. *J. Genet. Genomics* **43**, 513–523 (2016).
37. Volpe, T. & Martienssen, R. A. RNA Interference and Heterochromatin Assembly. *Cold Spring Harb. Perspect. Biol.* **3**, a003731 (2011).
38. Song, X. & Cao, X. Transposon-mediated epigenetic regulation contributes to phenotypic diversity and environmental adaptation in rice. *Curr. Opin. Plant Biol.* **36**, 111–118 (2017).
39. Gent, J. I. *et al.* CHH islands: de novo DNA methylation in near-gene chromatin regulation in maize. *Genome Res.* **23**, 628–637 (2013).
40. Lippman, Z. *et al.* Role of transposable elements in heterochromatin and epigenetic control. *Nature* **430**, 471–476 (2004).
41. Galindo-González, L., Sarmiento, F. & Quimbaya, M. A. Shaping Plant Adaptability, Genome Structure and Gene Expression through Transposable Element Epigenetic Control: Focus on Methylation. *Agronomy* **8**, 180 (2018).
42. Corem, S. *et al.* Redistribution of CHH Methylation and Small Interfering RNAs across the Genome of Tomato ddm1 Mutants. *Plant Cell* **30**, 1628–1644 (2018).
43. Jones, P. A. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat. Rev. Genet.* **13**, 484–492 (2012).
44. Gelfman, S., Cohen, N., Yearim, A. & Ast, G. DNA-methylation effect on cotranscriptional splicing is dependent on GC architecture of the exon–intron structure. *Genome Res.* **23**, 789–799 (2013).
45. Luco, R. F., Allo, M., Schor, I. E., Kornblihtt, A. R. & Misteli, T. Epigenetics in Alternative Pre-mRNA Splicing. *Cell* **144**, 16–26 (2011).
46. Lister, R. *et al.* Human DNA methylomes at base resolution show widespread epigenomic differences. *Nature* **462**, 315–322 (2009).
47. Bianchi, M., Crinelli, R., Giacomini, E., Carloni, E. & Magnani, M. A potent enhancer element in the 5'-UTR intron is crucial for transcriptional regulation of the human ubiquitin C gene. *Gene* **448**, 88–101 (2009).
48. Juneau, K., Miranda, M., Hillenmeyer, M. E., Nislow, C. & Davis, R. W. Introns Regulate RNA and Protein Abundance in Yeast. *Genetics* **174**, 511–518 (2006).
49. Laxa, M. Intron-Mediated Enhancement: A Tool for Heterologous Gene Expression in Plants. *Front. Plant Sci.* **7**, (2017).
50. Shaul, O. How introns enhance gene expression. *Int. J. Biochem. Cell Biol.* **91**, 145–155 (2017).
51. Callis, J., Fromm, M. & Walbot, V. Introns increase gene expression in cultured maize cells. *Genes Dev.* **1**, 1183–1200 (1987).
52. Rose, A. B. Requirements for intron-mediated enhancement of gene expression in Arabidopsis. *RNA* **8**, 1444–1453 (2002).
53. Nam, A.-R. *et al.* Alternative methylation of intron motifs is associated with cancer-related gene expression in both canine mammary tumor and human breast cancer. *Clin. Epigenetics* **12**, 110 (2020).
54. Walter, K. *et al.* DNA Methylation Profiling Defines Clinically Relevant Biological Subsets of Non-Small Cell Lung Cancer. *Clin. Cancer Res.* **18**, 2360–2373 (2012).

55. Grene, R. Oxidative Stress and Acclimation Mechanisms in Plants. *Arab. Book* **1**, e0036 (2002).
56. Demidchik, V. Mechanisms of oxidative stress in plants: From classical chemistry to cell biology. *Environ. Exp. Bot.* **109**, 212–228 (2015).
57. Krishnamurthy, A. & Rathinasabapathi, B. Oxidative stress tolerance in plants: Novel interplay between auxin and reactive oxygen species signaling. *Plant Signal. Behav.* **8**, e25761 (2013).
58. Tognetti, V. B., Mühlhennock, P. & Van Breusegem, F. Stress homeostasis - the redox and auxin perspective: Stress homeostasis. *Plant Cell Environ.* **35**, 321–333 (2012).
59. Arteca, R. N. Influence of IAA, NAA and 2, 4-D on ethylene production by potato discs (*Solanum tuberosum* L. cv. Red Pontiac). *Am. J. Potato Res.* **59**, 267–274 (1982).
60. Vendrell, M. Dual effect of 2, 4-D on ethylene production and ripening of tomato fruit tissue. *Physiol. Plant.* **64**, 559–563 (1985).
61. Gunawardena, A. H., Pearce, D. M., Jackson, M. B., Hawes, C. R. & Evans, D. E. Characterisation of programmed cell death during aerenchyma formation induced by ethylene or hypoxia in roots of maize (*Zea mays* L.). *Planta* **212**, 205–214 (2001).
62. Drew, M. C., He, C.-J. & Morgan, P. W. Programmed cell death and aerenchyma formation in roots. *Trends Plant Sci.* **5**, 123–127 (2000).
63. Vanderstraeten, L., Depaepe, T., Bertrand, S. & Van Der Straeten, D. The Ethylene Precursor ACC Affects Early Vegetative Development Independently of Ethylene Signaling. *Front. Plant Sci.* **10**, (2019).
64. Friml, J., Vieten, A., Sauer, M., Weijers, D., & others. Efflux-dependent auxin gradients establish the apical-basal axis of *Arabidopsis*. *Nature* **426**, 147 (2003).
65. Křeček, P. *et al.* The PIN-FORMED (PIN) protein family of auxin transporters. *Genome Biol.* **10**, 249 (2009).
66. Ruzicka, K. *et al.* Ethylene Regulates Root Growth through Effects on Auxin Biosynthesis and Transport-Dependent Auxin Distribution. *PLANT CELL ONLINE* **19**, 2197–2212 (2007).
67. Solano-Sánchez, W. Embriogenesis somatica en clones superiores de cacao (*Theobroma cacao* L.) obtenidos en el programa de mejoramiento genético del CATIE. (2008).
68. Fischer, E. R., Hansen, B. T., Nair, V., Hoyt, F. H. & Dorward, D. W. Scanning Electron Microscopy. in *Current Protocols in Microbiology* (eds. Coico, R., Kowalik, T., Quarles, J., Stevenson, B. & Taylor, R.) (John Wiley & Sons, Inc., 2012).
69. Williams, D. B. & Carter, C. B. *Transmission electron microscopy: a textbook for materials science*. (Springer, 2008).
70. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).
71. Xi, Y. & Li, W. BSMAP: whole genome bisulfite sequence MAPping program. *BMC Bioinformatics* **10**, 232 (2009).
72. Murtagh, F. & Legendre, P. Ward's Hierarchical Agglomerative Clustering Method: Which Algorithms Implement Ward's Criterion. *J. Classif.* **31**, 274–295 (2014).
73. Karolchik, D., Hinrichs, A. S. & Kent, W. J. The UCSC Genome Browser. in *Current Protocols in Bioinformatics* (eds. Baxevanis, A. D., Petsko, G. A., Stein, L. D. & Stormo, G. D.) (John Wiley & Sons, Inc., 2009).

Figures

Figure 1

Zygotic and somatic embryos comparison. (A) Abnormal type 1 (AT1) has fused more than two embryo axes, (B) AT2 are embryos without root and cotyledons formation, (C) AT3 are translucent embryos with curved axis, a single cotyledon and without apical shoot formation, (D) AT4 are translucent embryos with more than two cotyledons, axis without root and shoot apex formation, (E) AT5 are embryos with long and slender axis and poor development of cotyledons without root and shoot formation, (F) AT6 are translucent embryos with a single amorphous cotyledons and amorphous axis, (G) AT7 are embryos with long and slender cotyledons and curved and amorphous axis, (H) normal somatic embryo and (I) zygotic embryo in torpedo stage. Scale bar, 2 mm (Zygotic and abnormal somatic embryos); 1 mm (normal somatic embryos).

Figure 2

Morphological comparison between normal and abnormal somatic embryos. Scanning electron microscopy of AT4 SE transverse (A-B) and longitudinal cuts with detail, where it is observed cavities formation along of their body in these kinds of abnormal SE (C-D). Scanning electron microscopy of normal SE transverse (E-F) and longitudinal cuts with detail, in difference, normal SE do not have cavities formation in their bodies (G-H). Scale bar in the Figures A, E and D 1mm; C and G 2mm; B 200 μm; F 400μm; H 300μm.

Figure 3

Electron micrographs of zygotic and somatic embryo cells. (A) Zygotic, (B) normal somatic and (C) abnormal somatic cell. Different from zygotic and normal somatic cell, some cells from abnormal SE had a higher number of vesicles, large lytic vacuoles and the nonexistence of the nucleus. n, nucleus; lv, lytic vacuole; v, vesicles. Scale bar, 2 μm (zygotic and abnormal somatic cells (A and C)); 5 μm (normal somatic cell (B)).

Figure 4

Histological and fluorescence micrographs of normal and abnormal somatic embryos. (A) Histological analysis of apical meristem of normal SE. (B) Cell death analysis in apical meristem of normal SE, where there is no presence of fluorescence stained in meristematic cells, but superficial cell layers in cotyledons show fluorescence. It is typically characteristic of positive reaction to the kit "*In situ*" cell death detection fluorescein, product of cell death. (C) Histological analysis of apical meristem of abnormal SE. (D) Cell death analysis in cotyledon and apical meristem of abnormal SE, the nucleus in meristematic and cotyledon's regions are highly fluorescence stained. (E) Histological analysis of root meristem of normal SE. (F) Cell death analysis in root meristem of normal SE, where there are no cells-stained fluorescence. (G) Histological analysis of root meristem of abnormal SE. (H) Cell death analysis in root meristem of abnormal SE shows red fluorescence typically characteristic when the DNA in the cell is fragmented. Scale bar, 400 μm (histological micrographs); 100 μm (fluorescence micrographs). DNA strand breaks in cells with cell death

are stained bright red color. Arrows in blue color are showing apoptotic nuclei inside the cells-stained fluorescence, product of cell death.

Figure 5

Cell death detection through fluorescence analysis using the TUNEL reaction in normal and abnormal SE. Fluorescence area calculated in cells in normal and abnormal SE with cell death, where abnormal SE showed the high values in cotyledons and root meristem cells compared with normal SE. Error bars are 95% confidence interval, and outliers are shown as transparent dots.

Figure 6

Methylation distribution analysis in the CHG context of ZE, normal and abnormal SE by WGBS. (A) Hierarchical clustering of the samples by Euclidian distances, where normal and abnormal SE were grouped in the same cluster while ZE was grouped separately showing the similarities between samples in the methylation profile. (B) Frequency bar chart of methylation levels between ZE and normal SE where the highest frequencies were in regions with low levels of methylation for both samples. (C) Same analysis was done for ZE and abnormal SE. (D) Same analysis was done for normal and abnormal SE, showing the same behavior. (E) Frequency histogram representing the functional regions differentially methylated between ZE and normal SE, (F) ZE and abnormal SE and (G) normal and abnormal SE, where intronic and intergenic regions have the highest frequencies. Color code blue (ZE) and red (Normal SE) in B; blue (ZE) and red (abnormal SE) in C; blue (normal SE) and red (abnormal SE) in D. Codes in ZE (MCCS_TC4), Normal SE (MCCS_TC5) and Abnormal SE (MCCS_TC6).

Figure 7

Methylation distribution analysis in the CpG context of the ZE, normal and abnormal SE by WGBS. (A) Histogram of CpG coverage for ZE, normal and abnormal SE showing normal distribution for all the samples. (B) Histogram of frequencies of CpG percentages in ZE, normal and abnormal SE showing bimodal distribution in all the samples. (C) Plot of percentages of significative regions with hypo and hypermethylation in the ten chromosomes comparing the DNA methylomes between ZE, normal and abnormal SE. (D) Hierarchical clustering of the samples by ward method, where normal and abnormal SE were grouped in the same cluster while ZE is grouped separately. (E) Bar chart of methylation levels proportion between ZE and normal and abnormal SE showing similar proportion for hypomethylation and hypermethylation in whole epigenome. Codes T4 (ZE), T5 (Normal SE) and T6 (Abnormal SE) in E.

Figure 8

Methylation distribution analysis in the CpG context of ZE, normal and abnormal SEs by WGBS. (A) Frequency bar chart of methylation levels between ZE and normal SE shows bimodal distribution for both samples. (B) Same analysis was done for ZE and abnormal SE (C) and between normal and abnormal SE showing the same bimodal distribution. (D) Frequency histogram representing the functional regions differentially methylated between ZE and normal SE, (E) between ZE and abnormal SE and (F) between normal and abnormal SE where intronic, intergenic and exonic regions have the highest frequencies. Color code blue (ZE) and red (Normal SE) in A; blue (ZE) and red (abnormal SE) in B; blue (normal SE) and red (abnormal SE) in C. Codes ZE (MCCS_TC4), Normal SE (MCCS_TC5) and Abnormal SE (MCCS_TC6).

Figure 9

Genes differentially methylated related with SE abnormalities. (A) Circular heatmap showing differentially methylated regions in ZE, normal and abnormal SE (ASE = Abnormal SE and NSE = normal SE) across of ten chromosomes (873 genes). (B) Hierarchical heatmap grouping the differentially methylated genes related with similarities between ZE, normal and abnormal SE where 4 groups were formed. For ZE and abnormal SE there were more genes with similar methylation levels (TC4 = ZE and TC6 = Abnormal SE), normal SE present high differences when is compare with ZE (TC5 = normal SE). (C) Gene differentially methylated distribution across of ten chromosomes with important functions in stress response, defense response, auxin transport, cell polarity, PCD, cell wall biogenesis, and embryo shoot and root development. Gene regions differentially methylated between ZE (D), normal SE (E) and abnormal SE (F) with important functions in stress response, defense response, auxin transport, cell polarity, PCD, cell wall biogenesis, and embryo shoot and root development. Color code blue (hypomethylation), white (unmethylated) and red (hypermethylation) in A and B. Protein, chromosome position and function of genes with code (gene ID) are presented in the table 2.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFigures1.docx](#)
- [Supplementarytable5CGsigMethylation.xltx](#)
- [Supplementarytable6CG45.csv](#)
- [Supplementarytable7CG46.csv](#)

- [Supplementarytable8CG56.csv](#)
- [Supplementarytable9CHG45.csv](#)
- [Supplementarytable10CHG46.csv](#)
- [Supplementarytable11CHG56.csv](#)
- [Supplementarytable12CHH45.csv](#)
- [Supplementarytable13CHH46.csv](#)
- [Supplementarytable14CHH56.csv](#)