

Association Between Cigarette Smoking and Ovarian Reserve Among Women Seeking Fertility Care

Islamiat Oladipupo (✉ miyaah82@yahoo.com)

Department of Epidemiology and Population Health, University of Louisville, School of Public Health and Information Sciences <https://orcid.org/0000-0001-8783-403X>

T'shura Ali

University of Louisville School of Public Health and Information Sciences

David W Hein

University of Louisville School of Medicine

Kelly Pagidas

University of Louisville Department of Medicine

Henry Bohler

University of Louisville School of Medicine

Mark A Doll

University of Louisville School of Medicine

Merry Lynn Mann

University of Louisville School of Medicine

Adrienne Gentry

University of Louisville School of Medicine

Chiang L Jasmine

University of Louisville School of Medicine

Rebecca C Pierson

University of Louisville School of Medicine

Sashia Torres

University of Louisville School of Public Health and Information Sciences

Emily Reece

University of Louisville School of Public Health and Information Sciences

Kira C Taylor

University of Louisville School of Public Health and Information Sciences

Research article

Keywords: Anti-Müllerian hormone (AMH), ovarian reserve, polycystic ovary syndrome (PCOS), NAT2, smoking

Posted Date: November 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1075418/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

This study examined the association of smoking with ovarian reserve in a cross-sectional study of 207 women enrolled in the Louisville Tobacco Smoke Exposure, Genetic Susceptibility, and Infertility (LOUSSI) Study and assessed effect modification by NAT2 acetylator phenotype.

Methods

Information on current smoking status was collected using a structured questionnaire and confirmed by cotinine assay. Serum anti-Müllerian hormone (AMH) levels were used to assess ovarian reserve. Diminished ovarian reserve (DOR) was defined as AMH < 1 ng/mL. Single nucleotide polymorphisms in the *NAT2* gene, which metabolizes toxins found in cigarette smoke, were analyzed to determine NAT2 acetylator status. Linear and logistic regression were used to determine the effects of smoking on ovarian reserve and evaluate effect modification by NAT2. Regression analyses were stratified by polycystic ovary syndrome (PCOS) status and adjusted for age.

Results

Current smoking status, either passive or active, was not significantly associated with DOR. For dose-response assessed using self-report, the odds of DOR increased significantly for every additional cigarette currently smoked (odds ratio, OR:1.08; 95% confidence interval, 95%CI:1.01–1.15); additionally, every 1 pack-year increase in lifetime exposure was associated with an increased odds of DOR among women without PCOS (OR: 1.08 95%CI: 0.99–1.18). These trends appear to be driven by the heavy or long-term smokers. Effect modification by NAT2 genotype was not established.

Conclusion

A history of heavy smoking, but not current smoking status, may indicate increased risk of diminished ovarian reserve.

Background

Infertility and impaired fecundity among women of reproductive age remain important public health issues. Cigarette smoking in women has been associated with adverse reproductive outcomes such as poorer IVF outcome and increased adverse pregnancy outcomes [1–4]. However, studies of the association of smoking with ovarian reserve have yielded inconsistent results [1, 2, 5–15].

Some studies have reported significantly lower anti-Müllerian hormone (AMH) levels, a measure of ovarian reserve, among smokers [1, 2, 5–9]. Schuh-Huerta et al. however, found that serum AMH levels were significantly higher among smoking women compared to non-smoking women [10]. Other studies found no significant association between smoking and measures of ovarian reserve [11–15]. Previous results are difficult to reconcile as several of the studies [1, 2, 8, 9, 15, 16] were conducted among infertile or sub-fertile populations with none of the studies controlling for polycystic ovary syndrome (PCOS). Women with PCOS diagnosis have been shown to have significantly elevated AMH levels [17, 18].

Genetic heterogeneity is another factor that may explain the inconsistent results of previous studies on the association of smoking with ovarian reserve. N-acetyltransferase2 (NAT2) acetylator status (rapid, intermediate, or slow) has been shown to modify the association of smoking with different disease outcomes, with slow acetylators often more susceptible [19–24]. Taylor et al. reported reduced fecundability among current smokers who are slow acetylators [20]; however, it remains to be established whether these effects on fecundability are modulated through a reduction in ovarian reserve. No studies have assessed the effect of NAT2 acetylator status or the potential interaction with smoking on ovarian reserve.

This study assessed the association of smoking with ovarian reserve in a clinical population of women seeking fertility counseling and used urinary cotinine to validate current smoking status and a smoking questionnaire to estimate cumulative lifetime exposure. In addition, this study explored genetic heterogeneity by assessing whether NAT2 acetylator status influenced the association of smoking with ovarian reserve. Analyses were further stratified by PCOS status to account for higher baseline AMH levels.

Methods

Study Population

The Louisville Tobacco Smoke Exposure, Genetic Susceptibility, and Infertility Study (LOUSSI study) is an observational study that is focused on the effects of tobacco exposure and NAT2 acetylator status on ovarian reserve and IVF outcomes. All women 21 years and older seeking infertility treatment in the University of Louisville Reproductive Endocrinology and Infertility (REI) Division were eligible for enrollment into the study. Excluded were women with ongoing pregnancies and patients who could not communicate in English or were unable to understand and complete the informed consent and questionnaire. Women over 45 years of age at the time of initial consult visit were excluded. A total of 261 women were recruited between September 2016 and June 2018. The University of Louisville Institutional Review Board approved this study for human subjects, and all subjects provided written consent prior to participation (IRB number- 16.0063).

Smoking Assessment

Current smoking status was assessed based on urinary cotinine levels. Cotinine, a metabolite of nicotine, has a half-life that ranges from 7 to 40 hours and levels can be used to assess recent exposure to nicotine (past 3-5 days) [25-27]. Urine was collected at enrollment and stored at -80°C until time of assay. Cotinine ELISA assays (Calbiotech, Spring Valley, CA) were used to estimate urinary cotinine levels with maximum detectable level of cotinine for the assay of 100ng/mL. Urinary cotinine levels were estimated to the nearest 0.50 ng/mL. All self-reported smokers except for one had cotinine levels >100ng/ml; therefore, based on preliminary analysis examining the distribution of cotinine levels in the study population and evidence from literatures [25, 28, 29], the cotinine level used to discriminate smokers from non-smokers was set at ≥ 14 ng/mL. A “current active smoker” was defined as a person with urinary cotinine levels of ≥ 14 ng/mL [25]. Women with cotinine levels between 0.5-13.9 ng/mL were classified as “passive smokers” [25]. A “nonsmoker” was defined as a woman with cotinine levels <0.5 ng/mL [25].

Dose-response was assessed using response to a self-administered smoking questionnaire (SSQ) developed for the study (Supplementary File 1). Cumulative lifetime smoking was based on pack-years smoked, calculated for current and former smokers (women with history of smoking at least 1 cigarette/week, who reported quitting more than one month prior to enrollment) by multiplying the number of packs smoked per day by the years of smoking, as reported on the questionnaire.

NAT2 Genotyping

Genomic DNA was isolated from urine samples using the ZR Urine DNA Isolation Kit™ (Zymo Research, Irvine, CA, USA) according to the manufacturer’s instructions. NAT2 alleles, haplotypes, genotypes, and deduced phenotype were determined using a NAT2 four-SNP genotype panel of rs1801279 (191G>A), rs1801280 (341T>C), rs1799930 (590G>A) and rs1799931 (857G>A). The accuracy of the four-SNP panel in determining NAT2 acetylator status is 98.4% and comparable to the seven-SNP panel [30]. The assay uses SNP-specific PCR primers and fluorogenic probes designed using Primer Express™ (Applied Biosystems, CA, USA). The presence of the four SNPs was determined using a predeveloped TaqMan® SNP Genotyping Assay (Applied Biosystems) according to the manufacturer’s instructions. Positive and negative (no DNA template) controls were run to ensure that there was no amplification of contaminating DNA, and between 10% and 20% of samples from each plate were run in duplicate. Individuals possessing two of the NAT2 alleles associated with rapid acetylation (*NAT2*4*) were classified as rapid acetylators; individuals possessing one of the alleles associated with rapid acetylation and one allele associated with slow acetylation (*NAT2*5*, *NAT2*6*, *NAT2*7*, and *NAT2*14*) were classified as intermediate acetylators; and those individuals who possessed two slow acetylation alleles (*NAT2*5*, *NAT2*6*, *NAT2*7*, and *NAT2*14*) were classified as slow acetylators.

Assessment of Ovarian Reserve

Ovarian reserve can be measured using serum AMH levels or antral follicle count. Serum AMH and antral follicle count (AFC) have strong and similar linear relationship with the size of the primordial follicle pool

and ovarian reserve [31-34]. Studies have shown that AMH level and AFC are highly correlated [15, 35]. The limited intra- and inter-cycle variation, objectivity, and potential standardization of AMH assays, makes it the preferred biomarker of ovarian reserve in women.

Assessment of ovarian reserve was based on baseline serum AMH levels extracted from patients' medical records. Women seen in the University of Louisville REI Division routinely provide a serum sample for analysis of reproductive hormones, including AMH, as part of the initial infertility workup. AMH assays were performed by Quest Diagnostics (Louisville, KY) using chemiluminescence. The limit of detection (LOD) for the assay was 0.03ng/mL; AMH values below the LOD were assigned a value of 0.03ng/mL. For this study, diminished ovarian reserve was defined as baseline serum AMH level less than 1ng/mL [13, 36].

Statistical Analysis

The effect of current active smoking on ovarian reserve was assessed by comparing current active smokers with never smokers and passive smokers. The dose of exposure for current active smoking was based on number of cigarettes smoked per day reported on the SSQ (Supplementary File 1), irrespective of the urinary cotinine values. The dose-response relationship between current active smoking and ovarian reserve was modeled as both a continuous exposure variable and a categorical variable: non-smoker (0 cigarettes/day), moderate smoker (1– <10 cigarettes/day), or heavy smoker (≥ 10 cigarettes/day).

Cumulative lifetime exposure was based on number of packs smoked per day and duration of smoking in years reported on the SSQ by both current active smokers and former smokers. The effect of cumulative lifetime years of active smoking was assessed with both a continuous exposure variable and categorical variable, coded as never smoker, 1-5 pack-years: or >5 pack-years.

Covariates extracted from the medical record included age; race; weight and height at enrollment for calculation of body mass index (BMI); age at menarche; parity; and PCOS status. Age at enrollment was categorized as 21–25, 26–30, 31–35 and 36–45 years. Parity, defined as the number of live births, was put into 3 categories: nulliparous, one live birth, or two or more live births. PCOS status was extracted from the medical record, documented as a current diagnosis or as a previous diagnosis self-reported by the patient. BMI at enrollment was categorized as normal weight (17.5 – 24.9), overweight (25.0 – 29.9), obese (30.0 – 34.9), or morbidly obese (≥ 35). Differences in demographics and covariates between smokers and non-smokers were tested using a Chi-square/Fisher's exact test for categorical variables, Student's t-test for continuously normally distributed variables, and the Wilcoxon rank-sum test for continuous variables that were not normally distributed.

Linear regression models using the natural logarithm (ln) of AMH as the dependent variable were generated to estimate the effects of the different smoking variables on ovarian reserve, adjusted for covariates. Percent differences and 95% confidence intervals (CIs) in serum AMH between exposure groups were calculated from the parameter estimates as follows: $([\exp(\beta) - 1] * 100)$ and presented with

corresponding 95% confidence intervals (CIs) and p-values [13]. Unconditional logistic regression models were used to estimate odds ratios (ORs) and 95% CIs for the association of smoking on DOR. Regression analyses were adjusted for variables that were significantly associated with ovarian reserve (age and PCOS).

Effect modification by *NAT2* genotype was assessed by including an interaction term between *NAT2* and the smoking variables in the multivariable models. The rapid acetylator phenotype was combined with the intermediate acetylator phenotype to form the referent group for comparison with the slow acetylator group which has been used in other studies [30, 37-39]. Interaction models were further adjusted for race, in addition to other covariates. Ethnic differences in the frequency of rapid and slow acetylator *NAT2* alleles or haplotype have been reported [20, 40-42]. A P-value <0.05 was considered statistically significant. All regression models were also run restricted to women without PCOS. All analyses were two-sided and conducted using Statistical Analysis Software version 9.4 (SAS Institute, Cary, NC).

Results

Population

Out of 261 women who were enrolled into the LOUSSI study, 54 were excluded due to incomplete records (52 were missing serum AMH levels and 2 were missing PCOS status) (Fig. 1). Thus, 207 women were included in the final analysis of the questionnaire data (self-reported dose data such as pack-years). For the analysis of current smoking status using cotinine measurement, 8 of the 207 were excluded as they did not provide urine for cotinine assays (Fig. 1).

Descriptive Statistics and Bivariable Analyses

Of the remaining 199 participants, 29% were current active smokers with 55% of these reporting smoking 10 or more cigarettes per day, 21% smoking between 5 to 9 cigarettes per day and 24% reported smoking fewer than 5 cigarettes per day. Current active smokers were more likely to be of self-reported black race ($p < 0.001$), obese ($p = 0.03$) and single (67% vs. 22%, $p < 0.001$), compared with non-smokers (Table 1). There were no significant differences between current active smokers and non-smokers for the other characteristics examined (Table 1).

Table 1

Characteristics of women in the LOUSSI study, stratified by smoking status (N = 199)^a

	Nonsmokers (Cotinine < 0.5 ng/mL) N (%) (n = 71)	Passive smokers (Cotinine 0.5–13.9 ng/mL) N (%) (n = 70)	Active smokers (Cotinine ≥ 14 ng/mL) N (%) (n = 58)	P- value
Age (years)				0.06
21–25	6 (8.4)	10 (14.2)	15 (25.9)	
26–30	13 (18.3)	13 (18.6)	14 (24.1)	
31–35	23 (32.4)	27 (38.6)	12 (20.7)	
36–45	29 (40.9)	20 (28.6)	17 (29.3)	
Race				< 0.001
White	38 (53.5)	37 (53.6)	29 (50.0)	
Black	12 (16.9)	20 (29.0)	27 (46.6)	
Other	21 (29.6)	12 (17.4)	2 (3.4)	
<i>Missing</i>		1 (1.4)		
Age at menarche(years) mean (SD)^b	12.7 (2.0)	12.4 (1.7)	12.2 (1.5)	0.33
<i>Missing</i>	2 (2.8)	4 (5.7)	1 (1.7)	
Body Mass Index (kg/m²)				0.03
17.5–24.9	22 (31.4)	15 (21.8)	14 (24.1)	
25-29.9	26 (37.1)	19 (27.5)	12 (20.7)	
30-34.9	13 (18.6)	12 (17.4)	18 (31.1)	
≥ 35	9 (12.9)	23 (33.3)	14 (24.1)	
<i>Missing</i>	1 (1.4)	1 (1.4)		
Polycystic Ovary Syndrome				0.24

^a Women with cotinine measurements.^b SD = Standard Deviation

	Nonsmokers (Cotinine < 0.5 ng/mL) N (%) (n = 71)	Passive smokers (Cotinine 0.5–13.9 ng/mL) N (%) (n = 70)	Active smokers (Cotinine ≥ 14 ng/mL) N (%) (n = 58)	P- value
No	47 (68.1)	38 (54.3)	36 (62.1)	
Yes	2 (31.9)	32 (45.7)	22 (37.9)	
<i>Missing</i>	2 (2.8)			
Parity				0.11
None	26 (36.6)	35 (51.5)	16 (28.1)	
One	17 (23.9)	14 (20.6)	17 (29.8)	
Two or more	28 (39.5)	19 (27.9)	24 (42.1)	
<i>Missing</i>		2 (2.9)	1 (1.7)	
Marital Status				< 0.001
Single	15 (21.7)	17 (26.6)	37 (67.3)	
Married	54 (78.3)	47 (73.4)	18 (32.7)	
<i>Missing</i>	2 (2.8)	6 (8.6)	3 (5.2)	
Serum AMH categories (ng/mL)				0.32
< 1	19 (26.8)	19 (27.1)	9 (15.5)	
1–2	14 (19.7)	10 (14.3)	14 (24.1)	
2–3	20 (28.1)	18 (25.7)	22 (38.0)	
> 3	18 (25.4)	23 (32.9)	13 (22.4)	
Diminished ovarian reserve (DOR) (AMH < 1 ng/mL)				0.23
<i>No</i>	52 (73.2)	51 (72.9)	49 (84.5)	
<i>Yes</i>	19 (26.8)	19 (27.1)	9 (15.5)	
^a <i>Women with cotinine measurements.</i>				
^b <i>SD = Standard Deviation</i>				

	Nonsmokers (Cotinine < 0.5 ng/mL)	Passive smokers (Cotinine 0.5–13.9 ng/mL)	Active smokers (Cotinine ≥ 14 ng/mL)	P- value
	N (%) (n = 71)	N (%) (n = 70)	N (%) (n = 58)	
NAT2 Acetylator Status				
Rapid/ Intermediate Acetylators	40 (61.5)	39 (61.9)	18 (62.1)	0.99
Slow Acetylators	25 (38.5)	24 (38.1)	11 (37.9)	
<i>Missing</i>	6 (8.5)	7 (10)	1 (3.3)	
^a <i>Women with cotinine measurements.</i>				
^b <i>SD = Standard Deviation</i>				

We also examined the association of each potential covariate with ovarian reserve, as measured by either DOR or % change in AMH. Only age and PCOS status were significantly associated with ovarian reserve in bivariable analyses and were considered as potential confounders in the multivariable analyses. A 48% decrease (95%CI: -71.0 to -8.5) in AMH level was reported among women between ages 31 to 35 years and a 67% decrease (95%CI: -81.0 to -40.6) among women older than 35 years of age, relative to women less than 31 years old. A similar inverse association was found between age and DOR, with women between the ages of 36 to 45 years having 6 times the odds of DOR (OR = 6.2; 95% CI: 1.7–22.3) compared to women 21 to 25 years of age. A PCOS diagnosis was associated with a significant increase in AMH levels (177.0%; 95% CI: 89.0–306.1) and a decrease in the odds of DOR (OR = 0.3; 95%CI: 0.1–0.7). Neither serum AMH levels nor DOR was found to be significantly associated with race, BMI, age at menarche, or parity.

Multivariable Modeling Results

Current active smoking, based on cotinine measurement, was not significantly associated with DOR (OR = 0.62; 95% CI: 0.24–1.56); neither was passive smoking: (OR = 1.32; 95% CI: 0.59–3.0) (Table 2). Results were similar among women without PCOS. The results were similar when assessment of current smoking was based only on self-report (OR = 1.22; 95% CI: 0.45–3.27; Non PCOS: OR = 1.87; 95% CI: 0.56–6.33).

Table 2

Multivariable model for the association of cigarette smoking with AMH levels and diminished ovarian reserve (DOR)

	All women ^a			Women without PCOS ^b		
	N	% change in AMH (95% CI)	OR (95% CI) for DOR	N	% change in AMH (95% CI)	OR (95% CI) for DOR
Current smoking status, using cotinine measurement						
Nonsmoker	71	Referent	Referent	47	Referent	Referent
Passive smoker	70	-14.2 (-45.3– 34.4)	1.32 (0.59– 3.0)	38	1.5 (-44.1– 84.4)	0.85 (0.33– 2.19)
Active smoker	58	5.2 (-34.5– 68.9)	0.62 (0.24– 1.56)	36	60.1 (-14.0– 198.3)	0.52 (0.18– 1.52)
Self-reported cigarettes/day						
Treated continuously	207	-2.3 (-5.2– 0.7)	1.05 (0.997– 1.11)	125	-1.3 (-5.1– 2.7)	1.08 (1.01– 1.15) ^c
Categories:						
None	174	Referent	Referent	107	Referent	Referent
1–9	15	9.9 (- 45.7– 122.1)	0.27 (0.03– 2.27)	8	-9.5 (-67.6– 152.6)	0.44 (0.05– 4.27)
≥ 10	18	-38.6 (- 67.9– 17.4)	2.64 (0.81– 8.64)	10	-17.0 (-67.3– 110.8)	4.58 (0.91– 23.02)
Self-reported cumulative lifetime exposure in pack-years						
Treated continuously	207	-1.5 (-5.70– 3.0)	1.05 (0.97– 1.13)	125	-1.2 (-6.4– 4.3)	1.08 (0.99– 1.18)
Categories:						

a-Models adjusted for age and polycystic ovary syndrome

b-Models adjusted for age

c- p-value = 0.04

	All women ^a			Women without PCOS ^b		
Never	150	Referent	Referent	93	Referent	Referent
1–5	35	9.6 (-33.3– 80.2)	0.67 (0.24 – 1.82)	21	-12.8 (-56.0– 72.7)	1.09 (0.36– 3.34)
> 5	22	-27.6 (- 61.4– 35.6)	1.48 (0.49– 4.47)	11	-7.2 (-63.5– 136.1)	2.51 (0.61– 10.23)
<i>a-Models adjusted for age and polycystic ovary syndrome</i>						
<i>b-Models adjusted for age</i>						
<i>c- p-value = 0.04</i>						

However, in the analysis of current cigarettes reported per day on the questionnaire, which was analyzed regardless of cotinine levels, current cigarette smoking was associated with an increased odds of diminished ovarian reserve among women without PCOS. On average, the odds of DOR increased by 8% for every additional cigarette currently smoked (OR = 1.08; 95% CI:1.01–1.15). There was also suggestion of a dose-response with self-reported lifetime exposure. For every 1 pack-year increase in lifetime exposure, the odds of DOR increased by 8% (OR = 1.08; 95% CI: 0.99 – 1.18). Results were similar but not statistically significant when analyses were limited to women with validated cotinine levels and without PCOS (Cigs/day: OR = 1.06; 95% CI: 0.99–1.14; pack-year: OR = 1.08; 95% CI: 0.99–1.18).

We also examined cigarettes per day and pack-years of exposure in categories (Table 2). In these models, the effects of smoking on DOR were only observed in the highest exposure groups (at least 10 cigarettes/day or more than 5 pack-years of exposure), suggesting that the observed associations may be driven by those with high exposure levels.

NAT2 Interaction Analysis

A total of 11 *NAT2* genotypes were identified. The concordance of genotype duplicates was 100% for all four SNPs. DNA extraction or *NAT2* genotyping was unsuccessful in 14 participants, yielding a sample size of 155 for the *NAT2* interaction analysis. About 39% of women in the study were slow acetylators, 44% were intermediate, and 17% were rapid acetylators. The distribution of *NAT2* acetylator status did not differ by smoking status (Table 1) or self-reported race (data not shown).

After adjusting for age, race and PCOS status, there was no statistically significant interaction with *NAT2* phenotype (Table 3). Although there was a suggestion that the odds of DOR was higher among slow acetylators (OR = 1.59; 95% CI: 0.26–9.70), the sample size was insufficient to provide adequate statistical power.

Table 3
Interaction of current cigarette smoking and *NAT2* acetylator phenotype on diminished ovarian reserve (DOR)

	All women ^a				Women without PCOS ^b			
	Rapid/Intermediate		Slow		Rapid/Intermediate		Slow	
	N	OR (95% CI)	N	OR (95% CI)	N	OR (95% CI)	N	OR (95% CI)
Nonsmoker	38	1.00	25	0.42 (0.05–3.44)	26	1.00	16	0.35 (0.02–5.14)
Passive smoker	39	1.16 (0.30–4.50)	24	2.42 (0.37–15.97)	25	1.43 (0.31–6.66)	9	1.92 (0.11–33.78)
Active smoker	18	1.25 (0.16–9.54)	11	1.59 (0.26–9.70)	13	1.71 (0.17–17.12)	3	2.50 (0.15–42.6)
<i>a-Adjusted for age, race and polycystic ovary syndrome; p-value for interaction = 0.67</i>								
<i>b-Adjusted for age and race; p-value for interaction = 0.69</i>								

Discussion

In this study of women seeking fertility counseling, current smoking was not associated with diminished ovarian reserve. However, other results and trends observed in this study were consistent with the hypothesis that heavy smoking, or smoking for a long duration, may decrease ovarian reserve. Although most results were not statistically significant at the $\alpha = 0.05$ level, one statistically significant association was observed, between cigarettes smoked per day and odds of diminished ovarian reserve among women without PCOS. We also identified other trends consistent with the hypothesis that smoking reduces ovarian reserve. While the 95% confidence intervals crossed the null value for the other analyses, the magnitude and direction of effect suggested that heavy smoking (> 10 cigarettes per day or more than 5 pack-years over the lifespan) reduces ovarian reserve. The effect sizes among women without PCOS were larger, perhaps because the AMH levels among these women are a more accurate reflection of ovarian reserve. Although it is possible that variation in AMH levels among women with PCOS may also reflect ovarian reserve, women with and without PCOS represent two qualitatively different populations with regard to AMH levels.

The validity of this study is supported by its use of a targeted smoking questionnaire, which assessed both current and lifetime smoking habits, and the use of a urinary cotinine assay to validate current smoking. In addition, both diminished ovarian reserve and AMH levels were assessed as outcomes, and results using either outcome were consistent with a dose-response effect of cigarette smoking (current and lifetime) with decreased ovarian reserve. Current active smoking, assessed using cotinine levels, was not significantly associated with DOR. However, there is an established relationship between smoking

and earlier age at menopause [43–47], which further increases the plausibility that smoking is associated with decreased ovarian reserve, possibly through accelerated ovarian aging and follicle atresia.

Constituents of tobacco smoke known to have toxic effects on reproductive health include carbon monoxide, nicotine, cadmium, and polycyclic aromatic hydrocarbons (PAHs) such as benzo[a]pyrene (B[a]P) [48, 49]. Some of the proposed biological mechanisms through which tobacco smoke influences ovarian reserve include inhibition of follicular development; premature luteinization of the preovulatory follicle; reduction of oocyte vascularization and maturation; atresia of oocytes in primordial and small primary follicles; impaired steroidogenesis; increased chromosomal errors; and cytotoxicity [49–52].

Several previous studies have suggested increased risk for diminished ovarian reserve with smoking [1, 2, 5–10], but results have been inconsistent. However, in a cross-sectional study of 913 premenopausal women, White et al. (2016) found a significant association between self-reported smoking and serum AMH among women who smoked 20 or more cigarettes/day (-56.2% , 95%CI: $-80.3, -2.8\%$) [5]. Similar findings with number of pack-years smoked were reported by Dolleman et al., where a significant association between smoking and age-specific serum AMH was found for 10–15 pack-years ($\beta = -7.0$, $p = 0.003$) and 15–20 pack-years ($\beta = -8.5$, $p = 0.001$) pack-years but not for 5–10 pack-years ($\beta = -3.1$, $p = 0.15$) [6]. Therefore, like the present study, prior studies also found that a clinically meaningful reduction in ovarian reserve was only detected among heavy or long-term smokers.

While it is true that women and their partners trying to conceive have been counseled to quit smoking for decades, smoking was still highly prevalent (58/199, or almost 29%) among these women who were actively attempting to conceive and seeking treatment for fertility counseling. Other studies that reported significant inverse associations between smoking and ovarian reserve (as measured by AMH) also reported moderate [5, 7, 10] to high [1, 2, 8, 9] prevalence of smoking in the study population, emphasizing the continued importance of research in the area of smoking and fertility.

Studies on the association of smoking with ovarian reserve varied in the covariates selected for adjustment. None of the other studies controlled for PCOS, even though several of the studies [8, 9, 1, 2, 16] were conducted among infertile or sub-fertile populations. Women with PCOS diagnosis have been shown to have significantly elevated AMH levels [17, 18]. In analyses that excluded women with PCOS diagnosis, effect sizes were generally stronger, suggesting PCOS is a confounder and/or an effect modifier of the association of smoking with ovarian reserve. Therefore, controlling for and stratifying on PCOS was a major improvement in the study design compared to prior studies.

This study further explored population differences in NAT2 genotype frequencies as a possible reason for the inconsistency in the association of smoking with ovarian reserve. Results from this study, though not statistically significant, suggest that the risk of diminished ovarian reserve may be increased for current smokers who are slow NAT2 acetylators. Polymorphisms in the NAT2 gene modify susceptibility to harmful heterocyclic and aromatic amines constituents in tobacco smoke [40, 53, 54]. High levels of DNA adduct of 4-aminobiphenyl, a carcinogenic aromatic amine in tobacco smoke, have been reported in tissues of smokers such as larynx [55], liver [56], bladder [57, 58], breast [59–61] and sputum [62].

Additionally, it has been shown that slow NAT2 acetylators have variable reductions in catalytic activities that make them more susceptible to the effect of some toxins [40, 53, 54, 63]. Given the significantly reduced fecundability reported among current smokers who are slow acetylators by Taylor et al. [20], it is plausible that the effect is modulated through a reduction in ovarian reserve. Slow NAT2 acetylator phenotypes result from different mechanisms based on the presence of single nucleotide polymorphisms [63, 64], and subsequent studies have shown that the slow phenotype is not homogenous [65, 66]. Therefore, treating all slow acetylators as a single group may be diluting any effects present in a particular haplotype. Genetic heterogeneity of the slow NAT2 phenotype was investigated by assessing the interaction of smoking with individual NAT2 alleles and genotypes on ovarian reserve, but this analysis was limited by small sample size (data not shown). More studies, with a large sample size, are needed to clarify the role of NAT2 acetylator phenotype as well as specific NAT2 genotypes in the association of smoking with ovarian reserve.

This study has a number of limitations common to cross sectional studies, including the potential for recall bias (e.g., pack-years smoked), resulting in possible misclassification of exposure; potential selection bias; and residual confounding. Recruiting participants from a single clinic may result in lack of generalizability, though patients who seek care at this site have varied payor mix. In addition, women excluded from this analysis for missing information on important variables (e.g., AMH) may be different in meaningful ways from women who had complete data. Finally, we presented a parsimonious model, and there could be residual confounding. As the associations of socioeconomic status and other covariates such as alcohol and hormonal contraceptive use on ovarian reserve have not been well-established, it is uncertain how these unmeasured covariates may have influenced the effect estimates obtained from this study. Previous diagnosis of endometriosis or a history of ovarian surgery that may diminish ovarian reserve were not assessed in this study.

This study was limited by the small sample of heavy smokers in the study population which reduced the power to detect the main effect of smoking or its potential interaction with NAT2 on ovarian reserve. Consequently, it cannot be certain whether the null associations are attributable to an actual lack of association or to inadequate statistical power. Future research with a similar study design should plan to expand recruitment to multiple gynecological and fertility clinics and should conduct power calculations based on the expected number of heavy smokers.

Strengths of this study include the novel investigation of the impact of NAT2 acetylator status on the relationship of smoking with ovarian reserve; the use of questionnaires that permitted a more detailed characterization of both current and lifetime exposure than typical patient intake questionnaires; and validation of current exposure status with cotinine.

Conclusions

Current smoking was not associated with diminished ovarian reserve. However, the trends and directions of effect observed in this study suggest that heavy smoking, or smoking for a long duration, may reduce

ovarian reserve. Additional data are needed to better define the role of NAT2 polymorphisms with diminished ovarian reserve following exposure to toxins such as those found in cigarette smoke.

Abbreviations

AFC Antral Follicle Count

AMH Anti-Müllerian Hormone

DOR Diminished Ovarian Reserve

IVF In Vitro Fertilization

LOD Limit of Detection

NAT 2 N-acetyltransferase 2

PCOS Polycystic Ovary syndrome

REI Reproductive Endocrinology and Infertility

SNP Single Nucleotide Polymorphisms

SSQ Self-administered Smoking Questionnaire

Declarations

Ethics approval and consent to participate

The University of Louisville Institutional Review Board approved this study for human subjects (IRB number- 16.0063). All subjects provided written consent prior to participation.

Consent to publish

Written consent was obtained from all subjects regarding publication of their data.

Availability of data and material

Anonymized data that support the findings of this study are available from the Principal Investigator of LOUSSI study upon reasonable request.

Competing interests

The authors have no conflicts of interest to declare that are relevant to the content of this article.

Funding

The LOUSSI study is funded by the National Institutes of Health (NIH), Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD)– 1R15HD087911-01 (PI: Kira Taylor). Portions of this work constitute partial fulfillment for the award of PhD in Epidemiology to I.O at the University of Louisville.

Authors' contributions

I.O. and K.C.T were involved with all aspects of this manuscript including study design, execution, and all genetic and statistical analyses. I.O. drafted the initial manuscript. D.W.H helped to obtain funding, provided advice regarding study design, NAT2 analyses and interpretation, and revised the final manuscript. M.A.D provided additional laboratory support for the DNA extraction, NAT2 genotyping and interpretation of results. I.O, T.A, S.T, K.C.T and E.R collected all clinical data including patient questionnaire, cotinine assay, medical record review for variables of interest, DNA extraction and NAT2 genotyping. K.P and H.B provided access to their patients, helped with medical record extraction, and assessed the initial eligibility of participants with help from M.M, A.G and J.L.C. R.C.P assisted with medical record extractions and reviewed the manuscript for intellectual content. All authors participated in work group meetings where results were reviewed and interpreted, and analyses were refined accordingly. All authors reviewed the manuscript and provided critical feedback and discussion. K.C.T conceived and initiated the LOUSSI study with the goals of understanding the role of gene-environment interactions play in female fertility.

Acknowledgements

The authors express their sincere thanks to the LOUSSI study participants who made this work possible. Portions of this work constitute partial fulfillment for the award of PhD in Epidemiology to I.O at the University of Louisville.

References

1. Freour T, Masson D, Mirallie S, Jean M, Bach K, Dejoie T, et al. Active smoking compromises IVF outcome and affects ovarian reserve. *Reprod Biomed Online*. 2008;16(1):96–102.
2. Freour T, Masson D, Dessolle L, Allaoua D, Dejoie T, Mirallie S, et al. Ovarian reserve and in vitro fertilization cycles outcome according to women smoking status and stimulation regimen. *Archives of gynecology obstetrics*. 2012;285(4):1177–82. doi:10.1007/s00404-011-2172-7.
3. El-Nemr A, Al-Shawaf T, Sabatini L, Wilson C, Lower AM, Grudzinskas JG. Effect of smoking on ovarian reserve and ovarian stimulation in in-vitro fertilization and embryo transfer. *Hum Reprod*. 1998;13(8):2192–8.
4. Radin RG, Hatch EE, Rothman KJ, Mikkelsen EM, Sorensen HT, Riis AH, et al. Active and passive smoking and fecundability in Danish pregnancy planners. *Fertil Steril*. 2014;102(1):183 – 91.e2. doi:10.1016/j.fertnstert.2014.03.018.

5. White AJ, Sandler DP, D'Aloisio AA, Stanczyk F, Whitworth KW, Baird DD, et al. Antimullerian hormone in relation to tobacco and marijuana use and sources of indoor heating/cooking. *Fertil Steril*. 2016;106(3):723–30. doi:10.1016/j.fertnstert.2016.05.015.
6. Dolleman M, Verschuren WM, Eijkemans MJ, Dolle ME, Jansen EH, Broekmans FJ, et al. Reproductive and lifestyle determinants of anti-Mullerian hormone in a large population-based study. *J Clin Endocrinol Metab*. 2013;98(5):2106–15. doi:10.1210/jc.2012-3995.
7. Plante BJ, Cooper GS, Baird DD, Steiner AZ. The impact of smoking on antimullerian hormone levels in women aged 38 to 50 years. *Menopause (New York NY)*. 2010;17(3):571–6. doi:10.1097/gme.0b013e3181c7deba.
8. Fuentes A, Munoz A, Pommer R, Arguello B, Galleguillos A, Torres A, et al. Decreased anti-Mullerian hormone concentration in follicular fluid of female smokers undergoing artificial reproductive techniques. *Chemosphere*. 2012;88(4):403–6. doi:10.1016/j.chemosphere.2012.02.054.
9. Fuentes A, Escalona J, Cespedes P, Repetto V, Iniguez G. [Effects of smoking on plasma antimullerian hormone concentrations among infertile women]. *Revista medica de Chile*. 2013;141(1):23–7. doi:10.4067/s0034-98872013000100003.
10. Schuh-Huerta SM, Johnson NA, Rosen MP, Sternfeld B, Cedars MI, Reijo Pera RA. Genetic variants and environmental factors associated with hormonal markers of ovarian reserve in Caucasian and African American women. *Hum Reprod*. 2012;27(2):594–608. doi:10.1093/humrep/der391.
11. Jung S, Allen N, Arslan AA, Baglietto L, Brinton LA, Eggleston BL, et al. Demographic, lifestyle, and other factors in relation to antimullerian hormone levels in mostly late premenopausal women. *Fertil Steril*. 2017;107(4):1012–22.e2. doi:10.1016/j.fertnstert.2017.02.105.
12. Kline J, Tang A, Levin B. Smoking, alcohol and caffeine in relation to two hormonal indicators of ovarian age during the reproductive years. *Maturitas*. 2016;92:115–22. doi:10.1016/j.maturitas.2016.07.010.
13. Hawkins Bressler L, Bernardi LA, De Chavez PJ, Baird DD, Carnethon MR, Marsh EE. Alcohol, cigarette smoking, and ovarian reserve in reproductive-age African-American women. *Am J Obstet Gynecol*. 2016;215(6):758 e1- e9. doi:10.1016/j.ajog.2016.07.012.
14. Waylen AL, Jones GL, Ledger WL. Effect of cigarette smoking upon reproductive hormones in women of reproductive age: a retrospective analysis. *Reprod Biomed Online*. 2010;20(6):861–5. doi:10.1016/j.rbmo.2010.02.021.
15. Nardo LG, Christodoulou D, Gould D, Roberts SA, Fitzgerald CT, Laing I. Anti-Mullerian hormone levels and antral follicle count in women enrolled in in vitro fertilization cycles: relationship to lifestyle factors, chronological age and reproductive history. *Gynecol Endocrinol*. 2007;23(8):486–93. doi:10.1080/09513590701532815.
16. Caserta D, Bordi G, Di Segni N, D'Ambrosio A, Mallozzi M, Moscarini M. The influence of cigarette smoking on a population of infertile men and women. *Archives of gynecology obstetrics*. 2013;287(4):813–8. doi:10.1007/s00404-012-2643-5.

17. Tal R, Seifer DB, Khanimov M, Malter HE, Grazi RV, Leader B. Characterization of women with elevated antimullerian hormone levels (AMH): correlation of AMH with polycystic ovarian syndrome phenotypes and assisted reproductive technology outcomes. *Am J Obstet Gynecol*. 2014;211(1):59.e1-8. doi:10.1016/j.ajog.2014.02.026.
18. Pigny P, Merlen E, Robert Y, Cortet-Rudelli C, Decanter C, Jonard S, et al. Elevated serum level of anti-mullerian hormone in patients with polycystic ovary syndrome: relationship to the ovarian follicle excess and to the follicular arrest. *J Clin Endocrinol Metab*. 2003;88(12):5957–62. doi:10.1210/jc.2003-030727.
19. Dubinskaia ED, Gasparov AS, Fedorova TA, Lapteva NV. [Role of the genetic factors, detoxication systems and oxidative stress in the pathogenesis of endometriosis and infertility (review)]. *Vestnik Rossiiskoi akademii meditsinskikh nauk*. 2013(8):14–9.
20. Taylor KC, Small CM, Dominguez CE, Murray LE, Tang W, Wilson MM, et al. Alcohol, smoking, and caffeine in relation to fecundability, with effect modification by NAT2. *Ann Epidemiol*. 2011;21(11):864–72. doi:10.1016/j.annepidem.2011.04.011.
21. Wu H, Wang X, Zhang L, Mo N, Lv Z. Association Between N-acetyltransferase 2 Polymorphism and Bladder Cancer Risk: Results From Studies of the Past Decade and a Meta-Analysis. *Clin Genitourin Cancer*. 2016;14(2):122–9. doi:10.1016/j.clgc.2015.10.007.
22. Risch A, Wallace DM, Bathers S, Sim E. Slow N-acetylation genotype is a susceptibility factor in occupational and smoking related bladder cancer. *Human molecular genetics*. 1995;4(2):231–6.
23. Moslehi R, Chatterjee N, Church TR, Chen J, Yeager M, Weissfeld J, et al. Cigarette smoking, N-acetyltransferase genes and the risk of advanced colorectal adenoma. *Pharmacogenomics*. 2006;7(6):819–29. doi:10.2217/14622416.7.6.819.
24. Mendola P, Moysich KB, Freudenheim JL, Shields PG, Schisterman EF, Graham S, et al. Risk of recurrent spontaneous abortion, cigarette smoking, and genetic polymorphisms in NAT2 and GSTM1. *Epidemiology (Cambridge, Mass)*. 1998;9(6):666-8.
25. Benowitz NL, Schultz KE, Haller CA, Wu AH, Dains KM, Jacob P 3. Prevalence of smoking assessed biochemically in an urban public hospital: a rationale for routine cotinine screening. *Am J Epidemiol*. 2009;170(7):885–91. doi:10.1093/aje/kwp215. rd. ;) .
26. Benowitz NL. Biomarkers of environmental tobacco smoke exposure. *Environmental health perspectives*. 1999;107(Suppl 2):349–55.
27. Avila-Tang E, Al-Delaimy WK, Ashley DL, Benowitz N, Bernert JT, Kim S, et al. Assessing secondhand smoke using biological markers. *Tobacco control*. 2013;22(3):164–71. doi:10.1136/tobaccocontrol-2011-050298.
28. Benowitz NL, Bernert JT, Caraballo RS, Holiday DB, Wang J. Optimal serum cotinine levels for distinguishing cigarette smokers and nonsmokers within different racial/ethnic groups in the United States between 1999 and 2004. *Am J Epidemiol*. 2009;169(2):236–48. doi:10.1093/aje/kwn301.
29. Homa DM, Neff LJ, King BA, Caraballo RS, Bunnell RE, Babb SD, et al. Vital signs: disparities in nonsmokers' exposure to secondhand smoke—United States, 1999–2012. *MMWR Morbidity and*

- mortality weekly report. 2015;64(4):103-8.
30. Hein DW, Doll MA. Accuracy of various human NAT2 SNP genotyping panels to infer rapid, intermediate and slow acetylator phenotypes. *Pharmacogenomics*. 2012;13(1):31–41. doi:10.2217/pgs.11.122.
 31. Fleming R, Seifer DB, Frattarelli JL, Ruman J. Assessing ovarian response: antral follicle count versus anti-Mullerian hormone. *Reprod Biomed Online*. 2015;31(4):486–96. doi:10.1016/j.rbmo.2015.06.015.
 32. Tal R, Seifer DB. Ovarian reserve testing: a user's guide. *Am J Obstet Gynecol*. 2017;217(2):129–40. doi:10.1016/j.ajog.2017.02.027.
 33. Nelson SM. Biomarkers of ovarian response: current and future applications. *Fertil Steril*. 2013;99(4):963–9. doi:10.1016/j.fertnstert.2012.11.051.
 34. Jayaprakasan K, Campbell B, Hopkisson J, Johnson I, Raine-Fenning N. A prospective, comparative analysis of anti-Mullerian hormone, inhibin-B, and three-dimensional ultrasound determinants of ovarian reserve in the prediction of poor response to controlled ovarian stimulation. *Fertil Steril*. 2010;93(3):855–64. doi:10.1016/j.fertnstert.2008.10.042.
 35. van Disseldorp J, Lambalk CB, Kwee J, Looman CW, Eijkemans MJ, Fauser BC, et al. Comparison of inter- and intra-cycle variability of anti-Mullerian hormone and antral follicle counts. *Hum Reprod*. 2010;25(1):221–7. doi:10.1093/humrep/dep366.
 36. Kara M, Aydin T, Aran T, Turktekin N, Ozdemir B. Does dehydroepiandrosterone supplementation really affect IVF-ICSI outcome in women with poor ovarian reserve? *Eur J Obstet Gynecol Reprod Biol*. 2014;173:63–5. doi:10.1016/j.ejogrb.2013.11.008.
 37. Baumgartner KB, Schlierf TJ, Yang D, Doll MA, Hein DW. N-acetyltransferase 2 genotype modification of active cigarette smoking on breast cancer risk among hispanic and non-hispanic white women. *Toxicol Sci*. 2009;112(1):211–20. doi:10.1093/toxsci/kfp199.
 38. Ambrosone CB, Kropp S, Yang J, Yao S, Shields PG, Chang-Claude J. Cigarette smoking, N-acetyltransferase 2 genotypes, and breast cancer risk: pooled analysis and meta-analysis. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research. cosponsored by the American Society of Preventive Oncology*. 2008;17(1):15–26. doi:10.1158/1055-9965.Epi-07-0598.
 39. Kinzig-Schippers M, Tomalik-Scharte D, Jetter A, Scheidel B, Jakob V, Rodamer M, et al. Should we use N-acetyltransferase type 2 genotyping to personalize isoniazid doses? *Antimicrobial agents and chemotherapy*. 2005;49(5):1733–8. doi:10.1128/aac.49.5.1733-1738.2005.
 40. Walker K, Ginsberg G, Hattis D, Johns DO, Guyton KZ, Sonawane B. Genetic polymorphism in N-Acetyltransferase (NAT): Population distribution of NAT1 and NAT2 activity. *Journal of toxicology environmental health Part B Critical reviews*. 2009;12(5–6):440–72. doi:10.1080/10937400903158383.
 41. Lin HJ, Han CY, Lin BK, Hardy S. Slow acetylator mutations in the human polymorphic N-acetyltransferase gene in 786 Asians, blacks, Hispanics, and whites: application to metabolic

- epidemiology. *Am J Hum Genet.* 1993;52(4):827–34.
42. Gross M, Kruisselbrink T, Anderson K, Lang N, McGovern P, Delongchamp R, et al. Distribution and concordance of N-acetyltransferase genotype and phenotype in an American population. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology.* 1999;8(8):683–92.
43. Oboni JB, Marques-Vidal P, Bastardot F, Vollenweider P, Waeber G. Impact of smoking on fertility and age of menopause: a population-based assessment. *BMJ open.* 2016;6(11):e012015. doi:10.1136/bmjopen-2016-012015.
44. Schoenaker DA, Jackson CA, Rowlands JV, Mishra GD. Socioeconomic position, lifestyle factors and age at natural menopause: a systematic review and meta-analyses of studies across six continents. *Int J Epidemiol.* 2014;43(5):1542–62. doi:10.1093/ije/dyu094.
45. Sun L, Tan L, Yang F, Luo Y, Li X, Deng HW, et al. Meta-analysis suggests that smoking is associated with an increased risk of early natural menopause. *Menopause (New York NY).* 2012;19(2):126–32. doi:10.1097/gme.0b013e318224f9ac.
46. Parente RC, Faerstein E, Celeste RK, Werneck GL. The relationship between smoking and age at the menopause: A systematic review. *Maturitas.* 2008;61(4):287–98. doi:10.1016/j.maturitas.2008.09.021.
47. Midgette AS, Baron JA. Cigarette smoking and the risk of natural menopause. *Epidemiology (Cambridge Mass).* 1990;1(6):474–80.
48. Centers for Disease C, Prevention, National Center for Chronic. Disease P, Health P, Office on S, Health. Publications and Reports of the Surgeon General. How Tobacco Smoke Causes Disease: The Biology and Behavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General. Atlanta (GA): Centers for Disease Control and Prevention (US); 2010.
49. Dechanet C, Anahory T, Mathieu Daude JC, Quantin X, Reyftmann L, Hamamah S, et al. Effects of cigarette smoking on reproduction. *Hum Reprod Update.* 2011;17(1):76–95. doi:10.1093/humupd/dmq033.
50. Sadeu JC, Foster WG. Effect of in vitro exposure to benzo[a]pyrene, a component of cigarette smoke, on folliculogenesis, steroidogenesis and oocyte nuclear maturation. *Reprod Toxicol.* 2011;31(4):402–8. doi:10.1016/j.reprotox.2010.12.006.
51. Sadeu JC, Foster WG. The cigarette smoke constituent benzo[a]pyrene disrupts metabolic enzyme, and apoptosis pathway member gene expression in ovarian follicles. *Reprod Toxicol.* 2013;40:52–9. doi:10.1016/j.reprotox.2013.05.008.
52. Sadeu JC, Foster WG. Cigarette smoke condensate exposure delays follicular development and function in a stage-dependent manner. *Fertil Steril.* 2011;95(7):2410–7. doi:10.1016/j.fertnstert.2011.03.072.
53. Hein DW, Doll MA, Fretland AJ, Leff MA, Webb SJ, Xiao GH, et al. Molecular genetics and epidemiology of the NAT1 and NAT2 acetylation polymorphisms. *Cancer epidemiology, biomarkers &*

- prevention: a publication of the American Association for Cancer Research. cosponsored by the American Society of Preventive Oncology. 2000;9(1):29–42.
54. Sim E, Lack N, Wang CJ, Long H, Westwood I, Fullam E, et al. Arylamine N-acetyltransferases: structural and functional implications of polymorphisms. *Toxicology*. 2008;254(3):170–83. doi:10.1016/j.tox.2008.08.022.
 55. Flamini G, Romano G, Curigliano G, Chiominto A, Capelli G, Boninsegna A, et al. 4-Aminobiphenyl-DNA adducts in laryngeal tissue and smoking habits: an immunohistochemical study. *Carcinogenesis*. 1998;19(2):353–7.
 56. Wang LY, Chen CJ, Zhang YJ, Tsai WY, Lee PH, Feitelson MA, et al. 4-Aminobiphenyl DNA damage in liver tissue of hepatocellular carcinoma patients and controls. *Am J Epidemiol*. 1998;147(3):315–23.
 57. Curigliano G, Zhang YJ, Wang LY, Flamini G, Alcini A, Ratto C, et al. Immunohistochemical quantitation of 4-aminobiphenyl-DNA adducts and p53 nuclear overexpression in T1 bladder cancer of smokers and nonsmokers. *Carcinogenesis*. 1996;17(5):911–6.
 58. Airoidi L, Orsi F, Magagnotti C, Coda R, Randone D, Casetta G, et al. Determinants of 4-aminobiphenyl-DNA adducts in bladder cancer biopsies. *Carcinogenesis*. 2002;23(5):861–6.
 59. Ambrosone CB, Abrams SM, Gorlewska-Roberts K, Kadlubar FF. Hair dye use, meat intake, and tobacco exposure and presence of carcinogen-DNA adducts in exfoliated breast ductal epithelial cells. *Arch Biochem Biophys*. 2007;464(2):169–75. doi:10.1016/j.abb.2007.05.018.
 60. Faraglia B, Chen SY, Gammon MD, Zhang Y, Teitelbaum SL, Neugut AI, et al. Evaluation of 4-aminobiphenyl-DNA adducts in human breast cancer: the influence of tobacco smoke. *Carcinogenesis*. 2003;24(4):719–25.
 61. Zhu J, Chang P, Bondy ML, Sahin AA, Singletary SE, Takahashi S, et al. Detection of 2-amino-1-methyl-6-phenylimidazo[4,5-b]-pyridine-DNA adducts in normal breast tissues and risk of breast cancer. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2003;12(9):830–7.
 62. Besaratinia A, Van Straaten HW, Kleinjans JC, Van Schooten FJ. Immunoperoxidase detection of 4-aminobiphenyl- and polycyclic aromatic hydrocarbons-DNA adducts in induced sputum of smokers and non-smokers. *Mutat Res*. 2000;468(2):125–35.
 63. Walraven JM, Zang Y, Trent JO, Hein DW. Structure/function evaluations of single nucleotide polymorphisms in human N-acetyltransferase 2. *Curr Drug Metab*. 2008;9(6):471–86.
 64. Hein DW. N-acetyltransferase SNPs: emerging concepts serve as a paradigm for understanding complexities of personalized medicine. *Expert Opin Drug Metab Toxicol*. 2009;5(4):353–66. doi:10.1517/17425250902877698.
 65. Ruiz JD, Martínez C, Anderson K, Gross M, Lang NP, García-Martín E, et al. The differential effect of NAT2 variant alleles permits refinement in phenotype inference and identifies a very slow acetylation genotype. *PloS one*. 2012;7(9):e44629. doi:10.1371/journal.pone.0044629.
 66. Doll MA, Hein DW. Genetic heterogeneity among slow acetylators N-acetyltransferase 2 phenotypes in cryopreserved human hepatocytes. *Arch Toxicol*. 2017;91(7):2655–61. doi:10.1007/s00204-017-

Figures




Figure 1

Population Flow Diagram

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SmokingQuestionnaire.docx](#)

- [SupplementalTables.docx](#)