Bound States in the Continuum (BIC) Protected By Self-Sustained Potential Barriers in a Flat Band System

Yi-Cai Zhang (✉ zhangyicai123456@163.com)

Guangzhou University

Research Article

Keywords: BIC, self-sustained, barriers, flat band

Posted Date: November 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1082419/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Bound states in the continuum (BIC) protected by self-sustained potential barriers in a flat band system

Yi-Cai Zhang^{1*}

¹School of Physics and Materials Science, Guangzhou University, Guangzhou 510006, People's Republic of China

*zhangyicai123456@163.com

ABSTRACT

In this work, we investigate the bound states in the continuum (BIC) of a one-dimensional spin-1 flat band system with a potential of type III, which has a unique non-vanishing matrix element in basis $|1\rangle$. It is found that, for such a kind of potential, there exists an effective attractive potential well surrounded by infinitely high self-sustained barriers. Some bound states in the continuum (BIC) can appear for sufficiently strong potential. These bound states (BIC) are protected by the infinitely high potential barriers, which could not decay into the continuum. Taking a long-ranged Coulomb potential and a short-ranged exponential potential as two examples, the bound state energies are obtained. For a Coulomb potential, there exists a series of critical potential strength, near which the bound state energy can go to infinite. For a sufficiently strong exponential potential, there exists two different bound states with a same number of wave function nodes. The existences of BIC protected by the self-sustained potential barriers is quite a universal phenomenon in the flat band system under a strong potential. A necessary condition for existence of BIC is that the maximum value of potential is larger than two times band gap.

Introduction

For usual potential wells, the bound states usually appear outside the continuous spectrum. However, Neumann and Wigner constructed a bound state which is embedded in the continuous spectrum of scattering states¹ (the so-called bound states in the continuum (BIC)²). Later, it is found that BIC can also appear due to the interferences of two resonances, where one of the resonance widths becomes zero with the variations of a continuous parameter^{3,4}. BIC can also appear in the optical waveguides⁵⁻⁹, and condensed matter physics system¹⁰⁻¹⁴. In addition, BIC can also exist in bottomless potentials^{15,16}.

A lot of novel physics, for example, existences of localized flat band states¹⁷⁻¹⁹, the ferro-magnetism transition²⁰⁻²², super-Klein tunneling²³⁻²⁶, preformed pairs²⁷, strange metal²⁸, high T_c superconductivity/superfluidity²⁹⁻³⁹, ect., can appear in a flat band system. Due to the existence of infinitely large density of states in a spin-1 flat band system, a short-ranged potential, e.g., square well potential, can result in infinite bound states, even a hydrogen atom-like energy spectrum, i.e., $E_n \propto 1/n^2, n = 1, 2, 3, \dots$ ⁴⁰.

Furthermore, it is found that the existences of bound states also depend on the types of potentials. For example, a long-ranged Coulomb potential of type I (with three same diagonal elements in usual basis), an arbitrary weak Coulomb potential can destroy completely the flat band^{41,42}. In two-dimensional spin-1 systems, a strong Coulomb potential can result in a wave function collapse near the the origin^{41,43}. For one-dimensional case, an arbitrarily weak Coulomb potential also causes the wave function collapse⁴⁴. In addition, for a potential of type II, which has a unique non-vanishing potential matrix element in basis $|2\rangle$ ⁴⁰, a long-ranged Coulomb potential can cause a $1/n$ energy spectrum. For a Coulomb potential of type III, which has a unique non-vanishing potential matrix element in basis $|3\rangle$ ⁴⁵, there are also infinite bound states which are generated from the flat band. Near the the flat band, the energy is inversely proportional to the natural number, i.e., $E \propto 1/n$. Differently from the ordinary one-dimensional bound state energy which is parabolic function of potential strength, the bound state energy is linearly dependent on the potential strength as the strength goes to zero. For a given quantum number n , the bound state energy grows up with the increasing of potential strength α . There is a critical potential strength α_{cr} at which the bound state energy reaches the threshold of continuous spectrum. After crossing the threshold, these bound state may still exist and they would form the bound states in a continuous spectrum (BIC)⁴⁵.

In this work, we propose a new mechanism of the existence of BIC in a spin-1 flat band system with a strong potential of type III. To be specific, we would extend the above investigations (see Ref.⁴⁵) where the energy is in the continuum. It is found that for sufficiently strong potential, there exist an effective potential well which are surrounded by infinitely high potential barriers. Within the potential well, there may exist some bound states which are embedded in the continuous spectrum, i.e., BIC. The infinitely high barriers protect the BIC from decaying into the continuous spectrum. Taking a long-ranged Coulomb potential and a shorted-ranged exponential potential as two examples, we get the bound state (BIC) energies. Our results show that the existence of BIC is quite a universal phenomenon for a strong potential of type III in the spin-1 flat band system.

Figure 1. The energy spectrum of free particle Hamiltonian. The possible conventional bound states exist in the gaps **A** and **B**. Any possible bound states in the continuum (BIC) only exist outside these gaps.

Results

0.1 The model Hamiltonian with a flat band

In this work, we consider a spin-1 Dirac-type Hamiltonian⁴⁰ in one dimension, i.e.,

$$\begin{aligned} H &= H_0 + V_p(x) \\ H_0 &= -iv_F \hbar S_x \partial_x + m S_z, \end{aligned} \quad (1)$$

where $V_p(x)$ is potential energy, H_0 is the free-particle Hamiltonian, $v_F > 0$ is Fermi velocity, and $m > 0$ is energy gap parameter. S_x and S_z are spin operators for spin-1 particles⁴⁶, i.e.,

$$S_x = \begin{bmatrix} 0 & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & 0 \end{bmatrix}; \quad S_z = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \quad (2)$$

in usual basis $|i\rangle$ with $i = 1, 2, 3$. In the whole manuscript, we use the units of $v_F = \hbar = 1$. When $V_p(x) = 0$, the free particle Hamiltonian H_0 has three energy bands. One of them is flat band with eigen-energy $E_{k,0} \equiv 0$, and the other two of them are dispersion bands⁴⁰. Among the three band, there are two band gaps, whose sizes are given by parameter m (see Fig.1). Any possible BIC only exist in the two continuous spectrum, i.e., outside the gaps .

0.2 bound states in a potential of type III

In the following manuscript, we assume the potential energy V_p has following form in usual basis $|i = 1, 2, 3\rangle$, namely,

$$V_p(x) = V_{11}(x) \otimes |1\rangle\langle 1| = \begin{bmatrix} V_{11}(x) & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}. \quad (3)$$

In the whole manuscript, we would refer such a kind of potential as potential of type III⁴⁵. Adopting a similar process as Ref.⁴⁵, the Schrödinger equation

$$\begin{aligned} -i\partial_x \psi_2(x)/\sqrt{2} &= [E - m - V_{11}] \psi_1(x), \\ -i\partial_x [\psi_1(x) + \psi_3(x)]/\sqrt{2} &= E \psi_2(x), \\ -i\partial_x \psi_2(x)/\sqrt{2} &= [E + m] \psi_3(x). \end{aligned} \quad (4)$$

can be transformed into an effective Schrödinger equation (a second-order differential equation), i.e.,

$$-\partial_x^2 \psi(x) + \tilde{V} \psi(x) = \tilde{E} \psi(x). \quad (5)$$

where the auxiliary wave function

$$\psi(x) \equiv \frac{E - V_{11}/2}{E + m} \psi_1(x). \quad (6)$$

The effective total energy $\tilde{E}$ and effective potential $\tilde{V}$ are

$$\begin{aligned} \tilde{E} &= E^2 - m^2 > 0, \text{ for bound states in Continuum,} \\ \tilde{V} &= \frac{V_{11}}{2} \frac{(m + E)^2}{E - V_{11}/2}. \end{aligned} \quad (7)$$

In the following, we would solve the effective Schrödinger equation Eq.(5) to get the bound state (BIC) energies.

0.2.1 a long-ranged Coulomb potential

In this subsection, we assume the V_{11} is a Coulomb potential, i.e.,

$$V_{11}(x) = \frac{\alpha}{|x|}, \quad (8)$$

where α is the potential strength. The effective potential $\tilde{V}$ is

$$\tilde{V}(x) = \frac{V_{11}(x)}{2} \frac{(m + E)^2}{E - V_{11}(x)/2} = \frac{A}{|x| - x_0}. \quad (9)$$

In the above equation, we introduce parameter $A \equiv \frac{\alpha(m+E)^2}{2E}$ and $x_0 \equiv \frac{\alpha}{2E}$. It is shown that the effective potential $\tilde{V}$ is a shifted Coulomb potential⁴⁷ with an effective potential strength A , which depends on energy E . The Eq.(5) becomes

$$\partial_x^2 \psi(x) + [\tilde{E} - \frac{A}{|x| - \frac{\alpha}{2E}}] \psi(x) = 0. \quad (10)$$

Depending on the sign of α/E , there exists two kinds of effective potentials $\tilde{V}$ ⁴⁵. When $\alpha/E < 0$, the effective potential $\tilde{V}$ has a lowest point at $x = 0$. The bound state energies are in the gaps, $0 < E < m$. Hence there is no BIC for $\alpha/E < 0$.

When $\alpha/E > 0$ ($x_0 > 0$), the effective potential $\tilde{V}$ is negative in the interval $(-x_0, x_0)$ (a potential well), and positive in intervals $(-\infty, -x_0)$ and (x_0, ∞) (see Fig.2). There are two infinitely high potential barriers near two ends $x = \pm x_0$ of the interval $(-x_0, x_0)$. We see that the potential well width $w \equiv 2x_0 = \alpha/E$, and the position of potential barriers $x = \pm x_0$ depend sensitively on the bound state energy E . So in the whole work, we call the potential barriers as self-sustained potential barriers. In addition, it is found that the bound state energy can be larger than zero for $\alpha > 0$ or smaller than zero for $\alpha < 0$ ⁴⁵. In the following, we mainly focus on the case of $\alpha/E > 0$, where the bound states in the continuous spectrum (BIC) may appear.

Figure 2. The effective potential wells surrounded by two infinitely high barriers (the red solid lines). (a): The effective potential for Coulomb potential ($\alpha/E > 0$). The value of effective potential at $x = 0$, i.e., $\tilde{V}(0) = -(m + E)^2$. (b): The effective potential for exponential potential. The value of effective potential at $x = 0$, i.e., $\tilde{V}(0) = \frac{(m+E)^2}{\frac{2E}{V_0} - 1}$

When $x > 0$ and $|E| > m$ (for BIC), the equation Eq.(10) can be solved with some confluent hypergeometric functions. Its general solution is

$$\psi(x) = (x - x_0)e^{-i\sqrt{\tilde{E}}(x-x_0)} \{c_1 \times {}_1F_1[a, b, 2i\sqrt{\tilde{E}}(x-x_0)] + c_2 \times U[a, b, 2i\sqrt{\tilde{E}}(x-x_0)]\}, \quad (11)$$

where ${}_1F_1[a, b, z] = \sum_{k=0}^{\infty} \frac{(a)_k z^k}{k!(b)_k}$ is confluent hypergeometric function⁴⁸, $(a)_k = a \times (a+1) \times (a+2) \times \dots \times (a+k-1)$, and $c_1(c_2)$ are two arbitrary constants. $a = 1 + \frac{A}{2i\sqrt{\tilde{E}}}$, $b = 2$. $U[a, b, z]$ is a second linearly independent solution to the confluent hypergeometric equation (Tricomi function⁴⁹). When $z \rightarrow 0$, the two confluent hypergeometric functions behave as

$$\begin{aligned} {}_1F_1[a, b, z] &\simeq 1, \\ U[a, b, z] &= \frac{\Gamma(b-1)}{\Gamma(a)} z^{1-b} + O(|\log z|), \quad (b=2). \end{aligned} \quad (12)$$

When $\alpha/E > 0$, due to the existence of infinitely high potential barriers, the bound state in the continuum only exist in the interval $(-x_0, x_0)$ (see Fig.2). Outside the effective potential well, the wave function vanishes. At the two ends of the interval, the zero boundary conditions should be satisfied, i.e.,

$$\psi(\pm x_0) = 0. \quad (13)$$

Taking Eq.(12) into account, $U[a, b, 2i\sqrt{\tilde{E}}(x-x_0)]$ should be discarded. So the wave function is

$$\psi(x) = (x - x_0)e^{-i\sqrt{\tilde{E}}(x-x_0)} {}_1F_1[a, b, 2i\sqrt{\tilde{E}}(x-x_0)]. \quad (14)$$

Due to $\tilde{V}(-x) = \tilde{V}(x)$, the wave functions can be classified by parities. For odd parity states, the bound state energy equation is

$${}_1F_1\left[1 + \frac{\alpha(E+m)^2}{4iE\sqrt{E^2-m^2}}, 2, -\frac{i\alpha\sqrt{E^2-m^2}}{E}\right] = 0. \quad (15)$$

For even parity states, the bound state energy equation is

$$\begin{aligned} &[-4E(2E + i\alpha\sqrt{E^2-m^2})] \times {}_1F_1\left[1 + \frac{\alpha(E+m)^2}{4iE\sqrt{E^2-m^2}}, 2, -\frac{i\alpha\sqrt{E^2-m^2}}{E}\right] \\ &+ \alpha[4iE\sqrt{E^2-m^2} + \alpha(m+E)^2] \times {}_1F_1\left[2 + \frac{\alpha(E+m)^2}{4iE\sqrt{E^2-m^2}}, 3, -\frac{i\alpha\sqrt{E^2-m^2}}{E}\right] = 0. \end{aligned} \quad (16)$$

The results are reported in Fig.(3).

When $E^2 > m^2$ for BIC, with quasi-classical approximation method⁵⁰, the eigen-energy is given by

$$\frac{\alpha|m+E|}{2E} \left[\sqrt{\frac{2E}{m+E}} + \sqrt{\frac{m+E}{E-m}} \operatorname{arcsinh}\left(\sqrt{\frac{E-m}{E+m}}\right) \right] = (n + \Delta)\pi, \quad (17)$$

where $n = 1, 2, 3, \dots$, and

$$\begin{aligned} \Delta &= +\frac{1}{4}, \text{ for odd parity states,} \\ \Delta &= -\frac{1}{4}, \text{ for even parity states.} \end{aligned} \quad (18)$$

From the Fig. 3, we see that the the bound state (BIC) energy can be well described by the quasi-classical approximation Eq.(17).

For $\alpha > 0$, there exists a critical potential strength⁴⁵, which is determined by

$$\begin{aligned} \alpha_{cr1} &\simeq \frac{(1/4+n)\pi}{2}, \text{ for odd parity states,} \\ \alpha_{cr1} &\simeq \frac{(-1/4+n)\pi}{2}, \text{ for even parity states.} \end{aligned} \quad (19)$$

Figure 3. The bound state (BIC) energy of Coulomb potential in the case of $\alpha/E > 0$. The solid lines are the the exact results of Eqs.(15) and (16). The black dashed lines are given by the quasi-classical approximation formula Eq.(17).

After crossing these critical values, the bound states still exist and they form the bound states in continuum (BIC). For a given n , there exist another critical value α_{cr2} , near which the bound state energy goes to infinity, i.e., $E \rightarrow \pm\infty$. When $E \rightarrow \pm\infty$, the value is can be determined approximately with Eq.(17), i.e.,

$$\alpha_{cr2} \simeq \pm \frac{2(n+\Delta)\pi}{\sqrt{2} + \text{arcsinh}(1)}, \quad (20)$$

where sign $+$ ($-$) for $E >$ ($<$) 0 . For $\alpha < 0$, when $E < -m$ and $|\alpha| \gg 1$, for a given n , the energy approach the threshold of lower band, i.e., $E \rightarrow -m$.

The existence of the BIC can be understood qualitatively as follows. The width of effective potential well is $w \equiv 2x_0 = \alpha/E$. Based on the bound state energy formulas of infinitely deep square well potential, the bound state energy equation can be approximately written as

$$kw = n\pi, \quad (21)$$

where wave vector can be approximated by

$$k \sim \sqrt{\tilde{E} - \tilde{V}(0)} = \sqrt{2E(m+E)}. \quad (22)$$

So we find that the potential strength

$$\alpha = \frac{n\pi E}{\sqrt{2E(m+E)}}, \quad (23)$$

where $E^2 > m^2$ for BIC. Only when

$$\alpha > \alpha_{cr1} \simeq \pi/2, \text{ for } E > m, \quad (24)$$

BIC begin to appear. When $E \rightarrow \pm\infty$, the critical value

$$\alpha_{cr2} \simeq \pm n\pi/\sqrt{2} \simeq \pm 0.7071n\pi. \quad (25)$$

A more accurate formula Eq.(20) gives $\alpha_{cr2} \simeq \pm 0.6534(n+\Delta)\pi$.

0.2.2 short-ranged exponential potential

We should remark that the above mechanism of existence of BIC is quite a universal phenomenon for a sufficiently strong potential. As long as the self-sustained potential barriers can form, the BIC may exist. A critical condition for the existence of the infinitely high potential barriers is that the denominator of the effective potential has zero point [see Eq.(7)], i.e., the equation

$$E - V_{11}(x)/2 = 0 \quad (26)$$

has real solutions x . On the other hand, for BIC, the bound state energy should satisfy

$$|E| > m. \quad (27)$$

Taking the Eqs.(26) and (27) into account, we conclude that when the maximum value of potential is larger than two times gap parameter m , i.e.,

$$|V_{11}(x)|_{\max} > 2m, \quad (28)$$

there may exist BIC. We should emphasize that the above condition Eq.(28) just is a necessary condition for the existence of BIC. So when the potential is sufficiently strong, there may exist BIC in the flat band system.

In this subsection, we assume the potential have an exponential function form, i.e.,

$$V_{11}(x) = V_0 e^{-\beta|x|}, \quad (29)$$

where V_0 is the potential strength, $1/\beta$ describes decaying distance of exponential function. For the above sufficiently strong exponential potential, it is found that there exist BIC. The effective potential $\tilde{V}$ is

$$\tilde{V} = \frac{V_{11}}{2} \frac{(m+E)^2}{E - V_{11}/2} = \frac{(m+E)^2 e^{-\beta|x|}}{\gamma - e^{-\beta|x|}}, \quad (30)$$

where we introduce dimensionless parameter $\gamma \equiv \frac{2E}{V_0}$. When $\gamma \equiv \frac{2E}{V_0} < 1$, $\gamma - e^{-\beta|x|} = 0$, i.e., $x = \pm x_0 \equiv \mp \frac{1}{\beta} \log(\gamma)$, then the denominator of effective potential is zero. In the interval $(-x_0, x_0)$, the effective potential is negative, while outside the interval, the effective potential is positive. Near the two end of interval, i.e., $x = x_0$, there are also two infinitely high potential barriers (see Fig.2). So the bound states in continuum may exist in the interval.

The Eq.(5) becomes

$$\partial_x^2 \psi(x) + [\tilde{E} - \frac{\varepsilon e^{-\beta|x|}}{\gamma - e^{-\beta|x|}}] \psi(x) = 0. \quad (31)$$

where $\tilde{E} = E^2 - m^2$, $\varepsilon = (m + E)^2$. The general solution of Eq.(31) is

$$\psi(x) = c_1 (\gamma e^{\beta x})^{-\frac{\sqrt{-(\tilde{E}+\varepsilon)}}{\beta}} {}_2F_1[A_-, B_-; C_-, \gamma e^{\beta x}] + c_2 (\gamma e^{\beta x})^{\frac{\sqrt{-(\tilde{E}+\varepsilon)}}{\beta}} {}_2F_1[A_+, B_+; C_+, \gamma e^{\beta x}], \quad (32)$$

where ${}_2F_1[a, b; c, z] = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k z^k}{k! (c)_k}$ is hypergeometric function, $(a)_k = a \times (a+1) \times (a+2) \times \dots \times (a+k-1)$, and $c_1 (c_2)$ are two arbitrary constants. $A_{\mp} = \mp \frac{\sqrt{-(\tilde{E}+\varepsilon)}}{\beta} - \frac{\sqrt{-\tilde{E}}}{\beta}$, $B_{\mp} = \mp \frac{\sqrt{-(\tilde{E}+\varepsilon)}}{\beta} + \frac{\sqrt{-\tilde{E}}}{\beta}$, $C_{\mp} = 1 \mp \frac{2\sqrt{-(\tilde{E}+\varepsilon)}}{\beta}$. At two ends of interval $(-x_0, x_0)$, the wave function should vanish, i.e.,

$$\psi(x = x_0) = c_1 (\gamma e^{\beta x_0})^{-\frac{\sqrt{-(\tilde{E}+\varepsilon)}}{\beta}} \times {}_2F_1[A_-, B_-; C_-, \gamma e^{\beta x_0}] + c_2 (\gamma e^{\beta x_0})^{\frac{\sqrt{-(\tilde{E}+\varepsilon)}}{\beta}} \times {}_2F_1[A_+, B_+; C_+, \gamma e^{\beta x_0}] = 0. \quad (33)$$

Similarly, the bound state energy equation are given by

$$\begin{aligned} \psi(x=0) &= 0, \text{ for odd parity states,} \\ \psi'(x=0) &= 0, \text{ for even parity states.} \end{aligned} \quad (34)$$

To be specific, for odd parity states, the bound state energy equation is

$$c_1 \gamma^{-\frac{\sqrt{-(\tilde{E}+\varepsilon)}}{\beta}} \times {}_2F_1[A_-, B_-; C_-, \gamma] + c_2 \gamma^{\frac{\sqrt{-(\tilde{E}+\varepsilon)}}{\beta}} \times {}_2F_1[A_+, B_+; C_+, \gamma] = 0. \quad (35)$$

For even parity states, the bound state energy equation is

$$\begin{aligned} & \frac{\gamma^{-\frac{\sqrt{-(\tilde{E}+\varepsilon)}}{\beta}} c_1}{-\beta + 2\sqrt{-(\tilde{E}+\varepsilon)}} \{ [2(\tilde{E} + \varepsilon) + \beta \sqrt{-\varepsilon - \tilde{E}}] \times {}_2F_1[A_-, B_-; C_-, \gamma] + \varepsilon \gamma \times {}_2F_1[1 + A_-, 1 + B_-; 1 + C_-, \gamma] \} \\ & - \frac{\gamma^{\frac{\sqrt{-(\tilde{E}+\varepsilon)}}{\beta}} c_2}{\beta + 2\sqrt{-(\tilde{E}+\varepsilon)}} \{ [2(\tilde{E} + \varepsilon) - \beta \sqrt{-\varepsilon - \tilde{E}}] \times {}_2F_1[A_+, B_+; C_+, \gamma] + \varepsilon \gamma \times {}_2F_1[1 + A_+, 1 + B_+; 1 + C_+, \gamma] \} = 0 \end{aligned} \quad (36)$$

The results are reported in Fig.(4).

With quasi-classical approximation method, the eigen-energy is given by

$$\begin{aligned} & \frac{-2\sqrt{E^2 - m^2}}{\beta} \log \left[\sqrt{\frac{(V_0/2 - E)(E - m)}{V_0(E + m)/2}} + \sqrt{\frac{mE + V_0E - E^2}{V_0(E + m)/2}} \right] + \frac{\sqrt{2E(E + m)}}{\beta} \log \left[\frac{1 + \sqrt{(m + V_0 - E)/(V_0 - 2E)}}{-1 + \sqrt{(m + V_0 - E)/(V_0 - 2E)}} \right] \\ & = (n + \Delta)\pi, \end{aligned} \quad (37)$$

where $n = 1, 2, 3, \dots$, and

$$\begin{aligned} \Delta &= +\frac{1}{4}, \text{ for odd parity states,} \\ \Delta &= -\frac{1}{4}, \text{ for even parity states.} \end{aligned} \quad (38)$$

From Fig.4, it shows that for a given V_0 there exists two bound states with same quantum number n . When potential is very strong, i.e., $V_0 \rightarrow \pm\infty$, one of bound states $E \rightarrow V_0/2$ (see Fig.4). The other bound state energy

$$\begin{aligned} E &\rightarrow 0, \text{ for } E > 0 \\ E &\rightarrow -m, \text{ for } E < 0. \end{aligned} \quad (39)$$

Figure 4. The bound state (BIC) energy of exponential potential. The solid lines are the the exact results of Eqs. (33), (35), and (36) . The black dashed lines are given by the quasi-classical approximation formula Eq. (37). Here we take $\beta = m$.

Similarly, the above behaviors of bound state energy can be explained as follows. Based on the existence conditions of bound states of infinitely deep square well potential, when

$$kw = n\pi \quad (40)$$

where effective well width $w \equiv 2x_0 = -2\frac{1}{\beta}\log(\gamma)$, the bound states would exist. The wave vector can be approximated by

$$k \sim \sqrt{\tilde{E} - \tilde{V}(0)} = \sqrt{\frac{2E(m+E)(E-m-V_0)}{2E-V_0}} \quad (41)$$

Therefore,

$$\frac{2}{\beta} \sqrt{\frac{2E(E+m)(E-m-V_0)}{2E-V_0}} \log\left[\frac{V_0}{2E}\right] = n\pi \quad (42)$$

where $V_0/(2E) > 1$. When the potential is very strong, e.g., $V_0 \rightarrow \pm\infty$, for a given n , the asymptotic behaviors of bound state energy can be obtained with Eq.(42).

summary

In conclusion, we investigate the bound states in the continuum (BIC) of a one-dimensional spin-1 flat band system. It is found that BIC can exist for sufficiently strong potentials of type III. We get the bound state energies for a Coulomb potential and an exponential potential. For a Coulomb potential and a given quantum number n , when the potential strength reaches a critical value α_{cr1} , the BIC begin to appear. When the potential strength reaches α_{cr2} , the bound state energy goes to infinite. For exponential potential, there are two bound states with same number of wave function nodes. When the exponential potential is very strong, one of bound state energy approaches one half of the potential strength. For repulsive potential (positive V_0), the other bound state energy goes to zero. For attractive case, the other bound state energy approaches to the threshold of lower continuous spectrum. In addition, it is found that the bound state energies can be well described by the quasi-classical approximation.

A necessary condition for existence of BIC is that the maximum value of potential is larger than two times band gap. Our results shows that the existence of BIC is quite a universal phenomenon for a strong potential of type III in the flat band system.

References

1. von Neumann, J. & Wigner, E. On remarkable discrete eigenvalues. *Phys. Z.* **30**, 465, DOI: https://doi.org/10.1007/978-3-662-02781-3_19 (1929).
2. Stillinger, F. H. & Herrick, D. R. Bound states in the continuum. *Phys. Rev. A* **11**, 446, DOI: <https://doi.org/10.1103/PhysRevA.11.446> (1975).
3. Fano, U. Effects of configuration interaction on intensities and phase shifts. *Phys. Rev.* **124**, 1899, DOI: <https://doi.org/10.1103/PhysRev.124.1866> (1961).
4. Friedrich, H. & Wintgenand, D. Interfering resonances and bound states in the continuum. *Phys. Rev. A* **32**, 3231, DOI: <https://doi.org/10.1103/PhysRevA.32.3231> (1985).
5. Bendix, O., Fleischmann, R., Kottos, T. & Shapiro, B. Exponentially fragile p-t symmetry in lattices with localized eigenmodes. *Phys. Rev. Lett.* **103**, 030402, DOI: <https://doi.org/10.1103/PhysRevLett.103.030402> (2009).
6. Bulgakov, E. N. & Sadreev, A. F. Bound states in the continuum in photonic waveguides inspired by defects. *Phys. Rev. B* **78**, 075105, DOI: <https://doi.org/10.1103/PhysRevB.78.075105> (2008).
7. Longhi, S. Optical realization of relativistic non-hermitian quantum mechanics. *Phys. Rev. Lett.* **105**, 013903, DOI: <https://doi.org/10.1103/PhysRevLett.105.013903> (2010).
8. Vicencio, R. *et al.* Observation of localized states in lieb photonic lattices. *Phys. Rev. Lett.* **114**, 245503, DOI: <https://doi.org/10.1103/PhysRevLett.114.245503> (2015).
9. Fong, P. T. & Law, C. K. Bound state in the continuum by spatially separated ensembles of atoms in a coupled-cavity array. *Phys. Rev. A* **96**, 023842, DOI: <https://doi.org/10.1103/PhysRevA.96.023842> (2017).
10. Zhang, J., Braak, D. & Kollar, M. Bound states in the continuum realized in the one dimensional two-particle hubbard model with an impurity. *Phys. Rev. Lett.* **109**, 116405, DOI: <https://doi.org/10.1103/PhysRevLett.109.116405> (2012).

11. Molina M.I. and, A. E., Miroshnichenko & Kivshar, Y. S. Surface bound states in the continuum. *Phys. Rev. Lett.* **108**, 070401, DOI: <https://doi.org/10.1103/PhysRevLett.108.070401> (2012).
12. Gorbatshevich, A. A. & Shubin, N. M. Unified theory of resonances and bound states in the continuum in hermitian tight-binding models. *Phys. Rev. B* **96**, 205441, DOI: <https://doi.org/10.1103/PhysRevB.96.205441> (2017).
13. Xiao, Y. X., Zhang, Z. Q. & Chan, C. A band of bound states in the continuum induced by disorder. *Sci. Reports* **8**, 5160, DOI: <https://doi.org/10.1038/s41598-018-23576-z> (2018).
14. Takeichi, M. & Murakami, M. Topological linelike bound states in the continuum. *Phys. Rev. B* **99**, 035128, DOI: <https://doi.org/10.1103/PhysRevB.99.035128> (2019).
15. Cho, H. & Ho, C. L. A novel quasi-exactly solvable model with total transmission modes. *J. Phys. A: Math. Theor.* **41**, 172002, DOI: <https://iopscience.iop.org/article/10.1088/1751-8113/41/17/172002/meta> (2008).
16. Ahmed, Z., Kumar, S., Ghosh, D. & Goswami. Solvable model of bound states in the continuum (bic) in one dimension. *Phys. Scr.* **94**, 105214, DOI: <https://iopscience.iop.org/article/10.1088/1402-4896/ab2751/meta> (2019).
17. Sutherland, B. Localization of electronic wave functions due to local topology. *Phys. Rev. B* **34**, 5208, DOI: <https://doi.org/10.1103/PhysRevB.34.5208> (1986).
18. Vidal, J., Mosseri, R. & Doucot, B. Aharonov-bohm cages in two-dimensional structures. *Phys. Rev. Lett.* **81**, 5888, DOI: <https://doi.org/10.1103/PhysRevLett.81.5888> (1998).
19. Mukherjee, S. *et al.* Observation of a localized flat-band state in a photonic lieb lattice. *Phys. Rev. Lett.* **114**, 245504, DOI: <https://doi.org/10.1103/PhysRevLett.114.245504> (2015).
20. Mielke, A. Ferromagnetism in single-band hubbard models with a partially flat band. *Phys. Rev. Lett.* **82**, 4312, DOI: <https://doi.org/10.1103/PhysRevLett.82.4312> (1999).
21. Zhang, H., S. and Hung & Wu, C. Proposed realization of itinerant ferromagnetism in optical lattices. *Phys. Rev. A* **82**, 035618, DOI: <https://doi.org/10.1103/PhysRevA.82.035618> (2010).
22. Raoux, A., Morigi, M., Fuchs, J. N., Piéchon, F. & Montambaux, G. From dia to paramagnetic orbital susceptibility of massless fermions. *Phys. Rev. Lett.* **112**, 026402, DOI: <https://doi.org/10.1103/PhysRevLett.112.026402> (2014).
23. Shen, R., Shao, L. B., Wang, B. & Xing, D. Y. Single dirac cone with a flat band touching on line-centered-square optical lattices. *Phys. Rev. B* **82**, 041410, DOI: <https://doi.org/10.1103/PhysRevB.81.041410> (2010).
24. Urban, D. F., Bercioux, D., Wimmer, M. & Häusler, W. Barrier transmission of dirac-like pseudospin-one particles. *Phys. Rev. B* **84**, 115136, DOI: <https://doi.org/10.1103/PhysRevB.84.115136> (2011).
25. Fang, A., Zheng, Z. Q., Louie, S. G. & Chan, C. T. Klein tunneling and supercollimation of pseudospin-1 electromagnetic waves. *Phys. Rev. B* **93**, 035422, DOI: <https://doi.org/10.1103/PhysRevB.93.035422> (2016).
26. Betancur-Ocampo, Y., Cordourier-Maruri, G., Gupta, V. & de Coss, R. Super-klein tunneling of massive pseudospin-one particles. *Phys. Rev. B* **96**, 024304, DOI: <https://doi.org/10.1103/PhysRevB.96.024304> (2017).
27. Tovmasyan, M., Peotta, S., Liang, L., Törmä, P. & Huber, S. D. Preformed pairs in flat bloch bands. *Phys. Rev. B* **98**, 134513, DOI: <https://doi.org/10.1103/PhysRevB.98.134513> (2018).
28. Volovik, G. Flat band and planckian metal. *Jetp Lett.* **110**, 352, DOI: <https://doi.org/10.1134/S002136401917003X> (2019).
29. Peotta, S. & Törmä, P. Superfluidity in topologically nontrivial flat bands. *Nat. Commun.* **6**, 8944, DOI: <https://www.nature.com/articles/ncomms9944> (2015).
30. Hazra, T., Verma, N. & Randeria, M. Bounds on the superconducting transition temperature: Applications to twisted bilayer graphene and cold atoms. *Phys. Rev. X* **9**, 031049, DOI: <https://doi.org/10.1103/PhysRevX.9.031049> (2019).
31. Cao, Y. & et al. Unconventional superconductivity in magic-angle graphene superlattices. *Nature* **556**, 43, DOI: <https://doi.org/10.1038/nature26160> (2018).
32. Wu, Y. R. & Zhang, Y. C. Superfluid states in $\alpha - t_3$ lattice. *Chinese. Phys. B* **30**, 060306, DOI: <https://iopscience.iop.org/article/10.1088/1674-1056/abea8a/meta> (2021).
33. Kopnin, N., Heikkilä, T. T. & Volovik, G. E. High-temperature surface superconductivity in topological flat-band systems. *Phys. Rev. B* **83**, 220503(R), DOI: <https://doi.org/10.1103/PhysRevB.83.220503> (2011).
34. Julku, A., Peltonen, T. J., Liang, L., Heikkilä, T. T. & Törmä, P. Superfluid weight and berezinskii-kosterlitz-thouless transition temperature of twisted bilayer graphene. *Phys. Rev. B* **101**, 060505(R), DOI: <https://doi.org/10.1103/PhysRevB.101.060505> (2020).

35. Iglovikov, V., Hebert, F., Gremaud, B., Batrouni, G. & Scalettar, R. Superconducting transitions in flat-band systems. *Phys. Rev. B* **90**, 094506, DOI: <https://doi.org/10.1103/PhysRevB.90.094506> (2014).
36. Julku, A., Peotta, S., Vanhala, T. I., Kim, D.-H. & Törmä, P. Geometric origin of superfluidity in the lieb-lattice flat band. *Phys. Rev. Lett.* **117**, 045303, DOI: <https://doi.org/10.1103/PhysRevLett.117.045303> (2016).
37. Liang, L. & et.al. Band geometry, berry curvature, and superfluid weight. *Phys. Rev. B* **95**, 024515, DOI: <https://doi.org/10.1103/PhysRevB.95.024515> (2017).
38. Iskin, M. Origin of fat-band superfluidity on the mielke checkerboard lattice. *Phys. Rev. A* **99**, 053608, DOI: <https://doi.org/10.1103/PhysRevB.99.053608> (2019).
39. Wu, Y. R., Zhang, X. F., Liu, C. F., Liu, W. & Zhang, Y. C. Superfluid density and collective modes of fermion superfluid in dice lattice. *Sci Rep* **11**, 13572, DOI: <https://doi.org/10.21203/rs.3.rs-418100/v1> (2021).
40. Zhang, Y. C. & Zhu, G. B. Infinite bound states and hydrogen atom-like energy spectrum induced by a flat band. . . ., DOI: <https://www.researchgate.net/publication/354669549> (2021).
41. Gorbar, E., Gusynin, V. & Oriekhov, D. O. Electron states for gapped pseudospin-1 fermions in the field of a charged impurity. *Phys. Rev. B* **99**, 155124, DOI: <https://doi.org/10.1103/PhysRevB.99.155124> (2019).
42. Pottelberge, R. V. Comment on “electron states for gapped pseudospin-1 fermions in the field of a charged impurity”. *Phys. Rev. B* **101**, 197102, DOI: <https://doi.org/10.1103/PhysRevB.101.197102> (2020).
43. Han, C. D., Xu, H., Huang, D. & Lai, Y. C. Atomic collapse in pseudospin-1 systems. *Phys. Rev. B* **99**, 245413, DOI: <https://doi.org/10.1103/PhysRevB.99.245413> (2019).
44. Zhang, Y. C. Wave function collapses and $1/n$ energy spectrum induced by a coulomb potential in a one-dimensional flat band system. *Chin. Phys. B* . . ., DOI: <https://iopscience.iop.org/article/10.1088/1674-1056/ac3653> (2021).
45. Zhang, Y. C. Infinite bound states and $1/n$ energy spectrum induced by a coulomb potential of type iii in a flat band system. . . ., DOI: <https://www.researchgate.net/publication/356190293> (2021).
46. Zhang, Y. C., Song, S., Liu, C. F. & Liu, W. M. Zitterbewegung effect in spin-orbit-coupled spin-1 ultracold atoms. *Phys. Rev. A* **87**, 023612, DOI: <https://doi.org/10.1103/PhysRevA.87.023612> (2013).
47. Downing, A., C & Song, M. E., Portnoi. One-dimensional coulomb problem in dirac materials. *Phys. Rev. A* **90**, 052116, DOI: <https://doi.org/10.1103/PhysRevA.90.052116> (2014).
48. Abramowitz, M. & Stegun, I. A. "confluent hypergeometric functions." ch. 13 in handbook of mathematical functions with formulas, graphs, and mathematical tables. New York: Dover (1972).
49. Wang, Z. X. & Guo, D. R. Special functions. World Scientific, Singapore (1989).
50. Landau, L. D. & Lifshitz, E. M. Quantum mechanics: non-relativistic theory. Pergamon Press, Third Revised Edition (1977).

Acknowledgements (not compulsory)

Y.C. Z. are supported by the NSFC under Grants No.11874127 and startup grant from Guangzhou University.

Author contributions statement

Y.C.Z. performed calculations. Y.C.Z. analyzed numerical results. Y.C.Z. contributed in completing the paper.

Additional information

Competing interests: The authors declare no competing interests