Liana functional assembly along the hydrological gradient in Central Amazonia

Elisangela da Xavier Rocha (✉ elisangelarocha.xavier@gmail.com)

Instituto Nacional de Pesquisas da Amazonia <https://orcid.org/0000-0002-2348-3872>

Anselmo Nogueira

Universidade Federal do ABC

Flávia Costa

INPA: Instituto Nacional de Pesquisas da Amazonia

Robyn Burnham

Michigan State University

Caian Gerolamo

Universidade de São Paulo: Universidade de Sao Paulo

Cibelly Honorato

INPA: Instituto Nacional de Pesquisas da Amazonia

Juliana Schietti

Universidade Federal do Amazonas

Research Article

Keywords: Functional traits, plant diversity, species composition, tropical forest, water-use efficiency, wood density

Posted Date: December 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1083529/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Soil hydrology, nutrient availability and forest disturbance determine the variation of tropical tree species composition locally. However, most habitat filtering is explained by tree species' hydraulic traits along the hydrological gradient. We asked whether these patterns apply to lianas. At the community level, we investigated: (i) whether the hydrological gradient, soil fertility and forest disturbance explain liana species composition; and (ii) whether differences in leaf and stem wood functional traits were linked to species composition along ecological gradients. We sampled liana species composition in 18 1-ha plots across a 64 km² landscape in Central Amazonia and measured ten leaves and stem wood traits across 115 liana species in 2,000 individuals. We correlated liana species composition summarized with PCoA with the functional composition summarized by PCA, considering the species mean values of traits at the plot level. We tested the relationship between ordination axes and the environmental gradients. Liana species composition was highly correlated with functional composition. Taxonomic (PCoA) and functional (PCA) composition were strongly associated with the hydrological gradient, with a slight impact of forest disturbance on functional composition. Species at valley areas had higher stomata size and higher proportions of self-supporting xylem than plateau. Differently, lianas on plateaus invest more in fast-growing leaves (higher SLA), although with a higher wood density. Our study reveals that lianas use different functional solutions in dealing with each end of the hydrological gradient and that the relationships between habitat preferences and traits explain lianas species distributions not straightforwardly as previously found for trees.

Introduction

Tropical climates support high richness and morphological diversity in different groups of woody plants. In tropical forests, woody climbing plants (lianas) account for about 25% of the abundance and species diversity of woody stems (Gentry 1991; Schnitzer and Bongers 2011), playing an essential role in forest dynamics (Schnitzer and Bongers 2002), carbon sequestration (Chen et al. 2015), and as food and canopy pathways for animals (Putz and Mooney 1991; Schnitzer and Bongers 2002). Despite this importance, biologists only recently began to unveil the liana's distribution pattern in tropical forests. On continental or global scales, high liana abundance and richness has been associated with lower mean annual precipitation or greater seasonality (Schnitzer 2005; DeWalt et al. 2015). In local and landscape scales, high liana abundance and richness are associated with lower soil sand content (Nogueira et al. 2011), higher availability of soil nutrients, such as phosphorus (Malizia et al. 2010; Ledo et al. 2016) and heterogeneous forest structure or disturbances such as large-sized canopy gaps (Schnitzer and Carson 2010; Nogueira et al. 2011).

These relationships are based on the collective of liana assemblages. However, little is known about species level responses, which may differ widely from average assemblage responses (Rocha et al. 2020). Moreover, surprisingly, few studies have investigated the role of the soil water availability on the spatial distribution of liana species in tropical forests (Schiatti et al. 2014), which may derive from the assumption that lianas have higher efficiency in obtaining and transporting water in comparison with

trees (Yong et al. 2012; Chen et al. 2015). In this case, lianas would establish themselves independently of the local water availability. However, the uneven distribution of liana species along a local water gradient (e.g., Schietti et al. 2014) suggests that lianas' hydraulic efficiency and security varies across species, as was found for trees (Oliveira et al. 2019). Soil water, nutrient availability, and forest disturbance rates should affect the distribution and maintenance of lianas at the landscape scale.

The liana life-form has unique morphological features functionally linked to its success and is distinct from other woody plants in tropical forests (Schenck 1983; Ewers 1985; Ewers et al. 1990; Angyalossy et al. 2015). For example, liana stems have a high proportion of soft tissues (vascular parenchyma and secondary phloem) and a low percentage of fibers (Angyalossy et al. 2015), generating a stem structure with a lower bending stiffness (Rowe et al. 2004; Gerolamo et al. 2020). These traits, in combination with a high growth rate in high light environments (Baars and Kelly 1996), is responsible for the remarkable capacity of some liana species to survive and resprout after falling from the canopy or suffering crown damage (Putz 1984; Rocha et al. 2020). In addition, many liana species have xylem with larger vessels that produce a highly conductive vascular system extending from the roots to distal branches and leaves via a long pathway (Ewers et al. 1991; Schnitzer 2005; Gerolamo and Angyalossy 2017). A well-developed root system in lianas has been argued to ensure deep-water access and a more efficient stomatal aperture control in leaves as a strategy to reduce the risk of collapse of xylem water transport (Campanello et al. 2016; Schnitzer 2018). High hydraulic efficiency is correlated with fast growth in lianas (Schnitzer 2005). Although as a collective life-form, lianas may be characterized as efficient resource exploiters and fast-growing, it is expected that their functional traits vary widely across species. For example, variability among liana species has been documented in the ability to regenerate axillary buds after a fall (Rocha et al. 2020), as well as in potential hydraulic conductivity (Gerolamo and Angyalossy 2017). We postulate that the considerable, yet underexplored, variation of functional traits within liana assemblages could explain the differential distribution of liana species in tropical forest landscapes (e.g., Vivek and Parthasarathy 2018).

In Central Amazonia, the vertical distance to the nearest drainage, a proxy for the water-table depth, is a robust hydrological predictor of plant species composition, explaining the variation of plant assemblages across the landscape (Schietti et al. 2014). The water-table depth gradient may provide a primary habitat axis of plant species differing in water use requirements (Silvertown et al. 1999; Balvanera et al. 2011). This gradient would lead to differentiation of local species composition according to the hydrological niches of species (Jirka et al. 2007; Silvertown et al. 2015; Cosme et al. 2017; Oliveira et al. 2019), and the expected trade-off between cavitation resistance and water transport efficiency (van der Sande et al. 2019), which should be reflected in species morphological and anatomical traits. If true, we expect this primary gradient to be strongly correlated with the variation in liana taxonomic and functional composition, followed by secondary ecological axes such as edaphic and forest disturbance gradients.

This study investigated the distribution pattern of liana species and their functional traits by sampling lianas in permanent plots across a 30 km² landscape in a hyperdiverse Amazonian rainforest. We evaluated leaf and stem wood functional traits related to light capture, water transport, nutrient

conservation, and mechanical support to address the following questions: (i) Are changes in the species composition of liana assemblages shaped by hydrological, edaphic, and forest disturbance gradients? (ii) Do changes in functional trait composition parallel changes in floristic composition along the ecological gradients? (iii) which traits covary along the dominant environmental gradients? We hypothesized that liana species composition is primarily explained by hydrological gradient and secondarily by soil fertility and forest disturbance gradients in this moist tropical forest. Since species-environment relationships are presumed to be mediated by functional traits (Suding et al. 2008), we hypothesize that the combination of liana species in portions of the three distinct ecological gradients is mainly caused by leaf and stem wood functional traits varying across species.

Materials And Methods

Study site

The study was conducted in the Reserva Florestal Ducke (hereafter Ducke Reserve) of the Instituto Nacional de Pesquisas da Amazônia (INPA) in Central Amazonia, located 26 km north of Manaus (2°55' S, 59°59' W at reserve headquarters), Brazil. Ducke Reserve is a Brazilian Long-Term Ecological Research (LTER) site where ecological data on topography, soil properties, and species abundance have been systematically collected in permanent plots (<https://ppbio.inpa.gov.br>). The average annual temperature from 1966 to 2018 was *c.* 26°C, and the annual rainfall was 2572 mm with a monthly maximum of 331.4 mm in April and a minimum of 98.8 mm in August (Ducke Reserve weather station data). The dry season occurs between July and September, but on average, only two months have rainfall less than 100 mm (Marques-Filho et al. 1981). Soils are derived from tertiary marine sediments of the Alter do Chão formation (Chauvel et al. 1987).

The predominant vegetation is terra firme forest. The well dissected relief is formed by different topographic levels, slopes, soil types, and hydromorphic conditions that range from plateau to valley (Chauvel et al. 1987). This well dissected relief is typical of the old soils that originated on the Guiana Shield in the Central Amazon region (Sombroek 2000). The plateaus are the highest areas (elevation above sea level between 90 and 120 m) with clayey, well-drained and generally more fertile soils (Hopkins 2005) with water table *c.* 20 m deep. In these portions of the forest, it is possible to find the largest and probably the oldest trees (30-40 meters), including emergent trees such as *Dinizia excelsa* Ducke, reaching 60 meters in height (Hopkins 2005). As the altitude decreases, there is a transition of environments and the clay content of the soil decreases while the sand gradually increases. Valleys are topographic portions of riparian areas with sandy and generally less fertile soils at lower altitudes, although phosphorus content may vary among valleys. Valleys are also mostly flat or gently sloping with the water table close to the surface (as shallow as 1 m depth) and seasonally waterlogged during the rainy season (Drucker et al. 2008; Tomasella et al. 2008). The height of the canopy in the valley is lower than in plateau regions and includes many palm trees, such as *Oenocarpus bataua* Mart. and *Mauritia flexuosa* L.f. (Kahn and Castro 1985, Hopkins 2005).

Sampling Design And Liana Species Composition

Lianas were sampled in 18 plots at least 1 km from one another, distributed over a 5 x 6 km grid (30 km²) and spanning a range of altitudes from 39 to 140 m (Ribeiro et al. 1999). Each plot is 250-m long and 40-m wide following the RAPELD sampling system (<https://ppbio.inpa.gov.br/en/Methods/RAPELD>). The center line follows the terrain level isoline to minimize soil and water-table depth variations inside each plot (Magnusson et al. 2005; Costa and Magnusson 2010; Costa et al. 2020).

Lianas were inventoried in 2004 and recensused in 2014 (Nogueira et al. 2011; Gerolamo et al. 2018). All lianas (individual stems) with a diameter (D) ≥ 5 cm at 1.3 m above their rooting point were sampled in the 1 ha plot (250 x 40). Thinner lianas ($1 \text{ cm} \leq D \leq 4.9 \text{ cm}$) were sampled in subplots of 0.25 ha (250 x 10 m). In a previous study, 32 species of the tribe Bignonieae (Bignoniaceae) were investigated for species composition in the permanent plots of the Ducke Reserve (Schiatti et al. 2014). In the present study, we expanded species identification to all individuals of lianas marked in the previous censuses. We collected vouchers of specimens for botanical identification between 2016 and 2019. Botanical specialists assisted plant identification mainly based on vegetative characters such as phyllotaxy, stem form, wood anatomy, bark odour, colour, texture, lenticel distribution, leaf type/parts, leaf form, venation, presence of exudate, and anchoring structures. Fertile specimens were also considered when available. Liana species composition data are available at ppbio.inpa.gov.br/en/home and at forestplots.net.

Functional plant traits

We selected ten leaf and stem wood traits related to light capture, water transport, and nutrient conservation (see below) to describe the functional variation of each liana species (see plant traits studied in Table 1). In the field, we chose one individual per species per plot to collect plant samples, and in total, 3 to 10 individuals were sampled per species. We carefully selected stems growing approximately 1 m from the plant apex and multiple leaves from the liana's crown on the sunnier side in each individual. Leaves and stems were sampled, placed in sealed plastic bags, and kept moist until processed in the Ducke Reserve laboratory.

Table 1

Description of functional plant traits used to characterize each liana species assembly at the Ducke Reserve (Brazil).

Trait	abb.	Description	Related function*
Leaf dry matter content	LDMC	Oven-dry mass of a leaf divided by its water saturated fresh mass (mg g ⁻¹)	Leaf tissue density, leaf physical resistance, stress tolerance
Leaf size	LS	Mean area-leaf (cm ²)	Photosynthetic capacity; leaf temperature
Leaf thickness	LT	Leaf blade thickness (μm)	Resistance and protection
Specific leaf area	SLA	One-sided area of a fresh leaf divided by its oven-dry mass (cm ² g ⁻¹)	Photosynthetic rate, leaf longevity, relative growth rate
Stomata density	SD	Stomata density (no. mm ⁻²)	Determine maximum leaf diffusive (stomatal) conductance of CO ₂ (g_{max}) to sites of assimilation
Stomata size	SS	Mean stomata length (μm)	
Vein density	VLA	Vein length per unit area (veins mm ²)	Protection and water transport in plants.
Lianescent xylem	Pxl	Fraction of the xylem area (or proportion) with the characteristics of the lianescent vascular syndrome (%)	Flexibility and greater water transport potential.
Self-supporting xylem	Pxs	Fraction of the inner area (or proportion) of the xylem with a small diameter vessel and more fibers developed in the self-supporting phase (%)	Water transport and mechanic support
Vessel density	VD	Vessel density (vessels mm ²)	Xylem cells concerning long distance water transport in plants.
Wood density	WD	Wood density (g cm ³)	Highly interactive trait of wood
* Carlquist (1975); Farquhar, Ehleringer and Hubick 1989; Grime et al. 1997; Weiher et al., 1999; McIntyre et al., 1999; Westoby et al., 2002; Cornelissen et al., 2003; Garnier and Navas 2012; Sack and Scoffoni 2013; Angyalosy et al., 2015.			

Leaf traits – We selected leaves that were mostly intact and without injury for measuring leaf functional traits. Three mature leaves per stem were scanned with a scanner (HP Scanjet 200) and processed to calculate leaf area (cm²) using the ImageJ software (National Institutes of Health, Bethesda, MD, USA; <http://rsb.info.nih.gov/ij/>). Leaf thickness (mm) was measured with a digital micrometer (accuracy 0.001

mm). We also measured the fresh leaf weight (g) and the dry leaf weight (g) on an electronic balance accurate to four decimal places. For dry weight, fresh leaves were dried at 60°C in a dehydration oven for at least three days until a constant weight was recorded. We also calculated the leaf water content (LWC; %), leaf dry matter content (LDMC; mg g^{-1}) and specific leaf area (SLA; $\text{cm}^2 \text{g}^{-1}$) following Cornelissen et al. (2003).

Two additional mature leaves were fixed in FAA (formaldehyde, acetic acid, and 50% ethanol, Johanson 1940) in the field, and after 48 hours, preserved in 70% alcohol. In the laboratory, we cut two one-centimeter squares from the leaves, avoiding the central vein, to evaluate stomatal density (SD; number mm^2), stomatal size (SS; μm), and vein length per unit area (VLA; also known as “vein density”). In one leaf square, the leaf epidermis was dissociated for stomata detection, with immersion in a solution of glacial acetic acid and hydrogen peroxide (1:1 v/v) at 60 °C for 48h for most species. Some species (e.g., *Davilla rugosa*) required more time (Franklin 1945 modified by Kraus and Arduin 1997). Then, leaf samples were washed with distilled water, stained with 1% safranin, and mounted as semi-permanent slides with glycerin (Bukatsch 1972 modified by Kraus and Arduin 1997). Photographs were taken using a light microscope (Leica DM500, Leica Microsystems, Wetzlar, Germany) under 200x magnification. The stomatal density (SD) and stomatal size (SS) were measured and counted in photos over an area of 0.45 mm^2 in each of the two leaves using the ImageJ software. Stomatal size was defined as the length in micrometers between the junctions of the guard cells at each end of the stomata. To determine leaf vein density, we immersed a five mm^2 leaf square of the second leaf cut in 20% sodium hydroxide solution until depigmentation occurred (Arnott, 1959). Then, we washed the samples in distilled water, photographed them in a stereomicroscope Leica (Leica M205 C, Leica Microsystems, Wetzlar, Germany), and estimated the vein density (VLA) using the leaf GUI software (<http://www.leafgui.org/>).

Stem wood traits – Using the same stems from which the leaves were obtained for each plant, we cut three stem samples of 0.5-2 cm in diameter and 2 cm in length from the basal-most stem. The first sample was used to measure the cross-sectional area of stem tissues, the second for microscopic analyses of secondary xylem, and the third sample to estimate wood density.

To estimate the percentage of each stem tissues' cross-sectional area relative to the total area of the stem, we initially polished each stem sample with a decreasing series of sandpaper particle sizes (Barbosa et al. 2021). Then, we scanned the cross-section area of stems on the high-resolution scanner indicated above and analyzed tissue distribution using ImageJ software. The cross-sectional stem area of each sample was categorized and measured as four different regions: pith, two parts of the secondary xylem (inner-formed dense xylem developed in juvenile and supposedly self-supporting phases, and lianescent xylem with wide vessels developed in climbing phases) and bark (all tissues outside of the vascular cambium). See figure 1 for details about the four tissues categorized in the cross-sectional stem area. Recognition and classification of stem tissues and anatomical patterns of lianas were based on the descriptions of Angyalossy et al. (2015).

For microscopic xylem measurements, the stem sample was fixed in FAA (formaldehyde, acetic acid, and 50% ethanol, Johanson 1940) for at least 12 hours and then preserved in 70% alcohol. In the laboratory, cross-sections of each sample's secondary xylem (three to five individuals per species) were obtained with a sliding microtome, stained in 1% astra blue and 1% safranin (Bukatsch 1972, as modified by Kraus and Arduin 1997) to document cells with cellulose and those with cellulose and lignin. Permanent anatomical slides were prepared with a synthetic resin and then photographed with a light microscope (Leica DM500, Leica Microsystems, Wetzlar, Germany) at 200x magnification. To characterize each individual's secondary xylem, we estimated the vessel lumen diameter (d in μm), measuring at least 30 vessel elements and the number of vessels per square millimeter (VD) using at least two areas of 1 mm^2 . Both traits were measured using ImageJ software for each individual and then averaged by species.

We estimated each stem sample's wood density by first removing the bark and determining the fresh xylem volume using the water displacement method. Then each sample was dried for at least 72 hours at 105°C and weighed until a constant weight was recorded. Wood density ($\text{WD} - \text{g cm}^3$) was determined as the ratio between the dry wood mass and the fresh wood volume (including pith).

Ecological Gradients: Hand, Soil Fertility And Tree Turnover

Height above the nearest drainage (HAND) in the Ducke Reserve quantitatively describes the vertical distance of each plot to the nearest drainage. HAND is an indirect estimate of the water-table depth derived from the algorithm of Rennó et al. (2008) from the Shuttle Radar Topography Mission (SRTM) digital elevation model, with 90 m of spatial resolution (Schietti et al. 2014). High HAND values correspond to high elevation plots, associated with ridges and plateaus and vertically far from the water table. In contrast, low HAND values represent plots in lowland areas closer to the water table in riparian (valley) areas. Soil texture and the sum of exchangeable cations (calcium, magnesium, potassium, and sodium) of each plot are highly correlated with HAND (Pearson $r = 0.58$, $P < 0.001$, $n = 30$; $r = 0.57$, $P = 0.01$, $n = 18$, respectively) at the Ducke Reserve. Plots far from the water table (plateaus) have more clayey and cation-rich soils than plots close to the water table. Therefore, the HAND gradient incorporates other correlated gradients, such as soil texture and fertility given by the sum of cations and the vertical access to water (<https://ppbio.inpa.gov.br>). Soil concentration of available phosphorus (P ; mg kg^{-1}) is not correlated to the HAND gradient ($r = -0.21$, $P = 0.39$, $n = 18$). The phosphorus limitation in tropical soils can be an important axis in niche segregation (Condit et al. 2013). We used available phosphorus (P) concentration to characterize the soil fertility independent of the HAND gradient and explain the compositional changes of liana species at the Ducke Reserve.

We also investigated the role of forest dynamics on the distribution of liana species. Our metric of forest dynamics was tree turnover, calculated as the average annual rate of tree mortality and recruitment (Phillips and Gentry, 1994). Tree turnover estimation was based on dead and recruited trees with $\text{DBH} > 10\text{ cm}$ over the five years of tree censuses between 2003 and 2009 (Castilho et al. 2010). Because we do not have tree turnover data that precisely match the liana census time interval, we assumed that our

estimates of tree turnover rates reflect the forest's intrinsic properties and tendency to maintain similar levels for a relatively long period.

Statistical Analyses

We reduced the dimensionality of the species composition of lianas using Principal Coordinate Analysis (PCoA). Ordinations were based on species occurrence data (presence or absence of species) and relative species abundance data. In the first ordination, we applied the Sørensen dissimilarity index using species occurrence data. In the second ordination, we applied the Bray-Curtis dissimilarity index. In both cases, only the first two axes of PCoA were used in subsequent analyses to test the relationship between the liana assemblage composition and the three environmental gradients (HAND, soil P, and tree turnover). These two axes captured most of the variation of the original species composition.

To explore the variation in the liana assemblage's functional traits relative to each plot, we also performed two Principal Component Analyses (PCA), one based on the species occurrence and the other on species relative abundances. In both multivariate analyses, we used the average of the eleven leaf and stem traits per species. The first PCA used the simple average of each trait across species found in each plot (qualitative functional PCA). The second PCA used average trait values per species weighted by each species' abundance in each plot (quantitative functional PCA). In both cases, only the first two axes of PCA were used in the subsequent analyses to test the relationship between the functional composition and the three environmental gradients (HAND, soil P and tree turnover). The two axes captured most of the variation in functional composition. A correlation matrix between trait means and the first PCA axes is available in Supporting Information (Table S1).

The first two axes of the taxonomic-based PCoA (PCoA-1 and 2) and functional-based PCAs (PCA-1 and 2) were used independently as response variables in our statistical modelling. We tested the relationship between the response variables (PCoA and PCA axes) and the three ecological gradients (HAND, soil P, and tree turnover). In all models, the HAND gradient was log-transformed to ensure linearity. We used a model selection approach for each response variable to evaluate the minimally adequate model from a set of models that included a null model (only the intercept without ecological gradients as fixed factors). We considered a set of models with all possible combinations of the three ecological gradients. The minimally adequate model had the lowest Akaike Information Criterion (AIC) value, and models with $\Delta AIC < 2$ are similarly plausible. We also inspected the three ecological gradients' multicollinearity using the Variance Inflation Factor (VIF, Dormann et al. 2012), and the VIF values were always less than two in all cases.

Finally, we evaluated whether functional composition changes parallel floristic composition changes by performing a simple regression between the PCoA and PCA axes.

All statistical analyzes were performed in R program version 3.4.0 (R Core Team, 2017). PCoA and PCA were run with *vegdist* and *prcomp* functions of the *vegan* package (Oksanen et al. 2015). Model selection

was performed with the *dredge* function of the Mumin package (Barton 2014).

Results

Species composition patterns along the ecological gradients

We sampled 2,000 individuals of 115 species of lianas distributed in 22 families, of which the richest families were (in order): Bignoniaceae, Fabaceae, Celastraceae, Connaraceae, and Dilleniaceae. The first two axes of the qualitative PCoA ordination captured 57% of the variance in species composition (Table 2; supporting information; Table S3), and the first two axes of the quantitative PCoA ordination captured 73% (supporting information; Table S4).

Table 2

Model selection results for liana species composition across 18 plots covering three distinct ecological gradients at Ducke Reserve, Central Amazonia. The response variable used here was the first axis of qualitative PCoA (PCoA-1), describing the liana species composition. The analysis of the second PCoA axis is available in S3 (supporting information). Gaussian error distribution was used in all models, including none or even the three fixed factors: log (HAND), phosphorus soil content, and tree turnover gradients.

Models	Response variable	Fixed factors	K	AIC	Δ AIC
1	Qualitative	HAND	3	-21.2	0.00
2	PCOA1	HAND + Tree turnover	4	-19.4	1.75
3	(47%) *	HAND + Phosphorus	4	-19.2	1.94
4		HAND + Phosphorus + Tree turnover	5	-17.5	3.65
5		1	2	6.9	28.08
6		Phosphorus	3	7.8	28.94
7		Tree turnover	3	8.5	29.67
8		Phosphorus + Tree turnover	4	9.2	30.36
*Proportion of variance					

Liana species composition represented by the qualitative PCoA-1 axis was only related to the HAND gradient (model 1 of Table 2, Fig. 2). There was a gradual substitution of species from valleys to plateaus, clearly described by the HAND gradient, as illustrated in Figure 3 for the two most representative liana families in the Amazon Forest (Fabaceae and Bignoniaceae). High HAND plots (plateaus) had several exclusive species such as the legumes *Mimosa guilandinae*, *Senegalia altiscandens*, and *Clitoria leptostachya* (Fig. 3A), the Bignoniaceae *Pleonotoma longiflora*, plus several *Adenocalymma* species (Fig. 3B). Exclusive species in the low HAND plots (valleys) were *Fridericia japurensis* and *Anemopaegma robustum* (Fig. 3B). The second composition axis was not associated with any of the ecological

gradients, and the null model was the minimum adequate model selected for the qualitative PCoA-2 axis (Table S3). The results based on the quantitative PCoA axes were similar to those presented here for qualitative data (supporting information; Table S4).

Functional Composition Patterns Along The Ecological Gradients

The first two axes of the qualitative functional PCA captured 49% of the functional species composition variation, while the first two axes of quantitative functional PCA captured 59%. Along the qualitative PCA-1 axis, the average community values of stomatal size and self-supporting xylem increased, while leaf size, specific leaf area, and wood density decreased. Along the qualitative PCA-2 axis, stomatal density, vessel density, and leaf dry matter content increased while leaf thickness decreased (Fig. S1 and Table S1). The quantitative PCA axes were correlated with leaf and stem functional traits similar to the qualitative PCA axes.

The qualitative functional PCA-1 axis was negatively related to the HAND gradient but not the other environmental predictors (model 1 of Table 3 and Fig. 4A). In contrast, the qualitative functional PCA-2 axis was not related to any predictor. Thus, in low HAND plots (valleys), liana species had higher mean values of the area of self-supporting xylem (Pxs, Fig. 4B) and stomata size (SS), and lower mean values of leaf size (LS, Fig. 4C), specific leaf area (SLA, Fig. 4D), and wood density (WD). The quantitative PCA axes had a similar relationship to the environmental predictors as the qualitative PCA axes.

Table 3

Model selection results for liana species composition across 18 plots covering three distinct ecological gradients at Ducke Reserve, Central Amazonia. The response variable used here was the first axis of qualitative PCA (PCA-1), describing the liana species composition. The analysis of the second PCA axis is available in S6 (Supporting information). Gaussian error distribution was used in all models, including none or even the three fixed factors: log (HAND), phosphorus soil content, and tree turnover gradients.

Models	Response variable	Fixed factors	K	AIC	Δ AIC
9	First Axis	HAND	3	63.2	0.00
10	PC1	HAND + Tree turnover	4	64.9	1.77
11	(27.5%) *	HAND + Phosphorus	4	65.2	2.00
12		HAND + Phosphorus + Tree turnover	5	66.9	3.76
13		1	2	73.3	10.12
14		Phosphorus	3	74.7	11.54
15		Tree turnover	3	75.3	12.12
16		Phosphorus + Tree turnover	4	76.7	13.54
*Proportion of variance					

These functional changes of the qualitative PCA-1 axis were strongly correlated with taxonomic species composition changes captured in the qualitative PCOA-1 axis ($r^2 = 0.53$, Fig. 5).

Discussion

We asked whether changes in the species composition of liana assemblages are shaped by hydrological, soil fertility, and forest disturbance gradients and if functional trait composition changes parallel the gradients. Following our predictions, we found that: (i) liana species composition was strongly correlated with the hydrological gradient represented by the terrain height above the nearest drainage (HAND), a proxy for the water-table depth; (ii) the turnover of liana species along the hydrological gradient is strongly correlated with differences in leaf and stem wood functional traits across species, i.e., the taxonomic compositional turnover of lianas in space was strongly linked to the functional change. Thus, leaf and stem wood trait differences explained the strong association between liana species composition and the hydro-edaphic gradient on the old dissected terrains distributed in the Central Amazonia rainforest.

Water-table depth explains landscape changes in liana species composition

Our results support the hypothesis that soil hydrology shapes local liana floristic patterns within Amazonian terra firme forests. We explained 81% of the species composition variation with only one predictor, the water-table depth gradient (HAND). Simultaneously, soil fertility (P content) and forest dynamics (tree turnover) had only marginal effects in secondary models. Our results expand previous findings for the tribe Bignoniaceae in this same forest (Schietti et al. 2014; Gerolamo et al. in review), increasing the generality of the distribution pattern along the water-table depth gradient to range of taxonomic identities among the liana assembly. Previous studies have detected the filtering effect of the water table depth on Amazonian tree species composition (Jirka et al. 2007; Schietti et al. 2014) and linked that to the selection of species-level functional traits (Cosme et al. 2017; Oliveira et al. 2019; Schmitt et al. 2020). To our knowledge, this is the first study to show that liana species are also filtered along a hydrological gradient.

The turnover of liana species composition from valleys to plateaus indicates that variation in the soil water conditions plays an essential role for liana species distribution, similar to the patterns observed for herbs, trees, and palms (Costa et al. 2005; Costa et al. 2009; Schietti et al. 2014). This pattern is surprising because lianas have been generically described as very efficient organisms in obtaining and transporting water, which has created the expectation that lianas would establish independently from the local water availability, the opposite of what we found here. However, in the Ducke Reserve, valleys have sandy soils close to the drainage and are strongly influenced by waterlogging during the rainy season (Castilho et al. 2006; Toledo et al. 2017). Despite the expected rapid drainage of sandy soils, the constant inflow of groundwater from higher topographic positions generates a superficial water-table in the valleys (Hodnett et al. 1997). Consequently, roots remain superficial and in contact with waterlogged soils during the wet season or year-round, depending on the micro-relief. Waterlogged soils have hypoxic to anoxic conditions that should restrict the colonization of plants without appropriate functional traits (Larcher 2000, Parolin et al. 2003). Thus, species composition in these environments is expected to be composed of hydro-edaphic specialist species (Kalliola and Puhakka 1988; Muscarella et al. 2019), capable of coping with these extreme soil conditions (e.g., Cosme et al. 2017; Oliveira et al. 2019). As soils become deeper and drainage better towards plateaus, the species composition changes and many liana species are found co-occurring. This pattern is also observed in herbs and woody plants with other life-habits (Schietti et al. 2014, Muscarella et al. 2019). Therefore, our results confirm that soil hydrology is an essential factor determining local plant assembly variation, independent of plant habit, in Amazonian terra firme forests.

Patterns of functional composition change: are species-environment relationships mediated by functional traits?

Our results revealed that changes in the floristic composition of lianas along the hydrological gradient paralleled changes in functional trait composition, highlighting that liana assemblies are composed of species with different ecological strategies.

The plant economic spectrum proposes an organization of global trade-offs between resource acquisitive and conservative plant strategies (Wright et al. 2004; Chave et al. 2009), which are mainly influenced by resource (soil water and nutrients) availability (Aerts and Chapin 2000; Sandel et al. 2010). Along the economic spectrum, plants with a fast-growing acquisitive strategy based on cheap tissue investments can dominate in moist and fertile areas (Grime et al. 1997; Reich et al. 1999). On the other side of the spectrum, plants with a conservative strategy grow slowly with expensive tissue investments (Wright et al. 2004; Reich 2014), minimizing nutrient loss and increasing competitive abilities in dry and nutrient-poor environments (Hobbie 1992; Aerts 1995). In Central Amazonia, previous studies revealed that tree species in valley areas had more acquisitive strategies (e.g., lower WD and higher SLA), while those on plateaus had a more conservative strategy (Cosme et al. 2017; Toledo et al. 2017). Similar results were found for tree species in French Guiana (Schmitt et al. 2020). We detected similar trends for liana stem traits, which were more acquisitive (lower WD) in valleys than plateaus.

We emphasize the importance of stem wood traits, still poorly investigated, to explain the liana species turnover patterns along the hydro-edaphic gradient. The lower investment in WD and higher proportions of self-supporting xylem in shallow water-table areas of valleys may guarantee the needed fast growth in height during the self-supporting phase of lianas in this more dynamic portion of the forest. In fact, lianas in this portion of the hydrological gradient (valleys) in the Ducke Reserve have a higher recruitment rate (Gerolamo et al. 2018). The sandy and waterlogged soils of valleys provide poor root anchorage, which, together with the shallow root systems (Fan et al. 2017), leads to higher tree uprooting and mortality than in the clayey slopes and plateaus (Toledo et al. 2011). Although a more dynamic environment is more beneficial for liana development (Schnitzer et al. 2021), valleys have a higher density of palms, including canopy palms (Kahn and Castro 1985), which are poor supports for lianas (Nogueira et al. 2011). Therefore, valleys are less suitable for establishing lianas both in the understory and the forest canopy. This inference is supported by the lower density and diversity of lianas found in this portion of the hydrological gradient (Gerolamo et al. 2018; Gerolamo et al. in review). Under such conditions, stem traits allowing a fast-growth strategy and a higher investment in self-supporting xylem are essential for liana success. Self-supporting xylem appears at early stages in liana stems' wood development in the understory and on searching branches when stems are still elongating before reaching climbing support (Ewers et al. 1991). After attaching to a support tree, the cambium activity produces a unique secondary tissue known as lianescence xylem that strongly decreases stem stiffness, allowing the unique bending and twisting properties of lianas to climb towards the canopy (Rowe et al. 2004; Angyalossy et al. 2012, 2015). In sum, we argue that the correlation between higher proportions of self-supporting xylem and lower WD in shallow water tables confirms the hypothesis that some liana species have hydraulic-biomechanical adaptations conferring success in a dynamic habitat where seasonal soil waterlogging and a scarcity of adequate host trees for liana's climbing may exist (Oliveira et al. 2019; Gerolamo et al. 2020).

Our results show pairing between stem wood traits and leaf anatomy in the liana assemblage along the hydrological gradient. In valley areas, we found larger mean assemblage stomata size than in plateaus, favoring higher conductivity of the hydraulic system (large stomata and lower WD) under more favorable

conditions. The high specific hydraulic conductivity of lianas could achieve rapid diffusive conductance, allowing higher photosynthetic rates and water use efficiency (Franks et al. 2009; Zhang et al. 2012). On the other hand, the opposite is expected in habitats with low water availability as plateau, with plant species having higher stomata density and smaller stomata size (Spence et al. 1986, Xu and Zhou, 2008). Small stomata may limit water loss, keeping xylem tension within a safe range and reducing the risks of xylem dysfunction (Brodribb and Holbrook 2003; Johnson et al. 2011; Zhang et al. 2014). This morpho-functional mechanism has been used to explain the greater abundance of lianas in more seasonal forests (Schnitzer, 2018; but see the review Amorin et al. 2018), and liana and tree ecological strategies within a wide range of soil water availability in tropical rainforests.

We did not detect correlations between foliar traits such as SLA and LA and stem wood traits in the liana assemblage as expected by the plant economic spectrum. In contrast, tree species coordinate more acquisitive foliar and stem traits at the same site in valleys (higher SLA and lower WD) and the opposite in plateaus (Cosme et al. 2017), while lianas had more acquisitive stem traits (lower WD) but more conservative traits (lower SLA and LA) in valleys. This decoupled pattern agrees with a correlational study in Central Panama, showing that the SLA was poorly correlated with hydraulic traits across lianas species (Van der Sande et al. 2019). In this study site, lianas have more acquisitive traits than trees generally associated with higher light and nutrient resource niches (Van der Sande et al. 2019; Liu et al. 2021). Decoupling may arise under ecological conditions acting differentially on plant organs as observed on the liana assemblage. Higher nutrient availability may sustain SLA values that are comparatively higher in plateaus than valleys given that more fertile soils are associated with high values of foliar functional traits (Liu et al. 2012; Trujillo et al. 2021). Therefore, it appears reasonable that the higher nutrient availability in plateaus could explain the more acquisitive leaves associated with this portion of the hydroedaphic gradient. Our results highlight that this unique combination of functional traits on each end of the hydroedaphic gradient is probably associated with the joint effect of multiple environmental factors in liana assemblages.

In sum, the stem and leaf functional trait decoupling at each extreme of the hydrological gradient highlights that local habitat may optimize trait composition that violates, at least partially, the principles of optimization of resource acquisition and conservation (Reich et al. 2014). At the local scale, this violation disappears when average values of traits observed by this research are compared with regional and global values (Ordoñez et al. 2009), indicating that lianas have a typical acquisitive strategy. The acquisitive strategy shared by many lianas species is composed of alternative sets of features (more or less acquisitive) depending on the portion of the ecological gradient where they are found. The local ecological conditions associated with this pattern may be explained by the high dynamism and low liana support availability in valleys versus the understory light limitation and seasonal drought of the plateaus. In the valleys, support limitation favors lianas with well-developed self-supporting xylem, contributing to upward growth of the stems until reaching adequate support. The absence of water limitation allows low wood density, permeable stems, and larger stomatal size, which may also respond to the high hydric dynamism in this habitat. At the other extreme, on the plateaus, light limitation in the dense understory may favor lianas with traits capable of rapid ascension, allowed by greater nitrogen availability (Luizão et

al. 2004) and exchangeable bases while the seasonal water deficit may favor denser wood and smaller stomata. Thus, contrary to global observations on plant strategies, the optimization of resource acquisition and conservation in liana species is not easily understood locally (see Liu et al. 2021) until variations in functional design, optimized under a complex set of environmental conditions, are revealed. Thus, the biophysical constraint coupling between traits indicated by Reich (1997, 2014) is more flexible once the larger-scale environmental filters (e.g., climate) have optimized the local range of trait variation, allowing plant survival.

Declarations

Acknowledgements

We are grateful to Jose Francisco for lab support and guidance on anatomical measurements. Erick Lavado and Lorena Rincón provided statistical support and valuable suggestions on the discussion. ER thanks José Raimundo for help in the field. INPA provided transport, and the logistical structure at the Ducke Reserve was provided by PELD and the Brazilian Biodiversity Research Program (PPBio).

Funding

This contribution is part of the doctoral thesis of E.R undertaken at the Instituto Nacional de Pesquisas da Amazônia (INPA), with a fellowship from Coordination of Superior Level Staff Improvement (CAPES). Financial support was received from the Brazilian Long Term Ecological Research Program (CNPq/PELD# 441282/.2016-4) and Project CENBAM- INCT Centro de Biodiversidade Amazônica (FAPEAM/FDB/INPA#003/2012). Additional funds were provided to A.N. by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) through a PDE scholarship (234000/2014-7).

Conflict of interest

The authors declare that they have no conflict of interest

Ethics approval

Not applicable for that section

Consent to participate

Not applicable for that section

Consent for publication

Not applicable for that section

Availability of data and material

The datasets generated for this study are available on request to the corresponding author.

Authors' contributions

ER, AN, JS, RJB and FRC conceived the ideas and methodological design; ER, CG and CH collected the data; AN and RJB identified the plant species. ER, AN, JS and FRC analyzed the data; ER led the writing of the manuscript. All authors contributed to the writing and gave final approval for publication.

References

1. Aerts R (1995) The advantages of being evergreen. *Trends in Ecology* 10:402–407
2. Aerts R, Chapin FS (2000) The mineral nutrition of wild plants revisited. A re-evaluation of processes and patterns. *Adv Ecol Res* 30:1–67
3. Angyalossy V, Pace MR, Lima AC (2015) Liana anatomy: a broad perspective on structural evolution of the vascular system.. In: In: Schnitzer SA, Bongers F, Burnham RJ, Putz FE (eds) *The Ecology of Lianas*. Wiley-Blackwell, Oxford, UK
4. Arnott HJ (1959) Leaf clearings. *Turtox News* 37:192–194
5. Baar R, Kelly D (1996) Survival and growth responses of native and introduced lianes in New Zealand to light availability. *N Z J Bot* 34:389–400
6. Balvanera P, Quijas S, Pérez-Jiménez (2011) Distribution patterns of tropical dry forest trees along a mesoscale water availability gradient. *Biotropica* 43:414–422
7. Barbosa AC, Gerolamo CS, Lima AC et al (2021) Polishing entire stems and roots using sandpaper under water: An alternative method for macroscopic analyses. *Applications in Plant Sciences* 9:15–21
8. Barton K (2014) Package ‘MuMIn’: multi-model inference. R package version 1.13.4. <https://cran.r-project.org/web/packages/MuMIn/index.html>
9. Brodribb TJ, Holbrook NM (2006) Declining hydraulic efficiency as transpiring leaves desiccate: two types of response. *Plant Cell Environ* 29:2205–2215
10. Castilho CV, Magunsson WE, Araújo RNO et al (2006) Variation in aboveground tree live biomass in a central Amazonian Forest: Effects of soil and topography. *For Ecol Manag* 234:85–96
11. Magnusson WE, FJ (2010) Short-term temporal changes in tree live biomass in a central Amazonian Forest, Brazil. *Biotropica* 42:95–103
12. Chauvel A, Lucas Y, Boulet R (1987) On the genesis of the soil mantle of the region of Manaus, Central Amazonia, Brazil. *Experientia* 43:234–241
13. Chave J, Coomes D, Jansen S et al (2009) Towards a worldwide wood economics spectrum. *Ecol Lett* 12:351–366
14. Chen YJ, Cao KF, Schnitzer SA et al (2015) Water-use advantage for lianas over trees in tropical seasonal forests. *New Phytol* 205:128–136
15. Condit R, Bettina MJ, Engelbrecht DP et al (2013) Species distributions in response to individual soil nutrients and seasonal drought across a community of tropical trees. – *Proc. Natl Acad. Sci. USA*

- 110: 5064–5068. <https://doi.org/10.1073/pnas.1218042110>
16. Cornelissen JHC, Lavorel S, Garnier E et al (2003) A handbook of protocols for standardised and easy measurement of plant functional traits worldwide. *Aust J Bot* 51:335–380
 17. Cosme LHM, Schiatti J, Costa FRC, Oliveira RS (2017) The importance of hydraulic architecture to the distribution patterns of trees in a central Amazonian forest. *New Phytol* 215:113–125
 18. Costa FRC, Magnusson WE, Luizão RC (2005) Mesoscale distribution patterns of Amazonian understory herbs in relation to topography, soil and watersheds. *J Ecol* 93:863–878
 19. Costa FRC, Guillaumet JL, Lima AP, Pereira OS (2009) Gradients within gradients: the mesoscale distribution patterns of palms in a central amazonian forest. *Journal Vegetation Science* 20:69–78. <https://doi.org/10.1111/j.1654-1103.2009>
 20. Costa FRC, Magnusson WE (2010) The need for large-scale, integrated studies of biodiversity – The experience of the Program for Biodiversity Research in Brazilian Amazonia. *Natureza & Conservação* 8:1
 21. de Azevedo Amorim T, Nunes-Freitas AF, Rosado BHP (2018) Revisiting the hypothesis for increasing liana abundance in seasonal forests: a theoretical review. *Plant Soil* 430:1–6
 22. DeWalt SJ, Schnitzer SA, Alves LF et al (2015) Biogeographical patterns of liana abundance and diversity., In: In: Schnitzer SA, Bongers F, Burnham RJ, Putz FE (eds) *Ecology of Lianas*. Wiley, Chichester, pp 131–146
 23. Dormann CF, Elith J, Bacher S et al (2013) Collinearity: A review of methods to deal with it and a simulation study evaluating their performance. *Ecography (Cop)* 36:27–46
 24. Drucker DP, Costa FRC, Magnusson WE (2008) How wide is the riparian zone of small streams in tropical forests? A test with terrestrial herbs. *J Trop Ecol* 24:65–74
 25. Ewers F (1985) Xylem structure and water conduction in conifer tree, dicot trees, and lianas. *Pap Knowl Towar a Media Hist Doc* 6:12–26
 26. Ewers FW, Fisher JB, Chiu ST (1990) A survey of vessel dimensions in stems of tropical lianas and other growth forms. *Oecologia* 84:544–552
 27. Ewers FW, Fisher JB, Fichtner K (1991) Water flux and xylem structure in vines.. in: In: Putz FE, Mooney HA (eds) *The biology of vines*. Cambridge university Press, Cambridge, pp 127–160
 28. Fan Y, Miguez-Macho G, Jobbágy EG et al (2017) Hydrologic regulation of plant rooting depth. *Proc Natl Acad Sci USA* 40:10572–10577
 29. Franks PJ, Leitch IJ, Ruszala EM, Hetherington AM, Beerling DJ (2012) Physiological framework for adaptation of stomata to CO₂ from glacial to future concentrations. *Philos Trans R Soc Lond Ser B Biol Sci* 367:537–546
 30. Gerolamo CS, Angyalossy V (2017) Wood anatomy and conductivity in lianas, shrubs and trees of Bignoniaceae. *IAWA J* 38:412–432
 31. Gerolamo CS, Nogueira A, Costa FRC et al (2018) Local dynamic variation of lianas along topography maintains unchanging abundance at the landscape scale in central Amazonia. *Journal*

32. Gerolamo CS, Nogueira A, Pace MR, Angyalossy V (2020) Interspecific anatomical differences result in similar highly flexible stems in Bignoniaceae lianas. *Am J Bot* 107:1622–1634
33. Gentry AH (1991) The distribution and evolution of climbing plants. In: Putz FE, Mooney HA (eds) *The biology of vines* pp. Cambridge University Press, Cambridge, UK, pp 3–49
34. Grime JP, Thompson K, Hunt R et al (1997) Integrated screening validates primary axes of specialization in plants. *Oikos* 79:259–281
35. Hobbie SE (1992) Effects of plant species on nutrient cycling. *Trends in Ecology and Evolution* 7:336–339
36. Hodnett MG, Vendrame I, Marques Filho AO et al (1997) Soil and groundwater behaviour in a catenary sequence beneath forest in Central Amazonia: I. Comparisons between plateau, slope and valley floor. *Hydrology and Earth Systems Sciences* 1:265–277
37. Hopkins MJG (2005) *Flora da Reserva Ducke, Amazonas, Brasil*. *Rodriguesia* 56:9–25
38. Jirka S, McDonald AJ, Johnson MS et al (2007) Relationships between soil hydrology and forest structure and composition in the southern Brazilian Amazon. *J Veg Sci* 18:183
39. Johansen DA (1940) *Plant microtechnique* New York. McGraw-Hill Book
40. Johnson DM, McCulloh KA, Meinzer FC et al (2011) Hydraulic patterns and safety margins, from stem to stomata, in three eastern US tree species. *Tree Physiol* 6:659–668
41. Kahn F, Castro A (1985) The Palm community in a forest of Central Amazonia, Brazil. *Biotropica* 17:210–216
42. Kalliola R, Puhakka M (1988) River dynamics and vegetation mosaicism: a case study of the River Kamajohka, Northern Finland'. *J Biogeogr* 15:703–719
43. Kraus J, Arduim M (1997) *Manual básico de métodos em morfologia vegetal* pp 1-98. EDUR, Seropédica
44. Larcher W (2000) Temperature stress and survival ability of Mediterranean sclerophyllous plants. *Plant Biosyst* 134:279–295. . doi: 10.1080/11263500012331350455
45. Ledo A, Illian JB, Schnitzer SA et al (2016) Lianas and soil nutrients predict fine-scale distribution of above-ground biomass in a tropical moist forest. *J Ecol* 104:1819–1828
46. Liu X, Swenson NG, Wright SJ et al (2012) Covariation in plant functional traits and soil fertility within two species-rich forests. *PLoS ONE* 7:e34767
47. Liu Q, Sterck FJ, Zhang JL et al (2021) Traits, strategies, and niches of liana species in a tropical seasonal rainforest. *Oecologia* 196:499–514
48. Magnusson WE, Lima AP, Luizão R et al (2005) RAPELD: A modification of the Gentry method for biodiversity surveys in long-term ecological research sites. *Biota Neotrop* 5:2
49. Malizia A, Grau H, Lichstein J (2010) Soil phosphorus and disturbance influence liana communities in a subtropical montane forest. *J Veg Sci* 21:551–560

50. Marques-Filho AO, Ribeiro MNG, Santos HM, Santos JM (1981) Estudos climatológicos da Reserva Florestal Ducke – Manaus-AM. IV. Precipitação. *Acta Amazônica* 11:759–768
51. Muscarella R, Bacon CD, Faurby S et al (2019) Soil fertility and flood regime are correlated with phylogenetic structure of Amazonian palm communities. *Ann Botany* 123:641–655
52. Nogueira A, Costa FRC, Castilho CV (2011) Liana abundance patterns: The role of ecological filters during development. *Biotropica* (Lawrence, KS) 43:442–449
53. Oksanen J, Blanchet FG, Friendly M et al (2015) Vegan: community ecology package. <https://cran.r-project.org/web/packages/vegan/index.html>
54. Oliveira RS, Costa FRC, van Baalen E et al (2019) Embolism resistance drives the distribution of Amazonian rainforest tree species along hydro-topographic gradients. *New Phytol* 221:1457–1465
55. Ordoñez JC, Van Bodegom PM, Witte JP et al (2009) A global study of relationships between leaf traits, climate and soil measures of nutrient fertility. *Glob Ecol Biogeogr* 18:137–149. <https://doi.org/10.1111/j.1466-8238.2008>
56. Pace MR, Lohmann LG, Angyalossy V (2011) Evolution of disparity between the regular and variant phloem in Bignoniaceae (Bignoniaceae). *Am J Bot.* <https://doi.org/10.3732/ajb.1000269>
57. Parolin P, Ferreira LV, Junk WJ (2003) Germination characteristics and establishment of trees from central Amazonian flood plains. *Trop Ecol* 44:155–167
58. Phillips OL, Gentry AH (1994) Increasing turnover through time in tropical forests. *Science* 263:954–958. <https://doi.org/10.1126/science.263.5149.954>
59. Putz FE (1984) The natural history of lianas on Barro Colorado Island, Panama. *Ecology* 68:1713–1724
60. Putz FE, Mooney HA (1991) The biology of vines. Cambridge University Press, Cambridge
61. Rennó CD, Nobre AD, Cuartas LA et al (2008) HAND, a new terrain descriptor using SRTM-DEM; mapping terra-firme rainforest environments in Amazonia. *Remote Sens Environ* 112:3469–3481
62. Reich PB, Walters MB, Ellsworth DS (1997) From tropics to tundra: global convergence in plant functioning. *Proc Natl Acad Sci USA* 94:13730–13734
63. Reich PB, Ellsworth DS, Walters MB et al (1999) Generality of leaf traits relationships: a test across six biomes. *Ecology* 80:1955–1969
64. Reich PB (2014) The world-wide ‘fast–slow’ plant economics spectrum: a traits manifesto. *J Ecol* 102:275–301
65. Ribeiro AJ, Hopkins MJG, Vicentini A et al (1999) Flora da Reserva Ducke. Guia de Identificação das plantas vasculares de uma floresta de terra-firme na Amazonia Central. Manaus, Amazonas, Brasil Great Britain
66. Rocha EX, Schietti J, Gerolamo CS et al (2020) Higher rates of liana regeneration after canopy fall drives species abundance patterns in central Amazonia. *J Ecol* 108:1311–1321
67. Rowe N, Isnard S, Speck T (2004) Diversity of mechanical architectures in climbing plants: an evolutionary perspective. *J. Plant Growth Regul.* 23:108–128

68. Sandel B, Goldstein LB, Kraft NJB et al (2010) Contrasting trait responses in plant communities to experimental and geographic variation in precipitation. *New Phytol* 188:565-575. <https://doi.org/10.1111/j.1469-8137.2010.03382.x>
69. Schenck H 1893. Beiträge zur Biologie und Anatomie der lianen, im Besonderen der in Brasilien einheimischen Arten. ii. Beiträge zur Anatomie der lianen. in: *Botanische Mittheilungen aus den tropen*. Verlag von Gustav Fischer, Jena
70. Schnitzer SA, Bongers F (2002) The ecology of lianas and their role in forests. *Trends Ecol Evol* 17:223–230
71. Schnitzer AS (2005) A mechanistic explanation for global patterns of liana abundance and distribution. *Am Nat* 166:262–276
72. Schnitzer SA, Carson WP (2010) Lianas suppress tree regeneration and diversity in treefall gaps. *Ecol Lett* 13:849–857
73. Schnitzer SA (2018) Testing ecological theory with lianas. *New Phytol* 220:366–380
74. Schnitzer AS, Filippis DM, Visser M, et al (2019) **Local canopy disturbance as an explanation for long-term increases in liana abundance.** *Ecology Letters* 00:1–13. <https://doi.org/10.1111/ele.13881>
75. Sombroek W (2000) Amazon landforms and soils in relation to biological diversity. *Acta Amaz* 30:81–81
76. Silvertown J, Dodd M, Gowing DJ, Mountford JO (1999) Hydrologically defined niches reveal a basis for species richness in plant communities. *Nature* 400:61–63
77. Silvertown J, Araya Y, Gowing D (2015) Hydrological niches in terrestrial plant communities: a review. *J Ecol* 103:93–108
78. Spence RD, Wu H, Sharpe PJH, Clark KG (1986) Water stress effects on guard cell anatomy and the mechanical advantage of the epidermal cells. *Plant, Cell Environ* 9:197–202
79. Suding KN, Lavorel S, Chapin FS et al (2008) Scaling environmental change through the community-level: a trait-based response-and-effect framework for plants. *Glob Chang Biol* 14:1125–1140
80. Toledo JJ, Magnusson WE, Castilho CV, Nascimento HEM (2011) How much variation in tree mortality is predicted by soil and topography in Central Amazonia? *For Ecol Manag* 262:331–338
81. Toledo JJ, Castilho CV, Magnusson WE et al (2017) Soil controls biomass and dynamics of an Amazonian forest through the shifting of species and traits. *Brazilian of Journal Botany* 40:451–461
82. Tomasella J, Hodnett MG, Cuartas LA et al (2008) The water balance of an Amazonia micro-catchment: the effect of interannual variability of rainfall on hydrological behaviour. *Hydrol Process* 13:2133–2147
83. Trujillo W, Rivera-Rondón CA, Balslev H (2021) Palm Functional Traits, Soil Fertility and Hydrology Relationships in Western Amazonia. *Front For Glob Chang* 4:1–10
84. van der Sande MT, Poorter L, Schnitzer SA et al (2019) The hydraulic efficiency–safety trade-off differs between lianas and trees. *Ecology* 100(5):e02666

85. Vivek P, Parthasarathy N (2014) Liana community and functional trait analysis in tropical dry evergreen forest of India. *J Plant Ecol* 8:501–512
86. Wright IJ, Reich PB, Westoby M et al (2004) The worldwide leaf economics spectrum. *Nature* 428:821–827. . doi: 10.1038/nature02403
87. Xu Z, Zhou G (2008) Responses of leaf stomatal density to water status and its relationship with photosynthesis in a grass. *J Exp Bot* 59:3317–3325
88. Yong T, Kitching R, Cao M (2012) Lianas as structural parasites: A re-evaluation. *Chinese Sci Bull* 57:307–312
89. Zhang X, Facette M, Humphries JA et al (2012) Identification of PAN2 by quantitative proteomics as a leucine-rich repeat-receptor-like kinase acting upstream of PAN1 to polarize cell division in maize. *Plant Cell* 24:4577–4589
90. Zhang J, Nieminen K, Serra JAA, Helariutta Y (2014) The formation of wood and its control. *Plant Biology* 17:56–63. <https://doi.org/10.1016/j.pbi.2013.11.003>

Figures

Figure 1

Stem cross sections of Bignoniaceae and Fabaceae species collected at Ducke Reserve along of the hydroedaphic gradient. (A) *Anemopaegma robustum* and (B) *Adenocalymma validum* species of Bignoniaceae; (C) *Machaerium multifoliolatum* and (D) *Machaerium macrophyllum* species of Fabaceae. In (A) and (C) species commonly found in lowland areas closer to the water table in riparian (valley). In (B) and (D) species commonly found in upland areas (plateaus). The self-supporting xylem (XS) is similar to the xylem of many trees and shrubs. The self-supporting xylem is produced in most liana

species during the initial phase of secondary xylem development and is recognized by having a predominance of fibers, fewer parenchyma and smaller vessels. The lianescent xylem (XL) is recognized by large vessels, abundant parenchyma, huge rays and fewer fiber. In the bark we can see the regular phloem region (PR) and in many lianas it is quite common to see cambial variants such as phloem wedges (Pw) in species (A) and (B).

Figure 2

Relationship between lianas species composition (first PCoA axes) and the water table depth (HAND - height above the nearest drainage) gradient in 18 permanent plots at Ducke Reserve. A: Liana species composition represented by the qualitative PCoA-1. B: Liana species composition represented by the quantitative PCoA-1. r^2 adj and significance level (***) $P < 0.001$ are shown.

Figure 3

Distribution of the liana species along the HAND gradient describing the water-table depth. A: Liana species of the family Fabaceae. B: Liana species of the family Bignoniaceae. Vertical bars are the relative abundance of each species along the HAND gradient in 18 permanent plots at Ducke Reserve.

Figure 4

Relationship between the liana mean traits composition and some functional traits and the HAND gradient in 18 permanent plots at Ducke Reserve. (A) Liana mean traits composition represented by the qualitative PCA 1 axis along the hydrological gradient (HAND). Along PCA-1, axis values of stomatal size and self-supporting xylem increased, while leaf size, specific leaf area and wood density decreased. (B-D) Functional traits correlated with three other traits directly related to the hydrological gradient: self-

supporting xylem (B), leaf size (C) and specific leaf area (D). r^2 adj and significance level (* $P < 0.005$; ** $P < 0.01$) are shown.

Figure 5

Relationship between liana mean traits composition represented by the qualitative PCA-1 and species composition represented by the qualitative PCoA 1 axis in 18 permanent plots at Ducke Reserve. r^2 adj and significance level (** $P < 0.001$) are shown.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterial.pdf](#)