

# Sexual Risk Behaviour Among School-going Adolescents in Sierra Leone and Liberia. A Secondary Analysis of the 2017 Global School Health Survey

Peter Bai James (✉ [jamepeb@yahoo.com](mailto:jamepeb@yahoo.com))

University of Sierra Leone

Augustus Osborne

Njala University

Abdulai Jawo Bah

University of Sierra Leone

Emmanuel Kamanda Margao

University of Sierra Leone

Mohamed Conteh-Barat

St. Joseph Catholic Hospital, Monrovia

---

## Research Article

**Keywords:** Adolescents, Sexual risk behaviour, Sierra Leone, Liberia

**Posted Date:** November 30th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-1087719/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Background:** There is very little information on sexual risk behaviour among Sierra Leone and Liberia school-going adolescents. The present study assessed the prevalence and determinants of sexual risk behaviours among school-aged adolescents in Sierra Leone and Liberia.

**Method:** We used publicly available nationally representative cross-sectional datasets of the 2017 Sierra Leone and Liberia Global school health survey. The sample consisted of 2798 and 2744 school-going adolescents from Sierra Leone and Liberia, respectively.

**Results:** The majority of adolescents in the two countries were involved in multiple sexual risk behaviour (80.2%), with higher prevalence observed in Sierra Leone (85.2%) than in Liberia (75.3%), Liberian adolescents showed lesser odds of indulging in multiple sexual risk behaviours than their Sierra Leonean counterparts (AOR=0.572; 95%CI: 0.345-0.946). Male compared to females were more likely to engage in multiple sexual risk behaviour (AOR=2.310;95%CI:1.543-3.458), with a similar pattern observed in both countries. Alcohol use was associated with multiple sexual risk behaviour (AOR=3.064; 95%CI: 2.137-4.392). Also, in Sierra Leone and Liberia, adolescents with one and two or more forms of psychological distress were more likely to have ever had sex than those who do show any form of psychological distress. Missing class/school was associated with multiple sexual risk behaviour (AOR=1.655; 95%CI:1.133-2.418). Peer support was only found to be a protective factor against no condom use among Liberian adolescents (AOR=0.608; 95%CI: 0.435-0.850). Less parental support was only associated with ever had sex more likely to have ever had sex as seen among adolescents in Sierra Leone (AOR=2.027; 95%CI: 1.322-3.107) but not Liberia (1.034(0.650-1.644).

**Conclusion:** Our study found a high sexual risk behaviour among school-going adolescents in Sierra Leone and Liberia. Our finding highlights the need to strengthen sexual and reproductive health education in schools and communities that incorporate mental health promotion activities tailored to this group.

## Introduction

Youths in Sub-Saharan Africa are prone to risky sexual activities, unwanted pregnancy, and sexual violence[1, 2]. Having unprotected sexual intercourse; having multiple sexual partners over one's lifetime; having intercourse with a casual partner; sexual initiation at a young age; sexual intercourse with commercial sex workers; bartering sex for money, goods, or other favours; engaging in sexual activity while under the influence of alcohol/drugs; and sexualism are all examples of risky sexual behaviours[3].

Adolescence is characterised by greater autonomy, peer influence, risk-taking behaviours such as initiation of sex and alcohol/drug usage, [1, 4]. Compared to adults, adolescents are more likely to have several sexual relationships, participate in unprotected sexual intercourse, and choose high-risk partners[5]. The study of teenage sexual behaviour is crucial because 60% of youths globally are afflicted with sexually transmitted infections (STIs), including HIV[6].

Earlier studies in Liberia reported a significant prevalence of risky sexual practices among in-school children and young adults. 78% of kids were found to be sexually active; 24.9% of those sexually active said they had sex for money, and 20.9% said they had never used a condom [7]. In the same study, males were also shown to have more sex, have many sexual partners, and start sex earlier than females. Another study among Liberian youths found that 34% of those who were sexually active did so before the age of 15 (early sexual debut), and 21% of those who were sexually active had several sexual partners, and 26% of sexually active teenagers had never used a condom, 11% had gotten pregnant or helped someone become pregnant one or more times, and 11% had been sexually assaulted [8]. A recent study reported that majority of sexually active Sierra Leone youths had condomless sex in their last sexual encounter [9]. A United Nations Population Fund report on the impact of Ebola on adolescent pregnancy in Sierra Leone found that close to half had their first pregnancy during the Ebola outbreak period and close to a third had used ever use of any kind of family planning[10].

Various sexually related risk behaviours have been observed among adolescents in African countries. In Ghana, 33.5% of adolescents ever had sex, 73.8% had not used a condom at last sex, and 32.5% had multiple sexual partners[11]; in Namibia in 2004, 33.2% ever had sex, and 17.1% had multiple sexual partners[12, 13]. Between 2015 and 2017, a community survey of adolescents (15–19 years) in Uganda, Tanzania, Nigeria, Ghana, Eswatini, Ethiopia, and Burkina Faso found that 25.9% had ever had sex. Among sexually active adolescents, the early sexual debut was 21% for girls and 28% for boys, unprotected last sex was 46% for girls and 40% for boys, and 37% for girls and 8% for boys had made someone pregnant[4].

In a study of 15-year-olds in 30 European countries, Israel, and Canada, 27% had had sexual intercourse, and 14% had not used the contraceptive pill or condoms at their most recent sex[14], and in a study of 15-year-olds in 10 European countries, the prevalence of sexual initiation was 18.8%, and among sexually active, 52.4% had >1 sexual partner[15].

Although the commonness of sexual behaviour varies by country and culture, the relationships between sexual and non-sexual risk behaviours and the function of psychosocial modulators may follow similar patterns. A previous study has identified factors associated with sexual risk behaviour among adolescents (ever had sex, early sexual debut, no condom use, and no contraceptive use), and they include male sex, older age, substance use, psychological distress, school truancy, and a lack of parental and peer support[16]. Substance abuse has been connected to a higher chance of youths participating in unsafe sexual practices. When comparing youths who use substances to those who do not, studies show that those who use substances are more likely to engage in early sexual intercourse, have many sexual partners, and use condoms at a lesser rate[17, 18]. Few studies have been undertaken among Liberian youths to investigate the link between risky sexual behaviours and substance abuse. One study revealed no link between alcohol consumption and transactional sex[19], while another identified a link between alcohol consumption and having several sexual partners-but no other substances were investigated[7]. Other drugs, such as marijuana, cocaine, stimulants such as methamphetamines, have been linked to risky sexual behaviours in adolescence[7, 20, 21].

Sierra Leone and Liberia are neighbouring countries that have shared unique history. Both countries populations have experienced civil war and, most recently, an Ebola disease outbreak leading to profound economic hardship, psychopathologies such as posttraumatic stress disorder, depression, and psychosis as well family disruption [20, 22–24]. These mental health morbidities and family disruption were more profound among adolescents and young people, making them vulnerable to indulge in at risk behaviours such as sexual risk behaviours like early sexual debut, having multiple sexual partners, and not using condoms[10, 25, 26]. Also, the adolescent birth rate in these countries is reported to be high and above the average sub-Saharan Africa [27]. Given such a unique background of these two countries, it is important to examine how personal, psychosocial, and protective factors influence sexual risk behaviour, especially among adolescents. Currently, there is limited national data on sexual risk behaviour and related risk factors among only adolescents. Most studies conducted in these countries are either community based or are among adolescents and adults combined [7, 19, 28, 29]. Knowing the prevalence of sexual behaviour and the risk factors associated with it among teenagers in Sierra Leone and Liberia can aid in developing intervention programs aimed at delaying sexual initiation and encouraging "safer sex." As a result, the goal of this study was to assess the prevalence and determinants of sexual risk behaviours among school-aged adolescents in Sierra Leone and Liberia using their 2017 Sierra Leone and Liberia Global school health survey (GSHS).

## Methods

### Sample and procedure

We used publicly available nationally representative cross-sectional datasets of the 2017 Sierra Leone and Liberia Global school health survey[30]. The Sierra Leone and Liberia GSHS employ a two-stage cluster sample design to obtain a nationally representative sample of school-going adolescents. The first stage involves the selection of schools with probability proportional to enrolment size, whilst the second stage involves randomly selecting classes for which all students have equal chances of being selected. In the Sierra Leone GSHS, the school response rate was 94%, the student response rate was 87%, and the overall response rate was 82%[30]. In the Liberia GSHS, the school response rate was 98%, the student response rate was 73%, and the overall response rate was 71%[30].

### Measures

The questionnaire used in this study and the definition of the variables is shown in Table 1. Sexual risk behaviour was considered as the outcome variable in our study, and it was assessed using the following questions ever having had sexual intercourse, age of sexual debut, number of people who have had sexual intercourse within a lifetime, condom use at last sexual intercourse, and birth control use at last sexual intercourse. Sexual risk behaviour was defined as ever having had sex, early sexual debut (<14 years), having had two or more sexual partners in a lifetime, non-condom use at last sex and nonbirth control use at last sex. Composite sexual risk behaviour was defined as having had sex, early sexual debut (<14 years), having had two or more sexual partners in a lifetime and non-condom use at last sex.

As reported in previous studies[31, 32], we excluded non-birth control use because of the overlap with non-condom use at last sex. The Independent variables considered in this study are in Table 1. As in a previous study[31], we considered no close friends, loneliness, anxiety, suicidal ideation, and suicide attempt as psychological distress items. Based on similar study by Pengpid and Pelzer[33], we summed these items into three groups – 0 = 0, 1 = 1 single and 2–5 = 2 multiple. School attendance, peer and parental or guardian support was considered protective factors and they groups. The four items that measure parental or guardian support were summed and divided into three groups - 0–1 as low, 2 as medium and 3–4 as high support.'

Table 1  
Questionnaire items and coding scheme

Indicator	Item	Responses (coding scheme)
<b>Outcome Variable (Sexual risk behaviour)</b>		
Ever sex	'Have you ever had sexual intercourse?'	'Yes, No' (coded yes=1, no=0)
Age of sexual initiation	'How old were you when you had sexual intercourse for the first time?'	'I have never had sexual intercourse  11 years old or younger to 18 years  old or older'
Number of sex partners	'During your life, with how many people have you had sexual intercourse?'	'I have never had sexual intercourse, 1  person to 6 or more people'
Condom use	'The last time you had sexual intercourse, did you or your partner use a condom?'	'I have never had sexual intercourse,  Yes, No, I do not know'
Birth control use	'The last time you had sexual intercourse, did you or your partner use any method of birth control, such as withdrawal, rhythm (safe time), birth control pills, or any other method to prevent pregnancy?'	'I have never had sexual intercourse,  Yes, No, I do not know'
<b>Independent variables</b>		
<b>Substance use</b>		
Current alcohol use	'During the past 30 days, on how many days did you have at least one drink containing alcohol?'	'1 = 0 days to 7 = All 30 days (coded 1 = 0, 2-7 = 1)'
Cannabis use	'During your life, how many times have you used marijuana'	'1 = 0 times to 5 = 20 or more times  (coded 1 = 0 and 2-5 = 1)'
Amphetamine use	'During your life, how many times have you used amphetamines or methamphetamine'	'1 = 0 times to 5 = 20 or more times  (coded 1 = 0 and 2-5 = 1)'

Indicator	Item	Responses (coding scheme)
<b>Psychological distress</b>		
No close friends	'How many close friends do you have?'	'1 = 0-4 = 3 or more (coded 1+=0, 0 = 1)'
Loneliness	'During the past 12 months, how often have you felt lonely?'	'1=never to 5=always (coded 1-3 = 0 and 4-5 = 1)'
Anxiety	'During the past 12 months, how often have you been so worried about something that you could not sleep at night?'	'1=never to 5=always (coded 1-3 = 0 and 4-5 = 1)'
Suicide ideation	'During the past 12 months, did you ever seriously consider attempting suicide?'	'Yes, No'
Suicide attempt	'During the past 12 months, how many times did you actually attempt suicide?'	'1 = 0 times to 5 = 6 or more times (coded 1 = 0 and 2-5 = 1)'
<b>Protective factors</b>		
School attendance	'During the past 30 days, on how many days did you miss classes or school without permission?'	'1 = 0 days to 10 or more days (coded 1 = 1)'
Peer support	'During the past 30 days, how often were most of the students in your school kind and helpful?'	'1=never to 5=always (coded 1-3 = 0 and 4-5 = 1)'
Parental supervision	'During the past 30 days, how often did your parents or guardians check to see if your homework was done?'	'1=never to 5=always (coded 1-3 = 0 and 4-5 = 1)'
Parental connectedness	'During the past 30 days, how often did your parents or guardians understand your problems and worries?'	'1=never to 5=always (coded 1-3 = 0 and 4-5 = 1)'
Parental bonding	'During the past 30 days, how often did your parents or guardians really know what you were doing with your free time?'	'1=never to 5=always (coded 1-3 = 0 and 4-5 = 1)'

Indicator	Item	Responses (coding scheme)
Parental respect for privacy	'During the past 30 days, how often did your parents or guardians go through your things without your approval?'	'1=never to 5=always (coded 1–3 = 0 and 4–5 = 1)'

## Ethical consideration

No formal ethical approval to conduct this study was necessary, given that our study is based on an analysis of a publicly available deidentified secondary dataset. Notwithstanding, ethics approval was obtained from the Ministries of Health in Sierra Leone and Liberia prior to conducting the surveys in the two countries.

## Data analysis

We analysed our pooled data from the two sets of surveys using SPSS version 27. We employed descriptive statistics to describe our sample. Chi-square statistics were used to compare the independent variables between Sierra Leone and Liberia. Binary regression statistics were used to determine the correlates of individual sexual risk behaviours (non-birth control use at last sex, non-condom use at last sex, multiple sexual partners, early sexual debut, and ever had sex) and a composite measure of multiple sexual risk behaviour. We excluded current tobacco use as an independent variable in our analysis because the data was not available in the Sierra Leone GSHS dataset. We employed complex samples analysis in all statistical procedures to account for the sampling weights and the multi-stage design. Statistical significance was set at  $p < 0.05$ .

## Results

### Characteristics of the sample and sexual risk behaviour in Sierra Leone and Liberia

Table 2 provides a summary of the characteristics of the sample and sexual risk behaviour in Sierra Leone and Liberia. The sample consisted of 2798 and 2744 school-going adolescents from Sierra Leone and Liberia, respectively. Close to half of them were above the age of 17 years (45.3%) and were females (48.3%). A similar percentage of females were observed in Sierra Leone and Liberia. Overall, close to half of the students have ever had sex (48.4%) but more than a third (38.9%) in Sierra Leone and close to two-thirds (61.9%) in Liberia. Among sexually active school-going adolescents, close to a third (31.3%) have had an early sexual debut ( $\leq 14$  years). However, 44.4% reported early sexual debut ( $\leq 14$  years) in Sierra Leone and less than a quarter (20.6%) in Liberia. Close to one in five (19.3%) had multiple sexual partners, 13.0% in Sierra Leone and 27.8% in Liberia. The majority (80.2%) of school-going adolescents exhibit



multiple sexual risk behaviour, although such risky behaviours were more common among adolescents in Sierra Leone (85.2%) than Liberia (75.2%).

Table 2  
 Characteristics of the sample and sexual risk behaviour in Sierra Leone and Liberia 2017 GSHS.

Study characteristics	variable	All N (%)	Sierra Leone N (%)	Liberia N (%)	P-value
Age in years	14 or less	1365(26.7)	973(35.6)	392(16.1)	<0.001
	15-16	1452(28.0)	939(34.6)	513(20.2)	
	≥17	2624(45.3)	868(29.8)	1756(63.7)	
Sex	Male	2640(51.7)	1258(51.6)	1382(51.8)	0.960
	Female	2737(48.3)	1484(48.4)	1253(48.2)	
<b>Sexual behaviour</b>					
Ever had sex	Yes	2181 (48.4)	917(38.9)	1264(61.9)	<0.001
Early sexual debut (<14 years)	Yes	536(31.3)	330(44.4)	330(20.6)	<0.001
Multiple sexual partners	Yes	1044(19.3)	360(13.0)	684(27.8)	<0.001
No condom use at last sex	Yes	848(48.7)	437(58.6)	411(39.8)	<0.001
No birth control use at last sex	Yes	892(48.5)	389(50.7)	503(46.6)	0.424
Multiple sexual risk behaviour	Yes	1142(80.2)	529(85.2)	613(75.3)	0.023
<b>Substance use</b>					
Current alcohol use	Yes	1011(18.6)	393(13.4)	618(24.7)	<0.001
Ever use cannabis	Yes	358(6.6)	165(5.2)	193(8.4)	0.032
Ever use amphetamine use	Yes	371(9.7)	191(7.9)	180(7.9)	0.999
<b>Psychological distress</b>					
Have No close friends	Yes	579(10.8)	249(9.2)	330(12.6)	0.003
Felt Lonely	yes	933(16.9)	566(19.4)	367(13.9)	0.014
Anxiety	Yes	1060(19.2)	520(18.5)	540(20.0)	0.339
Suicide ideation	Yes	1076(19.7)	394(14.2)	682(26.3)	<0.001
Suicide Attempts	Yes	1402(25.7)	541(19.1)	861(33.7)	<0.001
<b>Protective factors</b>					
Missed class/school	Yes	2121(40.5)	992(35.0)	1129(47.6)	<0.001

Study characteristics	variable	All N (%)	Sierra Leone N (%)	Liberia N (%)	P-value
Peer support	Yes	1741(32.6)	854(29.6)	887(36.4)	0.015
Parental Support	Low	2153(42.4)	1116(39.6)	1037(46.4)	0.066
	Medium	1348(28.4)	781(30.0)	567(26.1)	
	High	1377(29.2)	771(30.4)	606(27.5)	

## Associations with sexual risk behaviour

Table 3 provides correlates associated with ever had sex among school-going adolescents in Sierra Leone. Adolescents in Liberia were more likely than their Sierra Leonean counterparts to have ever had sex (AOR=1.548;95%CI:1.199- 1.999). Among those sexually active, adolescents in Liberia compared to those in Sierra Leone were more likely to have multiple sexual partners (AOR=1.587;95%CI:1.137-2.214). However, adolescents in Liberia were less likely to show multiple sexual risk behaviour than their Sierra Leonean counterparts (AOR=0.572;95%CI:0.345-0.946). See Tables 4, 5 and 8.

Table 3

Associations with ever had sex among school-going adolescents in Sierra Leone and Liberia 2017 GSHS

<b>Study characteristics</b>	<b>All AOR (95%CI)</b>	<b>Sierra Leone AOR (95%CI)</b>	<b>Liberia AOR (95%CI)</b>
Liberia vs Sierra Leone	1.548(1.199-1.999)		
<b>Age group</b>			
14 or less	1	1	1
15-16	1.432(0.967-2.120)	1.554(0.964-2.505)	1.257(0.735-2.149)
≥17	<b>3.469(2.323-5.179)</b>	<b>3.086(1.721-5.533)</b>	<b>4.048(2.314-7.080)</b>
<b>Sex (Male vs female)</b>	<b>1.578(1.202-2.071)</b>	<b>1.880(1.280-2.762)</b>	1.246(0.938-1.656)
<b>Current alcohol use (yes vs no)</b>	2.221(1.672-2.948)	1.801(1.063-3.050)	<b>2.519(1.753-3.621)</b>
<b>Ever use cannabis (Yes vs No)</b>	6.041(2.423-15.060)	5.606(1.792-17.540)	6.436(1.271-32.595)
Ever use amphetamine use (Yes vs No)	1.651(0.851-3.201)	1.594(0.738-3.442)	2.218(0.644-7.639)
<b>Psychological distress items</b>			
0	1	1	1
1	<b>1.586(1.134-2.219)</b>	<b>1.883(1.262-2.810)</b>	<b>1.273(0.847-1.913)</b>
2-5	<b>2.403(1.568-3.683)</b>	<b>3.346(1.831-6.117)</b>	<b>1.611(1.105-2.350)</b>
<b>Missed class/school (yes vs no)</b>	<b>1.312(1.084-1.588)</b>	<b>1.583(1.136-2.206)</b>	<b>1.105(0.822-1.487)</b>
<b>Peer support (Yes vs No)</b>	0.935(0.723-1.209)	0.937(0.629-1.395)	0.920(0.707-1.197)
<b>Parental support</b>			
Low	<b>1.492(1.120-1.987)</b>	<b>2.027(1.322-3.107)</b>	<b>1.034(0.650-1.644)</b>
Medium			
High	1.168(0.852-1.602)	1.212(0.797-1.843)	1.115(0.746-1.667)
	1	1	1

Table 4

Associations with Early sexual debut among school-going adolescents in Sierra Leone and Liberia 2017  
GSHS

Study characteristics	All AOR (95%CI)	Sierra Leone AOR (95%CI)	Liberia AOR (95%CI)
Liberia vs Sierra Leone	0.640(0.384-1.067)		
<b>Age group</b>			
14 or less	1	1	1
15-16	0.060(0.025-0.143)	<b>0.051(0.016-0.161)</b>	<b>0.080(0.025-0.253)</b>
≥17	0.021(0.009-0.050)	<b>0.017(0.005-0.057)</b>	<b>0.035(0.014-0.086)</b>
<b>Sex (Male vs female)</b>	<b>1.754(1.151-2.674)</b>	1.575(0.768-3.231)	<b>2.513(1.498-4.215)</b>
<b>Current alcohol use (yes vs no)</b>	1.048(0.625-1.757)	1.835(0.635-5.301)	0.635(0.308-1.308)
<b>Ever use cannabis (Yes vs No)</b>	1.364(0.566-3.290)	0.575(0.192-1.726)	<b>7.338(2.004-26.864)</b>
Ever use amphetamine use (Yes vs No)	1.475(0.502-4.337)	1.199(0.355-4.053)	0.987(0.208-4.695)
<b>Psychological distress items</b>			
0	1	1	1
1	1.242(0.718-2.149)	1.185(0.589-2.387)	1.311(0.743-2.311)
2-5	<b>2.217(1.311-3.748)</b>	<b>2.855(1.262-6.457)</b>	1.428(0.882-2.312)
<b>Missed class/school (yes vs no)</b>	0.906(0.609-1.347)	0.922(0.501-1.699)	0.877(0.544-1.414)
<b>Peer support (Yes vs No)</b>	0.879(0.536-1.441)	1.019(0.487-2.132)	0.751(0.397-1.422)
<b>Parental support</b>			
Low	1.183(0.782-1.790)	1.071(0.512-2.242)	1.310(0.669-2.565)
Medium	1.427(0.814-2.501)	1.621(0.723-3.633)	1.359(0.533-3.469)
High	1	1	

Table 5

Associations with multiple sexual partners among school-going adolescents in Sierra Leone and Liberia 2017 GSHS

<b>Study characteristics</b>	<b>All AOR (95%CI)</b>	<b>Sierra Leone AOR (95%CI)</b>	<b>Liberia AOR (95%CI)</b>
Liberia vs Sierra Leone	<b>1.587(1.137-2.214)</b>		
<b>Age group</b>			
14 or less	1	1	1
15-16	1.686(1.105-2.574)	<b>1.666(1.013-2.739)</b>	<b>1.778(0.800-3.949)</b>
≥17	<b>4.463(2.890-6.894)</b>	<b>3.288(1.974-5.476)</b>	<b>6.648(3.130-14.121)</b>
<b>Sex (Male vs female)</b>	<b>2.232(1.553-3.208)</b>	<b>2.542(1.551-4.166)</b>	<b>1.992(1.294-3.066)</b>
<b>Current alcohol use (yes vs no)</b>	<b>2.981(2.048-4.340)</b>	<b>3.219(1.563-6.630)</b>	<b>2.626(1.886-3.655)</b>
<b>Ever use cannabis (Yes vs No)</b>	1.812(0.904-3.632)	<b>2.372(1.095-5.136)</b>	1.316(0.542-3.197)
Ever use amphetamine use (Yes vs No)	1.257(0.695-2.273)	1.477(0.657-3.318)	0.785(0.335-1.836)
<b>Psychological distress items</b>			
0	1	1	1
1	1.245(0.810-1.915)	1.179(0.575-2.420)	1.262(0.764-2.087)
2-5	1.403(0.955-2.061)	1.324(0.650-2.697)	1.431(0.998-2.053)
<b>Missed class/school (yes vs no)</b>	<b>1.683(1.338-2.118)</b>	<b>1.854(1.304-2.637)</b>	<b>1.582(1.145-2.187)</b>
<b>Peer support (Yes vs No)</b>	1.146(0.889-1.477)	1.423(0.973-2.080)	0.965(0.710-1.312)
<b>Parental support</b>			
Low	1.178(0.845-1.640)	1.768(0.925-3.378)	0.859(0.604-1.221)
Medium	0.972(0.669-1.413)	1.098(0.658-1.834)	0.940(0.581-1.523)
High	1	1	

Regarding age, adolescents who were 17 years and older were more likely than those 14 years and younger to have ever had sex (AOR=3.469; 95% CI: 2.323-5.179). A similar relationship was observed among adolescents in Sierra Leone (AOR=3.086; 95%CI:1.721-5.533) and Liberia (AOR=4.048;95%CI:2.314-7.080). Also, among sexually active adolescents, a similar association was observed between age and not using other birth control methods other than condom (AOR=2.474;95%CI:1.359-4.505). Regarding sex, overall, male adolescents were more likely than their female counterparts to have ever had sex (AOR=1.578;95%CI:1.202- 2.071), to have had an early sexual debut (AOR=1.754;95%CI:1.151-2.674) had multiple sexual partners (AOR=2.232;95%CI:1.553-3.208), multiple sexual risk behaviour (AOR=2.310;95%CI:1.543-3.458). Male gender was found to be associated with ever had sex, and such association was observed among Sierra Leoneans (AOR= 1.880;95%CI:1.280-2.762) but not among Liberian adolescents (AOR=1.246;95%CI:0.938-1.656). Also, among sexually active adolescents, males were more likely than females to not use other birth control methods other than condom in Sierra Leone (AOR=1.827;95%CI:1.063-3.141) but not in Liberia (AOR=0.814;95%CI:0.559-1.186). However, male gender was associated with high-risk sexual behaviour in Sierra Leone (AOR=3.632;95%CI:1.852-7.122) and Liberia (AOR=1.913;95%CI:1.156-3.167). Adolescents who were current alcohol users were more likely than non-users to have ever had sex (AOR=2.221;95%CI:1.672-2.948), had multiple sexual partners (AOR=2.981;95%CI:2.048-4.340) and had high-risk sexual behaviour (AOR=3.064;95%CI:2.137-4.392). See Tables 3, 5,7 and 8.

Overall, adolescents with one (AOR=1.586;95%CI:1.134-2.219) and two or more (AOR= 2.403;95%CI:1.568-3.683) forms of psychological distress were more likely to have ever had sex than those who do not show any form of psychological distress with a similar pattern seen in Sierra Leone and Liberia. Also, adolescents that showed two or more form psychological distress were more likely to have had an early sexual debut (AOR=2.217;95%CI:1.311- 3.748) with a similar relationship observed only among Sierra Leonean adolescents (AOR=2.855;95%CI:1.262- 6.457). In addition, adolescents that exhibited one form of psychological distress were more likely not to use a condom in last sexual intercourse (AOR=1.346;95%CI:1.016-1.783). See Tables 3,4,6

Table 6  
Associations with Non-condom use among school-going adolescents in Sierra Leone and Liberia 2017  
GSHS

Study characteristics	All AOR (95%CI)	Sierra Leone AOR (95%CI)	Liberia AOR (95%CI)
Liberia vs Sierra Leone	0.614(0.362-1.041)		
<b>Age group</b>			
14 or less	1	1	1
15-16	0.919(0.524-1.613)	0.943(0.398-2.234)	0.732(0.344-1.555)
≥17	0.686(0.398-1.181)	0.722(0.351-1.485)	0.535(0.250-1.143)
<b>Sex (Male vs female)</b>	1.174(0.733-1.882)	1.060(0.502-2.242)	1.209(0.810-1.806)
<b>Current alcohol use (yes vs no)</b>	1.106(0.767-1.596)	0.954(0.366-2.485)	1.075(0.765-1.509)
<b>Ever use cannabis (Yes vs No)</b>	0.622(0.193-1.997)	0.418(0.093-1.877)	1.533(0.819-2.868)
Ever use amphetamine use (Yes vs No)	1.755(0.693-4.446)	2.544(0.634-10.206)	0.754(0.295-1.929)
<b>Psychological distress items</b>			
0	1	1	1
1	<b>1.346(1.016-1.783)</b>	1.775(0.961-3.282)	1.138(0.812-1.595)
2-5	0.772(0.505-1.180)	0.800(0.352-1.817)	0.809(0.522-1.254)
<b>Missed class/school (yes vs no)</b>	1.226(0.874-1.718)	1.244(0.747-2.074)	1.174(0.834-1.652)
<b>Peer support (Yes vs No)</b>	0.784(0.514-1.196)	1.019(0.499-2.080)	<b>0.608(0.435-0.850)</b>
<b>Parental support</b>			
Low	1.183(0.792-1.767)	1.565(0.778-3.151)	0.979(0.646-1.482)
Medium	0.885(0.605-1.294)	0.813(0.425-1.558)	1.003(0.637-1.580)
High	1	1	1



Adolescents who missed school were more likely to have had sex (AOR=1.312;95%CI:1.084-1.588), had multiple sexual partners (AOR=1.683;95%CI:1.338-2.118) and multiple sexual risk behaviour (AOR=1.655;95%CI:1.133-2.418). Peer support was not associated with any of the sexual risk indicators. Adolescents with less parental support were more likely to have ever had sex (AOR=1.492;95%CI:1.120-1.987). A similar relationship was seen among adolescents in Sierra Leone (AOR=2.027;95%CI:1.322-3.107) but not in Liberia (1.034(0.650-1.644). However, adolescents with less parental support were less likely to show multiple sexual behaviours (AOR=0.697;95%CI:0.498- 0.977). See Tables 3,5, and 8

Table 7

Association with Non-birth control use other than condom among school-going adolescents in Sierra Leone and Liberia 2017 GSHS

Study characteristics	All AOR (95%CI)	Sierra Leone AOR (95%CI)	Liberia AOR (95%CI)
Liberia vs Sierra Leone	1.074(0.665-1.735)		
<b>Age group</b>			
14 or less	1	1	1
15-16	1.660(0.898-3.069)	1.915(0.839-4.374)	1.312(0.512-3.365)
≥17	<b>2.474(1.359-4.505)</b>	<b>3.273(1.430-7.492)</b>	1.851(0.973-3.518)
<b>Sex (Male vs female)</b>	1.181(0.841-1.659)	<b>1.827(1.063-3.141)</b>	0.814(0.559-1.186)
<b>Current alcohol use (yes vs no)</b>	0.750(0.499-1.126)	0.331(0.098-1.122)	1.149(0.878-1.503)
<b>Ever use cannabis (Yes vs No)</b>	1.544(0.733-3.253)	3.023(0.949-9.632)	0.663(0.255-1.723)
Ever use amphetamine use (Yes vs No)	1.007(0.482-2.103)	1.326(0.508-3.460)	1.062(0.263-4.288)
<b>Psychological distress items</b>			
0	1	1	1
1	1.071(0.547-2.097)	1.357(0.472-3.901)	0.898(0.567-1.423)
2-5	1.283(0.730-2.256)	1.877(0.622-5.666)	1.131(0.716-1.786)
<b>Missed class/school (yes vs no)</b>	0.966(0.733-1.273)	1.146(0.698-1.881)	0.896(0.642-1.250)
<b>Peer support (Yes vs No)</b>	0.868(0.683-1.103)	<b>0.577(0.342-0.975)</b>	1.139(0.809-1.603)
<b>Parental support</b>			
Low	<b>0.697(0.498-0.977)</b>	0.510(0.250-1.042)	0.826(0.571-1.195)
Medium	1.102(0.704-1.726)	0.784(0.471-1.307)	1.188(0.723-1.950)
High	1	1	1



Table 8

Association with multiple sexual risk behaviour among school-going adolescents in Sierra Leone and Liberia 2017 GSHS

Study characteristics	All AOR (95%CI)	Sierra Leone AOR (95%CI)	Liberia AOR (95%CI)
Liberia vs Sierra Leone	<b>0.572(0.345-0.946)</b>		
<b>Age group</b>			
14 or less	1	1	1
15-16	0.305(0.089-1.046)	0.200(0.0340-1.156)	0.663(0.217-2.023)
≥17	0.188(0.063-0.558)	0.091(0.017-0.483)	0.650(0.224-1.883)
<b>Sex (Male vs female)</b>	<b>2.310(1.543-3.458)</b>	<b>3.632(1.852-7.122)</b>	<b>1.913(1.156-3.167)</b>
<b>Current alcohol use (yes vs no)</b>	<b>3.064(2.137-4.392)</b>	29.012(2.303-365.529)	<b>1.923(1.190-3.107)</b>
<b>Ever use cannabis (Yes vs No)</b>	0.278(0.073-1.056)	0.100(0.025-0.403)	2.123(0.445-10.134)
Ever use amphetamine use (Yes vs No)	1.739(0.444-6.801)	0.863(0.159-4.678)	0.959(0.172-5.338)
<b>Psychological distress items</b>			
0	1	1	1
1	1.188(0.745-1.894)	1.459(0.468-4.554)	1.129(0.644-1.982)
2-5	1.324(0.805-2.177)	1.451(0.507-4.148)	1.132(0.683-1.875)
<b>Missed class/school (yes vs no)</b>	<b>1.655(1.133-2.418)</b>	1.213(0.614-2.399)	<b>1.921(1.289-2.865)</b>
<b>Peer support (Yes vs No)</b>	1.350(0.799-2.280)	1.179(0.482-2.883)	1.238(0.687-2.231)
<b>Parental support</b>			
Low	1.322(0.898-1.946)	1.611(0.608-4.270)	1.131(0.728-1.758)
Medium	1.132(0.754-1.698)	1.100(0.440-2.746)	1.249(0.746-2.094)
High	1	1	1

## Discussion

Our study found a high prevalence of sexual risk behaviours (ever had sex, early sexual debut, multiple sexual partners, no condom, and no birth control use at last sex) among school-going adolescents with Liberian adolescents showing lesser odds of indulging in multiple sexual risk behaviours than their Sierra Leonean counterparts. Close to half of them have ever had sex (48.4%), with high prevalence observed among Liberians (61.9%) than Sierra Leoneans (38.2%) school-going adolescents. Our finding for both countries is lower than what was reported in Mozambique, Ethiopia [31, 34] but higher than what was reported in Ghana[11], and in four Caribbean countries[35] and five East Asia countries[36]. However, in individual countries, the reported prevalence in Mozambique was higher than what is found for Sierra Leone but low regarding our finding for Liberia.

Close to a third had an early sexual debut (<14 years) with a higher prevalence seen among Sierra Leoneans, and this was lower than what other similar studies reported in Caribbean and Asian countries[35, 36]. However, the prevalence of early sexual debut for Sierra Leone and Liberia was consistent and higher than what was reported in studies conducted in Ethiopia and Mozambique [31, 34], respectively. In addition, close to half (48.7%) did not use a condom in their last sex, with higher prevalence seen among Sierra Leoneans (58.6%) than Liberians (39.8%), and our findings for both countries are higher than the prevalence reported in studies conducted in Mozambique, Ghana and Ethiopia[11, 31, 34]. Our study's high prevalence of risky sexual behaviour is consistent with previous community-based studies conducted among adolescents and youths in both countries[7–9, 29] and similar studies in other African countries[1, 4, 11, 31]. The higher prevalence of risky sexual behaviour in our study may be attributed to increased trauma and economic hardships, changes in parenting styles, breakdown in the social fabric in our society and broken homes experienced by young people, which is due to civil war and lately the Ebola outbreak in these two countries [37–39].

Consistent with a previous studies conducted in Ghana[11], Mozambique[31], Fiji, Kiribati, Samoa, and Vanuatu[16], males were more likely to have ever had sex, had an early sexual debut (<14 years), multiple sexual partners and high sexual risk behaviour composite score. Similarly, being 17years and older was associated with being sexually active, having multiple sexual partners, and non-birth control uses other than condoms. Our findings suggest the need to develop and implement gender and age-specific interventions that will help prevent adolescents from indulging in risky sexual behaviours. Substance use (alcohol and cannabis use) was associated with sexual risk behaviour among adolescents in Sierra Leone and Liberia. Our finding aligns with previous studies conducted in Liberia[7] and some African countries[11, 31]. Substance abuse has been reported to be higher among adolescent in Sierra Leone and Liberia[20, 21, 40], and such behaviour have been reported to be associated with risky sexual behaviour and teenage pregnancy[7, 41].

In contrast to a Mozambican study [31] but consistent with a Ghanaian study[11] and in the Caribbean as well as Asian pacific island countries[16, 35, 36], psychological distress was associated with sexual risk behaviours such as ever had sex, early sexual debut and non-condom use. High mental health burden has

been reported among adolescents in Sierra Leone and Liberia, and such burden is linked with exposure to trauma during the civil war and Ebola outbreak in these two countries[22, 26]. Psychological distress contributes to adolescents being vulnerable, leading to risky sexual behaviour, violence, substance abuse, partly caused by economic hardship, and parental loss[42]. Although peer support was identified as a protective factor in only one of the sexual risk behaviour indicators (non-condom use) among Liberian adolescents, previous studies have concluded that negative peer influence affects adolescents' sexual risk behaviour[23, 43]. Low parental support was associated as with ever had sex but was not linked with the other sexual risk behaviour indicators, which is consistent with other studies in which parental support was a protective factor for non-condom use and non-birth control use at last sex but not multiple sexual risk behaviours[11, 31]. Our finding may reflect the significant trauma due to war, poverty and infectious disease outbreaks in these countries leading to mental health needs that have not been fully addressed. Such an unmet need makes it difficult for adolescent to benefit from protective factors such as peer and parental support.

## **Policy and Practice Implication**

Our findings underscored the need for strengthening sexual and reproductive health education in schools and communities. Such educational programs should integrate mental health promotion activities targeting adolescent specific health needs, such as increasing their access to counselling and education. Also, peer support and parental involvement in adolescent daily activities will help reduce adolescent sexual risk behaviour.

## **Study Limitations**

The 2017 Sierra Leone and Liberia GSHS employed a cross-sectional study design, and as such, we cannot infer causal relationships between our dependent and independent variables. Also, our findings are only applicable to school-going adolescents in these two countries. Future research should focus on both in-school and out-of-school adolescents. There is a tendency for recall bias as responses were based on self-report. GSHS does not provide a clear definition of 'sexual intercourse, which would have led to the possibility of some respondents misinterpreting some questions.

## **Conclusion**

Our study suggests that most school-going adolescents in Liberia and Sierra Leone have indulged multiple sexual risk behaviours although those in Liberia had fewer odds of being involved in multiple sexual risk behaviours than their Sierra Leone counterparts. Sex, substance use, psychological distress and missing classes were associated with multiple sexual risk behaviours. Peer and parental support were the only protective factors for no condom use among Liberian adolescents and being sexually active among Sierra Leonean adolescents. Our finding highlights the need to strengthen sexual and reproductive health education in schools and communities that incorporate mental health promotion activities.

# Abbreviations

CI: Confidential Intervals

GSHS: Global School Health Survey

OR: odds ratio

SPSS: Statistical Package for The Social Sciences

# Declarations

## **Ethics approval and consent to participate:**

No formal ethical approval to conduct this study was necessary, given that our study is based on an analysis of a publicly available de-identified secondary dataset. Notwithstanding, ethics approval was obtained from the Ministries of Health in Sierra Leone and Liberia prior to conducting the surveys in the two countries

## **Consent for publication:**

Not Applicable

## **Availability of data and materials:**

The dataset informing the findings of this study is and publicly available. It can be freely available via the WHO NCD Microdata Repository. <https://extranet.who.int/ncdsmicrodata/index.php/catalog/GSHS>

## **Competing interests:**

The authors declared no conflict of interest.

## **Funding:**

No funding was obtained for this study.

## **Authors' contributions:**

PBJ contributed to designing the study, conducted the analysis, interpreted the results, and wrote the first draft of the manuscript. AO contributed to designing the study, participated in the interpretation of the results, and contributed to writing the manuscript. ABJ, EKM and MCB participated in the interpretation of the results and edited the manuscript. All authors read and approved the final manuscript.

## **Acknowledgements:**

We want to thank the high school students who participated in the 2017 Sierra Leone and Liberian GSHS. This paper uses data from the Global School Health. GSHS is supported by the World Health Organization and the US Centre for Disease Control and Prevention.

## References

1. Doyle AM, Mavedzenge SN, Plummer ML, Ross DA: **The sexual behaviour of adolescents in sub-Saharan Africa: patterns and trends from national surveys.** *Tropical Medicine & International Health* 2012, **17**(7):796-807.
2. Ganle JK, Amoako D, Baatiema L, Ibrahim M: **Risky sexual behaviour and contraceptive use in contexts of displacement: insights from a cross-sectional survey of female adolescent refugees in Ghana.** *International Journal for Equity in Health* 2019, **18**(1):127.
3. Perera UAP, Abeysena C: **Prevalence and associated factors of risky sexual behaviors among undergraduate students in state universities of Western Province in Sri Lanka: a descriptive cross sectional study.** *Reproductive health* 2018, **15**(1):105-105.
4. Berhane Y, Canavan CR, Darling AM, Sudfeld CR, Vuai S, Adanu R, Bärnighausen T, Dessie Y, Bukenya JN, Guwatudde D: **The age of opportunity: prevalence of key risk factors among adolescents 10–19 years of age in nine communities in sub-Saharan Africa.** *Tropical Medicine & International Health* 2020, **25**(1):15-32.
5. Ritchwood TD, Ford H, DeCoster J, Sutton M, Lochman JE: **Risky Sexual Behavior and Substance Use among Adolescents: A Meta-analysis.** *Children and youth services review* 2015, **52**:74-88.
6. Da Ros CT, da Silva Schmitt C: **Global epidemiology of sexually transmitted diseases.** *Asian journal of andrology* 2008, **10**(1):110-114.
7. Quiterio N, Harris B, Borba C, Henderson D: **Substance use and sexual risk behaviours amongst in-school youth and young adults living in Liberia.** *African Journal of Drug and Alcohol Studies* 2013, **12**(2).
8. Kennedy SB, Atwood KA, Harris AO, Taylor CH, Gobeh ME, Quaqua M, Woods DV, Bee EM, Warlonfa M: **HIV/STD risk behaviors among in-school adolescents in post-conflict Liberia.** *The Journal of the Association of Nurses in AIDS Care : JANAC* 2012, **23**(4):350-360.
9. Ali MM, Merdad L, Bellizzi S: **Socioeconomic variations in risky sexual behavior among adolescents in 14 sub-Saharan Africa countries who report ever having had sex.** *International Journal for Equity in Health* 2021, **20**(1):11.
10. UNFPA Sierra Leone: **Recovering from the Ebola Virus Disease: Rapid Assessment of Pregnant Adolescent Girls in Sierra Leone.** <https://sierraleone.unfpa.org/en/publications/recovering-ebola-virus-disease-rapid-assessment-pregnant-adolescent-girls-sierra-leone> Accessed 29 October 2021. 2018.
11. Kugbey N, Ayanore MA, Amu H, Oppong Asante K, Adam A: **International note: Analysis of risk and protective factors for risky sexual behaviours among school-aged adolescents.** *Journal of*



*Adolescence* 2018, **68**:66-69.

12. Twa-Twa J, Oketcho S, Siziya S, Muula A: **Prevalence and correlates of condom use at last sexual intercourse among in-school adolescents in urban areas of Uganda.** *East African journal of public health* 2008, **5**(1):22-25.
13. Chinsembu KC, Siziya S, Muula AS, Rudatsikira E: **Prevalence and social correlates of sexual intercourse among schoolgoing adolescents in Namibia.** *SAHARA: Journal of Social Aspects of HIV/AIDS Research Alliance* 2008, **5**(3):129-135.
14. Gabhainn SN, Baban A, Boyce W, Godeau E: **How well protected are sexually active 15-year olds? Cross-national patterns in condom and contraceptive pill use 2002–2006.** *International Journal of Public Health* 2009, **54**(2):209-215.
15. Gambadauro P, Carli V, Wasserman C, Hadlaczky G, Sarchiapone M, Apter A, Balazs J, Bobes J, Brunner R, Cosman D: **Psychopathology is associated with reproductive health risk in European adolescents.** *Reproductive health* 2018, **15**(1):1-12.
16. Peltzer K, Pengpid S: **Risk and Protective Factors Affecting Sexual Risk Behavior Among School-Aged Adolescents in Fiji, Kiribati, Samoa, and Vanuatu.** *Asia Pacific Journal of Public Health* 2016, **28**(5):404-415.
17. Vasilenko SA, Lanza ST: **Predictors of multiple sexual partners from adolescence through young adulthood.** *Journal of Adolescent Health* 2014, **55**(4):491-497.
18. Tucker JS, Hu J, Golinelli D, Kennedy DP, Green Jr HD, Wenzel SL: **Social network and individual correlates of sexual risk behavior among homeless young men who have sex with men.** *Journal of Adolescent Health* 2012, **51**(4):386-392.
19. Okigbo CC, McCarragher DR, Chen M, Pack A: **Risk factors for transactional sex among young females in post-conflict Liberia.** *African journal of reproductive health* 2014, **18**(3):133-141.
20. Harris BL, Levey EJ, Borba CPC, Gray DA, Carney JR, Henderson DC: **Substance use behaviors of secondary school students in post-conflict Liberia: a pilot study.** *International Journal of Culture and Mental Health* 2012, **5**(3):190-201.
21. Olurishe TO: **Drug and substance abuse in Anglophone West Africa: a mini review.** *West Afr J Pharm* 2019, **30**(21):1-12.
22. Hopwood H, Sevalie S, Herman MO, Harris D, Collet K, Bah AJ, Beynon F: **The burden of mental disorder in Sierra Leone: a retrospective observational evaluation of programmatic data from the roll out of decentralised nurse-led mental health units.** *International journal of mental health systems* 2021, **15**(1):31.
23. Liebling-Kalifani H, Mwaka V, Ojiambo-Ochieng R, Were-Oguttu J, Kinyanda E, Kwekwe D, Howard L, Danuweli C: **Women war survivors of the 1989-2003 conflict in Liberia: the impact of sexual and gender-based violence.** *Journal of International Women's Studies* 2011, **12**(1):1-21.
24. Alleyne-Green B, Kulick A, Grocher KB, DeLoach McCutcheon KP, Betancourt TS: **The impact of war violence exposure and psychological distress on parenting practices among a sample of young**

- adults affected by war postconflict Sierra Leone. *Peace and Conflict: Journal of Peace Psychology* 2019, **25**(4):325.
25. Betancourt TS, Newnham EA, McBain R, Brennan RT: **Post-traumatic stress symptoms among former child soldiers in Sierra Leone: follow-up study.** *The British Journal of Psychiatry* 2013, **203**(3):196-202.
26. Hook K, Ando K, Ghebrehiwet S, Harris B, Ojediran B, Syeda H, Henderson D, Borba C: **Current state of the literature on mental health in Liberia: A systematic review.** *2020* 2020, **26**.
27. UNICEF: **Progress for Every Child in the SDG Era Country Profile.** <https://data.unicef.org/resources/progress-every-child-country-profiles/> Accessed 25 October 2021. 2018.
28. Odimegwu CO, De Wet N, Banda PC: **Risky sexual behaviour among women: Does economic empowerment matter? Case of Gabon, Mozambique, Sierra-Leone and Zambia.** *African Journal of AIDS Research* 2016, **15**(4):333-340.
29. Labat A, Medina M, Elhassein M, Karim A, Jalloh MB, Dramaix M, Zhang W-H, Alexander S, Dickson KE: **Contraception determinants in youths of Sierra Leone are largely behavioral.** *Reproductive Health* 2018, **15**(1):66.
30. WHO: **NCD Microdata Repository. Global School-based Student Health Survey.** <https://extranet.who.int/ncdsmicrodata/index.php/catalog/GSHS> Accessed 2nd November 2021. 2021.
31. Pengpid S, Peltzer K: **Sexual risk behaviour and its correlates among adolescents in Mozambique: results from a national school survey in 2015.** *SAHARA-J: Journal of Social Aspects of HIV/AIDS* 2021, **18**(1):26-32.
32. Carver JW, Dévieux JG, Gaston SC, Altice FL, Niccolai LM: **Sexual risk behaviors among adolescents in Port-au-Prince, Haiti.** *AIDS and behavior* 2014, **18**(8):1595-1603.
33. Pengpid S, Peltzer K: **Leisure-Time Sedentary Behavior Is Associated with Psychological Distress and Substance Use among School-Going Adolescents in Five Southeast Asian Countries: A Cross-Sectional Study.** *International journal of environmental research and public health* 2019, **16**(12):2091.
34. Yimer B, Ashebir W: **Parenting perspective on the psychosocial correlates of adolescent sexual and reproductive health behavior among high school adolescents in Ethiopia.** *Reproductive health* 2019, **16**(1):66-66.
35. Pengpid S, Peltzer K: **Prevalence and Correlates of Sexual Risk Behavior among School-Going Adolescents in Four Caribbean Countries.** *Behav Sci (Basel)* 2020, **10**(11):166.
36. Peltzer K, Pengpid S: **Sexual risk behaviour and its correlates among adolescents in Indonesia, Laos, Thailand and Timor-Leste: results from national school surveys in 2015.** *International Journal of Adolescent Medicine and Health* 2020(20190230).
37. Dominguez S, Borba CPC, Fatima B, Gray DA, Stinehart C, Murphy G, Wang EKS, Harris BL, Henderson DC: **Mental health and adaptation of young Liberians in post-conflict Liberia: a key**

- informant's perspective.** *International Journal of Culture and Mental Health* 2013, **6**(3):208-224.
38. Denney L, Gordon R, Ibrahim A: **Teenage Pregnancy after Ebola in Sierra Leone.** *London: Overseas Development Institute* 2015.
39. Zuilkowski SS, Thulin EJ, McLean K, Rogers TM, Akinsulure-Smith AM, Betancourt TS: **Parenting and discipline in post-conflict Sierra Leone.** *Child Abuse & Neglect* 2019, **97**:104138.
40. Bøås M, Hatløy A, Bjørkhaug I: **Alcohol and drugs in post-war Sierra Leone.** *African Journal of Drug and Alcohol Studies* 2008, **7**(1).
41. Yakubu I, Salisu WJ: **Determinants of adolescent pregnancy in sub-Saharan Africa: a systematic review.** *Reproductive Health* 2018, **15**(1):15.
42. Borba CPC, Ng LC, Stevenson A, Vesga-Lopez O, Harris BL, Parnarouskis L, Gray DA, Carney JR, Domínguez S, Wang EKS *et al.*: **A mental health needs assessment of children and adolescents in post-conflict Liberia: results from a quantitative key-informant survey.** *International journal of culture and mental health* 2016, **9**(1):56-70.
43. Betancourt TS, Simmons S, Borisova I, Brewer SE, Iweala U, De La Soudière M: **High hopes, grim reality: Reintegration and the education of former child soldiers in Sierra Leone.** *Comparative Education Review* 2008, **52**(4):565-587.