Bourneville-Pringle disease (Tuberous Sclerosis Complex - TSC) with bilateral renal angiomyolipoma and an epitheloid monomorf angiomyolipoma (EM-AML) in a 43-year-old female, a case report

Kinga Mate (✉ matekingabrigitta@gmail.com)

Péterfy Sándor Hospital and Traumatology Center <https://orcid.org/0000-0001-7791-1759>

Fanni Márványkövi

Péterfy Sándor Hospital and Traumatology Center, Budapest, Hungary

László Vrecenár

Péterfy Sándor Hospital and Traumatology Center, Budapest, Hungary

Tamás Zóber

Péterfy Sándor Hospital and Traumatology Center, Budapest, Hungary

István Buzogány

Péterfy Sándor Hospital and Traumatology Center, Budapest, Hungary

Tamás Beöthe

Péterfy Sándor Hospital and Traumatology Center, Budapest, Hungary

Research Article

Keywords: Bourneville-Pringle disease, tuberous sclerosis complex (TSC), angiomyolipoma (AML), fat-poor AML, epitheloid monomorf angiomyolipoma (EM-AML), nephrectomy, ablation, embolization, mTOR inhibitor

Posted Date: February 18th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1093873/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: The Bourneville Pringle disease, also called tuberous sclerosis complex (TSC) is a rare genetic disorder with dominant autosomal inheritance. It is a rare neurocutaneous disorder, which can affect numerous internal organs, for example the kidney(s), the heart, and the lungs. The formation of multiple hamartomas is causing the multiorgan systemic involvement.

Case presentation: A 43-year-old patient presents with tuberous sclerosis, an autosomal dominant inherited neurocutaneous disorder which variably affects different organs, among them the kidneys. During follow-up the abdominal computer tomography (CT) showed bilateral angioliipomatosis (AML) and a fat-poor mass with a different solid aspect on the left side. A fat-poor AML can be a challenge to diagnose. It is vital to exclude malignancy to decide on further management. A CT guided biopsy was performed. The results showed an atypical renal angiomyolipoma, a rare monomorph, epitheloid type. A genetic examination was performed, which showed a mutation of the TSC2 genes.

Conclusions: The Bourneville Pringle disease is a rare genetic, neurocutaneous disease, which can affect multiple organs by the formation of hamartomas, including renal manifestation as AML. In case of radiological findings of a fat poor CT component AML it is essential to keep in mind the possibility of a rare monomorph epitheloid AML and if there is any doubt, it is crucial to exclude malignancy by using biopsy. A fat-poor AML can be a challenge to diagnose. It is vital to exclude malignancy, to take the size of the kidney AML in consideration, the high risk of potentially massive hemorrhage to decide on further management.

Background

The Bourneville Pringle disease, also called tuberous sclerosis complex (TSC) is a rare genetic disorder with dominant autosomal inheritance.¹ The incidence of the disease is around 1:12 000, but this can vary in different countries. The name of the disease comes from a

French neurologist, Desire-Magloire-Bourneville, who described it first in 1880. Later on, Pringle supplemented the description of the syndrome with the cutaneous manifestations,

more precisely the facial angiofibromas, also called adenoma sebaceum. The hamartomatous neurocutan disorder was named after them. It is a rare neurocutaneous disorder, which can affect numerous internal organs, for example the kidney(s), the heart, and the lungs. The formation of multiple hamartomas is causing the multiorgan systemic involvement.

The tuberous sclerosis is caused by the mutation of the TSC1 (also called hamartin) and / or the mutation of TSC2 (also called tuberin) genes.²

The diagnostic criteria for the Bourneville Pringle disease were revised in 2012. To diagnose the tuberous sclerosis complex, 2 major criteria or 1 major and 2 minor criteria should be present.³ The major criteria

include hypomelanotic macules (≥ 3 of ≥ 5 mm in diameter), angiofibromas (≥ 3) or fibrous cephalic plaque, ungula fibromas (≥ 2), Shagreen patch, multiple retinal hamartomas, cortical dysplasias, subependymal nodules, subependymal giant cell astrocytomas, cardiac rhabdomyomas, lymphangioliomyomatosis (LAM), angiomyolipomas (≥ 2). The minor features include 'confetti' skin lesions, dental enamel pits (> 3), intraoral fibromas (≥ 2), retinal achromic patch, multiple renal cysts, non-renal hamartomas.

The formation of hamartomas in different organs cause the symptoms, and are responsible for manifestations of certain clinical features.⁴

Skin manifestations mainly include the formation of facial angiofibromas (adenoma sebaceum), periungual angiofibromas, fibrous cephalic plaques, shagreen patches and focal hypopigmentations.⁵⁶

The **neurological manifestations** include epilepsy, formation of subependymal giant cell astrocytomas, TSC-associated neuropsychiatric disorders, intellectual disability, autism spectrum disorder.⁸

The **pulmonary manifestation** is caused by lymphangioliomyomatosis (LAM) that most commonly cause dyspnea and, less commonly, cough, chest pain, and hemoptysis.

Among the **kidney manifestations** the angiomyolipomas and cysts appear more frequently, but rare cases of renal cell carcinomas, epitheloid monomorph angiomyolipoma and oncocytoma were also described.⁸⁷⁸

If the **heart** is affected, we can find cardiac rhabdomyomas.

Case Presentation

We present the case of a 43-year-old female affected by tuberous sclerosis complex, Bourneville Pringle disease, who was diagnosed at the age of 29.

The clinical manifestations started with skin symptoms. She firstly developed facial angiofibromas, adenoma sebaceum at a young age. Later, attended the dermatology department with periungual fibromas and with the appearance of hypopigmental patches on

the back. (Fig.1)

In her adolescent years the patient had epileptic episodes, in remission under the treatment of Pregabalin. In 2006, the abdominal ultrasonography revealed bilaterally enlarged kidneys with masses different in size and echogenicity, and a hypo echogenic mass on the upper pole of the left kidney and also an enlarged liver with 2 hyper echogenic, inhomogeneous masses.

An abdominal CT scan was requested, which described the above-mentioned renal lesions as possible bilateral angioliipomas. The hypo echogenic mass with a different solid aspect on the upper pole of the

left kidney was described as a fat-poor CT scan component AML and the malignancy could not be excluded either. The liver lesions were considered as hepatic angioliipomas. (Fig. 2)

In order to prove or exclude malignancy, a CT guided biopsy was performed, which was reviewed by 2 pathologists. The histological findings showed an atypical renal angiomyolipoma, a rare monomorph, epitheloid type, which is known as a benign lesion, but certain patterns are potentially malignant, that need further observation. To determine these

patterns' potential to become malignant, the histological findings are essential.

The brain computer tomography (CT scan) and the brain magnetic resonance imaging (MRI) revealed typical lesions of tuberous sclerosis.

The patient was referred to the ophthalmology department; a hamartoma was found during the fundusoscopic evaluation.

A genetic examination was performed from the patient's specimen, which showed a mutation of the TSC2 c.2062 del, p. (Leu 688 Cysfs 10).

The lung manifestation, the lymphangioliomyomatosis was diagnosed at the beginning of year 2020, starting with cough and dyspnea.

In the guidelines it is mentioned that in the case of a kidney AML, a lesion >4 cm or aneurysms >5 mm in diameter, there is a high risk of potentially massive hemorrhage.⁹

Despite the histological findings and immunohistochemical features which were referring to a benign form of EM-AML, following the guidelines, taking the size of the lesions in consideration and the high risk to a potentially massive hemorrhage, the possibility of the surgery was offered. The patient refused to undergo urological surgery and decided to have interventions only for her skin manifestations.

The patient is being observed for 14 years now. The international guidelines recommend assessing renal function and blood pressure at least annually and renal imaging every 1-3 years. In our case, she developed secondary hypertension, which is treated with beta-blocker,

Metoprolol (Betaloc) 50mg 2x1 and valsartan plus hydrochlorothiazide (Co-Valsacor) 80/12,5 mg 2x1/2 with favorable outcome. The results of the last renal function test (2020.08.18) were between the normal intervals: creatinine 76 U/L, carbamide 4,4 mmol/L, eGFR > 60 ml/min/1.73m². Concerning the renal imaging, MRI and CT scans were used, at the beginning every 6 months, then annually. The results of the renal imaging during the follow-up period, did not show significant progression. Her epilepsy is treated effectively with Pregabalin (Lyrica) 2x75 mg.

After a 14 year follow up the patient is stable, under remission, without any progression regarding the renal lesions, without any renal surgical intervention.

Since the diagnosis of the lung manifestation (LAM) at the beginning of the year 2020, our patient is waiting for the approval to begin the mTOR inhibitor, Everolimus therapy.

Discussion

The tuberous sclerosis is a genetic disorder, which variably affects the neurological system, skin and among other internal organs, the kidneys, lungs, heart. The clinical manifestations are due to the multiorgan systemic involvement.

The kidney manifestation in most of the cases are cysts and angiomyolipomas. The AML is a benign tumor, which is composed by adipose tissue, blood vessels and smooth muscle cells. The classic AML is characterized by a high fat tissue component. In the literature, in some rare cases renal cell carcinoma, oncocytoma and a rare monomorph epitheloid AML can be present.¹⁰

The fat poor AML is a challenge to diagnose by computer tomography. In most of the cases the malignancy cannot be excluded either. A CT guided biopsy and the histological findings can help to diagnose and to decide further management and follow-up. Immunohistochemical

studies, detecting melanin A and HMB-45, diagnostic markers for AML, can differentiate the rare, monomorph, epitheloid type from RCC.¹¹

In an epitheloid AML the epitheloid cells are predominant. In the literature there are no exact data concerning the required amount of epitheloid cells to differentiate a classic AML from an epitheloid AML. Aydin et al. investigated clinicopathologic features of 194 renal AML cases, highlighting the epitheloid component and they set a minimum proportion of 10% to define AML as an epitheloid AML.¹²

The renal angiomyolipomas are frequent clinical manifestations of TSC, which need careful surveillance to prevent life-threatening complications. In carefully selected patients conservative selective embolization, ablative therapies and mTOR inhibitor therapies and surgeries are the options.

Yamakado et al. reported that there was a strong correlation between the AML size and the aneurysm size and the risk of hemorrhage.¹³

In our case, taking into consideration the renal AML size and the recommendations of the guidelines (a lesion > 4 cm or aneurysms > 5 mm in diameter, with a high risk of a massive hemorrhage) a radical nephrectomy was offered, but the patient refused any surgical intervention. The anatomy of the lesion excluded the possibility of a conservative selective embolization. The patient is waiting for the approval for the mTOR inhibitor, Everolimus therapy.

Conclusions

The Bourneville Pringle disease is a rare genetic, neurocutaneous disease, which can affect multiple organs by the formation of hamartomas, including renal manifestation as AML.

In case of radiological findings of a fat poor CT component AML it is essential to keep in mind the possibility of a rare monomorph epitheloid AML and if there is any doubt, it is crucial to exclude malignancy by using biopsy. A fat-poor AML can be a challenge to diagnose.

It is vital to exclude malignancy, to take in consideration the size of the kidney AML, the high risk of potentially massive hemorrhage to decide on further management.

Abbreviations

TSC: Tuberous Sclerosis Complex

AML: angiomyolipoma

EM-AML: epitheloid monomorph angiomyolipoma

CT: computer tomography

mTOR inhibitor: mammalian target of rapamycin inhibitor

LAM: lymphangiomyomatosis

MRI: magnetic resonance imaging

RCC: renal cell carcinoma

Declarations

Conflict of interests: None

Ethics approval and consent to participate

Our case report was approved by our institutional ethics committee, with the following reference number 22/2021. Consent of the patient signed, uploaded.

Consent for publication

Consent of the patient signed, uploaded.

Availability of data and material

Data will not be shared, because of private informatical healthcare system of the hospital.

Competing interests

The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

Not applicable

Authors' contributions

MK has coordinated the study. Follow-up of the patient was coordinated by BI. MF, MK has analyzed the patient's previous history. MK, MF, ZT and VL drafted, proofread and edited the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank to Dr. Becker Krisztina for the photos of the patient's dermatological manifestations and for our radiologists, Dr. Markó Róbert and Dr. Volford Gábor for their cooperation. And we also thank to Dr. Buzogány István for the excellent technical support and to Dr. Vrecenár László language editor for proofreading the manuscript.

References

1. Alshaiji JM, Spock CR, Connelly EA, Schachner LA. Facial angiofibromas in a mosaic pattern tuberous sclerosis: A case report. *Dermatol Online J* 2012;18:8.
2. Rodrigues DA, Gomes CM, Costa IM. Tuberous sclerosis complex. *An Bras Dermatol* 2012;87:184-96.
3. Roach ES, Gomez MR, Northrup H. Tuberous sclerosis complex consensus conference: Revised clinical diagnostic criteria. *J Child Neurol* 1998;13:624-8.
4. Kwiatkowski DJ. Tuberous sclerosis: From tubers to mTOR. *Ann Hum Genet* 2003;67(Pt 1):87-96.
5. Webb DW, Clarke A, Fryer A, Osborne JP. The cutaneous features of tuberous sclerosis: A population study. *Br J Dermatol* 1996;135:1-5.
6. Sahar El Aoud et al. Tuberous Sclerosis Complex (Bourneville–Pringle Disease) in a 25-year-old Female with Bilateral Renal Angiomyolipoma and Secondary Hypertension, *Saudi J Kidney Dis Transpl* 2017;28(3):633-638
7. Rosser T, Panigrahy A, McClintock W. The diverse clinical manifestations of tuberous sclerosis complex: A review. *Semin Pediatr Neurol* 2006;13:27-36.
8. Ben Hamida F, Gorsane I, Gharbi C, et al. Renal manifestations in tuberous sclerosis. *Rev Med Interne* 2006;27:836-42
9. Anil Kapoor et al. Evolving Strategies in the Treatment of Tuberous Sclerosis Complex-associated Angiomyolipomas (TSC-AML) *Urology* 2016 Mar;89:19-26. doi: 10.1016/j.urology.2015.12.009. Epub

2015 Dec 23.

10. Henske, E. P., Jóźwiak, S., Kingswood, J. C., Sampson, J. R., & Thiele, E. A. 2016). Tuberous sclerosis complex. *Nature Reviews Disease Primers*, 2, doi:10.1038/nrdp.2016.35
11. John C. Hulbert, and John J. Bissler, Tuberous Sclerosis Complex Renal Disease, *Nephron Exp Nephrol*. 2010Bradley P. Dixon Nov; 118(1): e15–e20.
12. Aydin H, Magi-Galluzzi C, Lane BR, Sercia L, Lopez JI, Rini BI, Zhou M. Renal angiomyolipoma: clinicopathologic study of 194 cases with emphasis on the epithelioid histology and tuberous sclerosis association. *Am J Surg Pathol*. 2009 Feb;33(2):289-97. doi: 10.1097/PAS.0b013e31817ed7a6. PMID: 18852677.
13. Yamakado K, Tanaka N, Nakagawa T, Kobayashi S, Yanagawa M, Takeda K. Renal angiomyolipoma: relationships between tumor size, aneurysm formation, and rupture. *Radiology*. 2002;225:78-82.

Figures

Figure 1:

Figure 1: A:Facial angiofibromas, B:periungual fibromas, C,D: hypopigmental patches
Department of Dermatology, Semmelweis, Budapest (dr. Becker Krisztina's photos)

Figure 1

See image above for figure legend

Figure 2:

Figure 2: Bilateral renal angiomyolipoma

Figure 2

See image above for figure legend