Arabidopsis CPR5 Plays a Role in Regulating Nucleocytoplasmic Transport of mRNAs in Ethylene Signaling Pathway

Jiacai Chen

Qingdao Agricultural University

Xinying Sui

Qingdao Agricultural University

Binran Ma

Qingdao Agricultural University

Yuetong Li

Qingdao Agricultural University

Na Li

Qingdao Agricultural University

Longfei Qiao

Qingdao Agricultural University

Yanchong Yu

Qingdao Agricultural University

Chun-Hai Dong (✉ chunhai79@yahoo.com)

Qingdao Agricultural University <https://orcid.org/0000-0003-0568-0300>

Research Article

Keywords: CPR5, ethylene, signaling, nucleocytoplasmic transport, Arabidopsis

Posted Date: November 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1094076/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Plant Cell Reports on February 24th, 2022.
See the published version at <https://doi.org/10.1007/s00299-022-02838-1>.

Abstract

The ETR1 receptor plays a predominant role in ethylene signaling in *Arabidopsis thaliana*. Previous studies showed that both RTE1 and CPR5 can directly bind to the ETR1 receptor and regulate ethylene signaling. RTE1 was suggested to promote the ETR1 receptor signaling by influencing its conformation, but little is known about the regulatory mechanism of CPR5 in ethylene signaling. In this study, we presented data showing that both RTE1 and CPR5 bound to the N-terminal domains of ETR1, and regulated ethylene signaling via the ethylene receptor. On the other hand, the research provided evidence indicating that CPR5 could act as a nucleoporin to regulate the ethylene-related mRNAs export out of the nucleus, while RTE1 or its homolog (RTH) had no effect on the nucleocytoplasmic transport of mRNAs. Nuclear qRT-PCR analysis and poly(A)-mRNA *in situ* hybridization showed that defect of CPR5 restricted nucleocytoplasmic transport of mRNAs. These results advance our understanding of the regulatory mechanism of CPR5 in ethylene signaling.

Key Message

Arabidopsis CPR5 is involved in regulation of ethylene signaling via two different ways: interacting with the ETR1 N-terminal domains, and controlling nucleocytoplasmic transport of ethylene-related mRNAs.

Introduction

Ethylene is the simplest gaseous plant hormone and widely distributed in plant tissues and cells. Previous studies showed that ethylene, as an important plant hormone, played important roles in regulating plant growth and development, such as seed germination, fruit maturity, flower development, sex determination (Kamachi et al., 1997; Yamasaki et al., 2000), and leaf development, senescence and abscission (Bialeski and Reid, 1992; Kieber and Ecker, 1993; Guerrero et al., 1998). Ethylene is also involved in plant responses to biotic and abiotic stresses, including resistance to hypoxia (Fukao and Bailey-Serres, 2008; Guillaume and Margret, 2008; Magneschi and Perata, 2009; Justin and Armstrong, 2010), drought and salt responses, inhibition of hypocotyl elongation under dark condition, and relief of photooxidation stress (Roman and Ecker, 1995; Bleecker and Kende, 2000). When ethylene or ethylene precursor ACC was added to the medium, the dark-grown *Arabidopsis* seedlings exhibited typical characteristics, termed the ethylene “triple response” with exaggerated apical hook, inhibited root and hypocotyl elongation, and swelled hypocotyl (Bleecker et al., 1988; Guzmán and Ecker, 1990). Using the ethylene “triple response” assay, a large number of *Arabidopsis* mutants with altered ethylene sensitivity were isolated. Based on genetic studies of the ethylene responsive mutants, a linear ethylene signal transduction pathway emerged (Bleecker et al., 1988; Guzmán and Ecker, 1990; Kieber et al., 1993; Roman and Ecker, 1995; Guo and Ecker, 2004).

Ethylene signaling pathway starts with the binding of ethylene to receptors, and then transmits through downstream factors, and finally reaches the nucleus and activates expression of the ethylene-responsive genes, thereby resulting in various ethylene responses. So far, some of the key regulatory factors in the

pathway have been isolated, including the ethylene receptors (ETR1, ETR2, ERS1, ERS2, EIN4) (Chang et al., 1993; Hua et al., 1998; Hua and Meyerowitz, 1998; Sakai et al., 1998), the Raf-like kinase CTR1 (Kieber et al., 1993), the endoplasmic reticulum (ER)-localized EIN2 (Alonso et al., 1999), the transcription factors EIN3 and EIN3-LIKE1 (EIL1) (Chao et al., 1997; Solano et al., 1998), and the downstream ETHYLENE-RESPONSIVE FACTORS (ERFs) (Fujimoto et al., 2000).

Ethylene receptors mainly exist in the ER and Golgi in dimeric form, and their N-terminal hydrophobic domains need cuprous ion as cofactor when binding to ethylene (Chen et al., 2002; Schaller and Bleecker, 1995; Rodriguez et al., 1999; Binder et al., 2010). Among the five ethylene receptors in Arabidopsis, ETR1 was believed to play a predominant role in ethylene signaling (Hua et al., 1995; Hua and Meyerowitz, 1998; Hall and Bleecker, 2003; Qu et al., 2007; Liu et al., 2010). To explore the regulation mechanism of the ETR1 receptor, we and colleagues previously reported the isolation of the ETR1 receptor-associated protein RTE1 and CPR5 based on their regulatory functions in the ETR1 receptor signaling (Resnick et al., 2006; Zhou et al., 2007; Dong et al., 2008; Dong et al., 2010; Wang et al., 2017; Zheng et al., 2017).

The Arabidopsis *RTE1* encodes a protein containing 250 amino acids, mainly localized in the ER and Golgi (Zhou et al., 2007; Dong et al., 2008). There are three homologues of *RTE1* in tomato, one of them was reported to be involved in the regulation of ethylene responses (Barry and Giovannoni, 2006; Harry, 2006; Ma et al., 2012). The rice *RTE1* homologue *OsRTH1* and the carnation *RTE1*-like genes (*DeRTE1* and *DeRTH1*) were shown to be involved in ethylene regulated functions in seedling growth and flower senescence (Yu et al., 2011; Zhang et al., 2012).

RTE1 is a positive regulator of the ETR1 receptor, and they can physically interact (Resnick et al., 2006; Dong et al., 2010). Genetic analyses showed that RTE1 is essential for ETR1 to function in Arabidopsis, but not for the other ethylene receptors (Resnick et al., 2006; Resnick et al., 2008). It was suggested that RTE1 may promote ETR1 signaling by influencing the conformation of the ethylene binding domain and/or its equilibrium state (Resnick et al., 2008). The Arabidopsis RTE1-HOMOLOG (RTH), shared the similar gene expression pattern and protein subcellular localization with RTE1, was thought to regulate ethylene signaling via a physical interaction with its homologue (Zheng et al. 2017).

CPR5 was initially isolated from the research on plant systemic acquired resistance (Bowling et al., 1997; Boch et al., 1998). The study showed that CPR5 participated in different physiological and pathological processes in plants, such as K⁺ dynamic balance, ABA signal transduction, redox balance, programmed cell death, and ROS status and signal transduction (Kirik et al., 2001; Jing et al., 2005; Jing and Dijkwel, 2008; Jing H et al., 2010). CPR5 was also shown to be involved in the regulation of gene replication, cell division, cell proliferation and spontaneous cell death (Brininstool et al., 2008; Perazza et al., 2011; Bao and Hua, 2014). More recently, it was reported that CPR5 may act as a nucleoporin to play a role in controlling of triggering immunity and programmed cell death (Gu et al., 2016). Although the regulatory function of CPR5 in ethylene signaling was reported (Aki et al., 2007; Jing et al., 2007; Wang et al., 2017), the regulation mechanism of CPR5 in ethylene signaling is largely unknown.

In the present study, we provided evidence showing that CPR5 can directly interact with the N-terminal transmembrane domains of the ETR1 receptor. CPR5 may mediate the interaction with the ETR1 receptor to regulate ethylene signaling. By using poly(A)-mRNA *in situ* hybridization and nuclear qRT-PCR analysis, we detected the effect of CPR5 on the nucleocytoplasmic transport of ethylene-related mRNAs, suggesting that CPR5 may regulate nucleocytoplasmic transport of mRNAs in ethylene signaling pathway. These observations significantly advance our understanding of the regulation mechanism of CPR5 in ethylene signaling.

Results

CPR5 can bind to the N-terminal domains of ETR1

As both CPR5 and RTE1 can directly bind to the ETR1 receptor (Dong et al., 2010; Wang et al., 2017), we attempted to examine whether they bind to the same site on the receptor. At first, the ETR1 was divided into three segments with different domains: the ETR1 (1-368 aa) contains three transmembrane domains and the GAF domain, the ETR1 (308-596 aa) has the histidine kinase domain, and the ETR1 (604-739 aa) consists of the response regulatory domain (Figure 1A). Yeast split-ubiquitin assay was performed and results indicated that both CPR5 and RTE1 can interact with the ETR1 (1-368 aa), but not the other two fragments (Figure 1B, C).

We further divided the ETR1 (1-368 aa) into three segments: the ETR1 (1-78 aa) contains two transmembrane domains, the ETR1 (79-146 aa) contains one transmembrane domain, and the ETR1 (140-368 aa) has only the GAF domain (Fig. 1D). Yeast split-ubiquitin assay showed that CPR5 could not interact with any of the segments (Figure 1E). In contrast, RTE1 did interact with the ETR1 (1-78 aa), but not the other two segments (Figure 1F).

The interactions were further examined by a bimolecular fluorescence complementation (BiFC) assay in living plant cells. As shown in Figure 2, the BiFC results are consistent with those of the yeast split-ubiquitin experiments. These observations indicated that both CPR5 and RTE1 could bind to the N-terminal domains of ETR1. CPR5 could bind to a larger area on the receptor, while RTE1 could bind to a smaller region.

Synergistic effect may not exist between CPR5 and RTE1 in regulating the ETR1 receptor signaling

The fact that both CPR5 and RTE1 could bind to the ETR1 N-terminal domains prompted us to investigate whether there exists a synergistic effect in their regulation of the ETR1 receptor signaling. To answer this question, we generated the double mutant *cpr5-T3 rte1-3* by a genetic cross between *cpr5-T3* (Wang et al., 2017) and *rte1-3* (Resnick et al., 2006). Ethylene “triple response” assay showed that all the etiolated seedlings of *cpr5-T3*, *rte1-3* and *cpr5-T3 rte1-3* were ethylene hypersensitive (Figure 3A, B). The hypocotyl growth of the double mutant *cpr5-T3 rte1-3* appeared similar to that of *cpr5-T3* or *rte1-3*, suggesting that synergistic effect may not exist in their regulation of the ETR1 receptor signaling.

CPR5 affects the nucleocytoplasmic transport of the ETR1 mRNA

As CPR5 may act as a nucleoporin in controlling of triggering immunity and programmed cell death (Gu et al., 2016), we examined whether CPR5 could function in regulating the nucleocytoplasmic transport of the mRNAs in ethylene signaling pathway. By *in situ* poly(A)-mRNA hybridization, we detected strong nuclear poly(A) signals in *cpr5-T3*, but not in WT, *rte1-3* or *rth-1* (Figure 4). This observation indicated that the nuclear export of mRNAs was defected in the *cpr5* mutant.

We next examined the nuclear accumulation of the *ETR1* receptor mRNA in the mutant plants. Total RNA and nuclear RNA were extracted from the 10-day-old seedlings of WT, *cpr5-T3*, *rte1-3*, and *rth-1*, respectively. qRT-PCR analysis showed that accumulation of the *ETR1* receptor mRNA in the nucleus of *cpr5-T3* reached over 60% of its total level, while those in the nucleus of WT, *rte1-3* or *rth-1* were about 30%. As a control, the *Actin1* mRNA accumulation in the nucleus of *cpr5-T3* was almost the same as in WT, *rte1-3* or *rth-1* (Figure 5).

CPR5 may regulate the nucleocytoplasmic transport of the mRNAs in ethylene signaling pathway

We further examined nucleocytoplasmic transport of the other ethylene-related mRNAs in *cpr5-T3*, including the ethylene receptors genes (*ETR2*, *ERS1*, *ERS2* and *EIN4*), the downstream components (*CTR1*, *EIN2*, *EIN3*), and some of the *ETHYLENE RESPONSIVE FACTORS* (*ERFs*) in the pathway (Figure 5). To our surprise, it was observed that the mRNAs of the *ETR2*, *ERS1*, *ERS2*, *EIN4* and *CTR1* were significantly aggregated in the nucleus of *cpr5-T3*, and their accumulation levels were more than 60% of totals, being similar to that of the *ETR1*. In contrast, the mRNAs of the *EIN2*, *EIN3*, and *ERFs* (*ERF1*, *ERF2*, *ERF4*, *ERF11*, *ERF105*) in ethylene signaling pathway did not dramatically accumulate in the nucleus of *cpr5-T3* compared to that of WT. As controls, the abnormal mRNAs accumulations were not detected in the nucleus of *rte1-3* or *rth-1* (Figure 5).

In addition, we examined the mRNAs accumulations of the genes which were not the components of ethylene signaling pathway such as *ADF1*, an regulator of actin cytoskeleton (Dong et al., 2001). Experiments revealed a significant increase of the *ADF1* mRNA accumulation in the *cpr5-T3* mutant. Moreover, the nuclear mRNA accumulation assay was carried out for the genes mainly involved in flowering pathway such as *LEY*, *REM16*, *GI* and *AP1* (Yu et al., 2020), and the observations showed that increased nuclear aggregations of the mRNAs were detected for *AP1* and *GI* but not *LFY* and *REM16* in *cpr5-T3* (Figure 5). Taken together, these studies suggested that CPR5 regulated mRNA nucleocytoplasmic transport may exist in different pathways.

Discussion

Previous studies showed that the *rte1* mutations can restore the ethylene sensitivity of *etr1-2* but not *etr1-1* (Resnick et al., 2006), while the *cpr5* mutations can restore the ethylene sensitivity of both *etr1-2* and *etr1-1* (Wang et al., 2017), indicating that CPR5 may function differently from RTE1 in regulating the ETR1 receptor signaling. The *etr1-2* encodes an Ala102-to-Thr substitution in the putative ethylene binding

domain, while the *etr1-1* mutation (Cys65-to-Tyr) fully blocks ethylene binding (Schaller and Bleecker 1995; Hall et al., 1999). These studies suggested that CPR5 differed from RTE1 in regulating the ETR1 receptor signaling. As supported, the present study provided evidence indicating that CPR5 may act as a nucleoporin in regulating the nucleocytoplasmic transport of the mRNAs in ethylene signaling pathway, whereas RTE1 or its homolog (RTH) had no effect on the mRNAs nucleocytoplasmic transport (Figures 4, 5).

RTE1 is mainly localized in the ER and Golgi and required for the function of ethylene receptor ETR1 (Resnick et al., 2006; Zhou et al., 2007; Dong et al., 2008). Genetic analyses revealed that Arabidopsis RTE1 is required for the ETR1 receptor signaling, but not for the other receptors, suggesting that RTE1 may play a specific role in regulating the ETR1 signaling (Resnick et al., 2006). The genetic analyses using the double mutants of *cpr5-T3* with each of the ethylene insensitive mutants (*etr1-2*, *etr1-1*, *ers1-10*, *etr2-1*, *ein2-1*) showed that the ethylene insensitivity of the mutant *etr1-2* and *etr1-1* was suppressed by *cpr5-T3*, while those of the others (*ers1-10*, *etr2-1*, *ein2-1*) were not suppressed by *cpr5-T3* (Wang et al., 2017), suggesting that the ETR1 receptor may also be the main target of CPR5 in ethylene signaling pathway. As an ETR1 associated protein, CPR5 may regulate the ETR1 receptor signaling via protein-protein interaction with the receptor. As supported, both CPR5 and RTE1 proteins are localized in the ER and Golgi (Gu et al., 2016; Wang et al., 2017), and both of them can interact directly with the ETR1 receptor (Wang et al., 2017). In this study, we further analyzed the binding domains of ETR1 by CPR5 or RTE1, and the results revealed that RTE1 could bind to a smaller region which contains two transmembrane domains, and the CPR5 binding required at least three transmembrane domains and the GAF domain of ETR1 (Figures 1, 2).

It is worth to note that all the etiolated seedlings of *cpr5-T3*, *rte1-3* and *cpr5-T3 rte1-3* were ethylene hypersensitive, and the hypocotyl lengths of the double mutant *cpr5-T3 rte1-3* appeared similar to those of the mutant *cpr5-T3* or *rte1-3* (Figure 3), suggesting that synergistic effect may not exist in their regulation of the ETR1 receptor signaling. Nevertheless, it is suggested that the enhanced ethylene sensitivity of *cpr5-T3* may be largely attributed to the defected nucleocytoplasmic transport of the *ETR1* mRNA and the other ethylene-related mRNAs (*ETR2*, *ERS1*, *ERS2*, *EIN4*, *CTR1*) in the mutant, while the phenotype of *rte1-3* may not be caused by alteration of the nucleocytoplasmic transport of mRNAs (Figures 4, 5).

CPR5 is originally isolated from the research on plant pathogenesis (Bowling et al., 1997; Boch et al., 1998). Arabidopsis CPR5 has five transmembrane domains at its C-terminus and nuclear localization signal at the N-terminus. Subcellular localization analysis indicated that CPR5 is localized in the ER and Golgi membrane, and the nuclear membrane (Gu et al., 2016). CPR5 was reported to be able to act as a nucleoporin in controlling of triggering immunity (Gu et al., 2016). For example, overexpression of *CPR5* resulted in the retention of stress and hormone related NPR1, JAZ1 and ABI5 in the cytoplasm (Gu et al., 2016). However, whether CPR5 regulates ethylene signaling via controlling of nucleocytoplasmic transport of mRNAs has not been reported. The present study provided evidence showing that CPR5 may play a crucial role in regulation of the nucleocytoplasmic transport of the mRNAs in ethylene signaling

pathway. When the mutant *cpr5-T3* was examined for poly (A) – mRNAs by *in situ* hybridization, obvious mRNAs aggregation was detected in the nucleus of *cpr5-T3* (Figure 4). Further qRT-PCR analysis showed that the ethylene signaling-related mRNAs including those of *ETR1*, *ETR2*, *ERS1*, *ERS2*, *EIN4* and *CTR1* were dramatically accumulated in the nucleus of *cpr5-T3*, but not in the controls (WT, *rte1-3*, *rth-1*) (Figure 5). Interestingly, there were no significant accumulations of the mRNAs of *EIN2*, *EIN3* and some ethylene-induced *ERFs* in the nucleus of *cpr5-T3*, suggesting that CPR5 may regulate the nucleocytoplasmic transport of the mRNAs in ethylene signaling pathway. It is worth to note that the negative regulators of ethylene signaling (*ETR1*, *ETR2*, *ERS1*, *ERS2*, *EIN4* and *CTR1*) were obviously affected, whereas the positive factors (*EIN2*, *EIN3*) were barely affected in the mutant *cpr5-T3*. This observation may give a reason why the *cpr5-T3* mutant exhibited enhanced ethylene sensitivity phenotype. On the other hand, some downstream *ERFs* may function as both positive and negative regulators in ethylene signaling, none of them seem to be compromised in their export out of the nucleus. For example, it was reported that some EAR-domain containing *ERFs* can function as both positive or negative regulators of transcription and ethylene responses (Lyons et al, 2013).

Interestingly, the increased aggregations of the nuclear mRNAs were observed for *AP1* and *G1* but not for *LFY* and *REM16* in the mutant *cpr5-T3* in flowering pathway (Figure 5), suggesting that the CPR5-mediated regulation of mRNAs nucleocytoplasmic transport also existed in the other pathways. Further research about the regulation mechanism and target selectivity is needed in the future study.

Nucleoporin is a basic unit of the nuclear pore complex (NPC) which controls the two-way transport of RNA and protein between nucleus and cytoplasm. Yeast NPC contains 35 to 50 unique proteins (Yang et al., 1998; Rout et al., 2000), while mammalian NPC is a larger complex consisting of 80 to 100 unique proteins (Görlach and Kutay, 1999). In plants, at least 30 nucleoporins were identified, involved in different events including flowering, pathogen interaction, nodulation, cold response, and hormone signaling (Zhang and Li, 2005; Dong et al., 2006; Parry et al., 2006; Meier and Brkljacic, 2009; Tamura et al., 2010). CPR5 was thought to be a plant-specific transmembrane nucleoporin that may contribute to the stability of the NPC core scaffold (Gu et al., 2016). Nucleoporins play multiple roles in various biological processes. In this study, we provided evidence supporting that CPR5 may function as a nucleoporin in regulating ethylene response via affecting the nucleocytoplasmic transport of the mRNAs in ethylene signaling pathway (Figure 5).

Compared to the studies in yeast and metazoa, the nucleocytoplasmic transport of mRNAs remains less understood in plants (Ehrnsberger et al., 2019a). It was reported that the quadruple mutant (*aly1aly2aly3aly4*) showed defects in plant growth, flowering and seed production, and a nuclear accumulation of mRNAs (Pfaff et al., 2018). The UIF-like proteins bind to RNA and UAP56, and cooperate with ALY1-4 to regulate nucleocytoplasmic mRNA transport (Ehrnsberger et al., 2019b). UAP56 encodes a DEAD-box RNA helicase, and acts together with ALY and MOS11 in regulating nuclear accumulation of mRNAs. The *mos11* knockout plants show nuclear accumulation of mRNAs (Germain et al., 2010). Recently, it was reported that two CaMV proteins, the coat protein P4 and reverse transcriptase P5, play important roles in nuclear export of mRNAs (Julie et al., 2021). In the present study, we provided evidence

demonstrating that *Arabidopsis* CPR5 play a role in nuclear accumulation of mRNAs in ethylene signaling pathway. Although it was thought that pre-mRNA splicing may play an essential role for the nuclear export of mature mRNAs, the regulation mechanism needs further investigation.

Materials And Methods

Plant materials and ethylene response assays

Seeds of the wild type (WT, Col-0) or mutants were surface sterilized and then sowed on 1/2 MS (Murashige and Skoog) medium or in soil in a controlled environment growth chamber set at 21°C with 16 h light/8 h in darkness. The *Arabidopsis* ethylene response mutants were either obtained from previous studies (Dong et al., 2008; Wang et al., 2017; Zheng et al., 2017), or generated by genetic crossing. The F2 progeny from the crosses were screened by specific PCR markers, as previously described (Resnick et al., 2006; Wang et al., 2017).

The ethylene response of *Arabidopsis* seedling was examined in the presence of ACC at different concentrations (0, 0.5, 5, 10, 20, 100 µM) as previously described (Wang et al., 2017). After sterilization, seeds were sowed on 1/2 MS medium containing ACC, and treated at 4°C for 3 d. Thereafter, the plates were moved to a growth chamber for 8 h under white light, and then wrapped with aluminum foil and placed in a growth chamber for indicated periods. Measurement of hypocotyl length and statistical data were evaluated either by Student's *t*-test or Fischer's test for a multiple comparison.

Yeast split-ubiquitin assay

For yeast split-ubiquitin assay, the portions of *Arabidopsis* ETR1 (1-368 aa; 369-596 aa; 597-739 aa; 1-78 aa; 79-139 aa; and 140-368 aa) were each PCR-amplified from an existing template, cloned into the PMD18-T vector and then transferred into a bait vector pPR3-N or a prey vector pBT3-STE through the restriction sites shown in the supplementary table (Table S1). The constructions were verified by sequencing of the inserts. The bait or prey vector harboring a full-length of *ETR1*, *CPR5* or *RTE1* was obtained from a previous study (Wang et al., 2017).

Bimolecular fluorescence complementation (BiFC) assay

To generate the constructs harboring the fusions of *cYFP-ETR1* (1-368 aa) and *cYFP-ETR1* (1-78 aa), the coding sequences of *cYFP* (466-717 aa), *ETR1* (1-368 aa) and *ETR1* (1-78 aa) were each PCR-amplified from existing templates as previously described (Wang et al., 2016), and cloned into the pMD18-T vector. After sequencing verification of the inserts, the fusions were cloned into a binary vector, pCambia1300-3HA, through the restriction sites (Xba I and Kpn I for *cYFP*, Kpn I and BamH I for *ETR1* (1-368 aa) and *ETR1* (1-78 aa)). The constructs containing the fusions of *nYFP-CPR5* and *nYFP-RTE1* were obtained from a previous study (Wang et al., 2017). Transformation mediated by *Agrobacterium* (GV3101) using onion epidermal cells was according to a previous study (Xu et al., 2014). Fluorescent signal from transfected cells was examined under a fluorescence microscope (Nikon ECLIPSE Ti-S, Japan) for a preview and a

laser scanning confocal microscope (Leica TCS SP5, Germany) for photopictures. The primers used for the constructs of BiFC assay are listed in a supplementary table (Table S2).

RNA extraction and qRT-PCR analysis

Total RNA was extracted from the 21-day-old *Arabidopsis* seedlings according to TRNzol (TIANGEN, China). Half of the same sample was used for the nuclear RNA extraction essentially as described by Wang et al (2011) with minor modifications. 3 g of the fresh seedlings were ground into powder in liquid nitrogen, and 6 mL extract A (0.3 M sucrose, 50 mM Tris-HCl, 40 U/mL RNase inhibitor, pH 8.5) was added to the ground material. After further grinding, the residue was filtered by micracloth, and the filtrate was collected in the RNase free tube. The residue was added to extract A to continue grinding and filtering. The two filtrates were mixed and centrifuged at 500 g under 4°C for 5 min. The supernatant was discarded, and 1 mL of extract B (20 mM Tris-HCl, pH 8.5, 25% glycerol, 2.5 mM MgCl₂, and 0.2% Triton X-100) was added to the sediment and suspended. The samples were centrifuged at 1300 g under 4°C for 2 min. After repeating two times, 300 µL of extract C (20 mM Tris-HCl, pH 8.5, 0.25 M sucrose, 10 mM MgCl₂, 0.5% Triton X-100, and 350 U/mL RNase inhibitor) was added to the sediment and suspended. Meanwhile, 300 µL of extract D (20 mM Tris-HCl, pH 8.5, 1.7 M sucrose, 10 mM MgCl₂, 0.5% Triton X-100, and 350 U/mL RNase inhibitor) was added to a new RNase free tube. Then the heavy suspension from the extract C was slowly added to the upper layer of extract D, and centrifuged at 1300 g under 4°C for 10 min. The supernatant was discarded, and 1 mL TRNzol reagent was added to the sediment, and the nuclear RNA was extracted accordingly.

cDNA was synthesized with oligo(dT) primers using PrimeScript™ 1st strand cDNA synthesis kit (Takara, Japan). Quantitative RT-PCR analysis was performed on an Agilent Real-Time qPCR apparatus (Mx3000P system) using SYBR Premix ExTaq™ II (Takara, Japan). Biological replicates for each set of experiments were carried out three times, and the mean value of three replicates was normalized using *Tubulin8* as an internal control. Quantitative RT-PCR was conducted at 95°C for 10 min, followed by 40 cycles at 95°C for 30 s, 55°C for 30 s, and 72°C for 20 s. The primers used for quantitative RT-PCR are listed in supplementary table (Table S3).

Poly(A)-mRNA *in situ* hybridization

Examination of the Poly(A)-mRNA *in situ* localization was performed as previously described (Engler et al., 1994; Gong et al., 2005). Fresh leaves taken from the 21-day-old seedlings were put into a fixation cocktail containing 10 ml of 50% fixation buffer (120 mM NaCl, 7 mM Na₂HPO₄, 3 mM NaH₂PO₄, 2.7 mM KCl, 0.1% Tween 20, 80 mM EGTA, 10% DMSO, 5% formaldehyde) and 50% heptane. After shaking gently at room temperature for 30 min, the samples were transferred into anhydrous methanol for dehydration for 5 min, and repeated once. Then the samples were transferred into anhydrous ethanol for further dehydration for 5 min, and repeated twice. The samples were incubated in a mixture with 50% ethanol and 50% xylene for 30 min. After washing twice with anhydrous ethanol for 5 min at each time, the samples were washed twice with anhydrous methanol for 5 min each. The samples were transferred

into a mixture with 50% methanol and 50% fixation buffer (excluding 5% formaldehyde) for 5 min, and then transferred into a fixation buffer containing 5% formaldehyde for 30 min at room temperature. After fixation, the samples were washed twice with the fixation buffer without 5% formaldehyde, and washed once with 1 ml of perfect Hyb Plus hybridization buffer (h-7033, Sigma Aldrich). Then 1 ml of hybridization buffer was added to each sample for prehybridization at 50°C for 1 h. After prehybridization, 5 pmol of 45-mer oligo(dT) labeled with one molecule of 6-FAM fluorescein at N-terminus (synthesized by Sangon Biotech, China) at 50°C in darkness for more than 8 h. After hybridization, the samples were washed with 2 x SSC (0.3 M NaCl, 0.03 M sodium citrate) and 0.1% SDS for 60 min, and then washed with 0.2 x SSC and 0.1% SDS at 50°C for 20 min in darkness. The samples were observed immediately using a laser scanning confocal microscopy (Leica TCS SP5, Germany) with a 488 nm excitation laser, and photopictures were taken from the leaf mesophyll cells. Each experiment was repeated at least three times, and similar results were obtained.

Declarations

Acknowledgments

We thank the Experimental Center of Qingdao Agricultural University for confocal microscopy support (Leica TCS SP5). This work was supported by National Natural Science Foundation of China (#31870255; #31370322), Shandong Natural Science Foundation (ZR2019MC061) and Shandong Agricultural Variety Project (2019LZGC015) to CHD; and National Nature Science Foundation of China (#31600224) to Y.Y.

Author contributions

J.C., X.S. and B.M. performed most of the research and wrote the first draft of the manuscript. Y.L., N.L., L.Q. and Y.Y. performed mutant screening and some qRT-PCR analysis. C.H.D. designed the experiments and revised the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest for publication.

References

1. Aki T, Konishi M, Kikuchi T, Fujimori T, Yoneyama T, Yanagisawa S (2007) Distinct modulations of the hexokinase1-mediated glucose response and hexokinase1-independent processes by HYS1/CPR5 in Arabidopsis. *Journal of Experimental Botany* 58:3239-3248.
2. Alonso JM, Hirayama T, Roman G, Nourizadeh S, Ecker JR (1999) EIN2, a bifunctional transducer of ethylene and stress responses in Arabidopsis. *Science* 284: 2148-2152.

3. Bao Z, Hua J (2014) Interaction of CPR5 with cell cycle regulators UVI4 and OSD1 in Arabidopsis. PLoS One 9: e100347.
4. Barry CS, Giovannoni JJ (2006) Ripening in the tomato Green-ripe mutant is inhibited by ectopic expression of a protein that disrupts ethylene signaling. Proc Natl Acad Sci U S A 103: 7923-7928.
5. Bialeski RL, Reid MS (1992) Physiological changes accompanying senescence in the ephemeral daylily flower. Plant Physiology 98: 1042-1049.
6. Binder BM, Rodriguez FI, Bleecker AB (2010) The Copper Transporter RAN1 Is Essential for Biogenesis of Ethylene Receptors in Arabidopsis. Journal of Biological Chemistry 285: 37263-37270.
7. Bleecker AB, Estelle MA, Somerville C, Kende H (1988) Insensitivity to Ethylene Conferred by a Dominant Mutation in Arabidopsis thaliana. Science 241: 1086-1089.
8. Bleecker AB, Kende H (2000) Ethylene: a gaseous signal molecule in plants. Annual Review of Cell and Developmental Biology 16: 1-18.
9. Boch J, Verbsky ML, Robertson TL, Larkin JC, Kunkel BN (1998) Analysis of Resistance Gene-Mediated Defense Responses in *Arabidopsis thaliana* Plants Carrying a Mutation in *CPR5*. Molecular Plant-Microbe Interactions 11: 1196-1206.
10. Bowling SA, Clarke JD, Liu Y, Klessig DF, Dong X (1997) The cpr5 mutant of Arabidopsis expresses both NPR1-dependent and NPR1-independent resistance. Plant Cell 9: 1573-1584.
11. Brininstool G, Kasili R, Simmons LA, Kirik V, Hulskamp M, Larkin JC (2008) Constitutive Expressor Of Pathogenesis-related Genes5 affects cell wall biogenesis and trichome development. BMC Plant Biology 8: 58.
12. Chang C, Kwok SF, Bleecker AB, Meyerowitz EM (1993) Arabidopsis ethylene-response gene ETR1: similarity of product to two-component regulators. Science 262: 539-544.
13. Chao Q, Rothenberg M, Solano R, Roman G, Terzaghi W, Ecker JR (1997) Activation of the ethylene gas response pathway in Arabidopsis by the nuclear protein ETHYLENE-INSENSITIVE3 and related proteins. Cell 89: 1133-1144.
14. Chen YF, Randlett MD, Findell JL, Schaller GE (2002) Localization of the ethylene receptor ETR1 to the endoplasmic reticulum of Arabidopsis. Journal of Biological Chemistry 277: 19861-19866.
15. Dong CH, Hu X, Tang W, Zheng X, Kim YS, Lee BH, Zhu JK (2006) A putative Arabidopsis nucleoporin, AtNUP160, is critical for RNA export and required for plant tolerance to cold stress. Molecular and Cellular Biology 26: 9533-9543.
16. Dong CH, Jang M, Scharein B, Malach A, Rivarola M, Liesch J, Groth G, Hwang I, Chang C (2010) Molecular association of the Arabidopsis ETR1 ethylene receptor and a regulator of ethylene signaling, RTE1. Journal of Biological Chemistry 285: 40706-40713.
17. Dong CH, Rivarola M, Resnick JS, Maggin BD, Chang C (2008) Subcellular co-localization of Arabidopsis RTE1 and ETR1 supports a regulatory role for RTE1 in ETR1 ethylene signaling. Plant Journal 53: 275-286.

18. Dong CH, Xia GX, Hong Y, Ramachandran S, Kost B and Chua NH (2001) ADF proteins are involved in the control of flowering and regulate F-actin organization, cell expansion and organ growth in *Arabidopsis*. *Plant Cell* 13: 1333-1346.
19. Ehrnsberger HF, Grasser M, Grasser KD (2019a) Nucleocytoplasmic mRNA transport in plants: export factors and their influence on growth and development. *J Exp Bot* 70: 3757-3763.
20. Ehrnsberger HF, Pfaff C, Hachani I, Flores-Tornero M, Sørensen BB, Längst G, Sprunck S, Grasser M, Grasser KD (2019b) The UAP56-Interacting export factors UIEF1 and UIEF2 function in mRNA export. *Plant Physiol* 179: 1525-1536.
21. Engler JA, Van Montagu M, Engler G (1994) Hybridization in situ of whole-mount messenger RNA in plants. *Plant Mol Biol Rep* 12: 321–331.
22. Fujimoto SYM, Ohta M, Usui A, Shinshi H, Ohme-Takagi M (2000) *Arabidopsis* Ethylene-Responsive Element Binding Factors Act as Transcriptional Activators or Repressors of GCC Box-Mediated Gene Expression. *The Plant Cell* 12: 393-404.
23. Fukao T, Bailey-Serres J (2008) Ethylene—A key regulator of submergence responses in rice. *Plant Science* 175: 0-51.
24. Germain H, Qu N, Cheng YT, Lee E, Huang Y, Dong OX, Gannon P, Huang S, Ding P, Li Y, Sack F, Zhang Y, Li X (2010) MOS11: a new component in the mRNA export pathway. *PLoS Genet.* 6(12): e1001250.
25. Gong Z, Dong CH, Lee H, Zhu J, Xiong L, Gong D, Stevenson B, JK Z (2005) A DEAD Box RNA Helicase Is Essential for mRNA Export and Important for Development and Stress Responses in *Arabidopsis*. *The Plant Cell* 17: 256-267.
26. Görlich D, Kutay U (1999) Transport between the cell nucleus and the cytoplasm. *Annu Rev Cell Dev Biol* 15: 607-660.
27. Gu YN, Zebell SG, Liang ZZ, Wang S, Kang BH, Dong XN (2016) Nuclear Pore Permeabilization Is a Convergent Signaling Event in Effector-Triggered Immunity. *Cell* 166: 1526-1538.
28. Guerrero C, de la Calle M, Reid MS, Valpuesta V (1998) Analysis of the expression of two thiolprotease genes from daylily (*Hemerocallis* spp.) during flower senescence. *Plant Molecular Biology* 36: 565-571.
29. Guillaume R, Margret S (2008) Ethylene biosynthesis and signaling in rice. *Plant Science* 175: 32-42.
30. Guo H, Ecker JR (2004) The ethylene signaling pathway: new insights. *Current Opinion in Plant Biology* 7: 40-49.
31. Guzmán P, Ecker JR (1990) Exploiting the Triple Response of *Arabidopsis* to Identify Ethylene-Related Mutants. *The Plant Cell* 2: 513-523.
32. Hall AE, Bleecker AB (2003) Analysis of combinatorial loss-of-function mutants in the *Arabidopsis* ethylene receptors reveals that the *ers1 etr1* double mutant has severe developmental defects that are EIN2 dependent. *Plant Cell* 15: 2032–2041.
33. Hall AE, Chen QG, Findell JL, Schaller GE, Bleecker AB (1999) The relationship between ethylene binding and dominant insensitivity conferred by mutant forms of the ETR1 ethylene receptor. *Plant*

Physiology 121: 291-300.

34. Harry K (2006) Highly conserved proteins that modify plant ethylene responses. *Proc Natl Acad Sci U S A* 103: 7537-7538.
35. Hua J, Chang C, Sun Q, Meyerowitz EM (1995) Ethylene insensitivity conferred by *Arabidopsis* ERS gene. *Science* 269: 1712–1714.
36. Hua J, Meyerowitz EM (1998) Ethylene responses are negatively regulated by a receptor gene family in *Arabidopsis thaliana*. *Cell* 94: 261-271.
37. Hua J, Sakai H, Nourizadeh S, Chen QG, Bleecker AB, Ecker JR, Meyerowitz EM (1998) EIN4 and ERS2 are members of the putative ethylene receptor gene family in *Arabidopsis*. *Plant Cell* 10: 1321-1332.
38. Jing H, Hebel R, Oeljeklaus S, Sitek B, Stuehler K, Meyer H, Sturre M, Hille J, Warscheid B, P D (2010) Early leaf senescence is associated with an altered cellular redox balance in *Arabidopsis cpr5/old1* mutants. *Plant Biology* 10: 85-98.
39. Jing HC, Anderson L, Sturre MJ, Hille J, Dijkwel PP (2007) *Arabidopsis* CPR5 is a senescence-regulatory gene with pleiotropic functions as predicted by the evolutionary theory of senescence. *Journal of Experimental Botany* 58: 3885-3894.
40. Jing HC, Dijkwel PP (2008) CPR5, a Jack of all trades in plants. *Plant Signal Behav* 3: 562-563.
41. Jing HC, Schippers JH, Hille J, Dijkwel PP (2005) Ethylene-induced leaf senescence depends on age-related changes and OLD genes in *Arabidopsis*. *Journal of Experimental Botany* 56: 2915-2923.
42. Julie Kubina, Angèle Geldreich, Jón Pol Gales, Nicolas Baumberger, Clément Bouton, Lyubov A Ryabova, Klaus D Grasser, Mario Keller, Maria Dimitrova (2021) Nuclear export of plant pararetrovirus mRNAs involves the TREX complex, two viral proteins and the highly structured 5'leader region. *Nucleic Acids Research* 49(15): 8900-8922.
43. Justin SHFW, Armstrong W (2010) Evidence for the involvement of ethene in aerenchyma formation in adventitious roots of rice (*Oryza sativa* L.). *New Phytologist* 118: 49-62.
44. Kamachi S, Sekimoto H, Kondo N, Sakai S (1997) Cloning of a cDNA for a 1-aminocyclopropane-1-carboxylate synthase that is expressed during development of female flowers at the apices of *Cucumis sativus* L. *Plant & Cell Physiology* 38: 1197-1206.
45. Kieber JJ, Ecker JR (1993) Ethylene gas: it's not just for ripening any more! *Trends in Genetics* 9: 356-362.
46. Kieber JJ, Rothenberg M, Roman G, Feldmann KA, Ecker JR (1993) CTR1, a negative regulator of the ethylene response pathway in *Arabidopsis*, encodes a member of the raf family of protein kinases. *Cell* 72: 427-441.
47. Kirik V, Bouyer D, Schöbinger U, Bechtold N, Herzog M, Bonneville J-M, Hülskamp M (2001) CPR5 is involved in cell proliferation and cell death control and encodes a novel transmembrane protein. *Current Biology* 11: 1891-1895.
48. Liu Q, Xu C, Wen CK (2010) Genetic and transformation studies reveal negative regulation of ERS1 ethylene receptor signaling in *Arabidopsis*. *BMC Plant Biol* 10: 60.

49. Lyons R, Iwase A, Gänsewig T, Sherstnev A, Duc C, Barton GJ, Hanada K, Higuchi-Takeuchi M, Matsui M, Sugimoto K, Kazan K, Simpson GG, Shirasu K (2013) The RNA-binding protein FPA regulates flg22-triggered defense responses and transcription factor activity by alternative polyadenylation. *Scientific Reports* 3: 2866.
50. Ma Q, Du W, Brandizzi F, Giovannoni JJ, Barry CS (2012) Differential control of ethylene responses by GREEN-RIPE and GREEN-RIPE LIKE1 provides evidence for distinct ethylene signaling modules in tomato. *Plant Physiology* 160: 1968-1984.
51. Magneschi L, Perata P (2009) Rice germination and seedling growth in the absence of oxygen. *Annals of Botany* 103: 181-196.
52. Meier I, Brkljacic J (2009) The nuclear pore and plant development. *Current Opinion in Plant Biology* 12: 87-95.
53. Parry G, Ward S, Cernac A, Dharmasiri S, Estelle M (2006) The Arabidopsis SUPPRESSOR OF AUXIN RESISTANCE proteins are nucleoporins with an important role in hormone signaling and development. *Plant Cell* 18: 1590-1603.
54. Perazza D, Laporte F, Balague C, Chevalier F, Remo S, Bourge M, Larkin J, Herzog M, Vachon G (2011) GeBP/GPL transcription factors regulate a subset of CPR5-dependent processes. *Plant Physiology* 157: 1232-1242.
55. Pfaff C, Ehrnsberger HF, Flores-Tornero M, Sørensen BB, Schubert T, Längst G, Griesenbeck J, Sprunck S, Grasser M, Grasser KD (2018) ALY RNA-Binding proteins are required for Nucleocytoplasmic mRNA transport and modulate plant growth and development. *Plant Physiol* 177: 226-240.
56. Qu X, Hall BP, Gao Z, Schaller GE (2007) A strong constitutive ethylene-response phenotype conferred on Arabidopsis plants containing null mutations in the ethylene receptors ETR1 and ERS1. *BMC Plant Biol* 7: 3.
57. Resnick J, Wen C, Shockey J, Chang C (2006) REVERSION-TO-ETHYLENE SENSITIVITY1, a conserved gene that regulates ethylene receptor function in Arabidopsis. *Proc Natl Acad Sci USA* 103: 7917-7922.
58. Resnick JS, Rivarola M, Chang C (2008) Involvement of RTE1 in conformational changes promoting ETR1 ethylene receptor signaling in Arabidopsis. *Plant Journal* 56: 423-431.
59. Rodriguez FI, Esch JJ, Hall AE, Binder BM, Schaller GE, Bleecker AB (1999) A copper cofactor for the ethylene receptor ETR1 from Arabidopsis. *Science* 283: 996-998.
60. Roman G, Ecker JR (1995) Genetic analysis of a seedling stress response to ethylene in Arabidopsis. *Philos Trans R Soc Lond B Biol Sci* 350: 75-81.
61. Rout MP, Aitchison JD, Suprpto A, Hjertaas K, Zhao Y, Chait BT (2000) The yeast nuclear pore complex: composition, architecture, and transport mechanism. *Journal of Cell Biology* 148: 635-651.
62. Sakai H, J. H, G. CQ, C. C, J. ML, B. BA, M. ME (1998) ETR2 is an ETR1-like gene involved in ethylene signaling in Arabidopsis. *Proc Natl Acad Sci USA* 95: 5812-5817.
63. Schaller GE, Bleecker AB (1995) Ethylene-binding sites generated in yeast expressing the Arabidopsis ETR1 gene. *Science* 270: 1809-1811.

64. Solano R, Stepanova A, Chao Q, Ecker JR (1998) Nuclear events in ethylene signaling: a transcriptional cascade mediated by ETHYLENE-INSENSITIVE3 and ETHYLENE-RESPONSE-FACTOR1. *Genes & Development* 12: 3703-3714.
65. Tamura K, Fukao Y, Iwamoto M, Haraguchi T, Hara-Nishimura I (2010) Identification and characterization of nuclear pore complex components in *Arabidopsis thaliana*. *Plant Cell* 22: 4084-4097.
66. Wang F, Wang L, Qiao L, Chen J, Pappa MB, Pei H, Zhang T, Chang C, Dong CH (2017) Arabidopsis CPR5 regulates ethylene signaling via molecular association with the ETR1 receptor. *J Integr Plant Biol* 59: 810-824.
67. Wang H, Sun Y, Chang J, Zheng F, Pei H, Yi Y, Chang C, Dong CH (2016) Regulatory function of Arabidopsis lipid transfer protein 1 (LTP1) in ethylene response and signaling. *Plant Molecular Biology* 91: 471-484.
68. Wang W, Ye RQ, Xin Y, Fang XF, Li CL, Shi HQ, Zhou XP, Qi YJ (2011) An importin β protein negatively regulates MicroRNA activity in Arabidopsis. *Plant Cell* 23: 3565-3576.
69. Yamasaki S, Fujii N, Takahashi H (2000) The ethylene-regulated expression of CS-ETR2 and CS-ERS genes in cucumber plants and their possible involvement with sex expression in flowers. *Plant & Cell Physiology* 41: 608-616.
70. Yang Q, Rout MP, Akey CW (1998) Three-dimensional architecture of the isolated yeast nuclear pore complex: functional and evolutionary implications. *Molecular Cell* 1: 223-234.
71. Yu Y, Qiao L, Chen J, Rong Y, Zhao Y, Cui X, Xu J, Hou X, Dong CH (2020) Arabidopsis REM16 acts as a B3 domain transcription factor to promote flowering time via directly binding to the promoters of SOC1 and FT. *Plant Journal* 103: 1386-1398.
72. Yu Y, Wang H, Liu J, Fu Z, Wang J, Liu J (2011) Transcriptional regulation of two RTE-like genes of carnation during flower senescence and upon ethylene exposure, wounding treatment and sucrose supply. *Plant Biol (Stuttg)* 13: 719-724.
73. Zhang W, Zhou X, Wen CK (2012) Modulation of ethylene responses by OsRTH1 overexpression reveals the biological significance of ethylene in rice seedling growth and development. *Journal of Experimental Botany* 63: 4151-4164.
74. Zhang Y, Li X (2005) A putative nucleoporin 96 Is required for both basal defense and constitutive resistance responses mediated by suppressor of npr1-1, constitutive 1. *Plant Cell* 17: 1306-1316.
75. Zheng F, Cui X, Rivarola M, Gao T, Chang C, Dong CH (2017) Molecular association of Arabidopsis RTH with its homolog RTE1 in regulating ethylene signaling. *Journal of Experimental Botany* 68: 2821-2832.
76. Zhou X, Liu Q, Xie F, Wen C-K (2007) RTE1 Is a Golgi-Associated and ETR1-Dependent Negative Regulator of Ethylene Responses. *Plant Physiology* 145: 75-86.

Figures

Figure 1

Analysis of the binding domains of CPR5 and RTE1 to the ETR1 receptor by yeast split-ubiquitin assay. (A & D) Diagram for the full length of CPR5, ETR1 and RTE1, and the truncated portions of ETR1. (B, C, E & F) Molecular interaction of CPR5 (B & E) and RTE1 (C & F) with the portions of ETR1 in the yeast split-ubiquitin assay. The bait protein CPR5 (B & E) and RTE1 (C & F) was paired with the full length or portions of ETR1. Positive interaction is indicated by growth on the medium lacking leucine, tryptophan, histidine and alanine (-LTHA). Undiluted and diluted liquid cultures were spotted on the plates and incubated for 5 days at 30°C.

Figure 2

BiFC assay for the molecular association of CPR5, RTE1 with the ETR1 portions in onion epidermal cells. The N-terminal half of YFP (nYFP) was fused to CPR5 or RTE1, and the C-terminal half of YFP (cYFP) was fused to ETR1 (1-368 aa) or ETR1 (1-78 aa), respectively, and the constructs were co-infiltrated into onion epidermal cells. Fluorescent signal of YFP was detected by laser scanning confocal microscopy at 505-530 nm. Bars: 50 μ m.

Figure 3

Ethylene “triple response” assays in double mutants. (A) Ethylene sensitivity analysis of the WT (Col-0) and mutants treated without ACC for 4 d. Bar=5 mm. (B) Ethylene sensitivity analysis of the WT (Col-0) and mutants treated with 20 μ M ACC for 4 d. Bar=5 mm. (C) Quantitative analysis of hypocotyl lengths for the wild type (WT) and mutants on 1/2 MS medium with 20 μ M ACC. For each genotype, the mean value $\pm$ SD is shown for >25 seedlings. ** P <0.01.

Figure 4

Defect of the *cpr5-T3* mutant plants in mRNA export. Poly(A)-mRNA export is blocked in the *cpr5-T3* mutant cells. WT (Col-0), *cpr5-T3*, *rte1-3* and *rth-1* mutant plants were grown at 22°C for 3 weeks. In situ hybridization with fluorescein-labeled oligo (dT) probe was performed using young leaf tissues. Fluorescent signal was examined by a laser scanning confocal microscopy at emission 505-530 nm. Bars: 25 μ m.

Figure 5

Analysis of the relative accumulation of mRNAs in the nucleus of WT (Col-0), cpr5-T3, rte1-3 or rth-1 by qPCR. Plants grown on 1/2 MS medium for 10 d were used for RNA extract. Relative accumulation of nuclear RNA/total RNA was calculated based on qPCR measurements. Black column, total RNA; white column, nucleus RNA. Values are mean $\pm$ SD; **p<0.01.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTables1s3.doc](#)