Diversity of active root-associated methanotrophs of three emergent plants in a eutrophic wetland in northern China

Jing Cui

Inner Mongolia University

Ji Zhao

Inner Mongolia University

Zheng Wang

Inner Mongolia University

Weiwei Cao

Inner Mongolia University

Shaohua Zhang

Inner Mongolia University

Jumei Liu

Inner Mongolia University

Zhihua Bao (✉ zhihua_bao@imu.edu.cn)

Inner Mongolia University

Original article

Keywords: active methanotrophs, high-throughput sequencing, emergent plant, microhabitat

Posted Date: January 9th, 2020

DOI: <https://doi.org/10.21203/rs.2.20409/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at AMB Express on March 14th, 2020. See the published version at <https://doi.org/10.1186/s13568-020-00984-x>.

Abstract

Root-associated aerobic methanotrophs play an important role in regulating methane emissions from the wetlands. However, the influences of the vegetation type on root-associated methanotrophic structures, especially on active flora, remain poorly understood. Transcription of the *pmoA* gene, encoding particulate methane monooxygenase in methanotrophs, was analyzed by reverse transcription PCR (RT-PCR) of mRNA isolated from root samples of three emergent macrophytes, including *Phragmites australis*, *Typha angustifolia*, and *Scirpus triqueter* from a eutrophic wetland. High-throughput sequencing of *pmoA* based on DNA and cDNA was used to analyze the methanotrophic community. Sequencing of cDNA *pmoA* amplicons confirmed that the structure of active methanotrophic was not always consistent with DNA. A type I methanotroph, *Methylobacter*, was the most active group in *P. australis*, whereas *Methylocystis*, a type II methanotroph, was the dominant group in *S. triqueter*. In *T. angustifolia*, these two types of methanotroph existed in similar proportions. However, at the DNA level, *Methylobacter* was predominant in the roots of all three plants. In addition, vegetation type could have a profound impact on root-associated methanotrophic community at both DNA and cDNA levels. These results indicate that members of the genera *Methylobacter* (type I) and *Methylocystis* (type II) can significantly contribute to aerobic methane oxidation in a eutrophic wetland.

Introduction

Wetlands can both produce and absorb greenhouse gases, which is a major component in the global climate change. Being the largest natural wetland at the same latitude of the earth, Wuliangsu (WLSH) is a typical eutrophication wetland in northern China and plays an important role in the earth's ecosystem. Methane, a greenhouse gas, accounts for 20–30% of the contribution of greenhouse gases to global warming (Conrad 2009). In nature, methane is normally produced by methanogens in deep anoxic soil, but is not directly released into the atmosphere. About 90% is consumed by methanotrophic bacteria when passing through the aerobic soil layer. As a biofilter, methanotrophs became a powerful biological weapon to combat global climate change (Hornibrook et al. 2009). Aerobic methanotrophic, with the help of a series of enzymes, can eventually convert methane into carbon dioxide, thus effectively reducing the greenhouse effect of methane. Methane monooxygenase (pMMO) enzyme plays an important role in this process. As a key gene encoding the β -subunit of pMMO, *pmoA* was found in almost all known aerobic methanotrophs. So far, the diversity of methanotrophs is typically assessed by the detection of the *pmoA* gene (Brabcova et al. 2015; McDonald et al. 2008; Semrau et al. 2010).

Eutrophication is a serious ecology problem in major aquatic ecosystems around the world, and aquatic macrophytes play critical roles in improving water quality (Dhote and Dixit 2009). *Phragmites* spp., *Typha* spp., and *Scirpus* spp. are three common emergent types of macrophyte vegetation that are present worldwide (Vymazal 2013), and they mediate CH₄ emissions from wetlands to the atmosphere (Grunfeld and Brix 1999). A well-developed aeration tissue inside the emergent plants can mediate the release of methane from the sediments into the atmosphere through the plants. In this process, the methanotrophs in the plant roots will have a decisive influence on the final methane gas emissions. As was observed in

constructed wetlands, the existence of plants play an important role in regulating the production, consumption and transportation of CH₄ (Sun et al. 2013). CH₄ flux is usually measured under the condition of constructed wetland, which is disturbed by human factors (Zhang et al. 2018). For CH₄ emissions, published results report lower (Bateganya et al. 2015; Maltais-Landry et al. 2009) or higher (Wang et al. 2013) in planted compared to unplanted constructed wetlands. However, the effect of different plant species on CH₄ fluxes remains controversial (Chen et al. 2019). Between 18% and 90% of the produced CH₄ in the root zone of emergent macrophytes in wetlands is consumed by aerobic methanotrophs (Grunfeld and Brix 1999; Laanbroek 2010). So far, many studies have focused on microorganisms in their rhizosphere sediment, including methylotroph- and heterotroph-mediated processes of carbon and other element cycles (Borruso et al. 2017). Through a 16S rRNA gene Illumina MiSeq sequencing, Pietrangelo et al. (2018) reported the bacterial community structure on the root surface of *P. australis* was indeed different from that of *Typha latifolia*. In addition, Fausser et al., (2012) have suggested that methylotrophic bacteria live in the root zones of *Phragmites australis* and *Typha latifolia*. However, the community structure of root-associated methanotrophs and the relationship between the root-associated methanotrophs (functional bacteria) and plant species remain poorly understood.

As more and more molecular biology techniques are applied to the environment, reverse transcription polymerase chain reaction (RT-PCR) is another useful tool to identify active methanotrophs in the environment (Burgmann et al. 2001; Chen et al. 2007; Esson et al. 2016; Griffiths et al. 2000). Different from the measurement of methane flux, studying the transcriptional activity of functional gene *pmoA* can help us understand the activities of aerobic methanotrophs directly. At the same time, transcriptional analysis from natural wetland samples without laboratory culture can more accurately reflect the community characteristics of aerobic methanogens in their natural state.

WLSH Lake is located near the city of Bayannur in the Inner Mongolia Autonomous Region in China. This lake is the largest freshwater lake in the Yellow River watershed. Recently, the lake has become eutrophic after having received industrial wastewater with high nitrogen and phosphorus content (Wu et al. 2017). *Phragmites australis* (common reed), *Typha angustifolia* (narrow leaf cattail), and *Scirpus triqueter* (bulrush) are the dominant macrophytes of WLSH (Duan et al. 2005).

Using the RT-PCR and MiSeq sequencing technique, we studied the structure of methanotrophic communities in the roots of three typical emergent plants (*P. australis*, *T. angustifolia*, and *S. triqueter*) in WLSH wetland, Inner Mongolia of China. This study was conducted to determine (1) whether any trends in active aerobic methanotrophs can be identified based on *pmoA* sequence analysis of cDNA, and (2) whether plant species influence the structure of methanotrophs. The results will be valuable for discussions and decisions related to the emission of greenhouse gas and the restoration of ecosystems by plants in WLSH wetland.

Materials And Methods

Sampling sites and plant materials

Three plants of each *Phragmites australis*, *Typha angustifolia*, and *Scirpus triqueter* were collected from WLSH wetland (N 40°52'36", E 108°51'16") in 15 July 2017 (Fig. 1). The physical and chemical properties of the sampling site are shown in the attached Table (Supplemental, Table.S1). The roots of the three plants were washed carefully with sterile water until all the soil was rinsed off. Some of the roots were carefully picked with sterilized forceps and divided into two equal parts which placed into 50 ml Falcon tubes (Bao et al. 2014a). One was quickly transferred to dry ice for DNA extraction, and the other was placed in a liquid nitrogen tank to extract RNA. We made three parallel lines for each sample, and quickly stored at -80°C prior to molecular analysis.

DNA and RNA extraction and cDNA synthesis

The tissues of the roots were ground to powder in liquid nitrogen, and stored at -80°C until molecular analysis (Bao et al. 2014a). Genomic DNA was extracted from 0.5 g root by using the Fast DNA SPIN for soil kit (MP Biomedicals, Solon, OH), while RNA was by RNeasy pure Plant Kit (Qiagen Biotech Co., Beijing), and both extraction processes were done according to the manufacturer's instructions.

In order to exclude the bacterial DNA from the extracted RNA, *pmoA* gene was used for PCR amplification with the template of the extracted total RNA. The PCR products were analyzed as negative results by agarose gel electrophoresis and NanoVue™ Plus (GE, USA) to ensure that there was no microbial DNA contamination in the total RNA. gDNA eraser reverse transcription kit (TaKaRa, Japan) was used to synthesize cDNA according to the manufacturer's instructions. All the DNA and cDNA samples were stored at -80°C until use.

High-throughput sequencing of *pmoA* genes based on both DNA and cDNA

The *pmoA* gene in DNA and cDNA samples from three plants root were sequenced using the Illumina MiSeq platform. The barcode primer pair A189f/A661r (Costello and Lidstrom 1999) and reagent kit (RR902A, Premix ExTaq™, Takara Bio Inc., Japan) were used for PCR amplification. A raw sequence file was processed by using the mothur software (version 1.33.3) for quality control and sample splitting (Schloss et al. 2009). The reads were processed using the online version of FunGene Pipeline (Fish et al. 2013), then high-quality *pmoA* sequences of each sample were classified as known *pmoA* groups or lineages as described (Cai et al. 2016). *PmoA* sequences were clustered into species-level operational taxonomic units (OTUs) using the FunGene Pipeline with a distance cutoff of 0.09 (Heyer et al. 2002).

Statistical analysis

The phylogenetic tree of the OTUs was drawn with the neighborhood joining method using MEGA5.2 (Tamura et al. 2011). The display and annotation of the tree were done with ITOL (<http://itol.embl.de/>). The representative sequence of OTUs were blasted in NCBI. Statistical analysis and data visualization were carried out in R (version 3.6.1). Mothur software (version 1.33.3) was used for calculating alpha diversity indices which were then analyzed using ANOVA (SPSS v16.0). Principal coordinates analysis

(PCoA) and analysis of similarities (ANOSIM) were done by the package *vegan*. The package *ggplot2* was used to draw the plot of PCoA. Heatmap were also plotted in R with the package *pheatmap*.

Nucleotide sequence accession numbers

All data from MiSeq sequencing of the *pmoA* have been deposited in the NCBI under the accession numbers: SRR10584604-SRR10584613.

Results

Comparison of methanotrophs community between DNA and cDNA amplicons

High throughput sequencing of *pmoA* gene was performed on root DNA and cDNA of three plants, and 109563 high quality reads (DNA 59742; cDNA 53344) were obtained. Fig. 2 showed the community structure at the genus level between DNA and cDNA. For DNA, more than 90% aerobic methanotrophic bacteria belonged to type I methanotrophs affiliated with *Methylomonas* in three plants root (Fig. 2a). The relative abundance of type II in all the samples was very low (0.2-8.9%). *Methylocystis* (8.8%) in *S. triqueter* was much more than the other two plants. Unlike DNA, *Methylomonas* was still dominant in *P. australis*, and *Methylocystis* had the highest abundance in the *S. triqueter* in cDNA (Fig. 2b). The relative abundance of four main genera in both DNA and cDNA were shown in Fig. 3. As *Methylomonas* of all three plants root had the highest abundance in DNA, it all decreased to some extent in cDNA; however, *Methylocystis* and *Methylosinus* had both increased in *S. triqueter* and *T. angustifolia*.

The alpha diversity was also different: in DNA, *S. triqueter* had the most OUTs, and *P. australis* had the highest diversity; while in cDNA, it was *P. australis* and *T. angustifolia* that had the most diversity (Table. 1, Fig. 4a). For beta-diversity, the same result was obtained: the aerobic methanotrophic bacteria of three roots clustered respectively (Fig. 4b), and the differences between groups were significant (Supplemental, Fig. S2, ANOSIM, $p < 0.01$).

Phylogeny of active methanotrophs associated with emergent plant roots

By studying the community of methanotrophs at the transcriptional level, we can reflect more directly on the role of functional flora in controlling methane emissions. The total sequences, obtained by high throughput sequencing of *pmoA* on root cDNA, were classified into 41 OTUs (Fig. 4; Supplemental, Table S2) assigned to 14 genera (Fig. 2b) and 5 classes (Fig. 4). The genus *Methylomonas* and *Methylococcaceae* which were type I methanotrophs in the *Gammaproteobacteria* and the genus *Methylocystis* and *Methylosinus* which were type II methanotrophs in the *Alphaproteobacteria* accounted for more than 80% of all samples.

The community structure varied greatly in different plants root (Fig. 2b). For *P. australis*, type I methanotrophs, including three main OTUs of *Methylomonas*, *Methyloglobulus morosus* (Lontoh et al. 2000) and *Methylomonas* sp. (QBB78506.1), represented 75.1%; however, in *S. triqueter*, the most OUT of *Methylocystis hirsute* (Lontoh et al. 2000) affiliated with type II accounted for 68.0%. The proportions of

type I and type II methanotrophs were almost equal in *T. angustifolia*. Two main OTUs, *Methylobomonas* of type I and *Methylocystis* of type II, accounted for 24.6% and 23.1%, respectively.

The alpha diversity is shown in Table 1, Fig. 5a and Supplemental, Fig. S1. The OTUs richness was evaluated through the Chao1 index whereas the OTUs evenness was evaluated through Shannon. *P. australis* had the most OTUs richness, while *T. angustifolia* had most diversity. In both alpha diversity measures, the methanotrophic diversity in roots of the three plants was significantly different from each other ($p < 0.05$). PCoA (Fig. 5b) and heatmap (Fig. 6) were used to analyze beta-diversity. It verified that samples from the same emergent plants tended to group together, while samples from different plant species were located far apart. The difference between groups was much larger than that within groups (anosim, $p < 0.01$).

Effects of plant host on community structure of methanotrophs

Although methanotrophs based on DNA and cDNA displayed many differences in alpha diversity and community structure, they revealed similar patterns in host dependence. As shown in PCoA (Fig. 5b) and heatmap (Fig. 6), beta-diversity of *pmoA* in both levels clustered with plant types. Plant host had significant influence on community structure of methanotrophs.

Discussion

Research has shown that type I methanotrophs are always found in soils with a limited CH_4 supply because they grow better than type II methanotrophs in a low- CH_4 environment (Hanson and Hanson 1996). On the other hand, type II methanotrophs such as *Methylocystis* are usually found in high- CH_4 systems (Shiau et al. 2018). Recently, Kits et al. (2015) have used complete genome analysis to show that the type I methanotrophs *Methylobomonas denitrificans* FJG1 possess related denitrification genes and demonstrate denitrification activity under hypoxic conditions. In addition, *Methylocystis* and *Methylosinus* of type II methanotrophs were found to be the predominant root-associated activity bacteria in rice paddy field (Bao et al. 2014b; Eller and Frenzel 2001; Qiu et al. 2009; Shinoda et al. 2019) and were identified as diazotrophic methanotrophs in rice root (Bao et al. 2014b; Shinoda et al. 2019). These results support that members of type I and type II methanotrophs inhabiting in aquatic plant in wetland. This study showed that type I methanotrophs dominates the root systems of the three species of emergent plants, which could easily be explained by the fact that the three species lived in the same water area and their aerenchyma was conducive to the proliferation of CH_4 , resulting in a low concentration of CH_4 in this water area. However, when analyzing the active methanotrophs in cDNA, we found that the structure was significantly different from the DNA level, especially *S. triqueter*, whose active communities were mainly *Methylocystis* of type II methanotrophs. This leads us to the question: what factors affect the transcriptional activity of different methanotrophs? *Methylobomonas* has been shown to be active in methane oxidation in the environments that are more neutral to alkaline in pH, such as a cave system, soda lake, and landfill cover soil (Cebren et al. 2007; Hutchens et al. 2004; Lin et al. 2004). On the other hand, Chen et al. (2008) determined that *Methylocystis* populations were

predominant in the active methanotrophs in a range of peatlands in the United Kingdom. These results showed that environmental factors (e.g., pH) affect the activity of different groups of methanotrophs. The pH of sediments in WLSH was 8.0 (Supplemental, Table. S1), and all three plants in this study grew in the same slightly alkaline environment which is suitable for the habitation of *Methylobacter*. Nevertheless, the main active groups of these plant roots varied with different plant species in this research, which may be due to the transcriptional activity of methanotrophs being more sensitive to the change of microenvironment (root secretion, etc.) of the plant root. Rhizodermis cells secrete a wide range of compounds, including organic acid ions, inorganic ions, phytosiderophores, sugars, vitamins, amino acids, purines, and nucleosides, and the root cap produces polysaccharide mucilage (Dakora and Phillips 2002). These exudates have a great influence on root microhabitats. The production of organic acids, for example, may alter the pH of plant roots. Though the root exudates of emergent plants might get diluted in the wetland environment, the ability of the microbial capacity to adhere to the root of different plants may vary (Pietrangelo et al. 2018). Root secretion of *P. australis* may have little effect on the pH of the environment, but to *S. triquetra* and *T. angustifolia*, it could change the pH of the root surface, thereby affecting the community of active methanotrophs. CH₄ emission may be related to the transcriptional activity of aerobic methanotrophs. As root-associated methanotrophs varies with plant species, it may affect the fluxes of plant mediated methane. Maltais-Landry et al. (2009) reported higher CH₄ emissions from constructed wetlands planted with *P. australis* and *P. arundinacea* than for constructed wetlands planted with *T. angustifolia*. Therefore, the community structure of active aerobic methanotrophs is more valuable for the study of natural wetland CH₄ methane emission.

As the microhabitat of bacterial life, the roots of emergent plants can be affected by various factors, such as the physical and chemical properties of sediments (Chen et al. 2018; Cui et al. 2017; Shiao et al. 2018), vegetation types (Chen et al. 2019; Yoshida et al. 2014), growth period, and the change of microenvironment is bound to lead the change of root-associated bacterial community. Bulgarelli et al. (2013) found that plant roots colonized by microbial communities have a distinctive phylogenetic structure. Is the aerobic methanotrophs affected by plant host? Studies of rhizosphere sediments have shown that vegetation types affected the community structure of methanotrophic. Zhang et al. (2018) reported that plant species had a profound impact on methanotrophic communities, and each plant species in constructed wetlands contained a specific group of methanotrophs. Yoshida et al. (2014) studied the communities of methanotrophs in the leaves, submerged part and emerged part of different aquatic plants, and drew the same conclusion. Little work was done on root-associated methanotrophs in natural wetland. In this study, beta-diversity based on both *pmoA* DNA and cDNA (Fig. 5b, Fig. 6) showed that the aerobic methanotrophic bacteria of three plants root vary with plant species. The root exudates of emergent plants in the wetland spread with water to other parts of the plant and surrounding sediments, which may be one reason why methanotrophic communities were different in rhizosphere and emerged part. In addition, Aerobic methanotrophs form an important bridge between the global carbon and nitrogen cycles including denitrification and nitrogen fixation (Bao et al. 2014b; Stein and Klotz 2011). Further investigation should be conducted, such as transcriptomic and/or metaproteomic analysis

of root-associated methanotrophs to clarify whether aerobic methanotrophs play a critical role to methane oxidation and denitrification or nitrogen fixation in natural wetland.

In this study, we demonstrated that the root-microhabitats of wetland emergent plants have significant influence on both the community structure and the active structure of aerobic methanotrophs.

Methylobacter of type I methanotrophs was predominant in root of three plant species in DNA level analysis. However, active root-associated *Methylobacter* of type II methanotrophs was predominant in root of *S. triquetra*, as well as in *T. angustifolia*. The diversity and composition of active methanotrophs were dependent on plants species. The community of active methanotrophs were affected by the host plant. These results will lay the foundation for further studies on plant-mediated methane emissions from natural wetlands.

Declarations

Acknowledgements

Authors acknowledge the member's input of the Microbial Ecology Research Team of School of Ecology and Environments of Inner Mongolia University during the sampling periods and sample analysis.

Funding

This study was funded by the National Natural Science Foundation of China grants 41563009 and 31160129, and the Science and Technology Major Project on Lakes of Inner Mongolia grant ZDZX2018054.

Author information

Affiliations

School of Life Sciences, Inner Mongolia University, Hohhot 010070, China

Jing Cui, Zheng Wang, Jumei Liu

Ministry of Education Key Laboratory of Ecology and Resource Use of the Mongolian Plateau & Inner Mongolia Key Laboratory of Grassland Ecology, School of Ecology and Environment, Inner Mongolia University, Hohhot, 010021, China

Ji Zhao, Weiwei Cao, Shaohua Zhang, Zhihua Bao

Inner Mongolia Key Laboratory of Environmental Pollution Control & Waste Resource Reuse, Inner Mongolia University, Hohhot 010021, China

Ji Zhao, Zhihua Bao

College of Chemistry and Environmental Engineering, Chongqing Key Laboratory of Environmental Materials & Remediation Technologies, Chongqing University of Arts and Sciences, Chongqing, 402160, China.

Jumei Liu

Contributions

ZB and JZ conceived and designed the research. JZ, ZB, JC, JL and WC collected samples. JC, ZW and WC conducted molecular biology experiments. SZ and JL measured the physical and chemical properties of sediment. JC and ZW analyzed data. JC and ZB wrote the manuscript. All authors read and approved the manuscript. ZB and JZ are joint corresponding authors.

Ethics declarations

Ethics approval and consent to participate

This article does not contain any studies with human participants or animals performed by any of the authors.

Consent for publication

All authors gave their consent for publication.

Competing interests

All authors declare that they have no conflict of interest.

Availability of data and material

We declared that materials described in the manuscript, including all relevant raw data, will be freely available to any scientist wishing to use them for non-commercial purposes, without breaching participant confidentiality.

References

Bao Z., Watanabe A., Sasaki K., Okubo T., Tokida T., Liu D., Ikeda S., Imaizumi-Anraku H., Asakawa S., Sato T., Mitsui H., Minamisawa K. (2014a) A rice gene for microbial symbiosis, *Oryza sativa* CCaMK, reduces CH₄ flux in a paddy field with low nitrogen input. *Appl Environ Microbiol* 80:1995-2003. doi:10.1128/AEM.03646-13

Bao Z., Okubo T., Kubota K., Kasahara Y., Tsurumaru H., Anda M., Ikeda S., Minamisawa K. (2014b) Metaproteomic identification of diazotrophic methanotrophs and their localization in root tissues of field-

grown rice plants. Appl Environ Microbiol 80:5043-5052. doi:10.1128/AEM.00969-14

Bateganya N. L., Mentler A., Langergraber G., Busulwa H., Hein T. (2015) Carbon and nitrogen gaseous fluxes from subsurface flow wetland buffer strips at mesocosm scale in East Africa. Ecol Eng 85:173-184. doi:10.1016/j.ecoleng.2015.09.081

Borruso L., Esposito A., Bani A., Ciccazzo S., Papa M., Zerbe S., Brusetti L. (2017) Ecological diversity of sediment rhizobacteria associated with *Phragmites australis* along a drainage canal in the Yellow River watershed. J Soil Sediment 17:253-265. doi:10.1007/s11368-016-1498-y

Brablcova L., Buriankova I., Badurova P., Chaudhary P. P., Rulik M. (2015) Methanogenic archaea diversity in hyporheic sediments of a small lowland stream. Anaerobe 32:24-31. doi:10.1016/j.anaerobe.2014.11.009

Bulgarelli D., Schlaeppi K., Spaepen S., Ver Loren van Themaat E., Schulze-Lefert P. (2013) Structure and functions of the bacterial microbiota of plants. Annu Rev Plant Biol 64:807-838. doi:10.1146/annurev-arplant-050312-120106

Burgmann H., Pesaro M., Widmer F., Zeyer J. (2001) A strategy for optimizing quality and quantity of DNA extracted from soil. J Microbiol Meth 45:7-20.

Cai Y., Zheng Y., Bodelier P. L., Conrad R., Jia Z. (2016) Conventional methanotrophs are responsible for atmospheric methane oxidation in paddy soils. Nat Commun 7:11728. doi:10.1038/ncomms11728

Cebon A., Bodrossy L., Chen Y., Singer A. C., Thompson I. P., Prosser J. I., Murrell J. C. (2007) Identity of active methanotrophs in landfill cover soil as revealed by DNA-stable isotope probing. FEMS Microbiol Ecol 62:12-23. doi:10.1111/j.1574-6941.2007.00368.x

Chen S., Chen J., Chang S., Yi H., Huang D., Xie S., Guo Q. (2018) Aerobic and anaerobic methanotrophic communities in urban landscape wetland. Appl Microbiol Biotechnol 102:433-445. doi:10.1007/s00253-017-8592-y

Chen X., Zhu H., Yan B., Shutes B., Xing D., Banuelos G., Cheng R., Wang X. (2019) Greenhouse gas emissions and wastewater treatment performance by three plant species in subsurface flow constructed wetland mesocosms. Chemosphere 239:124795. doi:10.1016/j.chemosphere.2019.124795

Chen Y., Dumont M. G., Cebon A., Murrell J. C. (2007) Identification of active methanotrophs in a landfill cover soil through detection of expression of 16S rRNA and functional genes. Environ Microbiol 9:2855-2869. doi:10.1111/j.1462-2920.2007.01401.x

Chen Y., Dumont M. G., Neufeld J. D., Bodrossy L., Stralis-Pavese N., McNamara N. P., Ostle N., Briones M. J., Murrell J. C. (2008) Revealing the uncultivated majority: combining DNA stable-isotope probing, multiple displacement amplification and metagenomic analyses of uncultivated *Methylocystis* in acidic peatlands. Environ Microbiol 10:2609-2622. doi:10.1111/j.1462-2920.2008.01683.x

- Conrad R. (2009) The global methane cycle: recent advances in understanding the microbial processes involved. *Environ Microbiol Rep* 1:285-292. doi:10.1111/j.1758-2229.2009.00038.x
- Costello A. M., Lidstrom M. E. (1999) Molecular characterization of functional and phylogenetic genes from natural populations of methanotrophs in lake sediments. *Appl Environ Microbiol* 65:5066-5074.
- Cui H., Su X., Wei S., Zhu Y., Lu Z., Wang Y., Li Y., Liu H., Zhang S., Pang S. (2017) Comparative Analyses of Methanogenic and Methanotrophic Communities Between Two Different Water Regimes in Controlled Wetlands on the Qinghai-Tibetan Plateau, China. *Curr Microbiol* 75:484-491. doi:10.1007/s00284-017-1407-7
- Dakora F. D., Phillips D. A. (2002) Root exudates as mediators of mineral acquisition in low-nutrient environments. *Plant Soil* 245:35-47. doi:10.1023/A:1020809400075
- Dhote S., Dixit S. (2009) Water quality improvement through macrophytes—a review. *Environ Monit Assess* 152:149-153. doi:10.1007/s10661-008-0303-9
- Duan X. N., Wang X. K., Mu Y. J., Ouyang Z. Y. (2005) Seasonal and diurnal variations in methane emissions from Wuliangsu Lake in arid regions of China. *Atmos Environ* 39:4479-4487. doi:10.1016/j.atmosenv.2005.03.045
- Eller G., Frenzel P. (2001) Changes in activity and community structure of methane-oxidizing bacteria over the growth period of rice. *Appl Environ Microbiol* 67:2395-2403. doi:10.1128/AEM.67.6.2395-2403.2001
- Esson K. C., Lin X., Kumaresan D., Chanton J. P., Murrell J. C., Kostka J. E. (2016) Alpha- and Gammaproteobacterial Methanotrophs Codominate the Active Methane-Oxidizing Communities in an Acidic Boreal Peat Bog. *Appl Environ Microbiol* 82:2363-2371. doi:10.1128/AEM.03640-15
- Fausser A. C., Hoppert M., Walther P., Kazda M. (2012) Roots of the wetland plants *Typha latifolia* and *Phragmites australis* are inhabited by methanotrophic bacteria in biofilms. *Flora* 207:775-782. doi:10.1016/j.flora.2012.09.002
- Fish J. A., Chai B., Wang Q., Sun Y., Brown C. T., Tiedje J. M., Cole J. R. (2013) FunGene: the functional gene pipeline and repository. *Front Microbiol* 4:291. doi:10.3389/fmicb.2013.00291
- Griffiths R. I., Whiteley A. S., O'Donnell A. G., Bailey M. J. (2000) Rapid method for coextraction of DNA and RNA from natural environments for analysis of ribosomal DNA- and rRNA-based microbial community composition. *Appl Environ Microbiol* 66:5488-5491. doi:10.1128/aem.66.12.5488-5491.2000
- Grunfeld S., Brix H. (1999) Methanogenesis and methane emissions: effects of water table, substrate type and presence of *Phragmites australis*. *Aquat Bot* 64:63-75. doi:10.1016/S0304-3770(99)00010-8
- Hanson R. S., Hanson T. E. (1996) Methanotrophic bacteria. *Microbiol Rev* 60:439-471.

- Heyer J., Galchenko V. F., Dunfield P. F. (2002) Molecular phylogeny of type II methane-oxidizing bacteria isolated from various environments. *Microbiology* 148:2831-2846. doi:10.1099/00221287-148-9-2831
- Hornibrook E. R. C., Bowes H. L., Culbert A., Gallego-Sala A. V. (2009) Methanotrophy potential versus methane supply by pore water diffusion in peatlands. *Biogeosciences* 6:1490-1504.
- Hutchens E., Radajewski S., Dumont M. G., McDonald I. R., Murrell J. C. (2004) Analysis of methanotrophic bacteria in Movile Cave by stable isotope probing. *Environ Microbiol* 6:111-120.
- Kits K. D., Campbell D. J., Rosana A. R., Stein L. Y. (2015) Diverse electron sources support denitrification under hypoxia in the obligate methanotroph *Methylobacterium album* strain BG8. *Front Microbiol* 6:1072. doi:10.3389/fmicb.2015.01072
- Laanbroek H. J. (2010) Methane emission from natural wetlands: interplay between emergent macrophytes and soil microbial processes. A mini-review. *Ann Bot-London* 105:141-153. doi:10.1093/aob/mcp201
- Lin J. L., Radajewski S., Eshinimaev B. T., Trotsenko Y. A., McDonald I. R., Murrell J. C. (2004) Molecular diversity of methanotrophs in Transbaikalian soda lake sediments and identification of potentially active populations by stable isotope probing. *Environ Microbiol* 6:1049-1060. doi:10.1111/j.1462-2920.2004.00635.x
- Lontoh S., DiSpirito A. A., Krema C. L., Whittaker M. R., Hooper A. B., Semrau J. D. (2000) Differential inhibition in vivo of ammonia monooxygenase, soluble methane monooxygenase and membrane-associated methane monooxygenase by phenylacetylene. *Environ Microbiol* 2:485-494. doi:10.1046/j.1462-2920.2000.00130.x
- Maltais-Landry G., Maranger R., Brisson J., Chazarenc F. (2009) Greenhouse gas production and efficiency of planted and artificially aerated constructed wetlands. *Environ Pollut* 157:748-754. doi:10.1016/j.envpol.2008.11.019
- McDonald I. R., Bodrossy L., Chen Y., Murrell J. C. (2008) Molecular ecology techniques for the study of aerobic methanotrophs. *Appl Environ Microbiol* 74:1305-1315. doi:10.1128/AEM.02233-07
- Pietrangeli L., Bucci A., Maiuro L., Bulgarelli D., Naclerio G. (2018) Unraveling the Composition of the Root-Associated Bacterial Microbiota of *Phragmites australis* and *Typha latifolia*. *Front Microbiol* 9:1650. doi:10.3389/fmicb.2018.01650
- Qiu Q., Conrad R., Lu Y. (2009) Cross-feeding of methane carbon among bacteria on rice roots revealed by DNA-stable isotope probing. *Environ Microbiol Rep* 1:355-361. doi:10.1111/j.1758-2229.2009.00045.x
- Schloss P. D., Westcott S. L., Ryabin T., Hall J. R., Hartmann M., Hollister E. B., Lesniewski R. A., Oakley B. B., Parks D. H., Robinson C. J., Sahl J. W., Stres B., Thallinger G. G., Van Horn D. J., Weber C. F. (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing

and comparing microbial communities. Appl Environ Microbiol 75:7537-7541. doi:10.1128/AEM.01541-09

Semrau J. D., DiSpirito A. A., Yoon S. (2010) Methanotrophs and copper. FEMS Microbiol Rev 34:496-531. doi:10.1111/j.1574-6976.2010.00212.x

Shiau Y.-J., Cai Y., Jia Z., Chen C.-L., Chiu C.-Y. (2018) Phylogenetically distinct methanotrophs modulate methane oxidation in rice paddies across Taiwan. Soil Biol Biochem 124:59-69. doi:10.1016/j.soilbio.2018.05.025

Shiau Y. J., Cai Y., Lin Y. T., Jia Z., Chiu C. Y. (2018) Community Structure of Active Aerobic Methanotrophs in Red Mangrove (*Kandelia obovata*) Soils Under Different Frequency of Tides. Microb Ecol 75:761-770. doi:10.1007/s00248-017-1080-1

Shinoda R., Bao Z., Minamisawa K. (2019) CH₄ oxidation-dependent ¹⁵N₂ fixation in rice roots in a low-nitrogen paddy field and in *Methylosinus* sp. strain 3S-1 isolated from the roots. Soil Biol Biochem 132, 40-46. doi: 10.1016/j.soilbio.2019.01.021

Stein L. Y., Klotz M. G. (2011) Nitrifying and denitrifying pathways of methanotrophic bacteria. Biochem Soc T 39:1826-1831. doi:10.1042/BST20110712

Sun H. Y., Zhang C. B., Song C. B., Chang S. X., Gu B. J., Chen Z. X., Peng C. H., Chang J., Ge Y. (2013) The effects of plant diversity on nitrous oxide emissions in hydroponic microcosms. Atmos Environ 77:544-547. doi:10.1016/j.atmosenv.2013.05.058

Tamura K., Peterson D., Peterson N., Stecher G., Nei M., Kumar S. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. Mol Biol Evol 28:2731-2739. doi:10.1093/molbev/msr121

Vymazal J. (2013) Emergent plants used in free water surface constructed wetlands: A review. Ecol Eng 61:582-592. doi:10.1016/j.ecoleng.2013.06.023

Wang Y., Yang H., Ye C., Chen X., Xie B., Huang C., Zhang J., Xu M. (2013) Effects of plant species on soil microbial processes and CH₄ emission from constructed wetlands. Environ Pollut 174:273-278. doi:10.1016/j.envpol.2012.11.032

Wu L. H., Wang X. Y., Li J. Y., Yu R. H., Wang L. X., Zhao J. (2017) Tag-Encoded Pyrosequencing Analysis of Bacterial Communities in the Sediments of a Eutrophic Lake on the Inner Mongolian Plateau. J Biobased Mater Bio 11:271-277. doi:10.1166/jbmb.2017.1669

Yoshida N., Iguchi H., Yurimoto H., Murakami A., Sakai Y. (2014) Aquatic plant surface as a niche for methanotrophs. Front Microbiol 5:doi:10.3389/fmicb.2014.00030

Zhang K., Luo H., Zhu Z., Chen W., Chen J., Mo Y. (2018) CH₄ flux and methanogen community dynamics from five common emergent vegetations in a full-scale constructed wetland. Environ Sci Pollut R 25:26433-26445. doi:10.1007/s11356-018-2692-9

Table

Table.1 The diversity index of high-throughput sequencing based on *pmoA* of three plant roots

	DNA			cDNA		
	PAR	TAR	STR	PAR	TAR	STR
Sequences	17618	20721	21403	18378	17350	17616
ChaoI	17.50±2.02	17.17±3.56	19.17±0.73	36.33±2.73	22.33±2.03	17±2.08
Simpson	0.37±0.010	0.57±0.012	0.63±0.004	0.22±0.008	0.49±0.012	0.17±0.003
Shannon	1.31±0.036	0.87±0.018	0.94±0.009	1.90±0.049	2.11±0.030	1.23±0.015

Note: PAR, *Phragmites australis* root; TAR, *Typha angustifolia* root; STR, *Scirpus triqueter* root

Figures

Fig. 1

Figure 1

Sampling. Localization(a) of the sampling site in the natural wetland of Wuliangsu Lake.

Fig. 2

Figure 2

Relative abundant of aerobic methanotroph community structure in different emergent plants at the genus level based on DNA (a) and cDNA (b) analysis of pmoA (PAR, Phragmites australis root; TAR, Typha angustifolia root; STR, Scirpus triqueter root).

Fig. 3

Figure 3

Relative abundant of four main genera in both DNA(a) and cDNA(b).

Fig. 4

Figure 4

Neighbor-joining tree of methanotrophic phylotypes detected in the emergent plants based on the cDNA of *pmoA* genes. The blue part of the inner circle represents type I; the pink part represents type II. The different colors in the outer circle represent the corresponding class of OUT. Multi value bar charts represent the relative abundance of OTU in different plants.

Fig. 5

Figure 5

Alpha diversity and beta-diversity of aerobic methanotroph calculation based on pmoA. The OTUs evenness of root microbiotas is shown by Shannon(a). The red boxplot based on DNA; the green boxplot based on cDNA. Error bars represent the standard error of the mean. Asterisks denote statistically significant differences between samples (**p < 0.01, *p < 0.05). The distance between samples of principal coordinates analysis (PCoA) (b) calculated using Bray–Curtis index.

Fig.6

Figure 6

Beta diversity and methanogenic community structure of different emergent plants at the genus level. Bray-cutis matrix heatmap of samples based on DNA(a) and cDNA(b).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterialAMB.pdf](#)