Health-related quality of life among cervical cancer survivors at a tertiary hospital in Kumasi, Ghana

Kwabena Amo Antwi (✉ amoantwikwabena@yahoo.com)

Komfo Anokye Teaching Hospital <https://orcid.org/0000-0002-9412-0673>

Ramatu Agambire

Faculty of Nursing, Garden City University College, Kumasi

Thomas Opkoti Konney

School of Medicine & Dentistry, Kwame Nkrumah University of Science & Technology, Kumasi

Yvonne Nartey

Division of Epidemiology and Public Health, University of Nottingham

Adu-Appiah Kubi

School of Medical Science, University of Health and Allied Sciences, Ho

Samuel Blay Nguah

School of Medicine & Dentistry, Kwame Nkrumah University of Science & Technology, Kumasi

Edward T. Dassah

School of Medicine & Dentistry, Kwame Nkrumah University of Science & Technology, Kumasi

Patrick K. Akakpo

School of Medical Science, University of Cape Coast, Cape Cape

Baafour K. Opoku

School of Medicine & Dentistry, Kwame Nkrumah University of Science & Technology, Kumasi

Frank Ankobea

School of Medicine & Dentistry, Kwame Nkrumah University of Science & Technology, Kumasi

Augustine Tawia

Komfo Anokye Teaching Hospital, Kumasi

Maxwell K. Nti

Komfo Anokye Teaching Hospital, Kumasi

Rex Mawuli Djokoto

School of Medicine & Dentistry, Kwame Nkrumah University of Science & Technology, Kumasi

Amponsah Peprah

School of Medicine & Dentistry, Kwame Nkrumah University of Science & Technology, Kumasi

Mavis Bobie Ansah

Komfo Anokye Teaching Hospital, Kumasi

Carolyn Johnston

University of Michigan, Ann Arbor

Research

Keywords: Cervical cancer survivors, Health-related quality of life, HRQoL, EORTC QLQ C-30, EORTC QLQ-CX24

Posted Date: January 27th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-110172/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Cervical cancer survivors often experience significant diminution in health-related quality of life (HRQoL). We aimed to investigate the overall HRQoL, determine the role of the stage of disease and type of treatment received on HRQoL, and evaluate predictors of HRQoL among cervical cancer survivors in Kumasi, Ghana.

Methods: A hospital-based cross-sectional study was conducted in 153 cervical cancer patients who completed curative treatment between January 2004 and December 2018 at Komfo Anokye Teaching Hospital. The European Organization for Research and Treatment of Cancer core questionnaire (EORTC QLQ C-30) supplemented with the cervical cancer-specific (EORTC QLQ-CX24) module was used. The Kruskal-Wallis test was used to determine the effect of the stage of cervical cancer and the type of treatment received on mean scores of the different domains of HRQoL. Multivariate logistic regression was performed to identify predictors of HRQoL. $P < 0.05$ was considered statistically significant.

Results: The mean global health score (GHS) was 79.7 (± 16.2), and it differed significantly with International Federation of Gynaecology and Obstetrics (FIGO) stage, with 84.1, 76.2, and 79.9, for stages I, II and III respectively ($p=0.012$). Financial burden was higher in participants with FIGO stage II compared to stage I disease (45.4 vs 20.8, $p=0.012$). The mean GHS scores for surgery, chemoradiation and radiation only were 85.2, 75.9, and 82.1 respectively ($p=0.027$). Compared to participants who were treated with Chemoradiation, those treated with surgery had lower financial difficulties (12.1 vs 41.6, $p=0.019$), better body image (95.7 vs 79.5, $p=0.039$) and better symptom experience (5.9 vs 12.0, $p=0.043$). The likelihood that survivors' GHS is affected is decreased with illiteracy (AOR = 0.30, 95% CI=0.09–1.00), and increased with complaints of pain (AOR=5.50, 95% CI=1.68-18.29), loss of appetite (AOR=13.24, 95% CI=2.71– 64.67) and diminution in body image perception (AOR = 6.04, 95% CI=1.67– 21.83).

Conclusion: Cervical cancer survivors in Kumasi, Ghana have overall satisfactory HRQoL. Surgical treatment is associated with improved HRQoL and less financial burden. Efforts to enhance HRQoL should also be focused on maximizing survivors' body image and managing pain and loss of appetite. Educating women about expected impacts on their quality of life, and strategies to manage and mitigate these impacts, is essential.

Background

Although the incidence of cervical cancer (CC) remains low in many countries, it is significantly higher in developing countries.^{1,2} Cervical cancer is the second most common cancer among women in Ghana. Treatment of CC is primarily by surgery or radiation therapy, with chemotherapy as an important adjunct.³ In Ghana, only 5.2% of women with CC present with Stage I disease, and are likely to benefit from surgery.⁴ The risk of nodal involvement in Stage IA1 disease is low, and surgical treatment by conization and simple hysterectomy is often adequate.³ Approximately, 7.4 % of patients with stage IA2

disease have pelvic lymph node involvement (PLNI), and modified radical hysterectomy and pelvic lymphadenectomy (PLND), with or without bilateral salpingo-oophorectomy, is standard treatment. Cone biopsy and radical trachelectomy plus PLND are options for women who desire future fertility.^{5,6} Radiotherapy may also be used to provide comparable treatment outcomes for early invasive disease in women with severe comorbidities, such as morbid obesity with attendant detrimental anaesthetic risk, uncontrolled hypertension, or diabetes mellitus.³ In the majority of low and middle income countries (LMICs) including Ghana, most patients present at a later stage where radiotherapy is the treatment of choice.^{4,7} The treatment of locally advanced disease (FIGO Stage IB3-IVA) has changed over the years from radiotherapy alone to concurrent chemoradiotherapy, with a number of studies demonstrating survival advantage of cisplatin-based chemotherapy in both adjuvant and definitive setting.^{8–10}

Cervical cancer survivors often experience physical symptoms and psychosocial distress that adversely affect their health-related quality of life (HRQoL). HRQoL encompasses domains of life which are directly affected by the presence of disease or treatments.¹¹ Several generic and disease-specific tools have been developed for the assessment of HRQoL. These are multidimensional, patient-reported tools, which assess aspects of life that are directly affected by changes in health.^{11,12} The European Organisation for Research and Treatment of Cancer (EORTC) HRQoL tool is a commonly used questionnaire to assess quality of life among all cancer patients (e.g. EORTC QLQ C30), and can be supplemented by the EORTC-QLQ CX-24 to focus specifically on cervical cancer patients or survivors.^{12,13}

In high-resource settings, the focus of cancer care has shifted from traditional treatment outcome measures (e.g. 5-year survival outcomes) to patients' symptoms and quality of life as they progress through treatment and into post-treatment life with cancer as a chronic disease.¹³ However, in low-resource settings like Ghana, there is limited data on HRQoL among cervical cancer patients and survivors. An assessment of the impact of patient- and treatment- related factors on the survivor's quality of life (QoL) will provide insight into whether the survival benefits are justifiable, when compared with limitations posed by patients and treatment characteristics. We aimed to investigate the overall HRQoL, HRQoL by stage and treatment received, and predictors of HRQoL among cervical cancer survivors in a single centre in Kumasi, Ghana. Results can help set targets for planners, treatment benchmarks for healthcare providers and create national benchmarks for care that correlate with improved quality of life for survivors.

Methods

Study setting

The study was conducted at Komfo Anokye Teaching Hospital (KATH), the second largest teaching hospital in Ghana. It is located in Kumasi, the capital city of Shanti Region. The hospital was established in 1942 and has more than 1000 beds providing diagnostic and treatment services for about 260,000 to 300,000 patients per year. The logistics and human resource base for gynaecologic oncologic services have developed through international cooperation and perform full multi-disciplinary care for gynaecologic cancer patients.¹⁴ The Gynaecologic Oncology Unit and Department of Oncology provide services for over 400 women with genital cancer annually. Ethical approval was obtained from the Committee on Human Research, Publications and Ethics (CHRPE) of the School of Medical Sciences, Kwame Nkrumah University of Science and Technology and KATH (**CHRPE/AP/661/19**) Written and verbal informed consent were obtained for participants interviewed via face-to-face interaction and telephone respectively.

Study design and participants

A hospital-based cross-sectional study was conducted from July to September 2019. Participants were defined as women diagnosed with cervical cancer who completed curative treatment between 2004 and December 2018, at Unit of Gynaecologic Oncology and the Department of Radiation Oncology of KATH. A total of 462 records of women with disease-free status after curative treatment were identified (Figure 1). An estimated sample size of 145 had adequate power to detect a global health score with a standard deviation of 21.5 and a desired margin of error of 3.5 scores.¹⁵ Considering a 10% of contingency for inappropriate and nonresponses, computer-generated simple random sampling without replacement was used to select 160 participants. A total of 153 women were willing to participate in the study and none of the patient data were incomplete. Participants were recruited consecutively until the required sample size was reached. Survivors were considered eligible if they were disease-free at least 6 months prior to the study. Women who were treated for malignancy of another anatomic site or who were critically ill were excluded from the study.

Data was collected by two trained oncology nurses working within the oncology clinic. Two days training was given for the oncology nurses focusing on; the contents of the questionnaire, the identification of patients based on the inclusion/exclusion criteria, and how to get consent. Participants were assured of anonymity and confidentiality of their information obtained in the study by excluding any personal identifier in the data collection form. They were also reassured that the report of the findings would not identify them and only the aggregate data would be reported. Telephone interviews or face-to-face interviews were conducted in Twi, a local language, or in English to administer the questionnaires. Telephone interview were conducted for patients who lived outside Kumasi city or were not scheduled/did not report for review within the study. Socio-demographic and clinical characteristics of the participants

were extracted from their medical records. All the collected data were checked for completeness by the principal investigator on a daily basis.

Study variables

- Dependent variables: global health score (GHS), functional scales and symptom scales
- Independent variables:
  - Socio-demographic characteristics such as age, time since completion of treatment, level of education, occupation, marital status, residence, proximity to treatment facility and tribe.
  - Clinical and treatment characteristics such as body mass index (BMI), parity, FIGO stage, primary treatment type, level of haemoglobin, units of blood transfused, and other comorbid conditions.

Instrument and scoring

The evaluation of the HRQoL was performed using the EORTC core questionnaire (EORTC QLQ C-30), supplemented with the cervical cancer-specific module EORTC QLQ-CX24.¹⁶ The 30-item EORTC QLQ-C30 is the most frequently used HRQoL assessment tool. Data on the use of the tool in Ghana is scarce, but it has been validated in many geographic and cultural settings.^{15,17} It is comprised of 15 subscales, 5 functional domains (physical, role, cognitive, emotional and social functioning); 3 multi-item symptom scales (fatigue, pain, and nausea/vomiting), 6 single items that assess additional symptoms commonly reported by cancer patients (dyspnoea, appetite loss, sleep disturbance, constipation, and diarrhoea), perceived financial difficulties and GHS. The EORTC QLQ-CX24 is a 24-item scale, grouped into 3 multi-item scales, 11 items with symptom experience domain, 3 items with body image domain, and 4 items with sexual/vaginal functioning domain. Further, it has single-item scales, that assess lymphoedema, peripheral neuropathy, menopausal symptom, sexual worry, sexual activity, and sexual enjoyment. All scores on the EORTC QLQ-C30 and QLQ-CX24 were transformed into 0 to 100 scale according to the EORTC QLQ scoring manual.¹⁸ A higher score on the GHS and functioning scale represented a better level of functioning. A higher score on the symptom scale represented worse levels of symptoms/ high level of problems or ill-health.

Data Analysis and interpretation

Descriptive statistics were used to present the QoL scores, and patient- and treatment -related variables. The Kruskal-Wallis test was used to determine the effect of stage of disease and type of treatment received on mean score of the different domains of HRQoL. The HRQoL domains were then dichotomized; functional domain and global health score/QoL score < 75 were considered affected (75 or above score indicated “no problem at all”) whereas on the symptom scale, a score 25 and above were

considered affected (below 25 indicated “no problem at all”).¹⁹ Stepwise multivariate logistic regression was performed to identify predictors of HRQoL. Only variables with $p < 0.25$ and clinically significant variables were included in the multivariable logistic regression model. All hypotheses involved were two-sided tests; $p < 0.05$ was considered statistically significant. An internal consistency of the multi-item scales was calculated using the Cronbach α coefficient; values of $\alpha \geq 0.70$ were considered acceptable for group comparisons. All analyses were performed using Stata 13 (StataCorp, College Station, TX, USA).

Results

Sociodemographic, clinical and treatment characteristics of cervical cancer survivors

A total of 153 women completed the questionnaires and seven declined participation on personal reasons. Nine survivors were critically ill (Figure 1). The mean ($\pm$ SD) age of the survivors was 58.3 ($\pm$ 11.4) years, with a median follow up time of 33.8 months (range: 6.3 to 180.5 months) after completion of treatment. (Table 1). Time since completion of treatment was more than one year in the majority (86.3%) with 29.4% being long-term survivors. Regarding level of education, 61 (39.9%) had no formal education and (11.1%) received senior high school or tertiary education. About fifty-seven percent had anaemia, necessitating haemotransfusion in a third (32.0%) of the cases, before or during treatment (Table 2). Most participants had FIGO stage III cervical cancer (37.3%). Primary chemoradiation was the treatment option in about half (47.7%) of the women. Twenty-eight (18.3%) participants underwent primary radical surgery with or without adjuvant radiotherapy.

Health-related quality of life (HRQoL) of cervical cancer survivors

The mean GHS ($\pm$ SD) score was 79.7 ($\pm$ 16.2). The EORTC QLQ-C30 multiple and single-item scales were also calculated and the mean ($\pm$ SD) score of the functional scales ranged from 88.2 ($\pm$ 14.1) to 91.8 ($\pm$ 18.4), with the highest being social functioning and the lowest being cognitive functioning (Table 3). For the symptom scales, the mean ($\pm$ SD) financial difficulty score was the highest 32.8 ($\pm$ 40.9) followed by pain 17.9 ($\pm$ 24.0) and fatigue 14.0 ($\pm$ 19.0). The EORTC QLQ-CX24 domains mean ($\pm$ SD) score ranged from 50.0 ($\pm$ 40.7) for sexual enjoyment to 85.3 ($\pm$ 26.0) for body image. On the cervical cancer-specific symptom scales, the lowest mean score 7.0 ($\pm$ 19.7) was for lymphoedema and the highest was sexual worry 21.1 ($\pm$ 35.4). The pattern of change in the EORTC QLQ C-30 and EORTC QLQ CX-24 score after treatment is displayed in Figures 2-4.

Mean score differences of EORTC QLQ-C30 and QLQ-CX24 scales with stage and treatment characteristics

On the EORTC QLQ-C30 scale, only GHS/Overall QoL and financial difficulty score differed significantly with regards to the FIGO stage and primary treatment type. (Tables 4). Participants who survived with stage I disease had a better GHS/Overall QoL score with lower financial difficulties compared to those who had FIGO stage II and III disease. More women with Stage II disease received chemoradiation. On the cervical cancer specific module, body image and symptoms differed significantly with respect to the treatment modality applied, with surgical treatment resulting in better GHS and body image and less symptoms compared to radiotherapy with or without chemotherapy. There was no difference in financial difficulties by treatment type.

Predictors of health-related quality of life

The outcome of the multivariate logistic model for HRQoL is displayed in table 5. For participants who were illiterate (AOR = 0.30, 95% CI = 0.09–1.00), they were 70% less likely to be classified as “affected” in the GHS dichotomized scale. For those who complained of pain (AOR = 5.50, 95%CI=1.68-18.29), loss of appetite (AOR = 13.24, 95% CI = 2.71– 64.67) or diminution in body image (AOR = 6.04, 95% CI = 1.67– 21.83) were more likely to have global health score affected.

Discussion

This is a maiden study conducted in Ghana to investigate HRQoL among cervical cancer survivors. The GHS/overall QoL was satisfactory. The score in functional domains were high, with highest scores seen in social functioning, followed by role functioning. Financial difficulties, pain and fatigue were the most frequently reported symptoms. Participants with FIGO stage II cervical cancer had lower GHS/overall QoL score and greater financial difficulties. Compared to other modes of treatment, surgery was associated with better overall QoL and body image. Chemoradiation was associated with more clinical complaints and financial difficulties. Illiteracy, body image, pain and loss of appetite were significantly associated with overall GHS/overall QoL. Cervical cancer and its treatment have significant impacts on the HRQoL, and thus its management should not be focused only on overall survival, but rather a holistic care also aimed at maximizing QoL.

Time since completion of treatment has effects on participants’ perception of HRQoL.²⁰ In our study, the pattern of change in GHS was highest at or after 15 years of treatment. The perceived increase in GHS was later when compared to the trend reported in a similar study in China.²⁰ This difference may be attributed to the mean age of the participants and the capacity to rehabilitate subjects after treatment. Similar recovery patterns were also noticed in all the EORTC functional domains with the exception of physical functioning, which was less brisk. Survivors reported worse score on this HRQoL domain with

increasing period of survivorship.²¹ The course of GHS appears to have an inverse relationship with the pattern of symptom complaints. Report of good GHS by survivors occur at the time of improved functioning, and they are unlikely to report symptoms during this period. This may explain the low symptom experience score. Comparable studies reported score range of 15.6 to 56 and 15.9 to 47 for and pain and fatigue respectively within a year of treatment completion.^{15,17,20,22} These scores compare to the score of 20.6 and 19.4 for fatigue and pain recorded respectively within a year of the treatment in this current study. Further, we noticed an upward trend in complaint of pain within 10 years following treatment. An earlier study by Brown et al., 2014, reported 5 to 10 in 100 cervical cancer survivors complained of pain.²³

Sexual function was a primary source of distress among cancer survivors.²⁴ It is an issues of paramount importance especially when a significant number of survivors are young in this study. At the time of the interview, only 56 (36.6%) were sexually active. This was comparatively higher than the rate reported among survivors in a similar study in Iran.²⁵ The level of sexual activity among survivor may have socio-demographic determinants, with increased sexual activity in a more liberal society. ²⁶ Higher score in sexual domains is also reported in participants reporting with early disease. Sexual worry (perception that sex will be painful) was higher within the first year of treatment, but thereafter, and then improved. Actual sexual activity decreased along the period of follow-up, although vaginal function and sexual enjoyment improved.²⁰

Female survivor populations treated for an early disease tend to have good GHS/overall QoL.²⁷ In our study, the global health status for the cervical cancer survivors was 79.7 ± 16.2 . This complements the findings of a similar in Taiwan which had few women surviving with stage IV disease.²⁸ A study that had relatively more women with advanced cervical cancer patient reported with a global health status of 59.5 ± 10.9 (after excluding critically ill patients).²⁹ The differences in the global health status scores between these studies is related to the differences in treatment options and time since completion of treatment.

Participants treated for FIGO stage I disease reported the highest GHS followed by that of stage III. This finding was consistent with that of an earlier study by Thapa et al., 2018 in Hubei, China.²⁰ However in a similar work in Malaysia, women treated for stage III reported the highest GHS.³⁰ No clear relationship exists between GHS and the FIGO stage of cervical cancer. The disparity in perceptions of GHS may be explained by the response shift theory, which postulates that perception of overall health is influenced by a change in ones' condition of health.³¹ Survivors who receive radiation with or without chemotherapy have worse HRQoL.^{20,26} Relatively more women who survived FIGO stage II disease had chemoradiation,

that may underline the poor perception of GHS in this group of survivors. The actual cost of treatment for stage II or III along with the cost accrued from loss of productivity is high with chemoradiation.³²

Treatment of cervical cancer is primarily by surgery or radiation therapy, with chemotherapy as valuable adjunct.³ In consonance with earlier studies, survivors who received surgery as a primary treatment modality in this study had lesser impact on their GHS, followed by survivors who received radiotherapy alone. The GHS, body image, financial difficulties and symptoms reported by survivors differed with the treatment option. In this study, surgery was associated with the highest GHS/overall QoL, which is consistent with earlier studies.^{17,20} Earlier studies also reported more symptoms in women treated with radiotherapy with or without chemotherapy.³³ The symptoms experienced after treatment may underline the significant alteration in body image observed across the various treatment modalities. The loss of reproductive organs and external scarring of the genitalia via radical surgery and radiotherapy respectively, have been reported to negatively impact survivors' psychophysical identity and cause distortion of the body image.³³ The absence of uterus and cervix, and the inability to reproduce traditionally, although often overlooked, is a major source of stress or discomfort to young women living with or surviving cancer.³⁴ These organs are seen by survivors as embodiment of womanhood, femininity and fertility. In most of the cultures in Sub-Saharan Africa, childbearing is a significant expression of femininity.

Prior studies demonstrate that women who had radiotherapy, were worse off financially, compared to those who received surgery alone or surgery with postoperative radiotherapy (PORT).^{17,26} These findings were mirrored in this current study. Longer treatment time, and the need to suspend work during and after treatment may explain this phenomenon.³⁵ Minimal use of PORT maximizes the gains after surgery, and multi-modal treatment may unnecessarily overburden the surgical and radiation oncology facilities which are already inadequate in low-resource countries.^{36,37} This places emphasis on diligent pre-treatment assessment especially in subjects with operable tumours. MRI should be considered in women being prepared for radical hysterectomy and pelvic lymph node dissection. Although accurate information about tumour size and lymph node involvement can be obtained from surgical pathology, it is desirable to estimate tumour volume and the presence of lymph node metastases or adverse prognostic findings before intended surgery. Women with node-positive disease or other surgical risk factors could then be managed appropriately, avoiding the potential drop in GHS associated with a combination of radical surgery and post-operative radiotherapy.³⁸

Disparities may exist in terms of demographic factors for cervical cancer patients and survivors. While cervical cancer patients tend to be less educated and often dwell in rural areas, survivors as reported by this study were better educated and only few had rural residence.³⁹ Surprisingly, survivors with no formal education were less likely to describe their GHS as "affected". An earlier study on cervical cancer

survivors did not report the role of level of education on HRQoL.²⁰ Pain was a major predictor of GHS among this group of survivors, which is consistent with an earlier study with similar clinical and treatment characteristics.²⁸ The cause of pain in the survivor may not necessarily be attributable to the cancer or its management.²³ Current interventions in cancer care may afford survivors the chance to live longer and experience pain that may be secondary to the aging process or senescence. Loss of appetite has been reported among subjects during and after treatment.^{17,25,40} Subjects who had lost appetite were likely to report an affected GHS in this study. Similar reports were recorded among survivors in Iran and Bangladesh.^{17,25} From these findings, good diet and nutrition extend well into the period of remission, and significantly influencing the QoL of the CCS. On body image, feeling less physically attractive, less feminine or been dissatisfied with the self-image were predictive of an affected GHS in this study. The absence of uterus and cervix, and the inability to reproduce traditionally, although often overlooked, is a major source of stress or discomfort to young women living with or surviving cancer.³⁴ For these survivors, the organs are traditionally seen as embodiment of womanhood, femininity and fertility, and for most of cultures in Sub-Saharan Africa, childbearing is a significant expression of femininity.

Limitations of this study include that the EORTC assessment scale has not been validated in Ghana. The closest locally relevant interpretation of questions was adopted by the researchers. The GHS of cancer survivors' changes over time. As this is a cross-sectional design, the assessment of GHS was not done over time, and the lack of the comparison of GHS score before and after treatment contribute to the limitations. Due to the retrospective nature of the treatment data, there is a possibility that some of clinical and treatment information may not have been documented. This was a single institution study so this population of cervical cancer survivors may not be representative of the cervical cancer survivor population in Ghana or in other LMICs. However, the study contributes to how to improve patient care and further research for women with CC in Ghana. Additional longitudinal and intervention studies with control groups may further evaluate the HRQoL of CCS.

Conclusion

Cervical cancer and its treatment affect survivors' HRQoL. Among cervical cancer survivors in Ghana, the HRQoL was satisfactory. Surgery offers the best prospects for HRQoL with the least financial burden. Efforts to enhance HRQoL should also be focused on maximizing survivors' body image and managing pain and loss of appetite. Educating women with cervical cancer about expected impacts on their quality of life, and strategies to manage and mitigate these impacts, is essential.

Abbreviations

CHRPE: Committee on Human Research, Publications and Ethics EORTC QLQ-CX24: European Organization for Research and Treatment of Cancer-Cervix Module; EORTC QLQ-C30: European

Organization for Research and Treatment of Cancer - core questionnaire; FIGO: International Federation of Gynaecology Obstetrics (FIGO) stage HRQoL: Health-related Quality of Life; KATH: Komfo Anokye Teaching Hospital; PLNl: pelvic lymph node involvement, PLND: Pelvic lymphadenectomy dissection, LMIC: low and middle income countries

Declarations

Competing interests

The authors declare no competing interests.

Funding

There were no external grants received for the study. All expenses related to the study were borne by the authors.

Author's contributions

KAA and YN conceived the research idea and designed the study; MBA was responsible for the data capture, KAA and SBN performed data analysis and interpretation; KAA and RA prepared the draft of the manuscript; KAA, YN, SBN, and PKA reviewed and edited the manuscript. All authors critically evaluated the final manuscript for important intellectual content and approved the final version of the manuscript.

Authors' information

KAA Gynaecologic Oncologist and Fellow of Ghana and West African Colleges of surgeons. Lecturer, School of Dental and Medical Sciences, Kwame Nkrumah University of Science and Technology.

RA is a lecturer and a nurse by profession with specialisation in public health, a Fellow of West African College of Nursing and Ghana College of Nurses and Midwives

YN is a Postdoctoral Research Fellow in Cancer Epidemiology, Division of Epidemiology and Public Health, School of Medicine, University of Nottingham, United Kingdom

TOK: Gynaecologic Oncologist and Fellow of Ghana and West African Colleges of surgeons. Lecturer, School of Dental and Medical Sciences, Kwame Nkrumah University of Science and Technology.

AAK: Gynaecologic Oncologist and Fellow of Ghana Colleges of Surgeons. Lecturer, School of Dental and Medical Sciences, Kwame Nkrumah University of Science and Technology.

SBN: Fellow of Ghana and West African Colleges of Physicians. Senior Lecturer, School of Dental and Medical Sciences, Kwame Nkrumah University of Science and Technology.

EDT: Public Health Physician and Fellow of Ghana and West African Colleges of surgeons. Senior Lecturer, School of Dental and Medical Sciences, Kwame Nkrumah University of Science and Technology.

FA is a Senior Lecturer and Head of Department, Department of Obstetrics and Gynaecology

AT is a Specialist, Department of Obstetrics and Gynaecology, Komfo Anokye Teaching Hospital

MKN is a Specialist, Department of Obstetrics and Gynaecology, Komfo Anokye Teaching Hospital

RMD: Fellow West African Colleges of surgeons and a Lecturer, School of Dental and Medical Sciences, Kwame Nkrumah University of Science and Technology

AP: Fellow Ghana Colleges of Surgeons and a Lecturer, School of Dental and Medical Sciences, Kwame Nkrumah University of Science and Technology

KAP is Pathologist and a Senior Lecturer, School of Medical Science, University of Cape Coast, Cape Coast-Ghana.

MBA is Specialist Oncology Nurse, Department of Radiation oncology, Komfo Anokye Teaching Hospital, Kumasi-Ghana

BKO is a Fellow of Ghana and West African Colleges of Surgeons. Medical Director, Komfo Anokye Teaching Hospital. Associate Professor of Obstetric of Obstetrics & Gynaecology, School of Dental and Medical Sciences, Kwame Nkrumah University of Science and Technology.

CJ is a Professor of Gynaecologic Oncology, University of Michigan

Acknowledgements

We would like to extend our sincere gratitude to all staffs of the Unit of Gynaecologic Oncology and the Department of Radiation Oncology of KATH, especially Abigail Osei-Mensah and Abigail Owusu Sekyere for their immense role during the interview. The authors acknowledge the support offered by the following people during the preparation of the manuscript: Dr. Abdul-Razak Abdul-Munin, Prof Alexandra T. Odoi Prof. Henry S. Opare-Addo and Dr. Ernest Bawuah Bonsu.

Availability of data and materials

For data protection the data set is not publicly accessible. However, data can be accessed from the primary or corresponding author upon reasonable request with a signature of data privacy form.

References

1. Cervical cancer control in developing countries: memorandum from a WHO meeting. Bull World Health Organ. 1996;74(4):345–51.
2. Laryea DO, Awuah B, Amoako YA, Osei-Bonsu E, Dogbe J, Larsen-Reindorf R, et al. Cancer incidence in Ghana, 2012: evidence from a population-based cancer registry. BMC Cancer. 2014 May 23;14:362.
3. Bhatla N, Aoki D, Sharma DN, Sankaranarayanan R. Cancer of the cervix uteri. Int J Gynecol Obstet. 143(S2):22–36.
4. Nartey Y, Hill PC, Amo-Antwi K, Nyarko KM, Yarney J, Cox B. Factors Contributing to the Low Survival Among Women With a Diagnosis of Invasive Cervical Cancer in Ghana. Int J Gynecol Cancer Off J Int

- Gynecol Cancer Soc. 2017;27(9):1926–34.
5. Du R, Li L, Ma S, Tan X, Zhong S, Wu M. Lymph nodes metastasis in cervical cancer: Incidences, risk factors, consequences and imaging evaluations. *Asia Pac J Clin Oncol*. 2018 May 31;
  6. Ware RA, van Nagell JR. Radical hysterectomy with pelvic lymphadenectomy: indications, technique, and complications. *Obstet Gynecol Int*. 2010;2010.
  7. Nkyekyer K. Pattern of gynaecological cancers in Ghana. *East Afr Med J*. 2000 Oct;77(10):534–8.
  8. Keys HM, Bundy BN, Stehman FB, Muderspach LI, Chafe WE, Suggs CL, et al. Cisplatin, radiation, and adjuvant hysterectomy compared with radiation and adjuvant hysterectomy for bulky stage IB cervical carcinoma. *N Engl J Med*. 1999 Apr 15;340(15):1154–61.
  9. Morris M, Eifel PJ, Lu J, Grigsby PW, Levenback C, Stevens RE, et al. Pelvic radiation with concurrent chemotherapy compared with pelvic and para-aortic radiation for high-risk cervical cancer. *N Engl J Med*. 1999 Apr 15;340(15):1137–43.
  10. Rose PG, Bundy BN, Watkins EB, Thigpen JT, Deppe G, Maiman MA, et al. Concurrent cisplatin-based radiotherapy and chemotherapy for locally advanced cervical cancer. *N Engl J Med*. 1999 Apr 15;340(15):1144–53.
  11. Cella DF. Quality of life: concepts and definition. *J Pain Symptom Manage*. 1994 Apr;9(3):186–92.
  12. Tax C, Steenbergen ME, Zusterzeel PLM, Bekkers RLM, Rovers MM. Measuring health-related quality of life in cervical cancer patients: a systematic review of the most used questionnaires and their validity. *BMC Med Res Methodol*. 2017;17:15.
  13. Wintner LM, Sztankay M, Aaronson N, Bottomley A, Giesinger JM, Groenvold M, et al. The use of EORTC measures in daily clinical practice-A synopsis of a newly developed manual. *Eur J Cancer Oxf Engl* 1990. 2016;68:73–81.
  14. Erem AS, Appiah-Kubi A, Konney TO, Amo-Antwi K, Bell SG, Johnson TRB, et al. Gynecologic Oncology Sub-Specialty Training in Ghana: A Model for Sustainable Impact on Gynecologic Cancer Care in Sub-Saharan Africa. *Front Public Health* [Internet]. 2020 [cited 2020 Dec 21];8. Available from: <https://www.frontiersin.org/articles/10.3389/fpubh.2020.603391/full>
  15. Khalil J, Bellefqih S, Sahli N, Afif M, Elkacemi H, Elmajjaoui S, et al. Impact of cervical cancer on quality of life: beyond the short term (Results from a single institution): Quality of life in long-term cervical cancer survivors: results from a single institution. *Gynecol Oncol Res Pract*. 2015;2:7.
  16. Greimel ER, Kuljanic Vlasic K, Waldenstrom A-C, Duric VM, Jensen PT, Singer S, et al. The European Organization for Research and Treatment of Cancer (EORTC) Quality-of-Life questionnaire cervical cancer module. *Cancer*. 2006 Oct 15;107(8):1812–22.
  17. Hossain N, Akter QM, Banu F, Mahmud S. Quality of life of cervical cancer patients after completion of treatment -A study among Bangladeshi women. *Bangladesh Med Res Counc Bull*. 2016 Oct 17;41(3):131–7.
  18. Fayers PM, Aaronson NK, Bjordal K, Groenvold M, Curran D, Bottomley A. EORTC QLQ-C30 Scoring Manual. In 1995.

19. Araya LT, Fenta TG, Sander B, Gebremariam GT, Gebretekle GB. Health-related quality of life and associated factors among cervical cancer patients at Tikur Anbessa specialized hospital, Addis Ababa, Ethiopia. *Health Qual Life Outcomes*. 2020 Mar 16;18(1):72.
20. Thapa N, Maharjan M, Xiong Y, Jiang D, Nguyen T-P, Petrini MA, et al. Impact of cervical cancer on quality of life of women in Hubei, China. *Sci Rep*. 2018 Aug 10;8(1):11993.
21. Quinten C, Coens C, Ghislain I, Zikos E, Sprangers MAG, Ringash J, et al. The effects of age on health-related quality of life in cancer populations: A pooled analysis of randomized controlled trials using the European Organisation for Research and Treatment of Cancer (EORTC) QLQ-C30 involving 6024 cancer patients. *Eur J Cancer Oxf Engl 1990*. 2015 Dec;51(18):2808–19.
22. Greimel E, Thiel I, Peintinger F, Cegnar I, Pongratz E. Prospective assessment of quality of life of female cancer patients. *Gynecol Oncol*. 2002 Apr;85(1):140–7.
23. Brown MR, Ramirez JD, Farquhar-Smith P. Pain in cancer survivors. *Br J Pain*. 2014 Nov;8(4):139–53.
24. Bergmark K, Avall-Lundqvist E, Dickman PW, Henningssohn L, Steineck G. Vaginal changes and sexuality in women with a history of cervical cancer. *N Engl J Med*. 1999 May 6;340(18):1383–9.
25. Torkzahrani S, Rastegari L, Khodakarami N, Akbarzadeh-Baghian A, Alizadeh K. Quality of life and its related factors among Iranian cervical cancer survivors. *Iran Red Crescent Med J*. 2013 Apr;15(4):320–3.
26. Greimel ER, Winter R, Kapp KS, Haas J. Quality of life and sexual functioning after cervical cancer treatment: a long-term follow-up study. *Psychooncology*. 2009 May;18(5):476–82.
27. Xie Y, Zhao F-H, Lu S-H, Huang H, Pan X-F, Yang C-X, et al. Assessment of quality of life for the patients with cervical cancer at different clinical stages. *Chin J Cancer*. 2013 May;32(5):275–82.
28. Huang H-Y, Tsai W-C, Chou W-Y, Hung Y-C, Liu L-C, Huang K-F, et al. Quality of life of breast and cervical cancer survivors. *BMC Womens Health [Internet]*. 2017 Apr 12 [cited 2017 Dec 7];17. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5389170/>
29. Dahiya N, Acharya AS, Bachani D, Sharma D, Gupta S, Haresh K, et al. Quality of Life of Patients with Advanced Cervical Cancer before and after Chemoradiotherapy. *Asian Pac J Cancer Prev APJCP*. 2016;17(7):3095–9.
30. Azmawati MN, Najibah E, Hatta MDAZ, Norfazilah A. Quality of life by stage of cervical cancer among Malaysian patients. *Asian Pac J Cancer Prev APJCP*. 2014;15(13):5283–6.
31. Bartoces MG, Severson RK, Rusin BA, Schwartz KL, Ruterbusch JJ, Neale AV. Quality of life and self-esteem of long-term survivors of invasive and noninvasive cervical cancer. *J Womens Health* 2002. 2009 May;18(5):655–61.
32. Cohn DE. Surgery or chemoradiation for stage IB cervical cancer? How cost effectiveness impacts a complex decision. *J Gynecol Oncol*. 2014 Apr;25(2):79–80.
33. Anderson B, Lutgendorf S. Quality of life in gynecologic cancer survivors. *CA Cancer J Clin*. 1997 Jul 1;47(4):218–25.

34. Kumbhaj PR, Sharma R, Bhatnagar A, Saini PK. SEXUAL FUNCTIONING & QUALITY OF LIFE IN CERVICAL CANCER SURVIVORS AFTER SURGERY AND RADIOTHERAPY -. Natl J Med Res. 2014;4(2):116–8.
35. Klee M, Thranov I, Machin D. Life after radiotherapy: the psychological and social effects experienced by women treated for advanced stages of cervical cancer. Gynecol Oncol. 2000 Jan;76(1):5–13.
36. Yeo RMC, Chia YN, Namuduri RPD, Yap SP, Soong YL, Yam PKL, et al. Tailoring adjuvant radiotherapy for stage IB-IIA node negative cervical carcinoma after radical hysterectomy and pelvic lymph node dissection using the GOG score. Gynecol Oncol. 2011 Nov;123(2):225–9.
37. Minig L, Patrono MG, Romero N, Rodríguez Moreno JF, Garcia-Donas J. Different strategies of treatment for uterine cervical carcinoma stage IB2-IIB. World J Clin Oncol. 2014 May 10;5(2):86–92.
38. Landoni F, Maneo A, Colombo A, Placa F, Milani R, Perego P, et al. Randomised study of radical surgery versus radiotherapy for stage Ib-IIa cervical cancer. The Lancet. 1997 Aug 23;350(9077):535–40.
39. Muliira R, Salas A, O'Brien B. Quality of life among female cancer survivors in Africa: An integrative literature review. Asia-Pac J Oncol Nurs. 2017;4(1):6.
40. Kamau RK, Osoti AO, Njuguna EM. Effect of diagnosis and treatment of inoperable cervical cancer on quality of life among women receiving radiotherapy at Kenyatta National Hospital. East Afr Med J. 2007 Jan;84(1):24–30.

Tables

Table 1 : Sociodemographic characteristics of cervical cancer survivors at KATH, Kumasi, Ghana

VARIABLE	N=153	PERCENTAGE (%)
Age at interview, years		
	Mean+/-SD 58.3-/ +11.4	
<40	9	5.9
40-49	27	17.7
50-59	48	31.4
60-69	40	26.1
>=70	29	18.9
Time since completion of treatment (TCT), years		
	Median 33.8 months	
< 1	21	13.7
1-5	87	56.9
>5	45	29.4
Education		
Illiterate	61	39.9
Elementary	45	29.4
Junior high	30	19.6
Senior high	13	8.5
Tertiary	4	2.6
Occupation		
Self-employed	103	67.3
Housewife	2	1.3
Civil Servant	3	2
Retired	13	8.5
Unemployed	32	20.9
Marital status		

Married	73	47.7
Widow	55	36
Divorced	22	14.4
Single	2	1.3
Cohabitant	1	0.7
Town of residence		
Urban	99	64.7
Rural	54	35.3
Region of residence		
Western Region	10	6.5
Ashanti Region	81	52.9
Brong Ahafo Region	28	18.3
Central Region	10	6.5
Other Regions*	24	15.6
Tribe/ethnicity		
Akan	123	80.4
Mole Dagbane	11	7.2
Gruni	9	5.9
Ewe	7	4.6
Others	3	2.0

Other Regions*

Table 2: Clinical characteristics of cervical cancer survivors at KATH, Kumasi, Ghana

CHARACTERISTICS	N=153	PERCENTAGE (%)
BMI (Kg/m2)	24.8-/±5.1	
Underweight (< 18.5)	10	7.9
Normal (18.5 - 24.9)	63	49.6
Over-weight (25 - 29.9)	35	27.6
Obese (> 30)	19	15
FIGO Stage		
stage I	42	27.4
Stage II	54	35.3
Stage III	58	37.3
Treatment Modality		
Surgery Alone	17	11.1
Surgery Plus Radiotherapy	11	7.2
Radiotherapy Alone	52	34.0
Chemoradiation	73	47.7
Presence of Comorbidities	Yes (%)	No (%)
Anaemia	87 (56.9)	66 (43.1)
Hypertension	49 (32.0)	104 (68.0)
*Others comorbidities	11 (11.1)	150 (88.9)
Haemoglobin level (g/dL)	Mean -/± SD 11.4 -/±1.6	
Severe Anaemia (< 8)	6	3.9
Moderate Anaemia (8 -10.9)	47	30.7
Mild Anaemia (11-11.9)	34	22.2
Normal (> or = 12)	66	43.1

Haemotransfusion		
No	111	72.6
Yes	42	27.4
Units of Blood transfused, N=42	Mean-/+ SD (3.1-/ +1.6)	
= < 2	18	11.8
3 – 4	19	12.4

*Other comorbidities: Deep vein thrombosis, diabetes, human immunodeficiency, cholelithiasis, haemorrhoid, left ventricular hypertrophy and paraparesis

Table 3: Mean score value of the EORTC QLQ-C30 and EORTC QLQ-CX24 Scales variables of cervical cancer survivors at KATH, Kumasi, Ghana

VARIABLES	Item Number	MEAN (SD)
QLQ-C30 Functional scale		
Global health status	29,30	79.7 (16.2)
Physical function	1-5	88.2 (14.1)
Role functioning	6,7	87.1 (19.5)
Cognitive functioning	20,25	80.5 (24.9)
Emotional functioning	21-24	87.8 (18.1)
Social functioning	26,27	91.8 (18.4)
QLQ-C30 Symptom scales		
Energy/fatigue	10,12,18	14.0 (19.0)
Nausea and vomiting	14,15	3.3 (10.3)
Pain	9,19	17.9 (24.0)
Short of breath	8	5.7 (15.2)
Sleep disturbance	11	9.8 (19.8)
Lack of appetite	13	7.4 (18.8)
Constipation	16	8.7 (19.0)
Diarrhea	17	3.2 (9.9)
Financial difficulty	18	32.8 (40.9)
QLQ-CX24 Functional scale		
Body image	45-47	85.3 (26.0)
Sexual activity	49	77.6 (34.0)
Sexual enjoyment, n=56	54	50.0 (40.7)
Sexual/vaginal functioning, n=56	50-53	71.7 (24.2)
QLQ-C24 Symptom scales		

Symptom experience	31-37,38,41-43	10.0 (10.3)
Lymphedema	38	7.0 (19.7)
Peripheral neuropathy	40	17.6 (26.2)
Menopausal symptoms	44	12.4 (25.0)
Sexual worry	48	21.1 (35.4)

Table 4: Kruskal Wallis Test Results: Global health score of cervical cancer survivors by FIGO Stage and Treatment options received at KATH, Kumasi, Ghana.

*Clin Rel.					
FIGO Stage					
	STAGE I	STAGE II	STAGE II		
EORTC_QLQ	N = 42	N = 54	N = 58	P	
Global Health Score	84.1 (15.7)	76.2 (16.0)	79.9 (16.2)	0.012	No
Final Difficulties	20.8 (37.4)	45.4 (41.9)	29.8 (39.7)	0.012	Yes
Primary Treatment					
*Clin Rel.					
Option					
	Surgery	Radiotherapy	Chemoradiation	p	
EORTC_QLQ	N = 28	N = 52	N = 73		
Global Health Score	85.2 (11.3)	82.1 (14.0)	75.9 (18.0)	0.027	No
Final Difficulties	12.1 (28.9)	31.7 (36.5)	41.6 (45.0)	0.019	Yes
Body Image	95.7 (15.6)	87.9 (21.9)	79.5 (30.1)	0.039	No
Symptom experience	5.9 (5.7)	9.4 (10.1)	9.4 (10.1)	0.043	No

*Clinical relevance ≥ 10 points differences

Table 5: Logistic regression analysis of global health score of cervical cancer survivors with socio-demographic, clinical characteristics, EORTC-QLQ-CX24 and EORTC-QLQ-C 30 subscale variables

Variables	GHS			
	Affected (%)	Unaffected (%)	COR (95%CI)	AOR (95%CI)*
Age at interview				
< 50	11 (28.2)	25 (21.3)	1	
50-59	11 (23.1)	37 (34.2)	0.68 (0.25-1.80)	
60-69	9 (23.1)	31 (27.2)	0.66 (0.24-1.84)	
> 70	8 (25.6)	21 (16.7)	0.87 (0.29-2.55)	
TCT				
> 5years	12 (30.7)	33 (29.1)	1	
1-5 years	20(51.3)	67 (58.7)	0.73 (0.33-1.66)	
< 1 year	6 (15.4)	15 (13.6)	0.98 (0.31-3.10)	
Level of education				
Literate	26 (66.7)	66 (57.9)	1	1
Illiterate	13 (33.3)	48 (42.1)	0.69 (0.32-1.47)	0.30 (0.09-1.00) *
Occupation				
Self-employed	24 (61.5)	79 (69.3)	1	
Civil Servant	0	3 (2.6)	1	
Housewife	1 (2.6)	1(0.9)	3.29 (0.20-54.63)	
Retired	5 (12.8)	8 (7.0)	2.06 (0.08-0.81)	
Unemployed	9 (23.1)	23 (20.2)	1.29 (0.34-2.17)	
Town of Residence				
Urban	28 (71.8)	71 (62.3)	1	
Rural	11 (28.2)	43 (37.7)	0. 65 (0.29-1.43)	
Proximity to facility				
Ashanti	20 (51.3)	61 (53.5)	1	
Brong Ahafo	6 (15.4)	22 (19.3)	0.88 (0.42-3.38)	
Central	2 (5.1)	8 (7.0)	0.82 (0.25-6.69)	
Western	6 (15.4)	4 (3.5)	4.89 (0.06-0.85)	

Northern	3 (7.7)	6 (5.3)	1.63 (0.15-2.87)
Other regions	2 (5.1)	13 (11.4)	0.82 (0.21 – 3.20)
Tribe			
Akans	32 (84.6)	91 (78.9)	1
Mole Dagbane	3 (7.7)	8 (7.0)	1.07 (0.26-4.27)
Gruni	2 (5.1)	7 (6.1)	0.81 (0.16-4.11)
Other tribes	2 (2.6)	8 (5.3)	0.71 (0.14-3.52)
Marital status			
Married	18 (46.2)	55 (48.3)	1
Widow	16 (41.0)	39 (34.2)	1.25 (0.60-2.78)
Other Status	5 (12.8)	20 (17.5)	0.76 (0.25-2.76)
Religion			
Christianity	34(87.2)	103(90.4)	1
Islam	5(12.8)	11(9.6)	1.98 (0.64-5.60)

Table 5 (Continuation I)

Variables	GHS			
	Affected (%)	Unaffected (%)	COR (95%CI)	AOR (95%CI)
BMI				
Normal weight	9 (33.3)	43 (43.4)	1	
Underweight	5 (17.9)	16 (16.2)	1.49 (0.43 - 5.13)	
Overweight	10 (35.7)	25 (25.3)	1.91 (0.22 - 1.63)	
Obese	4 (14.3)	15 (15.2)	1.27 (0.34 - 4.75)	
Parity				
≤ 4	15 (38.5)	55 (48.3)	1	
> 4	24 (61.5)	59 (51.8)	1.49 (0.71-3.13)	
Stage				
Stage I	6 (12.8)	36 (32.5)	1	
Stage II	16 (48.7)	38 (30.7)	2.52 (0.89-7.17)	
Stage III	17 (38.5)	40 (36.8)	2.55 (0.91-7.17)	
Primary Treatment				
Surgery	2 (5.1)	26 (22.8)	1	
Radiation alone	15 (33.3)	37 (34.2)	5.27 (1.11-25.04)	
Chemoradiation	22 (61.54)	51 (43.0)	5.61 (1.22-25.71)	
Haemoglobin Level				
Normal	15 (38.5)	51 (44.7)	1	
Mild	7 (18.0)	27 (23.1)	0.88 (0.33-2.42)	
Moderate	15 (38.5)	32 (28.1)	1.59 (0.69-3.70)	
Severe	2 (5.1)	4 (3.5)	1.70 (0.28-10.21)	
Units of Blood Transfused				
≤ 2	4 (33.3)	14 (46.7)	1	
> 2	8 (66.7)	16 (53.3)	1.75 (0.43-7.08)	
Physical functioning				
Unaffected	23 (59.0)	98 (86.0)	1	

Affected	16 (41.0)	16 (14.0)	4.2 (1.86-9.75)
Role Functioning			
Unaffected	23 (59.0)	96 (84.2)	1
Affected	16 (41.0)	16 (16.8)	3.71 (1.65-8.36)
Cognitive functioning			
Unaffected	17 (43.6)	84 (73.7)	1
Affected	22 (56.4)	30 (26.3)	3.62 (1.70-7.73)
Emotional functioning			
Unaffected	22 (56.4)	98 (86.0)	1
Affected	17 (43.6)	14 (14.0)	4.73 (2.08-10.80)
Social functioning			
Unaffected	24 (61.5)	105 (92.1)	1
Affected	15 (38.5)	9 (7.9)	7.29 (2.85-18.62)

Table 5 (continuation II)

Variables	GHS		AOR* (95%CI)	
			COR (95%CI)	
	Affected (%)	Unaffected (%)		
Fatigue				
Unaffected	36 (92.3)	104 (91.2)	1	
Affected	3 (7.7)	10 (8.8)	0.87 (0.23-3.33)	
Nausea & vomiting				
Unaffected	34 (87.2)	112 (98.3)	1	
Affected	5 (12.8)	2 (1.75)	8.24 (1.53 - 44.37)	
Pain				
Unaffected	14 (35.9)	90 (78.9)	1	1
Affected	25 (64.1)	24 (21.1)	6.70 (3.03-14.81)	5.5 (1.68 - 18.29) *
Dyspnea				
Unaffected	27 (69.2)	105 (92.1)	1	
Affected	12 (30.8)	9 (7.9)	5.19 (1.98-13.57)	
Insomnia				
Unaffected	20 (51.3)	101 (88.6)	1	
Affected	19 (48.7)	13 (11.4)	7.38 (3.14-17.32)	
Loss of appetite				
Unaffected	24 (61.5)	108 (94.7)	1	1
Affected	15 (38.5)	5 (4.3)	13.63 (4.52-41.11)	13.24 (2.71-64.67) *
Constipation				
Unaffected	24 (61.5)	99 (86.8)	1	
Affected	15 (38.5)	15 (13.2)	4.13 (1.78-9.59)	
Diarrhea				
Unaffected	32 (82.1)	107 (93.9)	1	
Affected	7 (17.9)	7 (6.1)	3.34 (1.09-10.24)	
Financial difficulty				
Unaffected	9 (23.1)	71 (62.3)	1	

Affected	30 (76.9)	43 (37.7)	5.5 (2.39-12.69)

Table 5 (continuation III)

Variables	GHS		COR (95%CI)	AOR* (95%CI)
	Affected (%)	Unaffected (%)		
Body image				
Unaffected	16 (41.0)	101 (88.6)	1	1
Affected	23 (58.0)	11 (11.4)	11.17 (4.72-26.41)	6.04 (1.67 – 21.83)*
Sexual activity				
Unaffected	26 (66.7)	71 (62.3)	1	
Affected	13 (33.3)	43 (37.7)	0.82 (0.38-1.78)	
Sexual enjoyment				
Unaffected	5 (38.5)	11 (25.6)	1	
Affected	8 (61.5)	32 (74.4)	0.55 (0.15-2.04)	
Sexual functioning				
Unaffected	4 (30.8)	19 (44.2)	1	
Affected	9 (69.2)	24 (55.8)	1.78 (0.47-6.69)	
Symptom experience				
Unaffected	30 (76.9)	110 (96.5)	1	
Affected	9 (23.1)	4 (3.5)	8.25 (2.37-28.65)	
Lymphoedema				
Unaffected	33 (84.6)	100 (87.7)	1	
Affected	6 (18.0)	14 (12.3)	1.70 (0.62-4.63)	
Peripheral neuropathy				
Unaffected	17 (43.6)	79 (69.3)	1	
Affected	22 (56.4)	35 (30.7)	2.92 (1.38-6.17)	
Menopausal symptom				
Unaffected	26 (66.7)	90 (79.0)	1	
Affected	13 (33.3)	24 (21.0)	1.88 (0.83-4.19)	
Sexual worry				

Unaffected	22 (56.4)	85 (74.6)	1
Affected	17 (43.6)	29 (25.4)	2.26 (1.06-4.84)

Other Status: Divorced, single/cohabitant

Other Tribes: Sissala, Gonja, Ewe and Nigerian

Other Regions: Greater Accra, Eastern, Upper East and West, Volta and Northern Regions of Ghana

COR: Crude odds ratio

*AOR: Adjusted odds ratio

TCT : Time since Completion of Treatment

Figures

Figure 1

Flow chart of the source of participants and sampling method used in the study.

Figure 2

The overall pattern of EORTC QLQ C30 functioning score in cervical cancer survivors at KATH, Kumasi, Ghana.

Figure 3

The overall pattern of EORTC QLQ C30 symptom score in cervical cancer survivors at KATH, Kumasi, Ghana.

Figure 4

The overall pattern of EORTC QLQ CX-24 functioning and symptom score in cervical cancer survivors at KATH, Kumasi, Ghana.