

The impact of parents' health behaviours on their preferences regarding vaccinations in Białystok, Poland

Jolanta Kraśnicka

Family Doctors Clinic

Elżbieta Krajewska-Kułał

Uniwersytet Medyczny w Białymstoku

Krystyna Klimaszewska

Uniwersytet Medyczny w Białymstoku

Mateusz Cybulski

Uniwersytet Medyczny w Białymstoku

Andrzej Guzowski

Uniwersytet Medyczny w Białymstoku

Jolanta Lewko

Uniwersytet Medyczny w Białymstoku

Cecylia Łukaszuk

Uniwersytet Medyczny w Białymstoku

Krystyna Kowalczyk

Uniwersytet Medyczny w Białymstoku

Halina Doroszkiewicz

Uniwersytet Medyczny w Białymstoku

Anna Baranowska

Uniwersytet Medyczny w Białymstoku

Katarzyna Krajewska-Ferishah

Uniwersytet Medyczny w Białymstoku

Hanna Rolka

Uniwersytet Medyczny w Białymstoku

Wojciech Kulak (✉ kneur2@wp.pl)

Medical Univeristy of Białystok <https://orcid.org/0000-0003-1118-9139>

Research article

Keywords: Vaccinations, health behaviours, parents' opinions

Posted Date: April 3rd, 2020

DOI: <https://doi.org/10.21203/rs.2.20442/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on July 25th, 2020. See the published version at <https://doi.org/10.1186/s12887-020-02235-1>.

Abstract

Background Currently, as the number of vaccinated children in Poland and throughout Europe is decreasing. Many factors impact on the rate vaccination and parents' health behaviours may affect the frequency of vaccinations.. The aim of the study was to assess the association of parents' health behaviors with children's vaccinations. **Methods** The study was carried out using a diagnostic survey method with an original questionnaire and the Inventory of Health Behaviours. Three hundred parents were recruited from the Pro Medica Family Medica Center in Bialystok Poland. **Results** Only 3.7% of respondents did not vaccinate their children. The level of health behaviours was average in 42.3% of the respondents, low in 33%, and high in 24.7%. Significant differences in health behaviours, mainly the level of normal eating habits ($p = 0.038$) and positive mental attitude ($p = 0.022$), were found in relation to views on the toxicity of vaccines. Participants who reported that vaccines can cause autism engaged in a higher level of prophylactic behaviours. Respondents who vaccinated their children with combined vaccines had a significantly higher level of health practices. **Conclusions** Parents preferred health behaviours did not effect on children vaccination. Parents who believed in the toxicity of vaccines were more concerned about proper nutrition, had a positive mental attitude, and engaged in a higher level of preventive behaviours and health practices. Parents who did not vaccinate their children had lower levels of normal eating habits. Parents who vaccinated their children with combined vaccines had a higher level of health behaviours, especially in terms of health practices. However, longitudinal studies are needed to confirm these results.

Background

There is a large amount of literature on health behaviours related obesity, physical activity, and diabetes [1-4]. But there are no studies evaluating the relationship between parental health behavior on vaccination of children.

Health behaviours comprise a very important element of human life that has been the subject of theoretical considerations and research by scientists from various fields for many years [5]. There are different concepts for defining and classifying health behaviours.

Health behaviours are a multifactorial set of reactions of social groups associated with treatment and healing agents and stimulated by the need to maintain health and life and prevent disease. Steptoe et al. [5] distinguished several classes of pro-health behaviours including stimulant avoidance, transportation safety, eating habits, positive health practices, and behaviours associated with preventive activities. Many factors that affect both the individual and the environment influence the formation of health behaviours [5-7].

'Vaccine hesitancy' is a new term used in last years to describe anyone who is doubtful about vaccinations or who chooses to delay or refuse immunizations even when they are readily available [8]. There are a wide range of factors that cause vaccine hesitancy including the adverse health outcomes, unfamiliarity with vaccine-preventable diseases, and lack of trust in public health agencies [9,10]. Reasons for refusing or delaying child vaccination reported by parents differ generally but can be classified into four categories: religious reasons, personal beliefs, safety concerns, and a desire for more information from healthcare providers.

Vaccinations included in the immunisation schedule are mandatory in Poland [11]. This means that every child residing in Poland can receive vaccines funded by the state, as well as that parents are obliged to show up on vaccination visits. At birth, each child receives an immunisation card that is stored at the general practitioner's (GP) office and used to monitor the immunisation schedule and progress. Based on this card, the GP calls parents for well-baby visits and administers scheduled vaccines as part of developmental monitoring. The current immunization schedule includes 11 mandatory vaccines: tuberculosis, hepatitis B, diphtheria, tetanus, pertussis, poliomyelitis, Haemophilus influenzae type b, pneumococci, measles, mumps, and rubella. The immunization schedule also includes a separate section describing which vaccines are recommended (against rotavirus, *Neisseria meningitidis*, hepatitis A, and human *papillomavirus*), but their cost must be paid by the parents

According to the Polish National Institute of Public Health [12], in Poland in 2017, 98.0% of 2-year-olds, 98.6% of 3-year-olds, 95.5% of 7-year-olds, 93.3% of 11-year-olds, 93.7% of 15-year-olds, 91.7% of 20-year-olds, and 93.3% of girls aged 14 (rubella) received the recommended *vaccines*. Despite the high vaccination rate of children and adolescents in Poland, the number of unvaccinated children has been increasing since 2010. At that time, 0.6% people had avoided their duty to vaccinate, while in 2018, about 5.4% parents refused to vaccinate their children [13]. 2011-2014, the number of people who avoided compulsory immunization more than doubled from approximately 5,000 to over 12,000 people [14,15].

A healthy lifestyle is comprised of the conscious or unconscious choices of a given individual, and vaccination is one of the most important health behaviours. It is worth noting that healthy lifestyle is greatly determined by the social and economic circumstances [16]. Parental vaccination decisions are complex and multi-dimensional.

The family is the first and the most important institution for shaping both attitudes towards maintaining health and the behavioural patterns necessary to maintain health [17]. Parents serve as fundamental role models, although they are not the only examples available to children.

Several studies described the relationship between parents' health communication behaviors and vaccinations for children [18]. Furthermore, parents with high levels of communicative and critical health literacy are less likely to vaccinate their children [19].

There is a large amount of literature in the immunisation space around parental attitudes [6,7,20-22], but there are no studies assessing the relationship between parental health behavior and vaccination of children.

The main purpose of the study was to assess the association of parents' health behaviors with children's vaccinations. We also evaluated parents' opinions on vaccination using an original questionnaire.

Methods

The study was carried out from July 2015 to June 2016 using the diagnostic survey method [23]. which is a method of gathering knowledge about structural and functional attributes and the dynamics of social

phenomena, opinions, and views of selected communities. The most common techniques used in surveys are interviews, questionnaires, and document analysis. An original questionnaire included demographic data: parents' age, gender, children's age, place of residence, education, financial status, and questions about vaccination was using. The vaccination questions were following: vaccination children, immunization is the most effective method of protection against infectious diseases; vaccine safety, compulsory vaccination source of information on vaccination (Details are shown in Table 2 and 3).

Parents were recruited from the Pro Medica Family Medica Center in Bialystok Poland. Children Center population in 2015 was 328. The 320 questionnaires were given for parents during doctor visits. The questionnaires were filled by parents in their homes.

The Polish version of the Inventory of Health Behaviour (IHB) [24], is use for testing different health practices. The IHB is intended for the study of healthy and ill adults. It contains 24 statements describing various types of health-related behaviours and enables the determination of both a general indicator of the level of health behaviours and specific indicators for the following categories of behaviours: proper nutrition (mainly taking into account the type of food consumed - including the primary types of food consumed, e.g., whole wheat bread, vegetables, and fruits), preventive behaviours (regarding compliance with health recommendations and obtaining information on health and disease), health practices (daily habits regarding sleep and rest), physical activity (daily exercise habits and recreation), and a positive mental attitude (avoiding strong emotions, stress, depressing situations) [24]. The internal reliability of the IHB scale, based on Cronbach's alpha, is 0.85 for the entire inventory; for the four subscales, internal reliability ranges from 0.60 to 0.65. According to the author's suggestions regarding the IHB scale, the scores for the health behaviour components were assessed as the average points for each subscale, while the total IHB score was determined by the sum of all obtained points. Thus, eating habits, preventive behaviours, positive mental attitude, and health practices can take values from 1 point (the worst result) up to 5 points (the best result), whereas the total scale score can range from 24 to 120 points, with higher scores indicating greater positive health behaviours.

Data analysis

The data management and analysis were conducted using Statistica version 13.0 (Statsoft, Tulsa, OK, USA). Descriptive analysis was performed by calculating frequencies and percentages of variables. Relationships between two variables were analysed using the chi-square independence test. Relationships among three groups were examined using the Kruskal-Wallis test. The critical level for all tests of significance was $p < 0.05$.

Results

The 320 questionnaires were delivered, and 314 (98%) were returned. But 300 (94%) surveys were properly completed.

The study included 300 parents; 46.3% were aged 31-40 years, 30% were aged 18 to 30 years, and 23.7% were aged 41 to 50 years. In the studied population, 83% were women, and 17% were men.

A total of 54.3% of the respondents had at least some higher education, 37.3% had a secondary education, 7.3% had a vocational education, and 1% had a primary education. Most had children aged 2 to 4 years (40%), followed by 0-23 months (25.3%), 7-10 years (22.7%), 11-15 years (22.3%), 5-6 years (21%), and 16-18 years (13.3%). Most of the respondents (57.3%) declared that their financial situation was good. The vast majority of the surveyed parents (85%) were not professionally involved in health care. Only 15% of the respondents were in the medical profession.

The vast majority of parents (96.3%) declared that they vaccinated their children. Only 3.7% of the respondents did not vaccinate their children. According to 68% of parents vaccination is the best method of preventing infectious diseases. Similarly 63% of parents reported that vaccines are safe and 65% reported that vaccinations should be compulsory. Most of respondents (84%) said that a family doctor should be a source of information on vaccinations.

In the surveyed population, health behaviours (proper eating habits, preventive behaviours, positive mental attitude and health practices) were practiced at similar levels, although the most frequently occurring health behaviour was a 'positive mental attitude'. Details are shown in Table 1.

The total IHB scale rates health behaviours on three levels: low, average, and high. This classification also takes into account the sex of the respondents, using slightly different criteria for women and men. Almost half of the parents (42.3%) had an average level of health behaviours; one in three respondents had a low level, and one in four had a high level. Details are provided in Table 1.

Next, the relationships among behaviour levels in the four areas specified on the IHB questionnaire was examined. The level of individual health behaviours was measured on a numerical scale. The analysis consisted of juxtaposing the mean IHB scores (with standard deviations) by group with the responses to the questions regarding vaccination and determining the significance of the relationship between them using the Kruskal-Wallis test. Opinions on immunity being infected with a disease were not related to the parents' health behaviours. Opinions on the need to vaccinate for all diseases, the vaccination system, and vaccine quality were not related to health behaviours. Details are shown in Table 2.

Significant differences in health behaviour levels, particularly eating habits ($p = 0.038$) and positive mental attitude ($p = 0.022$), were found among respondents with different opinions on the toxicity of vaccines. Parents who believed that vaccines were toxic cared about proper nutrition had positive mental attitudes and a higher level of preventive behaviours and health practices than other parents. Respondents who believed that vaccines can cause autism had a significantly higher level of preventive behaviours than other parents. Opinions on the general obligation to vaccinate were not related to health behaviour levels in any of the four areas identified in the IHB questionnaire. Additionally, opinions on the admission of unvaccinated children to crèches and kindergartens did not relate to the respondents' health habits. Details are shown in Table 2.

Parents who did not vaccinate their children with combined vaccines had a lower level of normal eating habits ($p = 0.058$). Parents who vaccinated their children with combined vaccines showed significantly higher levels of health practices (on average, 3.28 for this group compared with 3.12-3.13 in the other two

groups). Significant or near-significant differences in levels of various health behaviour were found between those who used recommended vaccinations and those who did not. Parents who vaccinated their children also had a higher level of health behaviours, especially in the field of health practices. Details are shown in Table 3.

Discussion

In the present study, parents preferred health behaviours did not effect on children vaccination. Only 3.7% of respondents did not vaccinate their children. Furthermore, parents with preferred health behaviours (proper eating habits and positive mental attitude) significantly more often reported that vaccines contain toxic ingredients. Also, parents with preventive behaviours more often reported that vaccines cause developmental disorders and autism. And, parents with positive mental attitudes significantly more often had doubts about vaccinating their children. Parents who vaccinated their children with combined and recommended vaccines showed significantly higher levels of health practices.

The present findings suggest less importance of parenteral health behaviour on children vaccination. It well known [25,26] that many factors impact on immunisation rates for example : social determinants such as young age of parents, level of parental education, family income, lack of health insurance, lack of periodic primary health care access, or pay for vaccines.

The role of the parents' perceptions health beliefs and attitudes toward childhood immunization [27] are risk factors for decreased vaccination. Other studies [28-30] suggest that socioeconomic factors play a more important role, and parents' beliefs may simply reflect their sociodemographic characteristics.

Our results on rate vaccination are consistent with a recent study from the United States [26]. Ninety-six percent of parents reported that their children had received all vaccines recommended for children up to their age. Moreover, 3.5% of all parents indicated they had decided not to have their child/children get a recommended vaccine. As in our study, some parents reported that vaccines have ingredients that are unsafe (35%) and 19% believed that vaccines may cause as autism.

Also, similar rate children immunisation was found in a study from Israel [31] where 90%-89% children had full immunisation in the years 2008 and 2016. However, a declining confidence of parents in official recommendations for vaccination from 87% to 72% was demonstrated.

It is believed that greater health behaviour awareness in the family results in better the health effects for its individual members [28]. The present and future health of the family system and all its members depends largely on the parents' actions, decisions, conduct, choices, and preferred lifestyle. It should be noted that parents raise children based not only on scientific and popular science knowledge but also on colloquial knowledge. These types of knowledge may have a positive or negative influence on the development of family behaviour patterns, the introduction of modifications and changes in behaviours and the consolidation of previously understood attitudes, including those related to health [14].

In the currently studied population, all categories of studied health behaviours (proper eating habits, preventive behaviours, a positive mental attitude, and health practices) were at a similar level; however, the most frequent behaviours was a positive mental attitude.

In the literature on the subject, health behaviours include the conscious undertaking of health-oriented actions (various behaviours related to physical health, mental health, self-management of health, preventive examinations, safe behaviours in everyday life, common sense behaviour regarding diseases) and the elimination of all activities that pose a threat to life and health, both directly and in the long term (e.g., abstaining from tobacco, alcohol, recreational drugs, and psychoactive substances) [3]. One of the most important health behaviours is vaccination, which is the most effective preventive method for combating diseases.

A survey of a representative sample of 990 adults in Poland showed that 79% of respondents considered vaccinations the most effective way to protect children from serious diseases. The vast majority of the surveyed parents (96.3%) declared that they vaccinated their children. The aforementioned survey of a representative random sample of 990 adults in Poland showed that 79% of respondents thought that vaccinating children causes more good than harm [15].

The present study showed that parents' health behaviours were not statistically correlated their beliefs regarding the vaccination system, the quality of the vaccines used in Poland, the general obligation to gain immunity through "sickness," the need to vaccinate against all diseases and the acceptance of unvaccinated children in nurseries and kindergartens.

In a recent study from 2020, including 5736 parents from eighteen country European, survey on parents' attitudes and behaviours regarding their children's immunization was performed. Fifty six percent respondents described themselves as "not at all hesitant", and 24% respondents "somewhat hesitant", respectively. Vaccine confidence was highest in Portugal and Cyprus, and lowest in Bulgaria and Poland [32].

At present, more and more parents avoid vaccinating their children. Negative opinions that undermine the effectiveness and safety of preventive vaccinations as a form of infection prevention are widely disseminated [8,9]. There are so-called anti-vaccine movements that aim to reduce mass vaccination. On online portals and forums, there is information regarding the harmfulness of preventive vaccinations, complications arising from vaccinations, and the impact of mercury on the emergence of autism, autoimmune diseases, or weakening of the body's resistance. Anti-vaccine content is also spread among those who use homeopathic and natural medicine [8].

In the present study, parents who believed that vaccines can cause autism showed a significantly higher level of prophylactic behaviour. Statistically, there were significant differences in health behaviour levels related to views on the toxicity of vaccines, mainly in terms of proper eating habits and a positive mental attitude. Furthermore, 8% of the respondents reported that vaccines cause autism, 45% had no opinion, and 47% reported that vaccines do not cause autism. But only 3.7% did not vaccinate their children. But, only 3.7 parents did not vaccinate their children.

Parents beliefs that vaccines can cause autism are complex [33]. Traditional approach to changing beliefs states that you need to first change values – the source of beliefs [34]. But values are strong, they are a consequence of our life experience and can not be changed so easily. Our behaviour is a direct result of our attitudes. Our attitudes are created based on our beliefs. And our beliefs come from our values [35]. A negative example of the dissemination of false information/ beliefs regarding vaccinations was the investigation by Wakefield, who put forward the concept of a relationship between the Measles, Mumps and Rubella Vaccine (MMR) vaccination and autism [36].

The most common reason for hesitancy or refusal for MMR is autism which was conformed in many previous studies [36,37]. Furthermore, vaccine-hesitant parents in Switzerland believed the risks of vaccination were worse than measles itself [38].

It is worth emphasizing the limitations of this study. First, this study involved relatively small group of parents, mainly woman. Secondly, parents were of different ages and had different educational level. Finally, 15% of the respondents were in the medical professions.

In summary, it is worth re-emphasizing that the family, as the most important, most basic social group on which society is based, should provide future generations with educational development, including vaccinations, based on cooperation with health care units and other educational units.

Further long-term studies involving a larger group of parents are needed to confirm these results.

Conclusions

In the present study, parents preferred health behaviours did not effect on children vaccination. Parents who believed in the toxicity of vaccines were more concerned about proper nutrition, had a more positive mental attitude, and had a higher level of preventive behaviours and health practices. Parents who did not vaccinate their children with combined vaccines had lower levels of normal eating habits, and those who vaccinated their children had a higher level of health behaviours, especially in the area of health practices.

Abbreviations

IHB: Inventory of Health Behavior

Declarations

Ethics approval and consent to participate

The study was approved by the bioethics committee of the Medical University of Bialystok, Poland R-I-002/196/2015 and the Directorate of Pro Medica Family Medicine Center in Bialystok, Poland. The informed consent obtained from study participants was written.

Consent for publication

Not applicable.

Availability of data and materials

The data analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

None

Authors' contributions

JK, EKK, K Klimaszewska made significant contributions to the conception and design of the study; MC AG, JL ,CŁ, K Kowalczyk, HD contributed to acquisition of data and initial analysis; AB, KKF and HR contributed to drafting of the manuscript. WK corrected the manuscript. All authors contributed to the interpretation of the data and revision of the manuscript. All authors approved the manuscript for publication.

Acknowledgements

We would like to thank Marek Sobolewski from the Quantitative Methods Department in Economics of the Faculty of Management and Marketing at the Rzeszów University of Technology, Poland for statistical analysis.

References

1. Phipps DJ, Hagger MS, Hamilton K. Predicting limiting 'free sugar' consumption using an integrated model of health behavior. *Appetite*. 2020 Mar 14;150:104668.
2. Teixeira PJ, Marques MM. Health Behavior Change for Obesity Management. *Obes Facts*. 2017;10(6):666-673.
3. Janssen E, van Osch L, de Vries H, Lechner L. Examining direct and indirect pathways to health behaviour: the influence of cognitive and affective probability beliefs. *Health Psychol* 2013;28(5):546-60.
4. Morgan PJ, Young MD, Smith JJ, Lubans DR. Targeted Health Behavior Interventions Promoting Physical Activity: A Conceptual Model. *Exerc Sport Sci Rev*. 2016;44(2):71-80.
5. Steptoe A, Gardner B, Wardle J. The role of behaviour in health. [in:] *Health Psychol*. French D, Kaptein A, Vedhara K, et al (eds). Blackwell, Oxford 2010: pp 12-32.
6. Suls J, Krantz DS, Williams GC. Three strategies for bridging different levels of analysis and embracing the biopsychosocial model. *Health Psychol*. 2013;2(5):597-601.

7. Łuszczynska A. Changing health behaviors. GWP, Gdańsk; 2004. (Polish)
8. Dubé E, Gagnon D, Nickels E, Jeram S, Schuster M. Mapping vaccine hesitancy–country-specific characteristics of a global phenomenon. *Vaccine*. 2014;32(49):6649-54.
9. Salmon DA, Dudley MZ, Glanz JM, Omer SB. Vaccine hesitancy: Causes, consequences, and a call to action. *Vaccine*. 2015;33 Suppl 4:D66-71.
10. Kennedy A, Lavail K, Nowak G, Basket M, Landry S. Confidence about vaccines in the United States: understanding parents' perceptions. *Health Aff (Millwood)*. 2011;30(6):1151-9.
11. Recommended vaccines. <https://gis.gov.pl/zdrowie/szczepienia-zalecane/> 01.2018
12. Vaccination of children and youths. <https://stat.gov.pl/en/topics/health/health/vaccinations-of-children-and-youth,5,1.html>. Accessed 22 Nov 2019.
13. The number of unvaccinated children in Poland is growing. Doctors urge parents not to believe in false theories about the harmfulness of <https://biznes.newseria.pl/news/liczba-nieszczepionych,p1691513164>, Accessed 22 Nov 2019.
14. Omyła-Rudzka M. Opinions on immunization. Raport BS/172/2013. Warszawa, 12, 2013. (Polish)
15. System szczepień ochronnych dzieci. Informacja o wynikach kontroli, 410.015. 00.2015, Nr ewid. 209/2015/P15/080/LKR, Warszawa, 2016, <https://www.nik.gov.pl/plik/id,10407,vp,12736.pdf>. Accessed 31 May 2018. (Polish)
16. Feinstein JS. The relationship between socioeconomic status and health: A review of the literature. *Milbank Q*. 1993;71(2):279–322
17. Poutiainen H, Hakulinen T, Mäki P, Laatikainen T. Family characteristics and parents' and children's health behaviour are associated with public health nurses' concerns at children's health examinations. *Int J Nurs Pract*. 2016;22(6):584-95.
18. Jung M, Lin L, Viswanath K. Associations between health communication behaviors, neighborhood social capital, vaccine knowledge, and parents' H1N1 vaccination of their children. *Vaccine*. 2013;31(42):4860-6.
19. Amit Aharon A, Nehama H, Rishpon S, Baron-Epel O. Parents with high levels of communicative and critical health literacy are less likely to vaccinate their children. *Patient Educ Couns*. 2017;100(4):768-75.
20. Rea CJ, Smith RL, Taveras EM. Associations of Parent Health Behaviors and Parenting Practices with Sleep Duration in Overweight and Obese Children. *J Clin Sleep Med*. 2016;12(11):1493-98. (Italian)
21. Garriguet D, Colley R, Bushnik T. Parent-Child association in physical activity and sedentary behaviour. *Health Rep*. 2017;28(6):3-11.
22. Plumtre I, Tolppa T, Blair M. Parent and staff attitudes towards in-hospital opportunistic vaccination. *Public Health*. 2020;182:39-44.
23. Metoda sondażu diagnostycznego. https://pl.wikipedia.org/wiki/Metoda_sondażu_diagnostycznego. (Polish) Accessed 22 Nov 2019.
24. Juczyński Z. Narzędzia pomiaru w promocji i psychologii zdrowia. Pracownia Testów Psychologicznych Polskiego Towarzystwa Psychologicznego, Warszawa, 2009. (Polish)

25. Zucs AP, Crispin A, Eckl E, Weitkunat R, Schlipkoter U. Risk factors for undervaccination against measles in a large sample of preschool children from rural Bavaria. *Infection* 2004; 32(3):127-33.
26. Rosenthal J, Rodewald L, McCauley M, Bermans S, Irigoyen M, Sawyer M, et al. Immunization coverage levels among 19- to 35-monthold children in 4 diverse, medically underserved areas of the United States. *Pediatrics* 2004; 113(4):E296-302.
27. Brenner RA, Simons-Morton BG, Bhaskar B, Das A, Clemens JD; NIH-D.C. Prevalence and predictors of immunization among inner-city infants: a birth cohort study. *Pediatrics* 2001;108(3):661-70.
28. Strobino D, Keane V, Holt E, Hughart N, Guyer B. Parental attitudes do not explain underimmunization. *Pediatrics* 1996;98 (6Pt1):1076-83.
29. Prislín R, Dyer JA, Blakely CH, Johnson CD. Immunization status and sociodemographic characteristics: the mediating role of beliefs, attitudes, and perceived control. *Am J Public Health* 1998; 88(12):1821-26.
30. Boyle J, Berman L, Nowak GJ, Iachan R, Middleton D, Deng Y. An assessment of parents' childhood immunization beliefs, intentions, and behaviors using a smartphone panel. *Vaccine*. 2020;38(10):2416-23.
31. Grossman Z, Hadjipanayis A, Degani A, Somekh E. Tracking Changes in Vaccine Attitudes and Decisions: Results from 2008 and 2016 Parental Surveys. *Pediatr Infect Dis J*. 2019;38(4):e75-e6.
32. Hadjipanayis A, van Esso D, Del Torso S, Dornbusch HJ, Michailidou K, Minicuci N, Pancheva R, Mujkic A, Geitmann K, Syridou G, Altorjai P, Pasinato A, Valiulis A, Soler P, Cirstea O, Illy K, Mollema L, Mazur A, Neves A, Završnik J, Lapii F, Efstathiou E, Kamphuis M, Grossman Z. Vaccine confidence among parents: Large scale study in eighteen European countries. *Vaccine*. 2020;38(6):1505-12.
33. Salwa A. Health education of parents in the field of immunization. *J Educ Health Sport* 2017;7(7):238-46.
34. Brown KF, Shanley R, Cowley NA, van Wijgerden J, Toff P, Falconer M, et al. Attitudinal and demographic predictors of measles, mumps and rubella (MMR) vaccine acceptance: development and validation of an evidence-based measurement instrument. *Vaccine* 2011; 29(8):1700-709.
35. Thomas C. How Underlying Beliefs Affect Behavior (The Effects of Coaching). 2012, 07.04. <https://coachcampus.com/coach-portfolios/research-papers/crystal-thomas-research-paper-how-underlying-beliefs-affect-behavior-the-effects-of-coaching/> Accessed 18 Mar 2020.
36. Offit PA, Coffin SE. Communicating science to the public: MMR vaccine and autism. *Vaccine*. 2003 Dec 8;22(1):1-6.
37. Newton P, Smith DM. Factors influencing uptake of measles, mumps and rubella (MMR) immunization in site-dwelling Gypsy, Roma and Traveller (G&T) communities: a qualitative study of G&T parents' beliefs and experiences. *Child Care Health Dev* 2017;43:504–10.
38. Gross K, Hartmann K, Zemp E, Merten S. 'I know it has worked for millions of years': the role of the 'natural' in parental reasoning against child immunization in a qualitative study in Switzerland. *BMC Public Health* 2015;15:373.

Tables

Table 1. Health behaviours assessed using the IHB scale

IHB		Me	<i>s</i>	<i>c</i> 25	<i>c</i> 75	min	max
proper eating habits	3.4	3.5	0.7	3.0	3.8	1.5	5.0
preventive behaviours	3.5	3.5	0.7	3.0	4.0	1.5	5.0
positive mental attitude	3.6	3.7	0.6	3.2	4.0	1.5	5.0
health practices	3.2	3.2	0.6	2.8	3.7	1.3	4.7
IHB	82.1	82	12.3	74	91	47	111
Level of health behaviours	Number		%				
low	99		33.0%				
average	127		42.3%				
high	74		24.7%				

Table 2. Selected opinions of parents depending on their preferred health behaviours.

	proper eating habits		preventive behaviour		positive mental attitude		health practices	
		s		s		s		s
It is better to develop resistance by being infected than by vaccination								
yes	3.47	0.64	3.57	0.67	3.78	0.61	3.45	0.61
no	3.42	0.64	3.54	0.66	3.58	0.60	3.20	0.61
hard to say	3.33	0.75	3.37	0.77	3.52	0.67	3.15	0.63
<i>p</i>	0.5632		0.2366		0.2531		0.1096	
You should be vaccinated against all diseases								
yes	3.38	0.69	3.50	0.66	3.57	0.64	3.10	0.61
no	3.42	0.69	3.45	0.69	3.58	0.60	3.29	0.58
hard to say	3.40	0.63	3.56	0.70	3.58	0.62	3.21	0.64
<i>p</i>	0.9595		0.2819		0.8752		0.1326	
The current vaccination programme takes into account the situation in Poland								
yes	3.40	0.67	3.50	0.75	3.55	0.66	3.19	0.64
no	3.49	0.61	3.59	0.61	3.58	0.49	3.18	0.58
hard to say	3.36	0.68	3.46	0.59	3.62	0.59	3.24	0.59
<i>p</i>	0.7630		0.6118		0.8036		0.7126	
The vaccines used in Poland are safe								
yes	3.40	0.64	3.52	0.71	3.58	0.62	3.19	0.63
no	3.60	0.67	3.58	0.56	3.71	0.56	3.00	0.59
hard to say	3.37	0.72	3.46	0.66	3.56	0.63	3.27	0.58
<i>p</i>	0.8341		0.7924		0.8125		0.3811	
Vaccines contain toxic ingredients								
yes	3.67	0.66	3.67	0.55	3.80	0.52	3.36	0.61
no	3.40	0.68	3.49	0.73	3.61	0.64	3.22	0.65
hard to say	3.32	0.64	3.46	0.68	3.49	0.61	3.16	0.58
<i>p</i>	0.0381*		0.3287		0.0223*		0.1416	
Vaccines cause developmental disorders and autism								
yes	3.63	0.56	3.80	0.61	3.90	0.64	3.42	0.62
no	3.39	0.71	3.41	0.73	3.57	0.65	3.18	0.63
hard to say	3.37	0.63	3.54	0.64	3.54	0.57	3.20	0.59
<i>p</i>	0.2963		0.0375*		0.0773		0.2425	
Vaccinations should be mandatory								
yes	3.43	0.63	3.50	0.71	3.55	0.63	3.20	0.63
no	3.32	0.68	3.53	0.66	3.71	0.58	3.27	0.56
hard to say	3.33	0.82	3.37	0.58	3.38	0.62	3.08	0.66
only some	3.62	0.77	3.67	0.51	3.60	0.44	3.21	0.37
<i>p</i>	0.5949		0.5182		0.1777		0.8082	
Children who are not vaccinated should not be admitted to kindergartens and crèches								
yes	3.40	0.64	3.48	0.70	3.52	0.67	3.16	0.68
no	3.43	0.68	3.62	0.63	3.60	0.57	3.21	0.63
hard to say	3.37	0.69	3.42	0.71	3.63	0.59	3.27	0.51
<i>p</i>	0.8455		0.3155		0.5410		0.5826	

I have doubts about vaccinating my child								
yes	3.40	0.77	3.52	0.65	3.63	0.55	3.28	0.58
no	3.40	0.61	3.55	0.66	3.62	0.62	3.19	0.61
hard to say	3.38	0.68	3.26	0.82	3.30	0.67	3.15	0.70
p	0.9613		0.0823		0.0156*		0.4125	

- Mean, *s* -Standard deviation

Table 3. Preference for child vaccination according to parents' preferred health behaviours

	proper eating habits		preventive behaviour		positive mental attitude		health practices	
		<i>s</i>		<i>s</i>		<i>s</i>		<i>s</i>
Has your child been vaccinated with combined vaccines ?								
yes	3.47	0.60	3.57	0.64	3.63	0.59	3.28	0.61
no	3.29	0.71	3.42	0.73	3.52	0.66	3.12	0.62
I do not remember	3.58	0.88	3.43	0.85	3.50	0.48	3.13	0.54
p	0.0584		0.2043		0.1620		0.0245*	
Has your child had the recommended vaccinations?								
		<i>s</i>		<i>s</i>		<i>s</i>		<i>s</i>
yes	3.47	0.65	3.59	0.69	3.62	0.67	3.28	0.66
no	3.31	0.62	3.40	0.68	3.56	0.56	3.12	0.57
I do not remember	3.38	0.95	3.39	0.63	3.36	0.52	3.14	0.48
p	0.1009		0.0783		0.0868		0.0380*	

- Mean, *s* -Standard deviation